

Additional file 2. Descriptive statistics of the study population

Characteristic	LTBI screening		LTBI diagnosis		Initiated LTBI treatment		Completed LTBI treatment	
	n	Column %	n	Row %	n	Row %	n	Row %
Total	257	(100%)	30	(12%)	29	(97%)	28	(97%)
Public Health Service (PHS)								
PHS 1	35	(14%)	2	(6%)	2	(100%)	1	(50%)
PHS 2	48	(19%)	8	(17%)	7	(88%)	7	(100%)
PHS 3	62	(24%)	12	(19%)	12	(100%)	12	(100%)
PHS 4	101	(28%)	8	(8%)	8	(100%)	8	(100%)
PHS 5	10	(3%)	0	(0%)	-	-	-	-
Gender								
Female	129	(50%)	10	(8%)	10	(100%)	10	(100%)
Male	128	(50%)	20	(16%)	19	(95%)	18	(95%)
Age category								
0-17 years	48	(19%)	1	(2%)	1	(100%)	1	(100%)
18-24 years	63	(25%)	5	(8%)	5	(100%)	4	(80%)
25-34 years	103	(40%)	12	(12%)	12	(100%)	12	(100%)
35-44 years	43	(17%)	12	(28%)	11	(92%)	11	(100%)
Household composition								
Alone	122	(45%)	16	(13%)	16	(100%)	15	(94%)
With family	123	(48%)	11	(9%)	10	(91%)	10	(100%)
Missing	12	(5%)	3	(25%)	3	(100%)	3	(100%)
Education level								
No formal / primary education	116	(45%)	12	(10%)	11	(92%)	11	(100%)
Secondary school	108	(42%)	13	(12%)	13	(100%)	12	(92%)
Higher education	28	(11%)	2	(7%)	2	(100%)	2	(100%)
Missing	5	(2%)	3	(60%)	3	(100%)	3	(100%)
Employment status								
Employed (payed job)	4	(2%)	0	(0%)	-	-	-	-
Unemployed (no payed job)	35	(14%)	6	(17%)	5	(83%)	5	(100%)
Student*	210	(82%)	22	(10%)	22	(100%)	21	(95%)
Missing	8	(3%)	2	(25%)	2	(100%)	2	(100%)
Duration of stay in the Netherlands								
<6 months	12	(5%)	1		0	(0%)		
≥6 months, <1 year	31	(12%)	2		2	(100%)	2	(93%)
≥1 year, <2 years	25	(10%)	1		1	(100%)	1	(100%)
≥2 years, <3 years	89	(35%)	14		14	(100%)	13	(93%)
≥3 years, <10 years	11	(4%)	4		4	(100%)	4	(100%)
Missing	90	(35%)	8		8	(100%)	8	(100%)

LTBI latent tuberculosis infection, PHS public health service, TB tuberculosis

*Includes: primary school, secondary school, college, integration courses