



Additional figures and tables



Additional Figure 1. Flow chart of subject selection.



Additional Figure 2. Antibiotic resistance genotypes over time. The percentage of subjects with the antibiotic resistance genotype are presented over time.

| Class | Drug Class | Type of antibiotics received during pregnancy | No. of subjects |
|-------------------------------|-----------------------|--|------------------------|
| Beta-lactam | Beta-lactam | Cefalexin capsule | 3 |
| Beta-lactam | Penicillin | Amoxicillin capsule | 6 |
| Beta-lactam | Penicillin | Amoxicillin capsule, Cloxacillin capsule | 1 |
| Macrolide, Beta-lactam | Macrolide, Penicillin | Erythromycin ESS, Amoxicillin | 3 |
| Macrolide | Macrolide | Azithromycin | 1 |

Additional Table 1. Maternal antibiotic exposures during pregnancy.

| Class | Drug Class | Type of antibiotics during labour | No of subjects |
|--------------------|------------------------------|--|-----------------------|
| Lincosamide | Lincosamide | Clindamycin | 1 |
| Beta-lactam | Penicillin | Penicillin G | 13 |
| Beta-lactam | Penicillin | Ampicillin/Amoxicillin | 7 |
| Beta-lactam | Penicillin, Cephalosporin | Penicillin G, Cefazolin | 1 |
| Beta-lactam | Cephalosporin | Cefazolin | 4 |
| Beta-lactam | Cephalosporin | Rocephine | 1 |

Additional Table 2. Maternal antibiotic exposure during labour.