

## RESEARCH

## QUESTIONNAIRE

### DATA COLLECTION

#### IDENTIFICATION:

NAME: \_\_\_\_\_

AGE: \_\_\_\_\_ DATE OF BIRTH: \_\_\_\_\_ SEX: M( ) F( ) Tel:

\_\_\_\_\_

Height: \_\_\_\_\_ Weight: \_\_\_\_\_ BMI:

\_\_\_\_\_

SMOKER: YES ( ) NO ( )

DIABETES: TYPE 1 ( ) TYPE 2 ( )

DURATION OF DIABETES IN YEARS: \_\_\_\_\_

MEDICATIONS: ORAL ANTI-DIABETIC DRUG ( )

INSULIN ( )

COMORBIDITY: HYPERTENSION ( )

HEART DISEASE ( )

KIDNEY DISEASE ( )

RETINOPATHY ( )

Prior STROKE ( )

Prior HEART ATTACK ( )

Prior AMPUTATION ( )

DIAMETER OF ULCER IN cm: \_\_\_\_\_ WOUND < 3 cm ( )

WOUND  $\geq$  3 cm ( )

DURATION OF ULCER IN MONTHS: \_\_\_\_\_

CLASSIFICATION OF ULCERS: WAGNER GRADE 2 ( )

WAGNER GRADE 3 ( )

WAGNER GRADE 4 ( )

LABORATORY EXAMS: HAEMOGLOBIN: \_\_\_\_\_

LEUCOGRAM: \_\_\_\_\_

ESR: \_\_\_\_\_

CRP: \_\_\_\_\_

FASTING GLUCOSE: \_\_\_\_\_

GLYCOSYLATED HAEMOGLOBIN: \_\_\_\_\_

CREATININE: \_\_\_\_\_

**DIAMETER OF THE ULCER IN cm (FOLLOW-UP)**

ON ADMISSION: \_\_\_\_\_

AFTER 10 SESSIONS / 2 WEEKS: \_\_\_\_\_

AFTER 20 SESSIONS / 4 WEEKS: \_\_\_\_\_

AFTER 30 SESSIONS / 6 WEEKS: \_\_\_\_\_

AFTER 30 SESSIONS / 7 WEEKS: \_\_\_\_\_

AFTER COMPLETING 6 MONTHS: \_\_\_\_\_

**ULTRASOUND FINDINGS (Doppler ultrasound)**

NAME: \_\_\_\_\_

**RIGHT LOWER LIMB**

<b>Arteries</b>	<b>Femoral Artery</b>	<b>Popliteal Artery</b>	<b>Anterior Tibial Artery</b>	<b>Posterior Tibial Artery</b>	<b>Fibular Artery</b>
<b>Peak Systolic Velocity</b>					
<b>Peak Diastolic Velocity</b>					
<b>Triphasic flow</b>					
<b>Biphasic Flow</b>					
<b>Monophasic Flow</b>					
<b>Occluded</b>					
<b>Calcification</b>					

## LEFT LOWER LIMB

<b>Arteries</b>	<b>Femoral Artery</b>	<b>Popliteal Artery</b>	<b>Anterior Tibial Artery</b>	<b>Posterior Tibial Artery</b>	<b>Fibular Artery</b>
<b>Peak Systolic Velocity</b>					
<b>Peak Diastolic Velocity</b>					
<b>Triphasic flow</b>					
<b>Biphasic Flow</b>					
<b>Monophasic Flow</b>					
<b>Occluded</b>					
<b>Calcification</b>					