

Effects of the grade/price of and thickness of cigarette on the PM2.5 concentration of smoke

CURRENT STATUS: POSTED



Shengzhe Ji
Shandong Agricultural University

Zhen Liu
Taian City Central Hospital

Yuxue Sha
Shandong Agricultural University

Yiwen Wang
Shandong Agricultural University

Qaing Wang
Taian City Central Hospital

Peng Liu
Shandong Agricultural University

✉ liupeng2003@sdau.edu.cn *Corresponding Author*

DOI:

10.21203/rs.2.19438/v1

SUBJECT AREAS

Health Policy

KEYWORDS

Effects of the grade, price of and thickness of cigarette on the PM2.5 concentration of smoke

Abstract

Background: Due to lower tar and nicotine, high grade/price and slim cigarette are believed as “safe cigarette” and cause less tobacco smoke pollution (TSP) advocated by cigarette producers. However, cigarette smokes comprise thousands of harmful constituents, only two substances cannot represent TSP. Then in this study, we wanted to discover if the grade/price and thickness of cigarette affect TSP.

Methods: Slim (S) and regular (R) cigarette with different grades/prices were selected to investigate TSP *via* measurement PM_{2.5} in a closed chamber (1 m³). Sidestream smoke was collected through cigarette combustion. Mainstream smoke was collected using piston-operated smoking machine.

Results: In sidestream smoke, cigarette grade/price did not affect PM_{2.5} of R or S cigarette. But in mainstream smoke, grade/price acted discrepancy roles in different thickness cigarette. In R cigarette, PM_{2.5} of lowest grade/price (R5) was lowest. While in S cigarette, PM_{2.5} of lowest grade/price (S5) was highest. Cigarette thickness obviously affected PM_{2.5}. PM_{2.5} of R cigarette was 116% higher than S cigarette in sidestream. In mainstream smoke, magnification was down to 31%, despite PM_{2.5} of R cigarette was also higher.

Conclusions: Irrespective of sidestream or mainstream smoke, the grade/price of cigarette did not obviously affect PM_{2.5}. But which was significantly influenced by cigarette thickness. Although S cigarette PM_{2.5} was significantly lower than R cigarette, it cannot mean that S cigarette was less harmful. Because harmfulness of smoke was not only affected by cigarette but also by smoker inhale habits. It needs further experiments to evaluate the harmfulness of slim cigarette.

Background

Tobacco smoke pollution (TSP), also known as “secondhand smoke” and “passive smoke”, in public places is a worldwide health problem. Tobacco smokes contain thousands of harmful constituents that come from cigarette combustion (“sidestream smoke”) and exhaled by the smoker (“mainstream smoke”) [1]. Accumulating evidence show that TSP may be the cause of many respiratory problems, acute cardiovascular effects, and lung cancer [2–4]. Due to the toxicity of TSP, > 600,000 people die every year worldwide [5].

The level of cigarette production and consumption in China is one of the highest in the world.

Cigarette smoking has an important role in Chinese social customs, such as at weddings and funerals.

Offering a cigarette to a friend or guest is regarded as a social courtesy. Nearly all these social courtesies happen in public places, and TSP is a serious problem [6].

In the past decade, the production and sales of cigarette in China has decreased, and this has been attributed to two main factors. The first is people's awareness of the hazards of cigarette smoking.

For example, the percentage of people who were aware that cigarette smoking increases the risk of lung cancer increased from 36% in 1996 to 77.5% in 2010 [7]. The other reason is that the Chinese government promoted multiple policies to reduce cigarette smoking, such as curbing cigarette advertisements and increasing the price of cigarette [8, 9].

The decline in cigarette sales severely affected the profits of State Tobacco Monopoly Administration (STMA) and China National Tobacco Corporation (CNTC). (Actually, the STMA and CNTC is the same organization, the different names represent the separation between the government and tobacco industry.) STMA/CNTC exploited new marketing tools to maintain their profits [10], one of which was the "safe cigarette", characterized by its low-tar and nicotine content. Some scientists attached to or funded by STMA/CNTC became engaged in reducing the harmfulness of smoking through improving cigarette-production technology or developing new types of cigarette.

New technologies were used to promote cigarette "safety". For instance, filters made from new materials, high-permeability cigarette papers, and ways to improve combustion were reported to decrease the content of hazardous substances [11-13]. These new technologies led to increases in manufacturing costs, which were passed onto the consumer. This increase in price could be misconstrued by consumers as being correlated with increased safety. That is, a high price of cigarette denotes high safety, and low price corresponds with low safety. A "slim" (S) cigarette (circumference of 17 ± 1 mm) contains less tobacco than a "regular" (R) cigarette (25 ± 1 mm). As the result, S cigarette has been advocated to be a "safe cigarette" to increase the profits of STMA/CNTC [14]. Under the impetus of STMA/CNTC, sales of S cigarette have shown tremendous growth. In 2014,

2.9 million cartons of S cigarette were sold, but in 2018 sales increased to 347 million cartons [15]. Apart from developing new products, cigarette producers have propagated the “safe cigarette” through introducing the work of tobacco researchers. A typical case was the “Tobacco Academician” who promoted low-tar tobacco products and was elected to the Chinese Academy of Engineering (the highest honor for scientists in China) [16]. All the propagation and research conducted by the tobacco industry created an illusion to people that TSP from “safe cigarette” did not or reduced harm to their health.

Cigarette smokes comprise thousands of harmful compounds; low-tar and nicotine content cannot fully represent TSP. “PM2.5” refers to atmospheric particulate matter (PM) of diameter < 2.5 µm. The PM2.5 concentration is a suitable and reliable airborne marker to evaluate emission and exposure to TSP [17].

In the present study, we selected six types of cigarette to evaluate the effects of grade/price, cigarette thickness, and grade/price × thickness on the PM2.5 concentration after cigarette smoking.

Methods

STMA/CNTC divides cigarette into five grades according to price in China [18]. To clarify the relationship between cigarette price and PM2.5, we selected three grades/prices: 1 (≥ ¥100), 3 (¥30 ≤ and <¥70) and 5 (< ¥16.5). According to their thickness, there are two types of cigarette: R and S (Table 1).

Table 1
Materials selected in the present study

Thickness	Grade	Tar (mg)	Nicotine (mg)
Regular	1	8	0.6
	3	10	1.0
	5	12	1.1
Slim	1	6	0.6
	3	7	0.7
	5	8	0.8

Measurement of the PM2.5 concentration

Measurement of the PM2.5 concentration was undertaken according to the method developed by Sendzik and colleagues [19]. A Dustrak aerosol monitor 8532 (TSI, St. Paul, MN, U.S.A.) was placed in a closed chamber (1 m × 1 m × 1 m) to measure the PM2.5 concentration. The monitor was calibrated before each experimental session according to manufacturer specifications. A customized calibration

factor of 0.32 was applied to the device, determined by calibrating the device in the present study with other light-scattering photometers measuring TSP [19, 20]. Sidestream smoke was collected in the chamber through cigarette combustion. Mainstream smoke was collected with a piston-operated smoking machine under the following conditions: puff volume of 40 mL, puff duration of 2 s, and interval between puffs of 10 s. According to this method, we could simulate the smoking of tobacco by a human [17]. Before data collection, we operated exhaust equipment to ventilate the PM2.5 of the chamber down to the air level.

Statistical analyses

Data were subjected to multivariate analysis of variance with general linear models. If thickness × grade/price had an interactive effect, further analyses were conducted. Data analysis were undertaken using SAS v8.0 (SAS, Cary, NC, USA).

Results

The cigarette grade/price had no significant effect on the PM2.5 concentration in either group (Fig. 1). However, the mean PM2.5 concentration for R cigarette was obviously higher than that for S cigarette (by 116%). Multivariate analysis of variance showed that cigarette thickness had main effect on the PM2.5 concentration of sidestream smoke (Table 2).

Table 2
Multivariate analyses of variance for PM2.5 in sidestream smoke

	df	F	P
Cigarette thickness	1	528.594	0.000
Grade/price	2	0.206	0.817
Thickness × grade/price	2	0.365	0.702

Table 3
Multivariate analyses of variance for PM2.5 in mainstream smoke

	df	F	P
Cigarette thickness	1	87.243	0.000
Grade/price	2	2.054	0.171
Thickness × grade/price	2	9.065	0.04

Just like for sidestream smoke, cigarette thickness had an effect on the PM2.5 concentration in mainstream smoke (Fig. 2). In addition, there was interaction effect between cigarette thickness and grade/price (Table 3). A simple effect analysis was conducted (Fig. 3). There was no obvious difference between R cigarette. The PM2.5 concentration of S1 and S3 were similar and lowest among all treatments. Interestingly, the PM2.5 concentration of S5 was significantly higher than that of S1

and S3, and close to that of R5.

Discussion

According to their annual report, STMA/CNTC had uninterrupted growth in recent years that reached ¥1155.6 billion (\$168 billion) in 2018 ^[21]. This “brilliant achievement” was supported by millions of smokers. Due to customs and habits, smokers in China can be found in most public places. As a result, many nonsmokers are affected by TSP. TSP-induced health risks burden the government and families. It has been reported that the total healthcare cost of TSP exposure in rural China accounted for 0.3% of China’s national healthcare expenditure in 2011 ^[22]. Fortunately, thanks to government policies, cigarette-carton sales decreased gradually from 260.98 billion in 2014 to 233.58 billion in 2018 according to the National Bureau of Statistics ^[23].

[21][22][23]

The discrepancy between the decrease in cigarette sales and enhancement of profit is noteworthy. The reason was due to the new sales strategy of STMA/CNTC. To align with the requirements for tobacco control from the government and society, STMA/CNTC decreased the output of low grade/price cigarette but slightly increased the output of high grade/price cigarette. Studies from several research institutions attached to or funded by STMA/CNTC indicated that high grade/price cigarette was likely to be much safer than low grade/price cigarette. The filters of cigarette made in China are composed of cellulose acetate (CA) or polypropylene (PP), of which CA has a higher filtration efficiency (by 6%) compared with PP ^[24]. PP filters are used in low grade/price cigarette ^[13]. Good-quality tobacco leaves in high grade/price cigarette have excellent flame properties, which can reduce the toxicity due to decrease tar and nicotine content ^[25]. In addition, some studies have indicated high-combustion cigarette papers and safer additive materials can decrease the toxicity of cigarette. However, these new technologies lead to much higher manufacturing costs and, consequently, price increases ^[11, 26]. Studies have created an illusion for smokers that high grade/price cigarette are harmless to health. However, we found that the PM2.5 concentration was not affected by the grade/price of cigarette in sidestream or mainstream smoke. The conflicting

results between our study and previous studies may be due to different research objectives. Several studies investigate a constituent of smoke (e.g., tar, nicotine, tobacco-specific nitrosamines) but we investigated the PM_{2.5} concentration. Tobacco smoke contains > 5000 constituents [27]. Smoke toxicity cannot be measured through investigating a few substances even if they have high toxicity. Conversely, PM_{2.5} almost contains all the constituents of tobacco smoke.

S cigarette is a relatively new type of cigarette, and attracts smokers rapidly upon their introduction. Their carton sales increased from 2.9 million to 347 million within 5 years. One of the important reasons for this massive increase in sales was that STMA/CNTC and their research institute transmitted a simple message: S cigarette can decrease harm to your health [28]. In the Chinese cigarette market, the tar and nicotine content of S cigarette annotated on cartons is 6–8 mg and 0.6–0.8 mg respectively, which is lower than that for R cigarette (Table 1). Ge and colleagues compared the components in mainstream smoke between S cigarette and R cigarette, and found that S cigarette released less nicotine, tar and carbon monoxide than R cigarette [29]. In the present study, the PM_{2.5} concentration was significantly lower in S cigarette in sidestream smoke and mainstream smoke. The mean PM_{2.5} concentration of sidestream smoke in S cigarette was lower by 54% compared with that in R cigarette whereas, in mainstream smoke, it was lower by 24%. These results seemed suggested that S cigarette was safer than R cigarette. In fact, the harmfulness of S cigarette cannot be reflected by the tar and nicotine content labeled on the carton, a hypothesis supported in the present study. The PM_{2.5} concentration of S5 was similar to that of R cigarette (Fig. 2, 3). And more important should to be noted that experiment results merely represented one cigarette differences between two types. But for one smoker, the objective of tobacco smoking is the satisfaction elicited by nicotine. If smokers changed from R cigarette to S cigarette, to maintain the desired intake of nicotine, they could change the way they smoked (e.g., deeply inhale each draw of the cigarette, increase the number of cigarettes smoked) [30]. Thus, their exposure to tobacco may be higher or at least equal to that of R cigarette.

To our knowledge, this was the first study to investigate the effects of grade/price and thickness of

cigarette on the PM_{2.5} concentration. We found no significant difference between lower and higher grade/price of cigarette. S cigarette produced lower concentrations of PM_{2.5} compared with that of R cigarette. However, this result does not demonstrate that S cigarette is safer than R cigarette because the harmfulness of smoking is determined not only by the cigarette but also by inhalation habits. This was theoretical research undertaken using a closed chamber, and indoor studies are needed to verify our data.

Consent for publication

No application

Availability of data and materials

All data generated or analysed during this study are included in this published article

Declarations

Consent for publication

No application

Availability of data and materials

All data generated or analysed during this study are included in this published article

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by Innovation and Entrepreneurship Training Program for College Students of Shandong (201910434038), Shandong Provincial Natural Science Foundation, China (ZR2018BC040).

Authors' contributions

L.Z. and L.P. designed the experiment and produced the draft. J.S., S.Y. and W.Y. collected the data. W.Q. analyzed the data.

References

- 1 Sureda, X., Fernández, E., López, M. J., Nebot, M. Secondhand tobacco smoke exposure in open and semi-open settings: a systematic review. *Environ Health Persp*, 2013. 121(7): 766-773.
- 2 Shikata, S. and Y. Takemura. Secondhand smoke exposure and risk of lung cancer in Japan: a

systematic review and meta-analysis of epidemiologic studies. *Jpn J Clin Oncol*, 2017. 47(3): 282-282.

3 Dunbar, A., W. Gotsis, and W. Frishman. Second-hand tobacco smoke and cardiovascular disease risk: an epidemiological review. *Cardiol Rev*, 2013. 21(2): 94-100.

4 Vozoris, N. and M.D. Lougheed. Second-hand smoke exposure in Canada: prevalence, risk factors, and association with respiratory and cardiovascular diseases. *Can Respir J*, 2008. 15(5): 263-269.

5 Öberg, M., Jaakkola, M. S., Woodward, A., Peruga, A., Prüss-Ustün, A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *The lancet*, 2011. 377(9760): 139-146.

6 Yang, G., Wang, Y., Wu, Y., Yang, J., & Wan, X. The road to effective tobacco control in China. *Lancet*, 2015. 385(9972): 1019-1028.

7 Chinese Center of Disease Control and Prevention. Global Adults Tobacco Survey (GATS) China 2010 Country Report. http://www.who.int/tobacco/surveillance/survey/gats/en_gats_china_report.pdf?ua=1(accessed Jan 7, 2015), 2011.

8 Legislation Office of the State Council. The notice to hear public opinion and comments on the regulation on control smoking in public places (draft). <http://www.nhfpc.gov.cn/zhuzhan/zqyj/201411/16b813422bae4cfd93823102ee45b7f3.shtml> (Nov 24, 2014), 2014.

9 National Health and Family Planning Committee. The first nationwide regulation of smoking ban in public spaces (draft)., <http://www.nhfpc.gov.cn/zhuzhan/zcjd/201411/cc090d9b0be641ac942daf1b101982b4.shtml>(accessed Nov 24, 2014). 2014.

10 Fang J, Yang G, Wan X. 'Pro-tobacco propaganda': a case study of tobacco industry-sponsored elementary schools in China. *Tob Control*, 2019.

11 Case P.D., Loureau J.M, Baskevitch N. Systematic studies on cigarette paper. The influence of natural permeability, added permeability and burn additive on paper properties and mainstream ISO yields. Coresta, 2003. ST 27.

12 Ning X, Wei Z., Zhang H, Lin H. Research progress on cigarette reducing tar technology. *Mod Agr*

Sci &Tech, 2019. 8: 240-243.

13 Wu J, Huang Y., Zeng J, Tang D. Filtration structure progress of cigarette filter. *Light Ind Mach*, 2017. 35(5): 86-90.

14 Zhang X, Xu X, Xu M, Hu T. The impact of tobacco taxation policy on slim cigarette use and the growing popularity of slim cigarette in China since 2014. *Health*, 2019. 11(6): 711-720.

15 State Tobacco Monopoly Administration. Analysis of national cigarette market. 2014-2018: <http://www.tobacco.gov.cn/html/56.html>.

16 Hvistendahl M. Tobacco scientist's election tars academy's image. *Science*, 2012:153-154.

17 Czogala, J., Goniewicz, M. L., Fidelus, B., Zielinska-Danch, W., Travers, M. J., Sobczak, A. Secondhand exposure to vapors from electronic cigarette. *Nicotine Tob Res*, 2013. 16(6): 655-662.

18 Qiao D. An empirical analysis on the structure of Chinese cigarette market and the competitiveness of the brands. *Yunnan Univ*, 2016.

19 Sendzik, T., Fong, G. T., Travers, M. J., Hyland, A. An experimental investigation of tobacco smoke pollution in cars. *Nicotine Tob Res*, 2009. 11(6): 627-634.

20 Travers, M. J., Hyland, A., Higbee, C., Travers, M. B., Peppone, L., Cummings, K. M. Tobacco smoke pollution exposure in hospitality venues around the US and the effect of smokefree air policies. in Madison, WI, Society for Research on Nicotine and Tobacco, Proceedings of 13th Annual Meeting, 2007:21-24

21 State Tobacco Monopoly Administration. The state tobacco administration held a press conference on the economic performance of the tobacco industry in 2018. <http://www.scio.gov.cn/xwfbh/gbwxfbh/xwfbh/ycj/Document/1646503/1646503.htm>.

22 Yao T, Sung H, Mao Z, Hu T, Max, W. The healthcare costs of secondhand smoke exposure in rural China. *Economics of Tobacco Control in China: From Policy Research to Practice*. 2016: 43-57.

23 National Bureau of Statistics of China. 2014-2018: <http://www.stats.gov.cn/was5/web/search?channelid=288041&andsen=%E5%8D%B7%E7%83%9F>.

24 Shao J, Liu H, Xia Q, Xie F, Liu K. Determination of filtration efficiencies of filters to tobacco-specific nitrosamines in cigarette smoke. *Tob Chem*, 2010(7): 34-38.

- 25 Yu J. Cigarette manufacturing technology. 2009.
- 26 Liu H, Zhao C, and Ding C. Research advances in the mechanism of cigarette harm-reducing mediated by filling materials of medicinal plants. *Chin Agric Sci Bull*, 2015. 31(28): 270-279.
- 27 Hall M G, Ribisl K M, Brewer N T. Smokers' and nonsmokers' beliefs about harmful tobacco constituents: implications for FDA communication efforts. *Nicotine Tob Res*, 2013. 16(3): 343-350.
- 28 Yan N, Wu J, Liu W, et al. Contribution of slim cigarette to reduce the tobacco addiction. *Hubei Agr Sci*, 2018. 57(2): 99-102.
- 29 Ge C, Zhao M, Hu y, et al., Analysis of routine indexes and neutral aroma components in mainstream smoke of slim and normal cigarette. *Tob Sci &Tech*, 2017. 50(4): 43-50.
- 30 Stratton, K., Shetty, P., Wallace, R., & Bondurant, S. Clearing the smoke: the science base for tobacco harm reduction—executive summary. *Tob Control*, 2001. 10(2): 189-195.

Figures



Figure 1

The cigarette grade/price had no significant effect on the PM_{2.5} concentration in either group



Figure 2

Multivariate analysis of variance showed that cigarette thickness had main effect on the PM_{2.5} concentration of sidestream smoke



Figure 3

In addition, there was interaction effect between cigarette thickness and grade/price (Table 3). A simple effect analysis was conducted