

Economic impact of therapies on perioperative hemodynamic optimization in surgical patients from the perspective of developing country: Economic study and Meta-analysis

João M Silva-Jr (✉ joao.s@usp.br)

Universidade de São Paulo <https://orcid.org/0000-0003-1494-0784>

Pedro Ferro L Menezes

Universidade de São Paulo

Flavia Helena S Carvalho

IAMSPE

Mariana Augusta N Oliveira

IAMSPE

Francisco Nilson F Cardoso Filho

IAMSPE

Bruna N Fernando

IAMSPE

Maria Jose C Carmona

Universidade de São Paulo

Vanessa D Teich

Hospital Israelita Albert Einstein

Suzana M Lobo

Faculdade de Medicina de São José do Rio Preto

Luiz Marcelo S Malbouisson

Universidade de São Paulo

Research

Keywords: surgery, hemodynamic optimization, complications, economic, cost-effective, public health system

Posted Date: January 8th, 2020

DOI: <https://doi.org/10.21203/rs.2.20292/v1>

Abstract

Background: Health resources are limited and need to be rationally distributed. Studies suggest that hemodynamic optimization therapy can reduce length of hospital stay, complications and costs. However, Brazilian data are scarce. Therefore, the objective of this analysis was to evaluate whether the improvement demonstrated by hemodynamic optimization therapy in surgical patients could result in lower costs from the perspective of the Brazilian public unified health system (SUS).

Method: A meta-analysis was performed comparing surgical patients who underwent hemodynamic optimization therapy (intervention) with patients submitted to standard therapy (control) in terms of complications and hospital costs. The cost-effectiveness analysis evaluated the clinical and financial benefits of hemodynamic optimization protocols for surgical patients. The analysis considered the clinical outcomes of randomized studies published in the last 20 years that involved surgeries and hemodynamic optimization therapy. Indirect costs were not included in the analysis.

Results: A total of 21 clinical trials with a total of 4872 surgical patients were selected. Comparison of the intervention and control groups showed lower rates of infectious (RR=0.66; 95% CI=0.58-0.74), renal (RR=0.68; 95% CI=0.54-0.87), and cardiovascular complications (RR=0.87; 95% CI=0.76-0.99) and a tendency toward lower rates of respiratory complications (RR=0.82; 95% CI=0.67-1.02). There was no difference in mortality (RR=1.02; 95% CI=0.80-1.3) between groups. In the analysis of total costs, the intervention group showed a cost reduction of R\$396,024.83 for every 1000 patients treated compared to the control group. The patients in the intervention group showed greater effectiveness, with 1.0 fewer day in the ICU and hospital. In addition, there were 333 fewer patients with complications, with a consequent reduction of R\$1,630,341.47 for every 1000 patients treated.

Conclusions: Hemodynamic optimization therapy is cost-effective and would increase efficiency and decrease the burden of the Brazilian public health system.

Background

Millions of major surgeries are performed every year worldwide [1]. The mortality and morbidity rates of high-risk surgeries vary among countries but are considered high, making them a global problem [2–4]. In Brazil, in 2008, the mortality rate of patients admitted to intensive care units after major surgeries was 15%; after 90 days, it reached 20.3%. The main postoperative complication found in Brazil was sepsis (24.7%), and the main cause of death was multiple organ dysfunction [5].

Postoperative complications in high-risk patients undergoing major surgery are associated with low cardiorespiratory reserve and the inability to maintain adequate oxygen delivery (O₂) during surgical trauma to meet the increased metabolic demand [6–8]. As a consequence, there is an imbalance in the ratio of oxygen delivery to oxygen consumption, leading to hypoperfusion, multiple organ dysfunction and severe infections, which are important causes of postoperative mortality [4]. Perioperative hemodynamic optimization therapy (PHOT) aims to adjust cardiac function to meet the increased

demand during the perioperative period, thus avoiding hypovolemia or hypervolemia and, ultimately, tissue hypoperfusion and postoperative complications. This requires adequate hemodynamic monitoring to guide the early treatment of each patient, allowing for earlier identification of the need for fluid optimization, blood transfusion, and vasoactive drugs.

Studies and meta-analyses have shown that PHOT has a significant impact on the outcomes of high-risk patients undergoing major surgery, potentially decreasing morbidity and mortality, length of ICU stay, and length of hospital stay [9–11]. However, the risk-benefit ratio of this type of monitoring has been questioned because it is invasive and carries risks [12]. Monitoring techniques include pulmonary artery catheterization, transpulmonary thermodilution, echocardiography, transesophageal Doppler echocardiography, pulse contour analysis, partial carbon dioxide rebreathing and bioimpedance.

Cost-effectiveness is represented by a ratio between monetary cost, usually expressed in a national currency, in the numerator and a measure of health gain in the denominator. The poor adherence to PHOT in clinical practice combined with the need to improve the effectiveness of care for surgical patients in the face of increasing demand points to the urgent need for cost-effectiveness assessments that encourage the use of PHOT in this area, particularly in resource-limited settings.

Health resources are increasingly limited, and a lack of knowledge about a method is a barrier to the implementation of new therapies with proven effectiveness that could not only save lives but also lead to a reduction in resource use. Our hypothesis is that these interventions, due to their significant impact on the length of hospital stay and complications, would be cost-effective for the public health system.

This study aimed to evaluate whether the use of PHOT is cost-effective for patients undergoing major surgery from the perspective of the Brazilian Unified Health System (SUS) by evaluating the impact of a reduction in length of hospital stay, complication rates and mortality on hospital costs.

Materials And Methods

A systematic review of the clinical trials indexed in CENTRAL (PubMed), MEDLINE (OvidSP) and EMBASE (OvidSP) between 2001 and 2018 was performed. The search used the following keywords, which were expected to be present in the title and/or abstracts: "randomized studies" and "surgeries", and/or "perioperative", and/or "high-risk", and/or "complications", and/or "intraoperative" and/or "postoperative", and/or "cardiac output", and/or "cardiac index", and/or "hemodynamic monitoring", and/or "hemodynamic optimization therapy", and/or "hemodynamic intervention", and/or "cost-effectiveness", and/or "mortality". Methods identical to those recommended by the Cochrane systematic review of randomized controlled trials on increased blood flow to the organs, with explicitly defined goals and results after surgery, were used.

Only studies published in the English language were included. Two independent researchers identified the titles and abstracts of the potentially eligible studies. Disagreements between the investigators were resolved by consensus. Two other researchers extracted the following data from the full texts of

potentially eligible studies: study design, patient population, interventions and outcomes. Similar to the approach used for the selection of texts, any disagreements between researchers regarding the data extraction were resolved by consensus. Each included study was assessed independently by the first and second reviewer for risk of bias in random sequencing generation, allocation concealment, blinding of participants and personnel, blinding of outcomes assessment, incomplete outcome data, selective reporting, and other sources of bias using the Cochrane Risk of Bias Tool. [13] In the absence of appropriate published data, at least one attempt was made to contact the authors of eligible studies to obtain necessary data. The analysis was performed with the best available information when there was no response.

The following information and outcomes were recorded: number of patients with respiratory complications (i.e., the need for respiratory support for more than 24 hours after surgery, hypoxemia, acute changes in lung mechanics), cardiovascular complications (need for hemodynamic support, such as the use of inotropes and vasopressors during the postoperative period), renal complications (oliguria, unexpected increase in creatinine and need for dialysis) and infectious complications (infections that occurred during the postoperative period) based on the records of each study. Mortality was assessed throughout the longer follow-up period (primary outcome) or was examined as in-hospital mortality. Many studies have reported complications as the number of complications rather than number of patients with complications, but we only reported the last unit in the analysis, which was the number of complications per patient.

To calculate the costs of these complications, only the length of ICU stay was considered.

The inclusion criteria of the selected studies were as follows:

- 1) studies in adult patients (18 years or older);
- 2) studies with patients undergoing hemodynamic optimization therapy with some type of cardiac output monitor;
- 3) studies that related the costs of hemodynamic optimization therapy and its outcomes, such as the reduction of mortality or morbidity rates or the reduction of length of hospital or ICU stay;
- 4) interventional studies comparing the use of invasive or minimally invasive monitoring with the standard strategy for hemodynamic optimization to alter clinical outcomes. The intervention should meet the following criteria:

Perioperative period: The administration of fluids with or without inotropes/vasoactive drugs to increase blood flow (standard therapy group) was compared with goals measured explicitly with invasive or minimally invasive hemodynamic monitoring (intervention group). The perioperative period started at the beginning of surgery and lasted up to 24 hours after surgery. Explicit goals were defined for the cardiac index, oxygen delivery (DO_2), oxygen consumption, systolic volume, mixed or central venous oxygen saturation, oxygen extraction, or serum lactate.

The exclusion criteria were as follows:

- 1) animal studies;
- 2) studies published prior to 2001;
- 3) observational studies that did not use clinical intervention to change outcomes or case reports;
- 4) studies involving critically ill patients prior to intervention or with established sepsis who therefore with a higher probability of unfavorable outcomes and death regardless of intervention.

Cost assessment

For the cost-effectiveness analysis, the costs were separated into 10 categories and two periods: the intraoperative period (monitoring and the costs of fluid infusion, inotropes or vasopressors and blood transfusions) and the postoperative period in the ICU, which was maintained at fixed daily rates regardless of disease, clinical exam and procedures depending on postoperative complications, laboratory diagnosis, use of antimicrobial and other agents (cardiac support, renal support, physical therapy and imaging). To avoid confounding factors, the costs of the surgical procedure (considering that the surgeries would have the same magnitude), postoperative analgesia and preoperative state were excluded from the final sum, as were costs related to the hospital infrastructure (electricity, safety system, etc.).

The overhead costs were estimated from a social perspective, i.e., regardless of who will bear the cost. However, the unit costs of health resources and services were obtained from the national databases of the Brazilian public health system (SUS), and therefore, the direct costs represent the costs borne by the payer [14–19].

The incremental cost effectiveness analysis was based on the difference in costs divided by the difference in survival days in each group.

Statistical analysis

The analyses were performed in Review Manager (RevMan 5.2.8) using fixed effects models with random effects models for comparison. We applied the intention-to-treat method for all analyses. Treatment effects were reported as relative risk (RR) and confidence intervals (95% CI) for clinical variables or as differences in weighted averages (SD) or medians for the length of ICU and hospital stays. Empty cells, the result of studies in which no event was observed in one or both arms, were corrected by adding a fixed value (0.5) to all cells with an initial value of zero. The chi-square test was used to assess whether the differences observed in the results were due to chance. A large chi-square (I^2 statistic) provided evidence of heterogeneity of the intervention effects (indicating that the estimated effect was beyond chance).

Results

Initially, 52 potential articles were identified. After the inclusion and exclusion criteria were applied, 21 articles remained. (Fig. 1)

Of the selected studies, 21 used techniques for measuring cardiac output; these studies had a total of 4,872 patients at high surgical risk. (Table 1)

Table 1

Selected studies and the technology used for hemodynamic monitoring.

Author-year (ref)	Objective	Technology
Bonazzi et al., 2002 [33]	Evaluation of the impact of hemodynamic optimization using a pulmonary artery catheter on the outcome of patients undergoing vascular surgery.	Pulmonary artery catheter
Venn et al., 2002 [34]	Evaluation of hemodynamic optimization therapy in patients undergoing hip surgery.	Transesophageal Doppler
Conway et al., 2002 [35]	Randomized study to evaluate the influence of fluid titration using transesophageal Doppler during intestinal surgeries.	Transesophageal Doppler
Gan et al., 2002 [36]	Evaluation of the impact of hemodynamic optimization therapy on the reduction of length of hospital stay after major surgeries.	Transesophageal Doppler
Sandham et al., 2003 [37]	Randomized study evaluating the use of pulmonary artery catheters in high-risk surgical patients.	Pulmonary artery catheter
Wakeling et al., 2005 [38]	Evaluation of transesophageal echocardiography-guided hemodynamic optimization therapy for the reduction of length of hospital stay during the postoperative period of major abdominal surgeries.	Transesophageal Doppler
Pearse et al., 2005 [39]	Evaluation of the use of GDT in highly complex surgeries to reduce perioperative complications and length of hospital stay.	LiDCO monitoring system
Lobo et al., 2006 [40]	Investigation of the effects of optimization of oxygen delivery in elective surgeries for high-risk patients.	Pulmonary artery catheter
Noblett et al., 2006 [41]	Evaluation of transesophageal echocardiography-guided hemodynamic optimization therapy in terms of the outcomes of patients undergoing colectomy.	Transesophageal Doppler
Harten et al., 2008 [42]	Randomized study evaluating the effect of hemodynamic optimization on renal function in patients undergoing emergency laparotomy.	FloTrac Vigileo system
Kapoor et al., 2008 [43]	Evaluation of GDT in patients with moderate- to high-risk cardiac surgery.	FloTrac Vigileo system
Mayer et al., 2010 [44]	Evaluation of GDT based on the monitoring of the blood pressure wave in high-risk surgical patients.	FloTrac Vigileo system
Benes et al., 2010 [45]	Evaluation of hemodynamic optimization by fluid loading based on data obtained by Vigileo.	FloTrac Vigileo system

GDT: goal-directed therapy, O2: oxygen

Author-year (ref)	Objective	Technology
Cecconi et al., 2011 [11]	Evaluation of hemodynamic optimization therapy for patients undergoing total hip arthroplasty under regional anesthesia.	FloTrac Vigileo system
Lobo et al., 2011 [46]	Evaluation of restrictive or conventional strategies for crystalloid administration during GDT in high-risk surgical patients.	LiDCO monitoring system
Salzwedel et al., 2013 [47]	Randomized study evaluating GDT based on the variation of the radial arterial pulse and the cardiac index and the effects of GDT on postoperative complications of major abdominal surgeries.	FloTrac Vigileo system
van Beest et al., 2014 [48]	Evaluation of the effect of the tissue oxygenation optimization-based protocol on perioperative complication rates.	FloTrac Vigileo system
Pearse et al., 2014 [25]	Evaluation of the clinical effectiveness of perioperative use of the cardiac output-guided hemodynamic therapy algorithm.	LiDCO monitoring system
Cannesson et al., 2015 [49]	Evaluation of the effects of the systematic implementation of GDT on the length of hospital stay and the incidence of complications after high-risk abdominal surgeries.	EV 1000 (Edwards Lifesciences, Irvine, CA, USA)
Kumar et al., 2015 [50]	Randomized study evaluating the impact of GDT on the cardiac index and O ₂ extraction rate in patients undergoing abdominal surgery.	FloTrac Vigileo system
Calvo-Vecino et al., 2018 [51]	Randomized study evaluating the impact of GDT on the outcome in major surgeries compared to controls.	Transesophageal Doppler
GDT: goal-directed therapy, O ₂ : oxygen		

The risk-of-bias assessment for each of the included studies can be visualized (Figs. 2a and 2b). Most studies have problems with blinding.

The mortality rate in the intervention group was 2.6% (125); in the control group, it was 2.5%. (122) There were no statistically significant differences in mortality rate. (Fig. 3)

The rates of infectious complications (RR = 0.66, 95% CI = 0.58-0.74), renal complications (RR = 0.68, 95% CI = 0.54-0.87) and cardiovascular complications after surgery (RR = 0.87, 95% CI = 0.76-0.99) were significantly higher in the control group. Regarding respiratory complications, there was no significant difference in the random effect analysis, but the fixed effect analysis yielded statistically significant differences (RR = 0.81; 95% CI = 0.66-0.98). (Fig. 4)

In the comparison between groups, the patients in the intervention group had shorter lengths of hospital and ICU stay. (Table 2)

Table 2

Comparison of the variables analyzed in determining the costs of the intervention and standard therapy groups.

Variables	Number of studies involved (total number of patients)	Intervention group	Standard therapy group	RR (95% CI)
Intraoperative period				
Use of vasoactive drugs	15 (N = 4104)	N = 785 (19.1%)	N = 609 (14.8%)	1.24 (1.13– 1.37)
Crystalloid fluids (mL); median (min-max)	17 (N = 2681)	3000 (1000– 6713)	2558 (1286– 6200)	
Colloid fluids (mL); median (min-max)	18 (N = 4393)	1188 (0– 2426)	817 (0– 2236)	
Blood products (mL); median (min-max)	15 (N = 4420)	244.5 (0– 825)	267 (0– 975)	
Complications (postoperative period)	21 (N = 4872)	1017 (20.9%)	1350 (27.7%)	0.75 (0.70– 0.81)
Infectious complications	17 (N = 4530)	N = 384 (8.4%)	N = 581 (12.8%)	0.66(0.58– 0.74)
Respiratory complications	19 (N = 4655)	N = 160 (3.4%)	N = 193 (4.1%)	0.82(0.67– 1.02)
Renal complications	19 (N = 4655)	N = 108 (2.3%)	N = 158 (3.4%)	0.68(0.54– 0.87)
Cardiovascular complications	21 (N = 4872)	N = 365 (7.4%)	N = 418 (8.5%)	0.87(0.76– 0.99)
Length of ICU stay (days); median (min-max)	14 (N = 1637)	1.9 (0–5)	2.9 (0–5)	
Mortality rate	20 (N = 4712)	N = 125 (2.6%)	N = 122 (2.5%)	1.02(0.80– 1.31)
Length of hospital stay (days); median (min-max)	20 (N = 4852)	10 (5–20)	11 (6–20)	
RR: relative risk compared to the intervention group; CI: confidence interval; ICU: intensive care unit; N: total number of patients; U: units				

Costs were calculated based on the costs for the intraoperative (anaesthesia, monitoring, infusions and blood products) and postoperative periods (patient care, clinical exams, routine procedures, routine laboratory and radiological exams and others). (Tables 3 and 4)

Table 3
Pattern of resource use by each patient treated with the intervention.

Resources	Quantity	Unit cost	Weighted cost
Intraoperative period			
Monitoring of cardiac output (average of prices considering only one-time-use disposable devices*)	1	R\$1363.00	R\$1363.00
Infusion of medications (risk of inotropes and vasopressors per patient)	1.24	R\$ 34.50	R\$42.78
Infusion of crystalloid fluids (per L)	3.0 L	R\$10.36	R\$31.08
Infusion of colloid fluids (per L)	1.188 L	R\$90.00	R\$106.92
Transfusions of blood products (per unit)	1.22 IU	R\$553.30	R\$675.02
Subtotal - intraoperative period	—	—	R\$2218.8
Hospital stay			
Hospital daily rate	10 days	R\$310.00	R\$3100.00
ICU daily rate	1.9 days	R\$1138.00	R\$2162.20
Subtotal - hospital stay	—	—	R\$5262.20
Treatment of complications (based on length of ICU stay)			
Infectious complications (treatment of sepsis)	1.9 days	R\$1868.00	R\$3549.2
Respiratory complications (including mechanical ventilation)	1.9 days	R\$1202.35	R\$2284.46
Renal complications (including days of dialysis)	1.9 days	R\$1449.42	R\$2753.89
Cardiovascular complications (including visits during the 1st postoperative period and the infusion of vasopressors)	1.9 days	R\$1252.70	R\$2380.13
Subtotal - treatment of complications	—	—	R\$10967.68
Total - intervention	—	—	R\$18448.68
* One-time-use or disposable devices: catheters and probes are acquired by the hospital in certain quantities, and the debt is borrowed by the hospital.			

Table 4
Pattern of resource use by each patient treated with standard therapy.

Resources	Quantity	Unit cost	Weighted cost
Intraoperative period			
Monitoring of cardiac output	0	R\$1363.00	0
Infusions of medications (risk of inotropes and vasopressors per patient)	1	R\$34.50	R\$34.50
Infusion of crystalloid fluids (per L)	2.558 L	R\$10.36	R\$26.50
Infusion of colloid fluids (per L)	0.817 L	R\$90.00	R\$73.53
Transfusions of blood products (per unit)	1.33 IU	R\$553.30	R\$735.88
Subtotal - intraoperative period	—	—	R\$870.41
Hospital stay			
Hospital daily rate	11 days	R\$310.00	R\$3410.00
ICU daily rate	2.9 days	R\$1138.00	R\$3300.2
Subtotal - hospital stay	—	—	R\$6710.20
Treatment of complications (based on length of ICU stay)			
Infectious complications (treatment of sepsis)	2.9 days	R\$1868.00	R\$5417.2
Respiratory complications (including mechanical ventilation)	2.9 days	R\$1202.35	R\$3486.81
Renal complications (including dialysis)	2.9 days	R\$1449.42	R\$4203.31
Cardiovascular complications (including visits during the 1st postoperative period and the infusion of vasopressors)	2.9 days	R\$1252.70	R\$3632.83
Subtotal - treatment of complications	—	—	R\$16740.15
Total - standard therapy	—	—	R\$24320.76

The amount was extracted from the findings of the studies (Table 2) and extrapolated to represent the costs of 1 patient from a Brazilian cost perspective.

The intervention group was monitored hemodynamically during the intraoperative period and was managed according to the data measured. The control group did not undergo hemodynamic monitoring and was treated according to the standard procedure.

Regarding complications in terms of only the length of ICU stay, those who received standard therapy showed higher costs when they developed infectious, renal, cardiovascular and respiratory complications. (Fig. 4)

Based on these calculations, when estimating costs for the two types of treatment per 1000 patients, the highest costs were observed for the standard treatment in the presence of complications. On the other hand, the benefits were not sustained for patients without complications or those who do not survive because the costs are higher for patients who are treated with goal-directed therapy. (Fig. 5)

The cost and final effectiveness of the intervention group (optimization therapy) were lower than those of the control group (conventional therapy). Although there was no significant difference in mortality rates, the cost of patients in the intervention group based on 1,000 treated patients was R\$396,024.83 less than that of patients in the standard therapy group (incremental). In addition, the intervention group showed a reduction of R\$1,630,341.47 per 1000 patients with complications. (Fig. 6)

Discussion

This study presented the first cost analysis of PHOT in Brazil. Despite the increase in costs due to the acquisition of materials and equipment and the provision of greater care in the perioperative period, PHOT showed more benefits and lower final costs for patients compared to the standard therapy for high-risk surgical patients.

The benefits of perioperative hemodynamic monitoring, especially in terms of complications and length of hospital stay are known [20, 21], but few studies have shown benefits in terms of mortality rates [22]. The data used to perform the calculations did not show differences in mortality rate, but when we performed a cost-effectiveness analysis according to the complications, we observed a significant reduction in costs (total cost of standard therapy, R\$24320.76, versus R\$1,488.68 for the intervention) and gains in effectiveness (1.0 days in the ICU and hospital) in terms of reduction in the length of stay. These facts translate to an incremental cost-effectiveness ratio of R\$5872.08 for each patient who receives only standard therapy for surgical patients if he/she presents with complications.

The results of this analysis are consistent with the economic health analyses performed in previous studies. A cost-effectiveness analysis using data from two recent clinical trials of perioperative hemodynamic therapy suggested that this treatment resulted in a net reduction in health costs and therefore was cost-effective [23, 24]. However, more recent data from clinical trials suggested that this treatment could only offer modest improvements in patient outcomes and, in some cases, provided evidence of borderline clinical effectiveness [25].

The first economic study conducted during the perioperative period was performed by Guest et al. [24] High-risk patients undergoing a goal-directed hemodynamic optimization strategy protocol experienced not only survival benefits but also lower total hospital costs (median of \$10,968.50 vs \$13,084.90). This analysis did not include the clinical benefit and therefore could be considered a pure cost minimization

study. Likewise, subsequent studies have shown cost savings when goal-directed therapy is initiated in the preoperative period [26] or the perioperative period [27]. Fenwick et al. [23] performed a cost-effectiveness analysis of a preoperative optimization regimen. In their analysis, the incremental costs were compared to the survival benefits among the different treatment groups; however, quality of life was not included in the model. Previous data from other countries have shown similar estimates in elderly patients undergoing hip fracture repair, which is considered a high-cost surgery [28].

Although the results of economic health simulations continue to suggest that perioperative hemodynamic therapy may be economical and, therefore, cost effective, these findings are sensitive to the size of the treatment effect [26, 27]. Curiously, the economic evaluations of early hemodynamic therapy in patients with severe sepsis also indicate that treatment can be cost-effective [29, 30]. However, these analyses are based on the assumption of a strong treatment effect, while the results of recent randomized clinical trials suggest that such protocols have little or no clinical benefit in patients with severe sepsis [31].

Thus, while the findings of the current study are consistent with those of previous analyses, definitive evidence provided by a large clinical effectiveness trial is necessary to confirm the economic impact of perioperative hemodynamic cardiac output optimization therapy. However, we emphasize that the findings are important because they show that perioperative hemodynamic intervention represents an excellent expenditure of public money from the SUS perspective.

However, this study has some limitations that we must discuss, such as the fact that the cost-effectiveness analysis was based on data from clinical trials with extrapolated results, despite the use of appropriate methods and long-term survival data. The cost of uncertainty in decision making was calculated along with traditional cost-effectiveness results. Cost data and results were collected only during ICU stays, and the cost-benefit analysis may involve other assumptions that could not be extrapolated with these data. Some follow-up data were missing and were resolved with an approximation by multiple imputation [32]. Furthermore, although economic health data were prospectively included in the dataset, the calculations were based on clinical results. These results should be interpreted considering that most tests are designed to detect differences in clinical effectiveness rather than cost-effectiveness. The costs of health care are more variable than the clinical results, and economic evaluations may therefore lack statistical power. The primary objective of economic evaluation is not to test hypotheses but rather to estimate incremental measures of cost-effectiveness and to provide an adequate representation of the uncertainty around these estimates.

Above all, these findings will inform a new clinical policy regarding priorities for future research.

Conclusions

Perioperative hemodynamic optimization therapy is indicated in high-risk surgical patients with greater chances of developing postoperative complications. The cost-effectiveness analysis is a useful tool for assisting in decision-making regarding the allocation of new resources for the appropriate treatment of

surgical patients. However, it is important to note that not only equipment and catheters but also trained anesthesiologists and intensivists are needed.

Abbreviations

SUS:Brazilian public unified health system; RR:Relative risk; CI:Confidence interval; PHOT:Perioperative hemodynamic optimization therapy; ICU:intensive care unit; DO2:oxygen delivery; SD:Standard deviation; GDT:goal-directed therapy; O2:oxygen; N:total number of patients; U:units; ICER:Incremental Cost-Effectiveness Ratio

Declarations

Acknowledgments

We would like to thank Vanessa D.T. Ferreira for technical advice and all CET SBA HSPE for secretarial assistance.

Author Contributions

Conceived and designed the study: JMSJr VDTF. Performed the study: JMSJr PFLM MJCC SML LMSM. Analyzed the data: JMSJr VDTF. Contributed data collect: PFLM FHSC MANO FNFCF BNF. Wrote the paper: JMSJr SML LMSM.

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

References

1. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, Gawande AA: **An estimation of the global volume of surgery: a modelling strategy based on available data.** *Lancet* 2008, **372**(9633):139-144.
2. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AH, Dellinger EP, Herbosa T, Joseph S, Kibatala PL, Lapitan MC *et al*: **A surgical safety checklist to reduce morbidity and mortality in a global population.** *N Engl J Med* 2009, **360**(5):491-499.
3. Juul AB, Wetterslev J, Gluud C, Kofoed-Enevoldsen A, Jensen G, Callesen T, Norgaard P, Fruergaard K, Bestle M, Vedelsdal R *et al*: **Effect of perioperative beta blockade in patients with diabetes undergoing major non-cardiac surgery: randomised placebo controlled, blinded multicentre trial.** *BMJ* 2006, **332**(7556):1482.
4. Khuri SF, Henderson WG, DePalma RG, Mosca C, Healey NA, Kumbhani DJ, Participants in the VANSQIP: **Determinants of long-term survival after major surgery and the adverse effect of postoperative complications.** *Ann Surg* 2005, **242**(3):326-341; discussion 341-323.
5. Lobo SM, Rezende E, Knibel MF, Silva NB, Paramo JA, Nacul F, Mendes CL, Assuncao M, Costa Filho RC, Grion CC *et al*: **Epidemiology and outcomes of non-cardiac surgical patients in Brazilian intensive care units.** *Rev Bras Ter Intensiva* 2008, **20**(4):376-384.
6. Bowdle TA: **Complications of invasive monitoring.** *Anesthesiol Clin North America* 2002, **20**(3):571-588.
7. Bennett-Guerrero E, Hyam JA, Shaefi S, Prytherch DR, Sutton GL, Weaver PC, Mythen MG, Grocott MP, Parides MK: **Comparison of P-POSSUM risk-adjusted mortality rates after surgery between patients in the USA and the UK.** *Br J Surg* 2003, **90**(12):1593-1598.
8. Bender JS, Smith-Meek MA, Jones CE: **Routine pulmonary artery catheterization does not reduce morbidity and mortality of elective vascular surgery: results of a prospective, randomized trial.** *Ann Surg* 1997, **226**(3):229-236.
9. Kern JW, Shoemaker WC: **Meta-analysis of hemodynamic optimization in high-risk patients.** *Crit Care Med* 2002, **30**(8):1686-1692.
10. Cecconi M, Corredor C, Arulkumaran N, Abuella G, Ball J, Grounds RM, Hamilton M, Rhodes A: **Clinical review: Goal-directed therapy-what is the evidence in surgical patients? The effect on different risk groups.** *Crit Care* 2013, **17**(2):209.
11. Cecconi M, Fasano N, Langiano N, Divella M, Costa MG, Rhodes A, Della Rocca G: **Goal-directed haemodynamic therapy during elective total hip arthroplasty under regional anaesthesia.** *Crit Care* 2011, **15**(3):R132.
12. Benes J, Zatloukal J, Simanova A, Chytra I, Kasal E: **Cost analysis of the stroke volume variation guided perioperative hemodynamic optimization - an economic evaluation of the SVVOPT trial results.** *BMC Anesthesiol* 2014, **14**:40.

13. Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JA *et al*: **The Cochrane Collaboration's tool for assessing risk of bias in randomised trials.** *BMJ* 2011, **343**:d5928.
14. **Informatics Department of the Brazilian Public Healthcare System**
15. **Management System of the Table of Procedures, Medications and Devices of SUS.**
16. **Brazilian Healthcare Database: Banco de Preços em Saúde**
[<http://portalsaude.saude.gov.br/index.php/cidadao/principal/banco-de-precos-em-saude?layout=edit&id=8667>]
17. Sogayar AM, Machado FR, Rea-Neto A, Dornas A, Grion CM, Lobo SM, Tura BR, Silva CL, Cal RG, Beer I *et al*: **A multicentre, prospective study to evaluate costs of septic patients in Brazilian intensive care units.** *Pharmacoeconomics* 2008, **26**(5):425-434.
18. **Price Index for Medical Procedures: CBHPM** [<http://cbr.org.br/wp-content/uploads/2013/05/COMUNICADO-CBHPM-2015-2016.pdf>]
19. **Development and Management. Prices Panel** [<http://paineldeprescos.planejamento.gov.br/analise-materiais>]
20. Hamilton MA, Cecconi M, Rhodes A: **A systematic review and meta-analysis on the use of preemptive hemodynamic intervention to improve postoperative outcomes in moderate and high-risk surgical patients.** *Anesth Analg* 2011, **112**(6):1392-1402.
21. Gurgel ST, do Nascimento P, Jr.: **Maintaining tissue perfusion in high-risk surgical patients: a systematic review of randomized clinical trials.** *Anesth Analg* 2011, **112**(6):1384-1391.
22. Aya HD, Cecconi M, Hamilton M, Rhodes A: **Goal-directed therapy in cardiac surgery: a systematic review and meta-analysis.** *Br J Anaesth* 2013, **110**(4):510-517.
23. Fenwick E, Wilson J, Sculpher M, Claxton K: **Pre-operative optimisation employing dopexamine or adrenaline for patients undergoing major elective surgery: a cost-effectiveness analysis.** *Intensive Care Med* 2002, **28**(5):599-608.
24. Guest JF, Boyd O, Hart WM, Grounds RM, Bennett ED: **A cost analysis of a treatment policy of a deliberate perioperative increase in oxygen delivery in high risk surgical patients.** *Intensive Care Med* 1997, **23**(1):85-90.
25. Pearse RM, Harrison DA, MacDonald N, Gillies MA, Blunt M, Ackland G, Grocott MP, Ahern A, Griggs K, Scott R *et al*: **Effect of a perioperative, cardiac output-guided hemodynamic therapy algorithm on outcomes following major gastrointestinal surgery: a randomized clinical trial and systematic review.** *JAMA* 2014, **311**(21):2181-2190.
26. Bartha E, Davidson T, Hommel A, Thorngren KG, Carlsson P, Kalman S: **Cost-effectiveness analysis of goal-directed hemodynamic treatment of elderly hip fracture patients: before clinical research starts.** *Anesthesiology* 2012, **117**(3):519-530.
27. Ebm CC, Sutton L, Rhodes A, Cecconi M: **Cost-effectiveness in goal-directed therapy: are the dollars spent worth the value?** *J Cardiothorac Vasc Anesth* 2014, **28**(6):1660-1666.

28. Bartha E, Davidson T, Brodtkorb TH, Carlsson P, Kalman S: **Value of information: interim analysis of a randomized, controlled trial of goal-directed hemodynamic treatment for aged patients.** *Trials* 2013, **14**:205.
29. Jones AE, Troyer JL, Kline JA: **Cost-effectiveness of an emergency department-based early sepsis resuscitation protocol.** *Crit Care Med* 2011, **39**(6):1306-1312.
30. Talmor D, Greenberg D, Howell MD, Lisbon A, Novack V, Shapiro N: **The costs and cost-effectiveness of an integrated sepsis treatment protocol.** *Crit Care Med* 2008, **36**(4):1168-1174.
31. Pro CI, Yealy DM, Kellum JA, Huang DT, Barnato AE, Weissfeld LA, Pike F, Terndrup T, Wang HE, Hou PC *et al*: **A randomized trial of protocol-based care for early septic shock.** *N Engl J Med* 2014, **370**(18):1683-1693.
32. McCleary L: **Using multiple imputation for analysis of incomplete data in clinical research.** *Nurs Res* 2002, **51**(5):339-343.
33. Bonazzi M, Gentile F, Biasi GM, Migliavacca S, Esposti D, Cipolla M, Marsicano M, Prampolini F, Ornaghi M, Sternjakob S *et al*: **Impact of perioperative haemodynamic monitoring on cardiac morbidity after major vascular surgery in low risk patients. A randomised pilot trial.** *Eur J Vasc Endovasc Surg* 2002, **23**(5):445-451.
34. Venn R, Steele A, Richardson P, Poloniecki J, Grounds M, Newman P: **Randomized controlled trial to investigate influence of the fluid challenge on duration of hospital stay and perioperative morbidity in patients with hip fractures.** *Br J Anaesth* 2002, **88**(1):65-71.
35. Conway DH, Mayall R, Abdul-Latif MS, Gilligan S, Tackaberry C: **Randomised controlled trial investigating the influence of intravenous fluid titration using oesophageal Doppler monitoring during bowel surgery.** *Anaesthesia* 2002, **57**(9):845-849.
36. Gan TJ, Soppitt A, Maroof M, el-Moalem H, Robertson KM, Moretti E, Dwane P, Glass PS: **Goal-directed intraoperative fluid administration reduces length of hospital stay after major surgery.** *Anesthesiology* 2002, **97**(4):820-826.
37. Sandham JD, Hull RD, Brant RF, Knox L, Pineo GF, Doig CJ, Laporta DP, Viner S, Passerini L, Devitt H *et al*: **A randomized, controlled trial of the use of pulmonary-artery catheters in high-risk surgical patients.** *N Engl J Med* 2003, **348**(1):5-14.
38. Wakeling HG, McFall MR, Jenkins CS, Woods WG, Miles WF, Barclay GR, Fleming SC: **Intraoperative oesophageal Doppler guided fluid management shortens postoperative hospital stay after major bowel surgery.** *Br J Anaesth* 2005, **95**(5):634-642.
39. Pearse R, Dawson D, Fawcett J, Rhodes A, Grounds RM, Bennett ED: **Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomised, controlled trial [ISRCTN38797445].** *Crit Care* 2005, **9**(6):R687-R693.
40. Lobo SM, Lobo FR, Polachini CA, Patini DS, Yamamoto AE, de Oliveira NE, Serrano P, Sanches HS, Spegiorin MA, Queiroz MM *et al*: **Prospective, randomized trial comparing fluids and dobutamine optimization of oxygen delivery in high-risk surgical patients [ISRCTN42445141].** *Crit Care* 2006, **10**(3):R72.

41. Noblett SE, Snowden CP, Shenton BK, Horgan AF: **Randomized clinical trial assessing the effect of Doppler-optimized fluid management on outcome after elective colorectal resection.** *Br J Surg* 2006, **93**(9):1069-1076.
42. Harten J, Crozier JE, McCreath B, Hay A, McMillan DC, McArdle CS, Kinsella J: **Effect of intraoperative fluid optimisation on renal function in patients undergoing emergency abdominal surgery: a randomised controlled pilot study (ISRCTN 11799696).** *Int J Surg* 2008, **6**(3):197-204.
43. Kapoor PM, Kakani M, Chowdhury U, Choudhury M, Lakshmy, Kiran U: **Early goal-directed therapy in moderate to high-risk cardiac surgery patients.** *Ann Card Anaesth* 2008, **11**(1):27-34.
44. Mayer J, Boldt J, Mengistu AM, Rohm KD, Suttner S: **Goal-directed intraoperative therapy based on autocalibrated arterial pressure waveform analysis reduces hospital stay in high-risk surgical patients: a randomized, controlled trial.** *Crit Care* 2010, **14**(1):R18.
45. Benes J, Chytra I, Altmann P, Hluchy M, Kasal E, Svitak R, Pradl R, Stepan M: **Intraoperative fluid optimization using stroke volume variation in high risk surgical patients: results of prospective randomized study.** *Crit Care* 2010, **14**(3):R118.
46. Lobo SM, Ronchi LS, Oliveira NE, Brandao PG, Froes A, Cunrath GS, Nishiyama KG, Netinho JG, Lobo FR: **Restrictive strategy of intraoperative fluid maintenance during optimization of oxygen delivery decreases major complications after high-risk surgery.** *Crit Care* 2011, **15**(5):R226.
47. Salzwedel C, Puig J, Carstens A, Bein B, Molnar Z, Kiss K, Hussain A, Belda J, Kirov MY, Sakka SG *et al*: **Perioperative goal-directed hemodynamic therapy based on radial arterial pulse pressure variation and continuous cardiac index trending reduces postoperative complications after major abdominal surgery: a multi-center, prospective, randomized study.** *Crit Care* 2013, **17**(5):R191.
48. van Beest PA, Vos JJ, Poterman M, Kalmar AF, Scheeren TW: **Tissue oxygenation as a target for goal-directed therapy in high-risk surgery: a pilot study.** *BMC Anesthesiol* 2014, **14**:122.
49. Cannesson M, Ramsingh D, Rinehart J, Demirjian A, Vu T, Vakharia S, Imagawa D, Yu Z, Greenfield S, Kain Z: **Perioperative goal-directed therapy and postoperative outcomes in patients undergoing high-risk abdominal surgery: a historical-prospective, comparative effectiveness study.** *Crit Care* 2015, **19**:261.
50. Kumar L, Kanneganti YS, Rajan S: **Outcomes of implementation of enhanced goal directed therapy in high-risk patients undergoing abdominal surgery.** *Indian J Anaesth* 2015, **59**(4):228-233.
51. Calvo-Vecino JM, Ripolles-Melchor J, Mythen MG, Casans-Frances R, Balik A, Artacho JP, Martinez-Hurtado E, Serrano Romero A, Fernandez Perez C, Asuero de Lis S *et al*: **Effect of goal-directed haemodynamic therapy on postoperative complications in low-moderate risk surgical patients: a multicentre randomised controlled trial (FEDORA trial).** *Br J Anaesth* 2018, **120**(4):734-744.

Figures

Figure 1- Flow Diagram




Figure 1

Flow Diagram. PRISMA diagram showing the inclusion and exclusion process used for the literature search and review.




Figure 2

Summary of risk-of-bias assessment. (A) Risk-of-bias summary for each included study. (B) Summary of domains for risk of bias for included studies.

Bonazzi, et al- 2002
Venn, et al- 2002
Conway, et al- 2002
Gan, et al- 2002
Sandham, et al- 2003
Wakeling, et al- 2005
Pearse, et al - 2005
Lobo, et al - 2006
Noblett, et al- 2006
Harten, et al- 2008
Kapoor, et al- 2008
Mayer, et al- 2010
Benes, at al- 2010
Ceconi, et al- 2011
Lobo, et al- 2011
Van Beest, et al- 2014
Pearse, et al-2014
Cannesson, et al- 2015
Kumar, et al- 2015
Calvo-Vecino, et al- 2018

Total (Fixed Effects)
Total (Random Effects)




Figure 3

Forest Plot for Mortality in the Intervention and Control Groups.




Figure 4

Forest Plot of Complications in the ICU; Comparison Between the Intervention and Control Groups. A - infectious; B - respiratory; C - renal; D - cardiovascular.




Figure 5

Incremental Cost-Effectiveness Ratio (ICER) per Patient Based on Complications.




Figure 6

Comparison of Costs per 1000 Patients Treated Between Groups in ICU Stay.




Figure 7

Comparison of Costs per 1000 Patients Treated Between Groups According to the Type of Complication.