

# Knowledge and Attitudes on First aid among Advanced Level Students in Gampaha Educational Zone, Sri Lanka.

**Prashan Alahakoon**

Management and Science University

**Kalpani Bandaranayaka** (✉ [kalpanee.b@gmail.com](mailto:kalpanee.b@gmail.com))

Management and Science Institute

**Pathirage Perera**

Institute of Indigenous Medicine, University of Colombo

**Chathura Wijesundara**

Ministry of Health, Nutrition and Indigenous Medicine

---

## Research Article

**Keywords:** first aid, school students, knowledge, attitudes

**Posted Date:** December 3rd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-112739/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# **Knowledge and Attitudes on First aid among Advanced Level Students in Gampaha Educational Zone, Sri Lanka.**

PrashanViranga Alahakoon<sup>1</sup>

Faculty of Health & Life Sciences, Management & Science University, University Drive, Off PersiaranOlahraga, 40100 Shah Alam, Selangor, Malaysia.

[prashanviranga@gmail.com](mailto:prashanviranga@gmail.com)

Pathirage Kamal Perera<sup>3</sup>

Institute of Indigenous Medicine, University of Colombo, Rajagiriya, Sri Lanka

[drpkamal@gmail.com](mailto:drpkamal@gmail.com)

Chatura Wijesundara<sup>4</sup>

Ministry of Health, Sri Lanka

[chaturawijesundara@gmail.com](mailto:chaturawijesundara@gmail.com)

KalpaniOmantha Bandaranayaka<sup>2</sup>

Management and Science Institute, Colombo 03, Sri Lanka

[kalpanee.b@gmail.com](mailto:kalpanee.b@gmail.com)

Corresponding author: KalpaniOmantha Bandaranayaka<sup>2</sup>

Management and Science Institute, Colombo 03, Sri Lanka

[kalpanee.b@gmail.com](mailto:kalpanee.b@gmail.com)

## **Abstract**

### **Background**

First aid is the initial care given in an emergency, to protect life and to prevent worsening of the condition of the victims, until they are undertaken by a professional medical person. It is essential to know basic first aid as a responsible person in society. The current study was conducted to assess the knowledge, attitudes and practices on first aid among advanced level students in government schools in Gampaha educational zone.

### **Methods**

A descriptive cross-sectional study was conducted among 510 grade 12 students in five randomly selected schools. A pre-tested self-administered questionnaire consisting of 25 multiple choice questions on knowledge and 10 statements formed as a Likert scale to assess attitudes. Chi-square test was used to assess the associations and a p-value <0.05 was considered as significant.

### **Results**

Student's mean first aid knowledge was  $57.4\% \pm 13.5\%$  according to the score grading and 1% (n=5) were categorized as having "inadequate knowledge", 30.6% (n=156) had "moderate knowledge", 61.4% (n=313) were having "good knowledge" and 7.1% (n=36) were categorized as having "excellent knowledge". Out of the sample, only 16.9% (n=86) students were previously trained in first aid. The overall attitudes towards first aid was positive and most of the students (98.8%, n=504) believed it is essential to have first aid knowledge. Their willingness to give first aid for emergencies was generally positive. There was only a slight correlation between knowledge and attitudes ( $r=0.134$ ). There was a statistically significant relationship between first aid knowledge and advanced level stream ( $\chi^2=31.7$ ,  $p<0.001$ ) where biology students were more knowledgeable than students in other streams. The students who had followed health science in their O/L had a significantly higher knowledge score when compared to others ( $\chi^2=12.4$ ,  $p=0.006$ ). However, previous training on first aid and gender had no significant relationship with their knowledge ( $p>0.05$ ).

### **Conclusions**

Addition of first aid as a part of the school curriculum and improving the standards of the existing first aid training programs is recommended. Further studies should be undertaken to assess the practical skills of students, as the current study mainly focused on the assessment of the theory knowledge.

**Keywords:** first aid, school students, knowledge, attitudes

## Background

First aid is the immediate care provided to a victim of injury or sudden illness in emergencies until they are undertaken by a professional medical person [1]. The person providing first aid should be able to assess the injured as soon as possible and give basic first aid correctly and then contact the proper medical facilities.

In 1990, there were more than 350 million new cases of injuries recorded globally and it has increased to 520 million new cases by 2017 [2]. Even though injuries are considered as an important cause of health loss worldwide, the mortality rate has declined by 2017 from 1079 to 738 per 100 000 individuals [2]. In 2013, 973 million people were subjected to injuries which required medical attention [2]. Most significant causes included car crashes (29%), self-harm (17.6%), falls (11.6%) and violence (8.5%). However, out of the total cases, only around 6% required hospital admission. Out of those cases, the most common reason (38.5%) for admission was having fractures [2].

Traumatic injuries are the main cause of hospitalization over the last ten years in Sri Lanka and more than 1 million people were hospitalized in 2016 due to injuries. It was ranked number 10 in terms of mortality with 1675 hospital deaths being reported in 2016 [3]. A community survey on injury occurrence among residents in a semi-urban area showed that the incidence of injuries among all age groups was 17.7 per 1000 population over six months. The common injuries were falling from a height, road accidents, animal bites, sports injuries, cuts and abrasions, poisoning, and burns [4]. When considering school-based injuries or accidents, the majority occur during 'free periods', within the classroom and are minor injuries [5].

Knowledge on first aid is important for everyone [5]. The person that will use first aid must know its rules and practices properly to achieve optimum results. A study conducted at Mosul University, Iraq on first aid knowledge using a "true/false" questionnaire among 124 students from the school of science and school of education revealed that the overall knowledge is 46%. The majority (n=57, 96%) of the students believed that they need to have proper training on first aid activities [6]. A similar study done in Turkey among 134 students of the Vocational School of Health Services at Kırklareli University revealed that 65.7% (n=88) of students had not received any formal first aid training. Of the sample, 65.7% (n=88) of students had not received any first aid knowledge before. However, 82.8% (110) knew the pulse rate of an adult while 56.7% (75) knew the respiratory rate of an adult. About one-fourth of the sample (n=35, 26.1%) stated that gasoline, bleaching liquid or

similar substances should not be used to induce vomiting if a patient has ingested poison [7]. Lack of local research makes it imperative to conduct large-scale research to assess knowledge, attitudes and practices on first aid to develop evidence-based interventions.

## **METHODS**

### **Study Aim:**

The current study was conducted to assess knowledge, attitudes and practices on first aid among advanced level students in government schools in Gampaha educational zone.

**Study design:** Descriptive cross-sectional study

### **Study setting:**

The study was conducted in Government schools located in Gampaha educational zone. According to the national school census 2016 – preliminary report, total government schools in Gampaha educational zone is 48, out of which only 14 schools had classes up to advanced level.

Table: 1 – Selected schools from the list of Schools at the Western Provincial Education Department

### **Study participants**

The study population was grade 12 and 13 students of both sexes (age range = 18-19 years) in government schools in Gampaha educational zone.

### **Sample size calculation**

Sample size determination was through the following equation [8] ;  $n = Z^2 P(1-P) / d^2$

Z = Critical value of specified confidence level, for this study the confidence level set at 95% and thus the critical value is 1.96. As there are no studies done in Sri Lanka to assess the knowledge, attitudes or practice levels regarding the first aid at the community level, the prevalence of poor knowledge was considered as 50% to maximize the sample size.

Applying the values to the equation;

$$n = (1.96 \times 1.96 \times 50 \times 50) / 5 \times 5 = 384$$

A non-response rate of 40% was considered. Therefore, the final sample size was increased to 540. The sample size achieved was 510.

### **Sampling techniques**

The list of schools with advanced level students (14) was obtained from the western provincial education department and rearranged according to the alphabetical order. Based on the logistical feasibility it was decided to select five (5) schools from the prepared sampling frame randomly. Accordingly, five (5) schools were selected based on computer-generated random numbers. The total sample size was divided equally among the selected schools.

### **Study instrument**

A self-administered questionnaire was used to collect data. It was developed in the English language (Annexure 1) and then translated to Sinhala and Tamil languages. The face and content validity was assessed by a panel of experts. The questionnaire contains the following parts; socio-demographic details, knowledge and attitudes on first aid. Pre-testing of the questionnaire was done among A/L students in a school which was not selected for the main study and some adjustments were made in the wording of the questionnaire. The data collection was done by the principal investigator. After explaining the details of the study, informed written consent was obtained.

### **Statistical analysis**

Primary data were entered to Microsoft Excel 2007 and exported to Statistical Package of Social Science for windows version 21 (SPSS Inc., Chicago, IL, USA). The level of knowledge and attitudes were considered as dependent variables. Numerical variables were described using mean and standard deviation. Categorical variables were expressed as percentages. The bivariate analysis was done using chi-square test. To calculate a total first aid knowledge score, one (1) mark was given to each correct answer and zero (0) marks were given for each wrong answer in the knowledge section. The total scores were then converted to percentages and were graded according to a scale where a score of  $\leq 24$  was considered as "Inadequate knowledge", scores between 25 to 49 were classified as "Moderate knowledge", scores between 50 and 74 were considered as "Good knowledge" and scores  $\geq 75$  were grouped as "Excellent knowledge". A p-value  $\leq 0.05$  was considered as significant.

## **Administrative and Ethical clearance**

Ethical clearance was obtained from National Institute of Health Science, Kalutara, Sri Lanka (ERC clearance No: NIHS/REC/17/05R). The information sheet was provided with all necessary information to the participants thus allowing asking questions as well. Informed written consent was obtained before the administration of the questionnaire. Administrative clearance was obtained from the Director, Department of Education - Western Province and principals of respective schools selected.

## **RESULTS**

### **Socio-demographic information of students**

The response rate was 94.4% and the final sample size achieved was 510. In the study sample, 185(36.3%) were males and all students were aged between 17(26.7%) to 18(74.1%)(Table 2). More than a quarter of the students (n= 144, 28.3%), were studying in biology stream while 122(23.9%) were in commerce stream, 144 (28.3%) were in mathematics stream and the rest were studying in the arts stream. Nearly half of the students (n=254, 49.8%) followed the health science subject during their ordinary level studies (O/L) and only 16.9% (n=86) of the students were previously trained on first aid (Table 2).

Table 2: Socio-demographic and education-related information of the respondents

### **Knowledge on first aid**

The mean knowledge score was 57.4 (SD=13.5) where the maximum was 89 and the minimum was 16 (Figure 1). According to the first aid knowledge grading, 1% (n=5) were categorized as having “inadequate knowledge”, 30.6% (n=156) had “moderate knowledge”, 61.4% (n=313) were having “good knowledge” and 7.1% (n=36) were categorized as having “excellent knowledge” (Table 3). The majority (n=486, 95.3%) were able to define “first aid” and 56.9% (n=290) knew first steps in managing a first aid situation. When considering the first aid knowledge for given emergencies, the majority was correct in following situations; 62.9% (n=321) knew the correct first aid for deep bleeding cut on the palm, 81.4% (n=415) were able to provide first aid for broken glass piece sticking in the leg, 89.2% (n=455) students were aware of first aid steps for electric shocks, 88.2% (n=450) knew correct first aid for snake bites before hospitalization and 87.6% (n=447) knew how to check for breathing in a person. In contrast, the majority was wrong in the following situations;

66.1%(n=337) were not aware of first aid to stop nasal bleeding, 74.3% (n=379) were unable to give the correct answer for first aid in sprains/strains, 72.4% (n=369) students were not aware of the correct first aid procedure for accidental ingestion of poisons and 82.7%(n=422) were not aware of correct CPR ratio. In following situations nearly half of the population were having the awareness; the first step in managing an emergency, first-aid for bone fracture in leg, first aid procedure for a choking person, first aid for burn injuries, first aid for seizures, steps in managing a heart attack and importance of CPR. More than three fourth of the sample(n=388, 76.1%) knew the shortcode (1990) number to contact “Suwasariya” free ambulance service (Table 4).

Figure 1: Distribution of the first aid knowledge scores among students.

Table 3: Students first aid knowledge according to the grading

Table 4: Knowledge score of the respondents

### **Attitudes on first aid**

The overall attitudes regarding first aid were positive and most of the students (n=504, 98.8%) believed it is essential to have first aid knowledge. Their willingness to give first aid in the following emergencies was high; heart attacks, drowning, severe bleeding, and road traffic accidents (Table 5). The common reasons to hesitate to provide first aid assistance for a victim of road traffic accident were lack of knowledge in first aid (n=202 39.6%), legal consequences (n=176, 34.5%), and fear of road traffic accidents (n=44, 8.6%). Nearly 20% (n=108) had a phobia of seeing blood (Table 5).

Table 5: Attitudes towards first aid

### **Correlation of first aid knowledge with attitudes**

There was only a slight correlation ( $r=0.134$ ,  $p=0.003$ ) between knowledge and attitudes indicating almost no relationship between knowledge and attitudes on first aid.

### **Factors associated with knowledge**

A statistically significant association was seen between first aid knowledge and advanced level stream ( $\chi^2=15.26$ ,  $p<0.001$ ; Table 6) where knowledge of students studying in the biology stream was higher than others. The students who had followed health science during their O/L studies had a significantly higher knowledge score when compared to others ( $\chi^2=12.4$ ,  $p=0.006$ ; Table 6). However, previous training on first aid had no significant



relationship with the first aid knowledge ( $\chi^2=7.2$ ,  $p=0.064$ ; Table 6). No statistical significance was found between first aid knowledge and gender ( $\chi^2=3.8$ ,  $p=0.284$ ; Table 6).

Table 6: Correlation of socio-demographic factors and first aid knowledge

## DISCUSSION

The study shows overall fair knowledge and positive attitudes towards first aid among advanced level students in Gampaha educational zone. More than half of the students were having a good knowledge while 7.1% were excellent in their first-aid knowledge. In contrast, a study done in Dehradun, India among grade 9 to 12 students showed that 33.3% had only a partial knowledge while only 23.1% had adequate knowledge on first aid [9]. Another study conducted in Pondicherry, India among grade 9 students shows that the majority (76.7%) have inadequate knowledge with only 23.3% having moderate knowledge on first aid [10]. A study in Telugana, India among undergraduate students showed that only 38% had inadequate knowledge [11].

In some specific first aid situations, students showed poor knowledge, such as nasal bleeding, sprain/strains, first aid for ingestion of poisons and correct CPR ratio. These observations were inconsistent with Semwal *et al.*, where it shows that nearly 99% of the students don't know first aid for nasal bleeding and sprain/strains [9]. In this study for the situation of accidental ingestion of poisons; 42.4% selected the option of "encourage the person to vomit" and 29.6% selected the "giving coconut milk" option. Both statements are wrong first aid measures. In Sri Lankan rural society, giving coconut milk for the cases of oral ingestion of poison is very popular as revealed in a study done in Sri Lanka about plant and kerosene oil poisoning among children, where 21.7% parents have practised forceful ingestion of coconut milk as a first-aid measure [12]. This clearly shows that some Sri Lankans still have a misconception regarding first-aid practices which could even lead to fatal outcomes [13]. In the above study, only 17.3% of the students knew the correct CPR ratio. In a similar study done in Sri Lanka by Priyangika *et al.*, in 2015 among senior school prefects in Galle education zone, only 26% of students knew the correct CPR ratio and those who knew the correct ratio were previously trained on first aid. The main reason for poor knowledge recorded in the current study is because the majority of the students (83.1%) had not received any formal training on first aid. A study done in Turkey by Metin and Mutlu in

2010 among university students shows that 53% were aware of the correct CPR ratio which is a comparatively high figure.

In the present study the overall attitudes towards first aid were positive (77.1%). A similar pattern was seen in a study done in New Zealand [14]. In this study, 95% of the students believed it is essential to have first aid knowledge with them however the knowledge and the attitudes were not significantly correlated ( $r = 0.134$ ). In another study done in Iraq by Makhlef, 2013 among university students, 96% of the respondents stated it's necessary to learn first aid [6]. Students in the current study showed a high willingness to provide first aid assistance for emergencies. A similar scenario is seen in other studies too [5][15]. However, in certain instances, people are hesitant to provide first aid assistance, for instance, road traffic accidents. The reasons for such hesitations were lack of knowledge in first aid (39.6%) and potential, legal consequences (34.5%). A study done in India by Patidar and Sharma in 2014 among school students on basic life support further illustrates the fear for legal actions leading to hesitancy in becoming a first aid provider. [16] Similar findings are reported in a study done in North Carolina among high school students, where legal issues and fear of infections are the main reasons for not intervening to provide first aid [17].

The evidence from this study indicates that a student's educational background influences their first aid knowledge with those studying in biology stream for the advanced level examinations and students who have studied health science during their O/Ls having a better first aid knowledge. This is because first aid knowledge is related to the health and life sciences fields. However, previous training in first aid did not influence their first aid knowledge. Similarly, this is noted in a nationwide survey conducted in Norway where most of the population were trained in first aid, but their theoretical first aid knowledge was poor than expected [15]. Therefore merely conducting training programmes on first aid is inadequate to improve and maintain the level of knowledge in the long run. A pilot study done in Norway suggests first aid education in preschools has a better outcome when the application of basic first aid for given situations is considered [18]. Therefore the addition of first aid education as a practical subject for the school curriculum for all grades will lead to better outcomes in terms of first aid knowledge among school students.

## **CONCLUSION AND RECOMMENDATION**

The majority (61.4%) of the students had “good” knowledge on first aid. A statistically significant association was seen between first aid knowledge and advanced level stream where knowledge of students studying in the biology stream was higher than others. The students who had followed health science during their O/L studies had a significantly higher knowledge score when compared to others. A majority showed positive attitudes towards providing first aid assistance in emergencies and their high willingness is observed.

It is recommended to add first aid as a part of the school curriculum and improve the standard of the all existing first aid training programs in the country. Further studies should be undertaken to assess the practical skills of students, as the current study mainly focused on the assessment of the theory knowledge.

## **DECLARATIONS**

### **Acknowledgements**

None.

### **Funding**

Self-funded

### **Authors' contributions**

The conception of the project hypothesis was by KOB, CW and PVA. The research proposal was designed by PKP, KOB, PVA and CW. Data collection tools were design & developed by KOB, CW and PVA. KOB, PKP, CW supervised the project. The manuscript was written by PVA. The manuscript was revised by KOB, PKP and CW. All authors read and accepted the final draft of the manuscript.

### **Competing interests**

The authors declare that they have no competing interests.

### **Consent for publication**

Not applicable.

## **Ethics approval and consent to participate**

Ethical clearance was obtained from National Institute of Health Science, Kalutara, Sri Lanka (ERC clearance No: NIHS/REC/17/05R). All methods were performed in accordance with the relevant guidelines and regulations being approved. Informed written consent was obtained by all study participants who were 18 years and above while for minors, consent was obtained from a parent and/or legal guardian.

## **Availability of data and materials**

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

## **REFERENCES**

1. National Safety Council. First aid: Taking action. New York: McGraw-Hill. 2007. <https://www.nsc.org/safety-training/first-aid/courses>. Accessed on August, 2020.
2. James SL, Castle CD, Dingels ZV, Fox JT, Hamilton EB, Liu Z. *et al.*, Global injury morbidity and mortality from 1990 to 2017: results from the Global Burden of Disease Study 2017. *Inj Prev: Journal of the International Society for Child and Adolescent Injury Prevention*. 2020;0:1-19 doi:10.1136/injuryprev-2019-043494
3. Annual Health Bulletin. Ministry of Health, Nutrition and Indigenous Medicine Sri Lanka. 2016. [http://www.health.gov.lk/moh\\_final/english/public/elfinder/files/publications/AHB/2017/AHB\\_2016.pdf](http://www.health.gov.lk/moh_final/english/public/elfinder/files/publications/AHB/2017/AHB_2016.pdf) [Accessed on July 2020]
4. Weerasinghe IE, Rajapaksa AW, Premaratne CS, Jayatilake JAMS, And Dharmaratne SD. Injury occurrence among residents in a semi-urban area in Sri Lanka; A community survey. *International Journal of Scientific and Research Publications*. 2015;5(3):1–8. Available at: [www.ijsrp.org/research-paper-0315/ijsrp-p3951.pdf](http://www.ijsrp.org/research-paper-0315/ijsrp-p3951.pdf).
5. Priyangika KGG, and Hettiarachchi M. Knowledge, Attitudes and Practices on First Aid Measures among Senior School Prefects in Galle Education Division, Sri Lanka. *Proceedings of 8<sup>th</sup> International Research Conference, KDU, 2015;(November):36–41*. Available at: <http://www.kdu.ac.lk/proceedings/irc2015/2015/ahs-008.pdf>.

6. Makhlef HH. Assessment of Mosul University Students' Knowledge about First Aid. *Kufa Journal for Nursing Sciences*. 2013;3(1):278–285. Available at: <http://www.iasj.net/iasj?func=fulltext&aId=75656>.
7. Metin I. and Mutlu C. Level of knowledge about first aid of the university students Std deviation. 2010;8:262–265
8. Lwanga S. and Lemeshow S. Sample Size Determination in Health Studies. A Practical Manual. 1991. [http://www.tbrieder.org/publications/books\\_english/lemeshow\\_samplesize.pdf](http://www.tbrieder.org/publications/books_english/lemeshow_samplesize.pdf)
9. Semwal J. Bakshi RK. Juyal R. et al., Study of knowledge and attitudes to first aid among school children of Doiwala block, Dehradun. *International Journal of Community Medicine And Public Health*. 2017;4(8):2934. doi: 10.18203/2394-6040.ijcmph20173348.
10. Geetha C. Knowledge on selected first aid measures among school children, Puducherry', *International journal of applied research*. 2016;2:504–506.
11. Rajakumari A. Knowledge attitude and practices on undergraduate students regarding first aid measures. *Indian Journal Scientific Research and Technologies*. 2015;3(3):22–25.
12. Dayasiri MBKC. Jayamanne SF. and Jayasinghe CY. Kerosene Oil Poisoning among Children in Rural Sri Lanka, *International Journal of Pediatrics*. 2017;1–7. doi: 10.1155/2017/8798610.
13. Dayasiri MBKC. Jayamanne SF. and Jayasinghe CY. Plant Poisoning among Children in Rural Sri Lanka, *International Journal of Paediatrics*. 2017;6187487. doi: 10.1155/2017/6187487.
14. Parnell MM. Pearson J. Galletly DC. and Larsen PD. Knowledge of and attitudes towards resuscitation in New Zealand high-school students. *Emergency Medicine Journal*. 2006; 23(12):899–902. doi: 10.1136/emj.2006.041160.
15. Kvåle BH. Tine S. Johan A. and Torben W. A nationwide survey of first aid training and encounters in Norway. *BMC Emergency Medicine*. 2017;17(1):6. doi: 10.1186/s12873-017-0116-7.
16. Patidar AB. and Sharma A. Attitude of School Children towards Basic Life Support in Punjab, India. *International Journal of Health Sciences and Research* 2014;4(June):193–201.
17. Michael H. Bachman W. Price W. Martin R. and Dennis NH. Willingness of high school students to perform cardiopulmonary resuscitation and automated external

defibrillation. Pre hospital Emergency Care. 2003;7(2):219–224. doi: 10.1080/10903120390936815

18. Bollig G. Myklebust AG. And Østringen K. Effects of first aid training in the kindergarten - a pilot study, Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine. 2011;19(1):13. doi: 10.1186/1757-7241-19-13.

## TABLES AND FIGURES

Table 1: Schools selected for the study

School	Type
Yasodara Devi Balika M.V	1AB
Thakshila M.V	1AB
Parakrama M.V	1C
Sri Siddhartha Kumara M.V	1C
Keppetipola M.V	1C

Table 2: Socio-demographic and education-related information of the respondents

Characteristics	Number of students (N=510)	Percentage (%)
Gender		
Male	185	36.3
Female	325	63.7
Age (years)		
≤16	4	0.8
17	126	24.7
18	378	74.1
≥19	2	0.4
Ethnicity		
Sinhala	510	100
Religion		
Buddhism	508	99.6
Catholic	2	0.4
A/L stream		
Biology stream	144	28.2
Commerce stream	122	23.9
Mathematics stream	144	28.2
Art stream	100	19.6
Health science subject for O/L		
Have followed	254	49.8
Not followed	256	50.2
First aid training		
Trained	86	16.9

Table 3: Students first aid knowledge according to the grading

Knowledge grading	Number of students (N=510)	Percentage (%)
Inadequate ( $\leq 24\%$ )	5	1.0
Moderate (25-49%)	156	30.6
Good (50-74%)	313	61.4
Excellent ( $\geq 75\%$ )	36	7.1

Table 4 Knowledge on First aid

Variable	Correct answers		Wrong answers	
	N=510	Percentage (%)	N=510	Percentage (%)
Define first aid	486	<b>95.3</b>	24	4.7
First step in a first aid situation	290	<b>56.9</b>	220	43.1
Bleeding cut on the palm	321	<b>62.9</b>	189	37.1
Nose bleeding	173	33.9	337	<b>66.1</b>
Glass sticking in the leg	415	<b>81.4</b>	95	18.6
First aid for a sprain or strain	131	25.7	379	<b>74.3</b>
Fracture in leg	241	47.3	269	52.7
First aid for choking	264	51.8	246	48.2
First aid for burning	292	57.3	218	42.7
First aid for drinking poison	141	27.6	369	<b>72.4</b>
First aid for seizures	210	41.2	300	58.8
First aid for electrical shock	455	<b>89.2</b>	55	10.8
First aid for snake bite	450	<b>88.2</b>	60	11.8
Managing heart attack situation	238	46.7	272	53.3
Checking for breathing on victim	447	<b>87.6</b>	63	12.4



What should commence if not breathing	272	53.3	238	46.7
Correct CPR ratio	88	17.3	422	<b>82.7</b>
Importance of CPR	263	51.6	247	48.4
Call number of free ambulance service	388	<b>76.1</b>	122	23.9

Table 5 Attitudes towards first aid

Statement	Extremely disagree (%)	Disagree (%)	Agree (%)	Extremely agree (%)
Do you think is it essential to have first aid knowledge	-	1.2	26.3	<b>72.5</b>
Do you think is it essential to have a first aid training to provide first aid assistance at an emergency?	0.8	4.4	<b>50.0</b>	44.1
Do you think is it essential to have first aid box to give first aid for some one?	10.0	<b>53.1</b>	29.8	6.3
Do you think you will help someone who is having a heart attack?	4.3	30.6	<b>39.0</b>	25.3
Do you think you will help someone who has drowned in a river?	0.8	15.9	<b>55.3</b>	27.3
Do you think you will help someone with a severe bleeding wound?	2.4	13.7	<b>46.3</b>	36.9
Do you think you will provide first aid assistance to a victim of an accident on the road?	1.6	17.5	<b>49.8</b>	28.8
Do you think you have a phobia to see others blood?	<b>45.9</b>	32.2	19.2	2.0

Table 6: Cross tabulation of first aid knowledge with selected factors (n=510)

Variable	IA/M n (%)	G/E n (%)	Statistical association	
			$\chi^2$ value	p value
Gender				
Male	52 (28.1)	133 (71.9)	3.8	0.284
Female	109 (33.5)	216 (66.5)		
A/L stream				
Non Biology subjects	134 (36.6)	232 (63.4)	15.26	<0.0001
Biology	27 (18.8)	117 (81.3)		
O/L health science				
Followed	71 (28.0)	183 (72.0)	12.4	0.006
Not followed	90 (35.2)	166 (64.8)		
First aid training				
Trained	36 (41.9)	50 (58.1)	7.2	0.064
Not trained	125 (29.5)	299 (70.5)		
(IA/M- Inadequate and moderate)(G/E- good and excellent)				




Figure 1: Distribution of the first aid knowledge scores among students.

# Figures




Figure 1  
Distribution of the first aid knowledge scores among students.