

# Amaryllidaceae alkaloids with anti-Trypanosoma cruzi activity

**Nieves Martinez-Peinado**

Instituto de Salud Global Barcelona

**Nuria Cortes-Serra**

Instituto de Salud Global Barcelona

**Laura Torras-Claveria**

Universitat de Barcelona

**Maria-Jesus Pinazo**

Instituto de Salud Global Barcelona

**Joaquim Gascon**

Instituto de Salud Global Barcelona

**Jaume Bastida**

Universitat de Barcelona

**Julio Alonso-Padilla** (✉ [julio.a.padilla@isglobal.org](mailto:julio.a.padilla@isglobal.org))

Instituto de Salud Global Barcelona <https://orcid.org/0000-0003-4466-7969>

---

## Research

**Keywords:** Chagas disease; Trypanosoma cruzi; alkaloids; Amaryllidaceae; hippeastrine; phenotypic assays; cytotoxicity.

**Posted Date:** January 22nd, 2020

**DOI:** <https://doi.org/10.21203/rs.2.21521/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Parasites & Vectors on June 10th, 2020. See the published version at <https://doi.org/10.1186/s13071-020-04171-6>.

# Abstract

**Background:** Chagas disease, caused by the protozoan *Trypanosoma cruzi*, is a neglected disease that affects ~7 million people worldwide. Development of new drugs to treat the infection remains a priority since those currently available have frequent side effects and limited efficacy at the chronic stage. Natural products provide a pool of diversity structures to lead the chemical synthesis of novel molecules for this purpose. Herein we analyzed the anti- *T. cruzi* activity of 9 alkaloids derived from plants of the Amaryllidaceae family. **Methods:** the activity of each alkaloid was assessed by means of a newly developed anti- *T. cruzi* phenotypic assay. We further evaluated the compounds that inhibited the parasite growth on two distinct cytotoxicity assays to discard those that were toxic to host cells and assure parasite selectivity. **Results:** we identified a single compound (hippeastrine 2 ) that was selectively active against the parasite yielding selectivity indexes of 12.7 and 35.2 against Vero and HepG2 cells, respectively. **Conclusions:** results reported here suggest that natural products are an interesting source of new compounds for the development of drugs against Chagas disease.

## 1. Background

Chagas disease (or American trypanosomiasis) is a neglected infectious disease caused by the protozoan parasite *Trypanosoma cruzi* (*T. cruzi*; order Kinetoplastida; family Trypanosomatidae). It is estimated that ~7 million people are affected by the disease, mainly in Latin America where *T. cruzi* infection is endemic and occur the majority of the ~20.000 deaths per year related to it (1).

The disease progresses in two phases. There is first a short acute phase that is usually asymptomatic and thus goes unnoticed. This is followed by a chronic phase characterized by absent or slow progression of clinical manifestations (2). Nonetheless, it is estimated that ~40% of those chronically infected will ultimately develop disruptive damage to the heart and/or digestive tract (esophagus and colon) tissues, which can lead to the formation of mega-syndromes and death if untreated (2,3).

Since the 1970s only two drugs are available to treat *T. cruzi* infections: benznidazole (BNZ) and nifurtimox (NFX) (1). Both have good efficacy and tolerability when administered to infected newborns (4). But their efficacy diminishes at the chronic stage, which is usually diagnosed at adulthood with serological tests that detect specific anti-*T. cruzi* type G immunoglobulins (1). Moreover, both drugs have long regimens of administration that entail the advent of frequent adverse events which often drive to treatment discontinuation (5–7). There is thus an urgent unmet need of safer and more efficacious drugs for the treatment of chronic Chagas disease, for which natural products may represent a promising approach to discover new lead compounds (8–10).

In this regards, members of the family Amaryllidaceae have attracted considerable attention in the last few years due to their unique alkaloid composition with multiple biological activities (11). Amaryllidaceae plants have been studied for their potential application as a source of anticancer, anti-inflammatory, antimicrobial, anti-parasitic and anticholinesterase activities (12). In fact they have been used for

centuries as part of traditional treatments for fever, swelling, cancer or malaria (12). Remarkably, in 2001 the Food and Drug Administration (FDA) approved the use of galanthamine (trade name Razadyne), an alkaloid identified from the Amaryllidaceae plant *Galanthus woronowi*, to treat Alzheimer's disease (13).

Alkaloid constituents found in these plants are classified in eight groups based on structure and biogenesis from the common precursor *O*-methylnorbelladine: galanthamine, lycorine, crinine, haemanthamine, homolycorine, narciclasine, tazettine and montanine (14). The unique structure of this set of alkaloids provides a viable platform for phytochemical-based drug discovery (8). With the aim to identify prospective drug development starting points that could eventually become new therapeutic solutions for Chagas disease we have developed an in vitro anti-*T. cruzi* phenotypic assay based on the parasite's Tulahuen strain engineered to express a bacterial  $\beta$ -galactosidase gene (15) and green monkey epithelial cells (Vero) as hosts. We evaluated the anti-*T. cruzi* activity of nine crystalized alkaloid compounds extracted from members of the Amaryllidaceae family: lycorine **1**, hippeastrine **2**, crinine **3**, haemanthamine **4**, narciclasine **5**, tazettine **6**, montanine **7**, sanguinine **8** and 1-*O*-acetylcaranine **9** (Figure 1) (16). In all the assays performed we always included the standard anti-parasitic drug BNZ for comparison.

In order to unveil the specific *T. cruzi* growth inhibitory capacity of those compounds that were found active in the anti-parasitic assay we further developed two secondary biological assays to determine the compounds' level of cytotoxicity. These were respectively based on the same host Vero cells and in the human hepatocellular carcinoma cell line HepG2. Results obtained were particularly promising for the compound hippeastrine **2** from *Narcissus* cv. Salome (17) and they are discussed herein.

## 2. Methods

### 2.1 Collection of purified alkaloid compounds from Amaryllidaceae plants.

Lycorine **1**, hippeastrine **2**, crinine **3**, haemanthamine **4**, narciclasine **5**, tazettine **6**, montanine **7**, sanguinine **8** and 1-*O*-acetylcaranine **9** alkaloids were isolated from extracts of different *Narcissus* species (16). In brief, alkaloids extraction process was as follows: plant material (60 mg) was macerated with MeOH, the mix was filtered and the solvent evaporated to dryness. After that, extracts were acidified with 500  $\mu$ L of H<sub>2</sub>SO<sub>4</sub> (2%, v/v). The neutral material was removed with Et<sub>2</sub>O and basified with 200  $\mu$ L NH<sub>4</sub>OH (25 %, v/v). Then, 750  $\mu$ L of Et<sub>2</sub>O were added to separate the organic phase two times and the solvent evaporated to dryness. Finally, extracts were subjected to a combination of chromatographic techniques and alkaloids identified by GC-MS and NMR (16, 17).

### 2.2 Host cells cultures.

Vero (green monkey kidney epithelial cells), LLC-MK2 (Rhesus monkey kidney epithelial cells) and HepG2 (human liver epithelial cells) cultures were kept with DMEM supplemented with 1% penicillin-streptomycin (P-S) and 10% heat inactivated fetal bovine serum (FBS) at 37°C, 5% CO<sub>2</sub> and >95% humidity as described (15). HepG2 were also supplemented with 10% non-essential aminoacids.

## 2.3 Culture of *cruzi* parasites.

*T. cruzi* parasites from the Tulahuen strain (Discrete Typing Unit VI) expressing  $\beta$ -galactosidase were kindly provided by Dr. Fred Buckner (University of Washington, Seattle, USA) and maintained using LLC-MK2 cells as hosts in DMEM supplemented with 2% FBS and 1% P-S as described (15). Free-swimming trypomastigotes were purified by spinning the supernatants for 7 min at 2,500 rpm with a low break speed and then letting them swim out of the pellet (18). Purified trypomastigotes were used to keep the parasite cycle in LLC-MK2 cells and for the performance of the anti-parasitic assays. In the last case, an extra round of spinning was done to clean out the phenol red from the maintenance DMEM and replace it by phenol red-free assay medium, which was supplemented with 1% P-S-glutamine, 2% FBS, 1 mM sodium-pyruvate and 25 mM HEPES (18).

## 2.4 Assay to detect *cruzi* growth inhibition in 96 well plates.

Our assay is based on Vero cells as hosts and infective trypomastigotes from Tulahuen strain that express the bacterial  $\beta$ -galactosidase enzyme as reporter activity (15). Firstly, alkaloids were added in the first column of a 96-well tissue culture treated plates at an initial concentration of 100  $\mu$ M and diluted in assay medium into the next columns of the plate to conform dose-response plate-maps following either a 1:2 or 1:3 fold pattern. Then, Vero cells were detached from their growing flasks, counted and diluted at a concentration of  $1 \times 10^6$  cells per mL. Trypan blue staining was used to check out their viability, which had to be >95% to proceed. On the other side, purified trypomastigotes were counted as well and diluted at a concentration of  $1 \times 10^6$  cells per mL. We directly mixed Vero cells and trypomastigotes in a falcon tube in sufficient volume so as to add 100  $\mu$ L of the mix per well (50,000 Vero cells and 50,000 trypomastigote cells per well; multiplicity of infection or MOI = 1). DMSO percentage in all wells was always kept below 0.5%.

The reference drug BNZ was used as control of drug growth inhibition in each run launched, whereas each plate contained its own negative (maximum parasite growth; Vero cells plus parasites without drugs) and positive (minimum parasite growth; trypomastigote forms alone marking an enzymatic zero time or baseline galactosidase activity). Note that trypomastigotes are unable to multiply in the absence of susceptible host cells. Plates were incubated for 4 days at 37 °C (18), and the readout was done adding 50  $\mu$ L per well of a PBS solution containing 0.25% NP40 and 500  $\mu$ M chlorophenol red-b-D-galactoside (CPRG) substrate as previously described (18). Upon addition of the substrate, plates were further incubated at 37 °C for another 4 h and the absorbance read out at 590 nm using an Epoch Gene5 spectrophotometer. All experiments were performed at least in triplicate.

## 2.5 Toxicity assays with Vero and HepG2 cells.

For the cell toxicity assays, compounds were added to tissue culture treated 96-well plates following a dose-response dilution pattern 1:2 or 1:3 with a starting concentration of 400 or 800  $\mu$ M per well in the first column of the plate. Cells viability was checked upon cell counting with Trypan blue staining and we only proceeded if it was >95%. Vero cells solution was diluted at a concentration of  $5 \times 10^5$  cells per mL

before adding 100  $\mu$ L per well. In the case of HepG2 cells, we used a dilution of  $3.2 \times 10^4$  cells per mL. Each test plate or run contained its own negative (untreated cells) and positive (medium alone) controls. Plates were incubated at 37° C for 4 days in the case of Vero cells and 2 days for HepG2 cells. We then added 50  $\mu$ L per well of a PBS solution containing 10% Alamar Blue reagent (ThermoFischer) and incubated the plates for 6 h at 37 °C before reading the fluorescence intensity in a Tecan Infinite M Nano<sup>+</sup> reader (excitation: 530 nm, emission: 590 nm). DMSO percentage in all wells was always kept below 0.5%. All experiments were performed at least in triplicate.

## 2.6 Data analysis.

Absorbance and fluorescence values respectively derived from the anti-*T. cruzi* and cell toxicity assays were normalized to the controls (19). IC50 and TC50 values were determined with GraphPad Prism 7 software (version 7.00, 2016) using a non-linear regression analysis model. These IC50 and TC50 values are the compounds concentration capable to inhibit by 50% the growth of parasites and cells, respectively.  $Z'$  values were calculated as described (20). Values provided are means  $\pm$  standard deviation (SD) of at least three independent experiments.

# 3. Results And Discussion

## 3.1 Quality assessment of the assays developed.

As part of the process of setting up the biological assays we calculated their  $Z'$  parameter to assess reproducibility and statistical robustness (20). In general, assays with a  $Z'$  between 0.5 and 1 are considered appropriate for the screening of compounds (20). Remarkably, our anti-parasitic assay had a very good performance and its  $Z'$  remained consistently  $>0.5$  with an average value of 0.89 (CI95% = 0.73 - 1.00) (**Figure 2A**). Regarding the two cytotoxicity assays developed for this work, we respectively retrieved  $Z' = 0.73$  (CI95% = 0.60 - 0.93) for the assay based on Vero cells, and a  $Z' = 0.71$  (CI95% = 0.65 - 0.82) for the assay based on HepG2 cells (**Figure 2B-C**).

Additionally, in every run of the *T. cruzi* growth inhibition assay and Vero cells toxicity assays performed, we included the reference drug BNZ as control, whereas the reference drug digitoxin (21) was included in all the HepG2 cells toxicity assays. Overall, averaged IC50 and TC50 values for BNZ were respectively 1.56  $\mu$ M (SD  $\pm$  0.39) and 173.4  $\mu$ M (SD  $\pm$  43.57) (**Figure 2A-B**), which correlate with previous reports (18, 22). Digitoxin TC50 mean value in the HepG2 cell assay was 0.29  $\mu$ M (SD  $\pm$  0.14).

## 3.2 Anti- *cruzi* activity of the alkaloids extracted from Amaryllidaceae.

Same as other widely used anti-*T. cruzi* assays (18, 19, 23) we relied on the genetically robust *T. cruzi*-Tulahuen strain expressing beta-galactosidase activity as a surrogate of parasite growth (15). However, being the amastigote replicative stage of *T. cruzi* obliged intracellular, our assay relies on Vero cells as hosts. Notably, Vero cells were described to be deficient in the interferon response due to a mutation in

the simian interferon beta gene (24), which makes them more susceptible to infections and thereby a good system for the discovery of active compounds against *T. cruzi*.

Results obtained from the phenotypic *T. cruzi* growth inhibition assay revealed that lycorine **1**, hippeastrine **2**, haemanthamine **4**, narciclasine **5** and montanine **7** were active, while crinine **3**, tazettine **6**, sanguinine **8** and 1-*O*-acetylcaranine **9** were inactive against the parasite (**Figure 3**). Tazettine **6**, sanguinine **8** and 1-*O*-acetylcaranine **9** have been described as poor antiprotozoal agents before (25, 26). Nonetheless, Machocho and co-workers described that compound 3-*O*-acetylsanguinine had some activity (IC<sub>50</sub> = 2.3 µg/mL; i.e. 7.29 µM) against trypomastigotes from *T. cruzi* strain Tulahuen C4 (25). Although different phenotypic anti-*T. cruzi* assays were performed in each case, the results obtained by those authors could suggest that the presence of an acetyl group might increase the anti-*T. cruzi* activity of sanguinine.

On the other hand, crinine **3** and haemanthamine **4** are crinane-type alkaloids that belong to the β-crinane and α-crinane subgroups, respectively (27). Other studies evaluating the anti-parasitic potential of alkaloids have reported that the presence of a methylene-dioxi group seemed to favour a more potent anti-parasitic activity (28). This could be the explanation for the different anti-*T. cruzi* activities observed by us between these two crinane-type alkaloids.

The compounds that showed the highest anti-*T. cruzi* activities were yielded by lycorine **1** (IC<sub>50</sub> = 0.70 µM; CI<sub>95%</sub> 0.65 - 0.73) and narciclasine **5** (IC<sub>50</sub> = 0.495; CI<sub>95%</sub> 0.46 - 0.54), which exceeded in potency that of the reference drug BNZ (**Table 1**). Lycorine **1** was the first alkaloid described from the Amaryllidaceae plant family and one of the most commonly found amongst different genus (29). Lycorine **1** extracted from *Crinum stuhlmannii*, *Zephyranthes citrina* and *Narcissus broussonetii* had been tested before against *T. cruzi* (30–32). However, in contrast to our results, it was described as inactive against the parasite; a feature that could be explained due to the use of different assays. It has been reported that major differences on compounds activity can be found depending on the host cell lines and the biological assays used (33). In this respect previous references relied on the use of L6 cells as hosts, whereas we have used Vero cells.

With respect to narciclasine **5**, which yielded the most potent activity against *T. cruzi* in this study (**Table 1**), to the best of our knowledge it has not been tested against *T. cruzi* before. Likewise, there were no previous reports on the anti-*T. cruzi* activity of montanine **7**, also shown for the first time in this study. Montanine **7** and haemanthamine **4** showed averaged IC<sub>50</sub> values similar to that of BNZ, respectively 1.99 µM (CI<sub>95%</sub> 1.82 - 2.1) and 1.59 µM (CI<sub>95%</sub> 1.47 - 1.42) versus the 1.56 µM of BNZ (CI<sub>95%</sub> 0.93-2.20; see **Figure 2B**). Osorio and co-workers had previously described the high activity of haemanthamine against *T. cruzi* (IC<sub>50</sub> = 1.8 µg/mL; i.e. 5.97µM) (26).

Finally, we found that hippeastrine **2** was the active alkaloid with the higher IC<sub>50</sub> value (IC<sub>50</sub> = 3.63 µM, CI<sub>95%</sub> 2.95 - 3.82). Its activity was poorer than that of BNZ but still fell within a range five times the IC<sub>50</sub> of the standard drug (**Table 1**). Hippeastrine **2** was first isolated from the Amaryllidaceae plant *Lycoris*

*radiata* and reported to exhibit activity against avian influenza virus H5N1 ( $IC_{50} = 47.5 \pm 0.37 \mu M$ ) (34). Moreover, promising results against ZIKA virus infection have been recently reported with hippeastrine hydrobromide (35). This was shown to remove ZIKA virus presence from infected human neural progenitors, recover a ZIKV-induced microcephaly phenotype in human forebrain organoids and even suppress virus propagation in infected adult mice (35). Antiviral (34, 35), antibacterial and antifungal (36) activities have been reported for hippeastrine **2**, despite little information is available about its anti-parasitic activity. Cedron et al. tested twenty one hippeastrine **2** derivatives that included functional group transformations, structural simplification and dimers formation against *Plasmodium falciparum* (strain F-32 Tanzania) (37). The anti-malarial activity increased by 10-fold when dimers were evaluated compared to the single alkaloid activity, suggesting an improved binding to the related target or the hydrolysis of the dimer onto two molecules (37). To our knowledge this is the first time that anti-*T. cruzi* activity is reported for hippeastrine **2**. Results reported by Cedron and co-workers suggest to further pursue research with hippeastrine **2** derivatives against *T. cruzi*.

### 3.3 Identification of alkaloid compounds with specific anti- *cruzi* activity.

With the aim of further selecting those alkaloids with specific activity against the parasite and discard those that were toxic to host cells we developed two secondary cell toxicity assays with monkey (Vero) and human (HepG2) cells. Since the compounds activity might vary depending on the characteristics of the cell line used, assaying cytotoxicity in two cell lines provided a more robust readout. Moreover, HepG2 cells are a widespread cell model to anticipate potential liver toxicity of drugs metabolism (35, 36). We determined a selectivity index (SI; or  $TC_{50}$  to  $IC_{50}$  ratio)  $>10$  to consider an alkaloid for further progression as described elsewhere (19).

Thereafter, all the alkaloids reported as active against *T. cruzi* were analysed through both cell toxicity assays. All of them were more toxic to Vero cells than to HepG2 cells (**Figure 4**). The cytotoxicity values registered against Vero and HepG2 cells indicated that narciclasine **5** activity was not specific against *T. cruzi* ( $TC_{50} = 0.66 \mu M$  against Vero cells; and  $TC_{50} = 2.73 \mu M$  against HepG2 cells), resulting in  $SI < 10$  in both cases (**Table 1**). In addition, montanine **7** with a  $SI = 2.5$  in relation to Vero cells was not specific to *T. cruzi* either (**Table 1**), even though it showed low toxicity to HepG2 cells ( $TC_{50} = 46.1 \mu M$ ; **Table 1**). Something similar occurred with lycorine **1**, which was weakly active against HepG2 cells ( $TC_{50} = 21.87 \mu M$ ), presenting a  $SI$  versus this cell line over ten times its registered anti-*T. cruzi* activity, but turned to be toxic to Vero cells ( $TC_{50} = 5.21 \mu M$ ) yielding a  $SI < 10$  and thus being discarded it from further progression.

Haemanthamine **4** had been reported to present a  $TC_{50} = 13 \mu g/ml$ ; i.e.  $43.14 \mu M$  for HepG2 cells (40) which correlate with the  $TC_{50}$  value we have retrieved in our study with this cell line ( $TC_{50} = 42.48 \mu M$ ; **Table 1**). But again, although it showed a  $SI > 10$  with respect to HepG2 cells, when evaluated on Vero cell its  $TC_{50}$  to  $IC_{50}$  ratio was below that threshold and thus its anti-parasitic activity could not be considered specific (**Table 1**).

In contrast to all the aforementioned results, hippeastrine **2** did show low toxicities against Vero (TC50 = 45.99  $\mu$ M) and HepG2 cells (TC50 = 128.1  $\mu$ M), and it indeed complied with the SI window > 10 against both cell lines (**Table 1**). It was the only compound that had a SI > 10 versus Vero cells (SI = 12.7; **Table 1**). In addition, hippeastrine **2** presented the highest SI against HepG2 cells (SI = 35.3), with a TC50 value similar to that previously reported by Weniger et al. (TC50 = 40  $\mu$ g/ml; i.e. 126.85  $\mu$ M) (40). Those cytotoxic results together with the in vivo assays reported by Zhou et al. (35) predict a low toxicity in future evaluation of hippeastrine **2** anti-*T. cruzi* activity in animal models, and thus lead us to be optimistic about the future progression of the compound.

**Table 1.** Table displaying the alkaloids' average IC50,TC50 and SI values for Vero and HepG2 cells.

No.	Alkaloid	IC50 ( $\mu$ M)	TC50 <sup>1</sup> ( $\mu$ M)	SI <sup>1</sup>	TC50 <sup>2</sup> ( $\mu$ M)	SI <sup>2</sup>
-	BNZ	1.56	173.4	111.15	168.76	108.18
<b>1</b>	Lycorine	0.70	5.21	7.5	21.87	31.2
<b>2</b>	Hippeastrine <sup>#</sup>	3.63	45.99	<b>12.7</b>	128.10	<b>35.2</b>
<b>3</b>	Crinine	57.93	-	-	-	-
<b>4</b>	Haemanthamine	1.59	11.52	7.3	42.48	26.7
<b>5</b>	Narciclasine	0.495	0.66	1.3	2.73	5.5
<b>6</b>	Tazettine	83.03	-	-	-	-
<b>7</b>	Montanine	1.99	5.04	2.5	46.10	23.1
<b>8</b>	Sanguinine	213.4	-	-	-	-
<b>9</b>	1- <i>O</i> -acetylcaranine	35.49	-	-	-	-

Superscripts <sup>1</sup> and <sup>2</sup> respectively refer to Vero and HepG2 cell toxicity assays TC50s and SIs. <sup>#</sup>Indicate the only alkaloid evaluated in this study that showed specific anti-*T. cruzi* activity. The standard drug BNZ is included in the first line as comparator.

## 4. Conclusions

Upon the evaluation of nine alkaloid compounds purified from extracts of different *Narcissus* species (family Amaryllidaceae) (16) we identified one compound that had specific anti-*T. cruzi* activity, proving that natural products are an interesting source to potentially identify new departing chemical structures for Chagas disease drug discovery. Our findings suggest that hippeastrine **2** (17) could be a relevant compound to progress into in vivo studies.

## 5. Declarations

**Ethics approval and consent to participate:** "Not applicable".

**Consent for publication:** "Not applicable".



**Availability of data and materials:** “Data and materials can be made available upon reasonable request to the authors”.

**Competing interests:** “The authors declare no competing interests”.

**Funding:** “We want to thank the support by the Departament d’Universitats i Recerca de la Generalitat de Catalunya, Spain (AGAUR; 2017SGR00924), and the funding by the Instituto de Salud Carlos III RICET Network for Cooperative Research in Tropical Diseases (ISCIII; RD12/0018/0010) and FEDER. JAP was funded by a Juan de la Cierva – Incorporación contract from the Spanish Science Ministry. LTC and JB (UB research group 2017SGR604) thank CYTED (416RT0511) for financial support. MJP research is supported by the Ministry of Health, Government of Catalunya (PERIS 2016-2010 SLT008/18/00132).

**Authors' contributions:** NMP, NCS, JB and JAP conceptualized the study; NMP performed all biological assays; NCS helped with the assays; LTC and JB purified the alkaloids and provided the collection; MJP and JG provided funds; NMP and JAP wrote the article; all authors read and approved the final manuscript, and declare that there is no conflict of interests.

**Acknowledgements:** “Not applicable”.

## References

1. WHO: Chagas disease (American trypanosomiasis). URL: [https://www.who.int/en/news-room/fact-sheets/detail/chagas-disease-\(american-trypanosomiasis\)](https://www.who.int/en/news-room/fact-sheets/detail/chagas-disease-(american-trypanosomiasis)). Last accessed 8<sup>th</sup> Jan 2020.
2. Pinazo MJ, Gascon J. Chagas disease: from Latin America to the world. *Reports Parasitol.* 2015;4:7–14.
3. Prata A. Clinical and epidemiological aspects of Chagas disease. *Lancet Infect Dis.* 2001;1:92–100.
4. Alonso-Padilla J, Gállego M, Schijman AG, Gascon J. Molecular diagnostics for Chagas disease: up to date and novel methodologies. *Expert Rev Mol Diagn.* 2017;17:699-710.
5. Crespillo-Andujar C, Venanzi-Rullo E, López-Vélez R, Monge-Maillo B, Norman F, López-Polín A, et al. Safety profile of benznidazole in the treatment of chronic Chagas disease: experience of a referral centre and systematic literature review with meta-analysis. *Drug Saf.* 2018. 41:1035-48.
6. Forsyth CJ, Hernandez S, Olmedo W, Abuhamidah A, Traina MI, Sanchez DR, et al. Safety profile of nifurtimox for treatment of Chagas disease in the United States. 2016;63:1056-62.
7. Pinazo MJ, Muñoz J, Posada E, López-Chejade P, Gállego M, Ayala E, et al. Tolerance of benznidazole in treatment of Chagas’ disease in adults. *Antimicrob Agents Chemother.* 2010;54:4896–99.
8. Cimmino A, Masi M, Evidente M, Superchi S, Evidente A. Amaryllidaceae alkaloids: absolute configuration and biological activity. *Chirality.* 2017;29:486-99.
9. Izumi E, Ueda-Nakamura T, Dias Filho BP, Veiga Júnior VF, Nakamura CV. Natural products and Chagas’ disease: a review of plant compounds studied for activity against *Trypanosoma cruzi*. *Nat*

- Prod Rep. 2011;28:809–23.
10. Zulfikar B, Jones AJ, Sykes ML, Shelper TB, Davis RA, Avery VM. Screening a natural product-based library against kinetoplastid parasites. *Molecules*. 2010; 22(10). pii: E1715.
  11. Nair JJ, Bastida J, Viladomat F, van Staden J. Cytotoxic agents of the crinane series of Amaryllidaceae alkaloids. *Nat Prod Commun*. 2012;7:1677-88.
  12. Presley CC, Krai P, Dalal S, Su Q, Cassera M, Goetz M, Kingston DGI. New potentially bioactive alkaloids from *Crinum erubescens*. *Bioorg Med Chem*. 2016;24: 5418–22.
  13. Sramek JJ, Frackiewicz EJ, Cutler NR. Review of the acetylcholinesterase inhibitor galanthamine. *Expert Opin Investig Drugs*. 2000;9:2393–402.
  14. Tallini LR, de Andrade JP, Kaiser M, Viladomat F, Nair JJ, Zuanazzi JAS, et al. Alkaloid constituents of the Amaryllidaceae plant *Amaryllis belladonna* L. *Molecules*. 2009;22(9). pii: E1437.
  15. Buckner FS, Verlinde CL, La Flamme AC, Van Voorhis WC. Efficient technique for screening drugs for activity against *Trypanosoma cruzi* using parasites expressing  $\beta$ -Galactosidase. *Antimicrob Agents Chemother*. 1996;40:2592-7.
  16. Bastida J, Lavilla R, Viladomat F. Chemical and biological aspects of *Narcissus* alkaloids. *Alkaloids Chem Biol*. 2006;63:87-179.
  17. Almanza GR, Fernández JM, Wakori EWT, Viladomat F, Codina, C. Alkaloids from *narcissus* cv. salome\*. *Phytochemistry*. 1996;43:1375–8.
  18. Bettiol E, Samanovic M, Murkin AS, Raper J, Buckner F, Rodriguez A. Identification of three classes of heteroaromatic compounds with activity against intracellular *Trypanosoma cruzi* by chemical library screening. *PLoS Negl Trop Dis*. 2009;3(2):e384.
  19. Peña I, Pilar Manzano M, Cantizani J, Kessler A, Alonso-Padilla J, Bardera AI, et al. New compound sets identified from high throughput phenotypic screening against three kinetoplastid parasites: an open resource. *Sci Rep*. 2015; 5:8771.
  20. Zhang JH, Chung TD, Oldenburg KR. Z' - a simple statistical parameter for use in evaluation and validation of high throughput screening. *J Biomol Screen*. 1999;4:67-73.
  21. Xiao Y, Yan W, Guo L, Meng C, Li B, Neves H, et al. Digitoxin synergizes with sorafenib to inhibit hepatocellular carcinoma cell growth without inhibiting cell migration. *Mol Med Rep*. 2017;15:941-7.
  22. Alonso-Padilla J, Cotillo I, Presa JL, Cantizani J, Peña I, Bardera AI, et al. Automated high-content assay for compounds selectively toxic to *Trypanosoma cruzi* in a myoblastic cell line. *PLoS Negl Trop Dis*. 2015;9:e0003493.
  23. Khare S, Nagle AS, Biggart A, Lai YH, Liang F, Davis LC, et al. Proteasome inhibition for treatment of leishmaniasis, Chagas disease and sleeping sickness. *Nature*. 2016;537:229–33.
  24. Mosca JD, Pitha PM. Transcriptional and posttranscriptional regulation of exogenous human beta interferon gene in simian cells defective in interferon synthesis. *Mol Cell Biol*. 2015;6:2279–83.
  25. Machocho AK, Bastida J, Codina C, Viladomat F, Brun R, Chabra SC. Augustamine type alkaloids from *Crinum kirkii*. *Phytochemistry*. 2004;65:3143-9.

26. Osorio EJ, Berkov S, Brun R, Codina C, Viladomat F, Cabezas F, et al., In vitro antiprotozoal activity of alkaloids from *Phaedranassa dubia* (Amaryllidaceae). *Phytochem Lett*. 2010;3:161–3.
27. Nair JJ, Wilhelm A, Bonnet SL, van Staden J. Antibacterial constituents of the plant family Amaryllidaceae. *Bioorg Med Chem Lett*. 2017;27: 4943–51.
28. Tallini LR, Osorio EH, Santos VDD, Borges WS, Kaiser M, Viladomat F et al. *Hippeastrum reticulatum* (Amaryllidaceae): alkaloid profiling, biological activities and molecular docking. *Molecules*. 2017;22(12). pii: E2191.
29. Nair JJ, van Staden J. Cytotoxicity studies of lycorine alkaloids of the Amaryllidaceae. *Nat Prod Commun*. 2014;9:1193-210.
30. Machocho A, Chhabra SC, Viladomat F, Codina C, Bastida J. Alkaloids from *Crinum stuhlmannii*. *Planta Med*. 1998;64:679-80.
31. Herrera MR, Machocho AK, Brun R, Viladomat F, Codina C, Bastida J. Crinane and lycorane type alkaloids from *Zephyranthes citrina*. *Planta Med*. 2001;67:191-3.
32. de Andrade JP, Pigni NB, Torras-Claveria L, Berkov S, Codina C, Viladomat F, et al. Bioactive alkaloid extracts from *Narcissus broussonetii*: mass spectral studies. *J Pharm Biomed Anal*. 2012;70:13-25.
33. Franco CH, Alcântara LM, Chatelain E, Freitas-Junior L, Moraes CB. Drug discovery for Chagas disease: impact of different host cell lines on assay performance and hit compound selection. *Trop Med Infect Dis*. 2019;4(2). pii: E82.
34. He J, Qi W, Wang L, Tian J, Jiao P, Liu G. Amaryllidaceae alkaloids inhibit nuclear-to-cytoplasmic export of ribonucleoprotein (RNP) complex of highly pathogenic avian influenza virus H5N1. *Influenza Other Respir Viruses*. 2013;7:922–31.
35. Zhou T, Tan L, Cederquist GY, Fan Y, Hartley BJ, Mukherjee S, et al. High-content screening in hPSC-neural progenitors identifies drug candidates that inhibit Zika virus infection in fetal-like organoids and adult brain. *Cell Stem Cell*. 2017;21:274-83.e5
36. Evidente A, Andolfi A, Abou-donia AH, Touema SM, Hammouda HM, Shawky E, et al. *Amaryllis belladonna* L . growing in Egypt. *Phytochemistry*. 2004;65:2113–8.
37. Cedrón JC, Gutiérrez D, Flores N, Ravelo ÁG, Estévez-Braun A. Preparation and antimalarial activity of semisynthetic lycorenine derivatives. *Eur J Med Chem*. 2013;63:722–30.
38. Choi JM, Oh SJ, Lee SY, Im JH, Oh JM, Ryu CS, et al. HepG2 cells as an in vitro model for evaluation of cytochrome P450 induction by xenobiotics. *Arch Pharm Res*. 2015;38:691-704.
39. Crouch SP, Kozlowski R, Slater KJ, Fletcher J. The use of ATP bioluminescence as a measure of cell proliferation and cytotoxicity. *J Immunol Methods*. 1993;160:81-8.
40. Weniger B, Italiano L, Beck JP, Bastida J, Bergoñón S, Codina C, et al. Cytotoxic activity of Amaryllidaceae alkaloids. *Planta Med*. 1995;61:77-9.

## Additional File Legend

**Figure A.** BNZ and DTX dose-response curves. Both reference drugs were included in every assay as a control of drug inhibition. Anti-*T. cruzi* assays are represented by circles while Vero and HepG2 cells toxicity assays by squares and triangles, respectively.

## Figures




Figure 1

Chemical structures of the alkaloids evaluated in this work. Structures were drawn with ChemDraw software (PerkinElmer).






Figure 2

Quality controls of the *T. cruzi* growth inhibition assay (A-B), toxicity assay with Vero cells (C-D) and toxicity assay with HepG2 cells (E-F). Z' values for each of the rounds launched are represented on the left (A, C, E); dashed line marks the 0.5 threshold. IC50 and TC50 values of the reference drugs BNZ and digitoxin (DTX) are represented on the right (B, D, F); straight lines indicate the average values, whereas the dashed lines indicate  $\pm 3$  SD limits.



**Figure 3**

Anti-T.cruzi phenotypic assay dose-response curves. Graphs represent mean results and SD of at least three biological replicas.



**Figure 4**

Dose-response curves yielded by the Vero and HepG2 cell toxicity assays. Vero cells toxicity assays are represented by circles and straight lines while HepG2 cell toxicity assays are represented by triangles and dashed lines. Graphs represent mean results and SD of at least three biological replicas.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Abstractgrafic.jpg](#)
- [Additionalfile1.pdf](#)