

# Characteristics of persistent hotspots of *Schistosoma mansoni* in western Côte d'Ivoire

**Rufin Kouassi Assaré** (✉ [hrufinass@yahoo.fr](mailto:hrufinass@yahoo.fr))

Universite Félix Houphouet-Boigny

**Roméo N. N'Tamon**

Universite Félix Houphouet-Boigny

**Louise G. Bellai**

University of Basel

**Judicaelle A. Koffi**

Universite Félix Houphouet-Boigny

**Tra-Bi I. Mathieu**

Universite Félix Houphouet-Boigny

**Mamadou Ouattara**

Universite Félix Houphouet-Boigny

**Eveline Hürlimann**

Swiss Tropical and Public Health Institut

**Jean T. Coulibaly**

Universite Félix Houphouet-Boigny

**Salia Diabaté**

Centre d'Entomologie Médicale et Vétérinaire

**Eliézer K. N'Goran**

Universite Félix Houphouet-Boigny

**Jürg Utzinger**


Swiss Tropical and Public Health Institut

---

## Research

**Keywords:** *Biomphalaria pfeifferi*, Côte d'Ivoire, persistent hotspot, preventive chemotherapy, *Schistosoma mansoni*, schistosomiasis, water, sanitation and hygiene (WASH)

**DOI:** <https://doi.org/10.21203/rs.2.21775/v3>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## Abstract

**Background** Preventive chemotherapy with praziquantel is the cornerstone of schistosomiasis control. However, in some social-ecological settings, the prevalence and/or intensity of *Schistosoma* infection does not lower meaningfully despite multiple rounds of preventive chemotherapy, a phenomenon termed persistent hotspot (PHS). We assessed the characteristics of PHS in a *Schistosoma mansoni*-endemic area of Côte d'Ivoire. **Methods** In October 2016, a cross-sectional parasitological survey was conducted in 14 schools in the western part of Côte d'Ivoire, one year after multiple rounds of preventive chemotherapy. In each school, 50 children aged 9-12 years were randomly selected and invited to submit two stool samples and one urine sample. Stool samples were subjected to triplicate Kato-Katz thick smears for *S. mansoni* diagnosis. Urine samples were examined by a filtration method for *S. haematobium* eggs. PHS was defined as failure to achieve a reduction in the prevalence and/or intensity of *S. mansoni* infection of at least 35% and 50%, respectively after multiple rounds of preventive chemotherapy targeting school-aged children. A questionnaire was administered in six schools to obtain data on demographic characteristics, including human-water contact behaviours, water, sanitation and hygiene and history of preventive chemotherapy. A malacological survey was conducted in December 2016. **Results** The overall prevalence of *S. mansoni* and *S. haematobium* in the 14 surveyed schools was 9.5% and 2.6%, respectively. There were four PHS among six schools that underwent detailed investigation. The *S. mansoni* prevalence in the four PHS was 10.9% (95% confidence interval (CI): 6.4-15.5%), compared to 6.6% (95% CI: 1.4-11.8%) in the remaining two schools. The arithmetic mean of *S. mansoni* eggs per gram of stool (EPG) among infected children was 123.8 EPG in PHS and 18.7 EPG in the remaining two schools. A statistically significant difference between PHS and the other schools regarding praziquantel coverage was found with higher coverage rate observed in PHS (98.4% versus 89.0%,  $\chi^2 = 3.7$ ,  $p = 0.001$ ). There was no significant difference with regard to access to clean water in the two groups of schools. Children bathing in open freshwater bodies were at higher odds of *S. mansoni* infection (odds ratio (OR) = 4.5, 95% CI: 1.6-12.6). A total of 76 human-water contact sites (53 in PHS and 23 in the other schools) were examined and 688 snails were collected, including 92 *Biomphalaria pfeifferi*, 25 *Bulinus forskalii*, 16 *Bu. globosus* and one *Bu. truncatus*. **Conclusion** Children in PHS schools bathed more frequently in open freshwater bodies, and hence, they were potentially more exposed to *S. mansoni*. Our findings call for an integrated control approach, complementing preventive chemotherapy with behavioural change implementation in PHS settings.

## Background

Although schistosomiasis affects more than 250 million people, it is considered a neglected tropical disease that caused an estimated burden of 1.4 million disability-adjusted life years in 2017 [1, 2]. Preventive chemotherapy with praziquantel is the mainstay of the global schistosomiasis control strategy [3]. In 2016, 52.8 million school-aged children requiring preventive chemotherapy received praziquantel, a coverage of 54% [4].

In order to strengthen the current evidence-base for decisions about preventive chemotherapy for gaining and sustaining control of schistosomiasis, the Schistosomiasis Consortium for Operational Research and Evaluation (SCORE) implemented a series of large-scale, multi-year trials in sub-Saharan Africa [5, 6]. In Côte d'Ivoire, a SCORE study pertaining to sustaining control of schistosomiasis *mansoni* was carried out in 75 schools in the western part of the country. Schools were included when the prevalence of *Schistosoma mansoni* was between 10% and 24%, as determined by duplicate Kato-Katz thick smears among 50 children aged 13-14 years [7]. The selected schools were randomly allocated to one of three treatment arms, including 25 schools per arm. Children in arm A received annual praziquantel treatment for four years, children in arm B received praziquantel in the first two years, followed by two years of "drug holidays", while children in arm C received treatment in years 1 and 3, alternated by "drug holidays" in years 2 and 4. The dynamics of schistosomiasis was heterogeneous, as revealed by the initial treatment follow-up 1 year after the first drug administration. Indeed, while in 43 of the 50 schools surveyed at this time point, a decrease in *S. mansoni* prevalence was observed, in seven schools, the prevalence increased [8]. Similar results have been reported elsewhere in sub-Saharan Africa [9, 10, 11, 12].

Social-ecological settings where the prevalence and/or intensity of *S. mansoni* increases or does not change meaningfully after preventive chemotherapy are termed persistent hotspots (PHS). There are different definitions of PHS. The approach adopted in the current study has been described elsewhere [13]. In brief, a village that failed to reduce the *S. mansoni* prevalence by at least 35% and/or failed to reduce the intensity of infection by at least 50% between baseline and year five testing is considered a PHS, while a village that fulfils these criteria is termed low-prevalence setting. Of note, PHS of schistosomiasis have been investigated in different parts of the world [14, 15, 16, 17] and in the gaining and sustaining control studies of SCORE [13]. Prior research showed that various factors can contribute to the existence of PHS, including close proximity of human habitation to open surface water bodies where intermediate host snails proliferate, human behaviours, particularly in terms of water, sanitation and hygiene (WASH) parameters [18, 19, 20] and socioeconomic status of the study population [21, 22].

In the western part of Côte d'Ivoire where a multi-year sustaining schistosomiasis *mansoni* control study was conducted, 45% (34 of 75) of the investigated villages were PHS [13]. However, the factors that drive PHS in this area are unknown. Hence, new research is needed to investigate factors, including human behaviours and environmental characteristics that might explain persistence of pockets of schistosomiasis transmission in western Côte d'Ivoire [13, 15, 17]. The specific objectives of this study were to (i) characterise PHS of *S. mansoni* in a SCORE sustaining control study in western Côte d'Ivoire; (ii) determine environmental and demographic factors and WASH indicators that might govern PHS, (iii) assess the awareness of the study population on schistosomiasis and praziquantel treatment; and (iv) identify intermediate host snails and determine whether they shed schistosome cercariae.

## Methods

### Ethics statement

Approval for this study was obtained from the ethics committee of the Ministry of Public Health in Côte d'Ivoire (reference no. 046/MSHP/CNER-kp; date of assignment: 30 May 2016). The purpose and procedures of the study were explained to the education and health authorities. The local communities were informed on the goal, potential risks and benefits of the study in their native language. Parents/guardians signed a written informed consent form on behalf of

the children. Participation was voluntary; hence, children could withdraw at any time without further obligations. Urine and stool samples may be sources of bacterial, parasitic and viral infections; hence, children were asked to clean their hands with soap after producing and transferring stool and urine samples. At the end of the study, all school-aged children were treated with praziquantel at a single oral dose of 40 mg/kg [3].

### Study area and population

The study was carried out between October and December 2016 in Cavally, Guemon, Haut-Sassandra and Tonkpi, four regions of western Côte d'Ivoire (geographical coordinates: 06°32'42.0" to 07°36'54.8" N latitude; 06°44'09.8" to 07°33'48.9" W longitude). Preventive chemotherapy targeting school-aged children was conducted in the study areas in late 2015 [7]. Infection with *S. mansoni* is highly endemic in this part of Côte d'Ivoire [23, 24, 25]. A longitudinal malacological survey carried out monthly from 1986 to 1988 in two villages in the Tonkpi region revealed the presence of *Biomphalaria pfeifferi*, the intermediate host snail of *S. mansoni* [26].

In the frame of a large SCORE-funded project, a parasitological survey was conducted in 2011 and 2012, in 75 purposely selected communities to determine the baseline situation and village characteristics before the implementation of a 4-year randomised controlled trial designed to evaluate the impact of three different school-based treatment schedules with praziquantel to sustain the control of *S. mansoni* infection [7, 27]. There was an overall prevalence of *S. mansoni* of 5.4% among first-graders and 22.1% in children aged 9-12 years [27]. One year after the initial administration of praziquantel, the overall prevalence of *S. mansoni* in children aged 9-12 years in 50 schools decreased significantly from 19.7% (95% confidence interval (CI): 18.5–20.8%) at baseline to 12.8% (95% CI: 11.9–13.8%) at the 1-year follow-up with considerable spatial heterogeneity in the dynamic of *S. mansoni* infection. While the prevalence of *S. mansoni* decreased in most of the schools, in some schools the prevalence remained unchanged or even increased.

In October 2016, a sub-study was launched in 14 schools, consisting of a parasitological survey in all 14 schools, and a malacological and questionnaire survey in six of the schools. Limited financial resources, challenges in accessing some of the schools and other logistic reasons precluded surveying all 14 schools. Among the six schools, three had a *S. mansoni* prevalence above 24%, and hence, were considered high endemicity according to a pre-set SCORE threshold. These three villages are Ziondrou (32.7%), Zê (27.0%) and Tobly Bangolo (26.0%). The other three schools were classified as moderate endemicity (10-24% prevalence): Gbadrou (19.0%), Zoukougbeu (15.2%) and Mona (14.0%).

### Parasitological survey

In a first step, a cross-sectional parasitological survey was conducted in the 14 schools. In brief, in each school, 50 children aged 9-12 years were randomly selected and invited to participate. We selected 50 children per school, according to guidelines put forth by the World Health Organization (WHO) [28]. Children aged 9-12 years were chosen in accordance to SCORE protocols and the fact that children at this age are particularly active, frequently contact open freshwater bodies and are thus exposed to schistosomiasis [28, 29]. On the first day of the survey, children with written parental informed consent received two empty plastic containers. They were asked to fill one container with a small portion of fresh morning stool and to collect a mid-day urine sample in the second container. On the second day, children provided an additional stool sample but no further urine sample. All stool samples were subjected to triplicate 41.7 mg Kato-Katz thick smears [30]. After a clearing time of at least 30 min, the thick smears were examined under a microscope by one of four experienced laboratory technicians. Urine samples were examined using a filtration method [31]. Eggs of *S. mansoni* and *S. haematobium* were counted and recorded for each individual separately.

### Questionnaire survey

A questionnaire was administered in November 2016 to those children who participated in the parasitological survey in the six villages chosen for in-depth analyses. The interviewers were well acquainted with the study setting and received prior training and specific instructions to conduct the interviews. There were two questionnaires; the first pertaining to schistosomiasis risk factors and the second to recent history of preventive chemotherapy campaigns. Both questionnaires has been used before.

The schistosomiasis risk factor questionnaire investigated demographic data, house construction materials, main source of water for drinking and cooking, socioeconomic factors, latrine access and use, defecation behaviours, hygiene, contact to open surface water and other water-related activities. For the current study, we defined stagnant water as a freshwater that does not flow (e.g. pond, lake or backwater). The history of preventive chemotherapy questionnaire has been described elsewhere [32]. In brief, the main focus was on children's participation at the last school-based treatment campaign in late 2015 and knowledge on schistosomiasis and praziquantel treatment. In case children did not take the medication, they were asked about the underlying reasons.

### Malacological survey

In December 2016, a malacological survey was conducted for the identification of schistosomiasis intermediate host snails. In each of the six target villages, all human-water contact sites located within a buffer of 4 km from the village centre were identified after discussion with local authorities. These human-water contact sites were grouped into ponds, small dams, swamps, rivers, lakes, paddy fields and streams. Snail sampling was conducted by two experienced field workers. Snails were collected during 15 min at each site by hand using soft tweezer. In addition, a long-handed kitchen scoop (1.5 m pole; 0.8-1.2 mm mesh

size) was used to reach under deeper vegetation and to sample where access was difficult. Collected snails were identified at genus and, whenever possible, at species level [33]. Potential intermediate host snails of schistosomiasis (i.e. *Biomphalaria* and *Bulinus*) were placed in screw top plastic containers with freshwater from the site and transferred to a nearby laboratory. The remaining snails were put back into the water body. Human-water contact sites were georeferenced using a hand-held phone (Infinix, Hot 2, X510; Guangzhou, People's Republic of China). Potential schistosomiasis intermediate host snails were reared in the laboratory and subjected to cercarial shedding at days 1, 15 and 30 after collection. The snails were exposed to artificial light from 11:00 to 14:00 hours.

### Snail habitat data

Human-water contact sites were characterised (e.g. accessibility, presence of water during the year, estimated canopy coverage, vegetation and substratum). Water flow, level and depth were also registered. The presence of domestic animals (e.g. cows, horses, goats, dogs, sheep, pigs and donkeys) and wild animals (e.g. monkey and birds) was documented through direct observations, by recording footprints and information given by local residents. Water contact activities such as laundry, swimming, playing, water collection, rice growing, crossing river, fishing, open defecation, washing bicycle and cars were determined by direct observations and interviews with key stakeholders (e.g. heads of households and mothers of young children). Water samples were collected in 1.5 l plastic bottles and transferred to a laboratory in Abidjan for determining physico-chemical parameters: total dissolved solids (mg/l), conductivity ( $\mu\text{S}/\text{cm}$ ) and pH, recorded using a portable multimeter (Hanna Instruments; Woonsocket, United States of America).

### Statistical analysis

Data were entered into Microsoft Excel 2010 (Microsoft Corporation; Redmond, United States of America) and cross-checked with EpiInfo version 3.5.4 (Centers for Disease Control and Prevention; Atlanta, United States of America). Statistical analyses were done with STATA version 13.1 (Stata Corporation; College Station, United States of America). Age was stratified into two groups: 9-10 and 11-12 years. Intensities of *Schistosoma* infection were classified according to WHO guidelines [34]. In brief, intensity of *S. mansoni* infection was categorised into light (1-99 eggs per gram of stool (EPG)), moderate (100-399 EPG) and heavy ( $\geq 400$  EPG). Intensity of *S. haematobium* infection was grouped into light (1-49 eggs/10 ml of urine) and heavy ( $\geq 50$  eggs/10 ml of urine). All statistical analyses involving comparisons or associations across schools took into account the clustered data structure treating schools as primary sampling units. Stata's survey methodology was used for this purpose. Proportions of outcomes were compared according to sex, age group and whether or not a school was considered PHS. Statistical significance was defined at the 5% level. Logistic regression models were used to examine the relationship between the odds of *S. mansoni* and potential risk factors.

## Results

### Prevalence of *Schistosoma* infection

Table 1 summarises the prevalence of *Schistosoma* infection, stratified by sex, age group and school. Schistosome eggs were detected in the stool and urine of 33 of the 274 children surveyed (12.0%) in October 2016. The prevalence of *S. mansoni* and *S. haematobium* were 9.5% and 2.6%, respectively. None of the children were concurrently infected with both *S. mansoni* and *S. haematobium*. From the initial 75 SCORE schools, six were included in the current study. In Gbadrou, Mona, Ziondrou and Zoukougbeu, *S. mansoni* prevalence failed to decrease by at least 35% and/or the intensity of infection failed to decrease by at least 50% when comparing the baseline situation in 2012 with the endline situation in 2016 after multiple rounds of treatment (Table 2). Hence, these four schools were considered PHS in the current analysis. The remaining two schools (Tobly Bangolo and Zè) achieved a *S. mansoni* prevalence reduction by at least 35% and/or the intensity of infection declined by at least 50%. Subsequently, these two villages were considered low-prevalence schools. In general, prevalence and intensity of *S. mansoni* infection decreased meaningfully in schools where children received four rounds of praziquantel treatment compared with schools where only two treatments were performed.

The average *S. mansoni* prevalence in the four PHS was 10.9% (95% CI: 6.8-16.4%), while it was 6.6% (95% CI: 2.5-13.8%) in the two low-prevalence schools. The highest *S. mansoni* infection prevalence was found in Ziondrou (22.7%, 95% CI: 11.5-37.8%). The arithmetic mean egg count among *S. mansoni*-positive individuals was 123.8 EPG in the four PHS schools and 18.7 EPG in the two low-prevalence schools. *S. mansoni* prevalence was significantly different between schools ( $\chi^2 = 14.3$ ,  $p = 0.014$ ). Infections with *S. mansoni* were mainly light (80.8%), while moderate (15.4%) and heavy (3.9%) infections were less common. The two age groups did not differ significantly in the *S. mansoni* prevalence ( $\chi^2 = 0.1$ ,  $p = 0.751$ ).

The average *S. haematobium* prevalence was 3.3% (95% CI: 1.2-7.0%) in the four PHS schools and 1.1% (95% CI: 0.03-6.0%) in the two low-prevalence schools. The highest prevalence of *S. haematobium* was found in Zoukougbeu (8.2%, 95% CI: 0.2-16.1%). Among *S. haematobium*-positive children in the four PHS schools, the intensity of the infection was 16.7 eggs/10 ml of urine (95% CI: 0.3-33.0 eggs/10 ml of urine). All *S. haematobium* infection were of light intensity ( $< 50$  eggs/10 ml of urine). *S. haematobium* prevalence differed significantly between the two age groups with children aged 11-12 years at higher odds of infection ( $\chi^2 = 11.0$ ,  $p = 0.001$ ).

### Demographic characteristics

Table 3 summarises the demographic characteristics of the study villages, including results from the questionnaire survey regarding the last round of preventive chemotherapy. Overall, 299 children participated in the parasitological survey. However, 25 children were absent during the questionnaire survey, resulting in 274 children with complete parasitological and questionnaire data. There were 150 (54.7%) boys. There was no significant sex difference between the two groups of schools. With regard to age, 60.6% of the children were in the younger age group (9-10 years) with no significant difference in the age groups between the PHS and low-prevalence schools ( $p = 0.805$ ) (Table 4).

Most of the children (97.5%) declared that they were present during the last treatment round and 95.3% of them had received praziquantel. Reasons for not taking the drugs were absence for at least five days (37.5%), unwillingness to be treated based on a dose pole (25.0%), feeling healthy (12.5%), feeling too sick (12.5%) or absence of a community health worker during treatment administration (12.5%). There was no statistical significant difference between PHS and low-prevalence schools in terms of praziquantel coverage ( $p = 0.055$ ). In Ziondrou and Zoukougbeu, all the surveyed children reported having received the drugs, while in the remaining schools some children were missed out. Two-third of the children (65.7%) said that they did not know which disease praziquantel cured. About one out of seven children (13.5%) was of the opinion that praziquantel cured stomach diseases. Slightly more than 10% of the children reported that praziquantel cured worms (10.6%) or other non-specified diseases (10.2%). There was no association between children's knowledge and the prevalence of *S. mansoni* ( $\chi^2 = 0.9$ ,  $p = 0.818$ ). There was no significant difference between PHS and low-prevalence schools in the proportion of children who reported that praziquantel cured worm disease or stomach ache ( $p = 0.714$ ).

### Socioeconomic status and WASH indicators

Figure 1 shows the characteristics of household materials in the study villages. Most of the children in Zoukougbeu lived in houses with walls made of bricks, stones or cement (91.8%) and floors done with cement, tiles or linoleum (93.9%). The majority of children's families had houses covered by metal roofs (91.8%). In Mona, houses were predominantly constructed with simple or natural materials for wall (52.2%) and roof (63.0%). Three-quarter of children's families in the study villages had electricity at home. In Tobly Bangolo, only half of the participants had access to the power grid. Mobile phones were available in the households of all children surveyed in Zoukougbeu and Zè. Electricity at home was significantly associated with households that had at least one mobile phone ( $\chi^2 = 25.6$ ,  $p < 0.001$ ), walls made of bricks, stones or cement ( $\chi^2 = 10.4$ ,  $p = 0.001$ ), floors done with cement, tiles or linoleum ( $\chi^2 = 5.2$ ,  $p = 0.023$ ) and roofs made of iron sheets ( $\chi^2 = 13.3$ ,  $p < 0.001$ ). We found a statistically significant association between households possessing at least one mobile phone and households with walls made of bricks, stones or cement ( $\chi^2 = 7.9$ ,  $p = 0.005$ ) and roofs made of cement, tiles or linoleum ( $\chi^2 = 13.7$ ,  $p < 0.001$ ). The odds of being infected with *S. mansoni* was higher in households with floor done with cement, tiles or linoleum compared with households having other types of floors (OR = 3.55, 95% CI: 0.56-22.60) (Table 5).

Table 6 summarises WASH indicators in the study area. There were three types of public water sources: wells, pumps and taps. In PHS schools, between 30 and 38 children per school (65.2-86.4%) mentioned that their parents collected water from safe water sources for drinking or cooking. The proportions of children who reported that their parents obtained water from safe water sources were 86.4%, 72.7%, 65.3% and 65.2% in Gbadrou, Ziondrou, Zoukougbeu and Mona, respectively. In Ziondrou (65.9%) and Gbadrou (52.3%) safe water was predominantly collected from taps. In Zoukougbeu (30.6%) and Mona (28.3%), wells were the most common source of water collection for domestic use.

In low-prevalence schools, the number of children who reported that they used safe water sources was 37 (82.2%) and 33 (71.7%) in Zè and Tobly Bangolo, respectively. In Zè, water was mainly collected from taps (42.2%), while in Tobly Bangolo, wells were the predominant source (39.1%). The proportion of children who said that their parents collected water from safe sources was 72.1% in PHS schools and 76.9% in low-prevalence schools with no statistically significant difference ( $\chi^2 = 0.72$ ,  $p = 0.396$ ). There was a statistically significant difference between PHS and low-prevalence schools in the proportion of children who reported receiving water for drinking ( $p = 0.005$ ) with a higher proportion of children receiving water for drinking in low-prevalence schools.

Three-quarter of the children (74.1%) had latrines at home. Slightly less than half of the children (48.9%) had no latrine at school. In PHS, latrines were available in more than 80% of children's houses. A high coverage of latrines at school was mainly recorded in Ziondrou (81.8%). In the low-prevalence school Zè, high coverage of latrines both at home (80.0%) and at school (91.1%) was reported. The logistic regression analysis showed that the odds of being infected with *S. mansoni* was significantly lower among children from households with a latrine compared with household without a latrine (OR = 0.26, 95% CI: 0.11-0.63).

Figure 2 shows latrine availability and open defecation practice, stratified by village. Open defecation was practiced in each of the six villages studied in greater depth. In general, open defecation was less practiced in villages with high number of latrines. In PHS, less than half of the children (43.2%) from Gbadrou claimed that they never used a latrine. Most of the children in Gbadrou (95.5%), and Zoukougbeu (65.3%) did not use latrines at school. Open defecation was widely practiced in Gbadrou (90.9%). There was no significant difference between PHS and low-prevalence schools in the proportion of children practicing open defecation ( $p = 0.893$ ).

In low-prevalence schools, the number of children who had latrines ( $n = 36$ ) and those practicing open defecation ( $n = 35$ ) were similar in Tobly Bangolo. Most of the children (80.4%) in the later school did not use latrine at school while defecating at school. Open defecation was commonly practiced in Tobly Bangolo (76.1%).

### Water-related activities

Table 7 summarises the water sources for playing, laundry, bathing and collecting water for domestic or agricultural use in the study area. Most of the children (61.7%) reported that they performed the aforementioned activities in lake, pond or stream water.

According to the questionnaire survey, only 7.7% of the responders played in backwater, while in Zê most of the children (97.8%) played in this type of water. More than 60% of the children did not play in rivers, while more than half of the children did so in Zoukougbeu (79.6%) and Gbadrou (52.3%) in the PHS, and in the low-prevalence school Zê (68.9%).

Most of the children did not do laundry in lakes or small dams, with the exception of children from Zoukougbeu. Most of the responders did not do laundry in a river, while the majority of children did so in Gbadrou (61.4%), Mona (69.6%) and Zê (71.1%). Only 25.2% of the children bathed in the river, but in Zê, half of the children did so (50.0%).

Significant differences between the four PHS and the two low-prevalence schools were found for doing the laundry, playing, bathing and collecting water for domestic or agriculture use, playing in small dams, playing in river, doing the laundry in small dams, bathing in backwater, collecting water from backwater, collecting stagnant water for domestic or agricultural use and using stagnant water for bathing ( $p < 0.05$ ). Statistically significant differences were not found for playing and doing laundry in pond ( $p = 0.059$ ). In addition, there were no significant differences for the remaining variables of water-related activities between the four PHS and the two low-prevalence schools.

Children bathing in stagnant water were at higher odds of *S. mansoni* infection compared with children not reporting this activity (OR = 3.67, 95% CI: 1.02-13.18) (Table 3). Children from PHS schools bathed in stagnant water more frequently compared to those from low-prevalence schools ( $\chi^2 = 5.1$ ,  $p = 0.024$ ).

### Results from the malacological survey

Table 8 summarises the results of a cross-sectional malacological survey carried out in the study area. Overall, 76 human-water contact sites were reported in the six villages. In PHS schools, the number of human-water contact sites ranged between 10 (Mona) and 19 (Gbadrou). In the two low-prevalence schools, 14 and 9 human-water contact sites were visited in Zê and Tobly Bangolo, respectively. Rivers were the most common human-water contact sites ( $n = 59$ ). Rice paddies constituted another 20 human-water contact sites. There were 41 easily accessible human-water contact sites. Water is present throughout the year in all the visited sites. Among the surveyed human-water contact sites, 100% open canopy and 100% covered canopy were reported in 45 and 12 sites, respectively. Vegetation was found in 74 sites. The predominant substrata were mud, sand and roots, observed in 76, 70 and 69 of the sites, respectively. Domestic animals were found in 31 human-water contact sites, including pig (31.3%), dogs (26.6%), cows (15.6%), chicken (10.9%), sheep (9.4%) and goats (6.3%). Birds were the only wild animals observed in all the surveyed human-water contact sites. The average pH was 6.2 (range: 5.0-7.1). The average total dissolved solids was 9.5 ppm (range: 4-51 ppm) and the average conductivity was 38.9  $\mu$ S (range: 9-103  $\mu$ S).

A total of 688 snails belonging to nine genera were found; namely, *Biomphalaria*, *Bulinus*, *Ferrissia*, *Gyraulus*, *Lanistes*, *Lymnaea*, *Melanooides*, *Pila* and *Physa*. *Melanooides tuberculata* (400 specimens) was the most abundant snail species. With regard to potential schistosomiasis intermediate host snails, there were 92, 25, 16 and one specimens of *Bi. pfeifferi*, *Bu. forskalii*, *Bu. globosus* and *Bu. truncatus*, respectively. However, none of the snails were shedding schistosome cercariae, while some snails were observed shedding xiphidio-cercariae.

## Discussion

The current study was carried out in six schools in the western part of Côte d'Ivoire, where *S. mansoni* is endemic [7, 23, 24]. The study was embedded in a cluster-randomised trial, implemented from 2012 to 2016, to determine the effect of different praziquantel treatment schedules for sustaining the control of schistosomiasis mansoni in moderate endemicity settings (*S. mansoni* prevalence at baseline ranging from 10% to 24%) [7, 8, 27]. Our results after multiple rounds of preventive chemotherapy with praziquantel revealed considerably lower prevalence and intensity of *S. mansoni* compared to the baseline situation. Yet, there were a few PHS schools, where the prevalence of *S. mansoni* did not reduce meaningfully or even increased despite repeated administration of praziquantel [8, 35].

Tobly Bangolo and Zê, the two low-prevalence schools included in our study, were subjected to four rounds of preventive chemotherapy with praziquantel. It is conceivable that the meaningful decrease in the prevalence and intensity of *S. mansoni* infection is due to the repeated administration of praziquantel, as part of the SCORE interventions from 2012 to 2016 [7]. However, in Mona, where schoolchildren also received four rounds of praziquantel, the intensity of *S. mansoni* infection failed to decrease by at least 50%. Continued transmission and rapid reinfection after the annual treatment campaign might explain this observation [10, 36].

We found that households in one of the PHS settings (i.e. Zoukougbeu) were economically better-off compared to the other villages. It is widely acknowledged that schistosomiasis is a disease of poverty with most cases reported from rural parts of low- and middle-income countries [22, 37]. Households with relatively higher wealth obviously have a higher ability to implement schistosomiasis preventive measures, such as WASH [21, 24]. The comparatively low prevalence of *S. mansoni* infection was probably due to high socioeconomic status.

We also found that open defecation was commonly practiced in the six study villages. As expected, open defecation was less common in villages with high numbers of latrines. Logistic regression analysis confirmed this observation; the odds of *S. mansoni* infection was lower in households with latrines compared with households without latrines, corroborating a systematic review and meta-analysis [38]. Hence, access to latrines is associated with a lower odds of *S. mansoni* infection. Nevertheless, the highest *S. mansoni* prevalence was observed in Ziondrou, where the highest latrine coverage was reported. This observation points to factors other than sanitation explaining specific patterns within PHS. The idiosyncrasy in the relationship between latrine coverage and

schistosomiasis prevalence has been reported before in the south-central part of Côte d'Ivoire [39]. A plausible explanation of this observation might lie in the behaviours of local residents.

Our questionnaire data showed that the proportions of households collecting water from safe water sources for domestic use were approximately the same in PHS and low-prevalence settings. In addition, no statistically significant differences were observed between PHS and low-prevalence schools with regard to the proportion of children with parents using water from safe water sources. These results indicate that the difference in schistosomiasis *mansoni* endemicity between PHS and low-prevalence schools was not explained by sources of water collection for drinking and cooking. Moreover, despite the availability of safe water sources, some households probably preferred using water from open surface sources. Hence, in the present study area, access to improved water supply has little impact on the transmission of schistosomiasis. Other factors that were not investigated in our study might also be at play. Our observation contradicts to a major review including data from 144 studies that concluded that improved water supply and sanitation is associated with lower prevalence and severity of schistosomiasis and other parasitic diseases [40]. One of Africa's largest population-based cohort study carried out in KwaZulu-Natal in South Africa demonstrated that high coverage of piped water could decrease schistosomiasis significantly [41]. A parasitological survey coupled with a questionnaire and direct observation methods, assessing access to, and use of, different water sources in the district 'des Savanes' in northern Côte d'Ivoire revealed that access to unimproved water for any activity (including crossing rivers) was significantly associated with schistosomiasis [42]. Plausible explanations of these conflicting results in the impact of safe water might be due to differences in study populations with regard to cultural habits and water contract behaviours.

We found that children bathing in stagnant water such as lakes, ponds, backwaters or dams were at high risk of *S. mansoni* infection. Lakes, ponds, backwaters and small dams offer suitable conditions for the development of *Bi. pfeifferi*, the intermediate host snail of *S. mansoni* [43]. Hence, human-water contact through bathing in stagnant water exposes children to the risk of schistosomiasis transmission. Children from PHS settings practiced bathing in stagnant water more frequently than children from low-prevalence schools. This suggests that children in PHS schools maintained schistosomiasis *mansoni* transmission through bathing in stagnant water, which might explain the apparent failure of praziquantel treatment in the four PHS schools, where cured children may have been rapidly reinfected after praziquantel administration [10, 36, 44].

Data from our questionnaire survey revealed low awareness and knowledge of schistosomiasis. Similar results were reported from communities in northern Senegal after several years of health education [36] and in high schistosomiasis prevalence settings in western Kenya [37]. Lack of knowledge pertaining to schistosomiasis might be associated with low rates of health-seeking and low compliance with preventive chemotherapy [45].

Malacological data confirmed the presence of *Bi. pfeifferi*, *Bu. globosus* and *Bu. truncatus* in western Côte d'Ivoire, corroborating previous reports [26]. A unique feature of our research was the observation of *Bu. forskalii* an intermediate host snail of *S. haematobium*. However, none of snails subjected to shedding indeed released schistosome cercariae. Similar results were reported from a 3-year study assessing the dynamics of freshwater snails and schistosomiasis prevalence in schoolchildren during the construction and operation of a multipurpose dam in the central part of Côte d'Ivoire [46]. In contrast, a previous study conducted from 1986 to 1989 in two villages in western Côte d'Ivoire found *Bi. pfeifferi* infection rates ranging from 3.8% to 12.7% [47]. The absence of snails shedding schistosome cercariae should be interpreted with caution, as we only pursued a single malacological survey. The presence of *Lymnaea* snails shedding xiphidio cercariae suggests that fascioliasis might be transmitted to human and domestic ruminants, such as cows, sheep and goats.

Our study has several limitations. First, a recent analysis using a dataset from Tanzania showed that PHS settings can be defined based on four different approaches, including prevalence and/or intensity of *Schistosoma* infection. The Tanzania study revealed that the same dataset yielded different numbers of PHS depending on the approach used to define them [48]. Hence, a PHS school can become a low-prevalence school and *vice versa* depending on the definition of PHS. We employed the most stringent definition of a PHS, namely that a village that did not decrease in *S. mansoni* prevalence by at least 35% and/or failed to reduce in intensity by at least 50% comparing baseline with year five testing. The second shortcoming pertains to limited malacological data. Indeed, only a single malacological survey was conducted during the dry season. Previous work in the western part of Côte d'Ivoire showed that intermediate host snails for schistosomiasis are more abundant during the rainy season and at least four malacological surveys should be conducted within a 12-month period to obtain meaningful data. In view of financial and human resource constraints, however, we were unable to conduct repeated malacological surveys. Yet, human-water contact sites in close proximity to the six study villages were characterised and georeferenced, and hence, these sites could be surveyed longitudinally in the future. A third shortcoming of the study pertains to the diagnostic approach. We collected two stool samples and a single urine sample, and hence, the true prevalence of infection was underestimated [49]. Fourth, we only included children aged 9-12 years, which might have biased questionnaire results. Indeed, 9- to 12-year-old children might not participate in household chores as their older counterparts. Hence, caution should be taken when interpreting our findings. However, the selection of children in this age range is in line with SCORE protocols and facilitated parasitological and questionnaire surveys because children in this age are easily found at school. Older children are more ashamed and thus less likely to participate in stool and urine collection.

## Conclusions

Bathing in stagnant water bodies and availability of latrines were the main factors characterising PHS of *S. mansoni* in the western part of Côte d'Ivoire. School-aged children become infected with *S. mansoni* primarily while bathing in open freshwater bodies, such as ponds, lakes and backwater. The availability and use of latrines at home decreased the risk of *S. mansoni* infection. Although *S. mansoni* is highly endemic in western Côte d'Ivoire, children had relatively little knowledge on schistosomiasis and they commonly practiced open defecation. Potential schistosomiasis intermediate host snails were present in open surface water bodies. Hence, an integrated approach for schistosomiasis control is warranted that includes information, education and communication to initiate behaviour change, provision and sensitization on the use of latrines and snail control, in order to complement preventive chemotherapy with praziquantel.

# Declarations

## Ethics approval and consent to participate

Not applicable

## Abbreviations

CI, confidence interval; EPG, eggs per gram of stool; PHS, persistent hotspot; OR, odds ratio; SCORE, Schistosomiasis Consortium for Operational Research and Evaluation; WASH, water, sanitation and hygiene; WHO, World Health Organization.

## Consent for publication

No applicable

## Availability of data and material

Data generated or analysed during the current study are not publicly available, but are available from the corresponding author on reasonable request.

## Competing interests

The authors declare that they have no competing interests.

## Funding

This work received financial support from the Rudolf Geigy Foundation for a 1-year post-doctoral fellowship granted to RKA. The study received additional financial support from the University of Georgia Research Foundation, Inc., funded by the Bill & Melinda Gates Foundation for the Schistosomiasis Consortium for Operational Research and Evaluation (SCORE). The funders had no role in study design, data collection and analysis, data interpretation, decision to publish or to write the manuscript.

## Authors' contributions

RKA, EKN and JU wrote the study protocol. RKA, TBIM and SD conducted the parasitological survey. RKA, LGB and JAK carried out the questionnaire survey. RKA, RNN and SD conducted the malacological survey. RKA managed, analysed and interpreted the data. RKA wrote the first draft of the manuscript. RNN, LGB, JAK, TBIM, MO, EH, JTC, SD, EKN and JU reviewed the manuscript. All authors read and approved the final version of the manuscript prior to submission.

## Acknowledgements

We are grateful to the secretariat of the Schistosomiasis Consortium for Operational Research and Evaluation (SCORE) for reviewing the study protocol and for useful comments on earlier drafts of the manuscript. We thank Prof. Christian Schindler from the Swiss Tropical and Public Health Institute for assistance in data analysis and interpretation. We thank Sadikou Touré, Raphael G. Diabré and Vincent A. Anzara for help with the field and laboratory work. We thank the team of the 'Laboratoire de Zoologie et de Biologie Animale' at the Université Félix Houphouët-Boigny for assistance with laboratory work. We thank the personnel of the Centre Suisse de Recherches Scientifiques en Côte d'Ivoire for administrative and technical support. We are grateful to Dr. Aboulaye Meïté, Director of the 'Programme National de Lutte contre les Maladies Tropicales Négligées à Chimiothérapie Préventive' of the Ministry of Health, and his team for the administrative support and for conducting the mass drug administration within the multi-year SCORE project. We thank the teachers, parents, guardians and children for their participation in the study. We thank the health, education and village authorities of the regions Cavally, Guémon, Haut-Sassandra and Tonkpi for their contributions.

## Authors' information

<sup>1</sup>Unité de Formation et de Recherche Biosciences, Université Félix Houphouët-Boigny, 22 BP 582, Abidjan 22, Côte d'Ivoire. <sup>2</sup>Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, 01 BP 1303, Abidjan 01, Côte d'Ivoire. <sup>3</sup>Swiss Tropical and Public Health Institute, P.O. Box, CH-4002 Basel, Switzerland.



## References

1. Hotez PJ, Alvarado M, Basáñez MG, Bolliger I, Bourne R, Boussinesq M, et al. The global burden of disease study 2010: interpretation and implications for the neglected tropical diseases. *PLoS Negl Trop Dis*. 2014;8 7:e2865.
2. GBD 2017 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1859–922.
3. WHO. Preventive chemotherapy in human helminthiasis: coordinated use of anthelmintic drugs in control interventions. Geneva: World Health Organization. 2006.
4. WHO. Schistosomiasis and soil-transmitted helminthiasis: number of people treated in 2016. *Wkly Epidemiol Rec*. 2017;92:749-60.
5. Colley DG. Morbidity control of schistosomiasis by mass drug administration: how can we do it best and what will it take to move on to elimination? *Trop Med Health*. 2014;42:25-32.
6. Ezeamama AE, He CL, Shen Y, Yin XP, Binder SC, Campbell CHJ, et al. Gaining and sustaining schistosomiasis control: study protocol and baseline data prior to different treatment strategies in five African countries. *BMC Infect Dis*. 2016;16:229.
7. Assaré RK, Knopp S, N'Guessan NA, Yapi A, Tian-Bi YNT, Yao PK, et al. Sustaining control of schistosomiasis mansoni in moderate endemicity areas in western Côte d'Ivoire: a SCORE study protocol. *BMC public health*. 2014;14:1290.
8. Assaré RK, Tian-Bi YNT, Yao PK, N'Guessan NA, Ouattara M, Yapi A, et al. Sustaining Control of schistosomiasis mansoni in western Côte d'Ivoire: results from a SCORE study, one year after initial praziquantel administration. *PLoS Negl Trop Dis*. 2016;10:e0004329.
9. Ernoult JC, Ba K, Sellin B. Increase of intestinal schistosomiasis after praziquantel treatment in a *Schistosoma haematobium* and *Schistosoma mansoni* mixed focus. *Acta Trop*. 1999;73:143-52.
10. N'Goran EK, Utzinger J, N'Guessan AN, Müller I, Zambé K, Lohourignon KL, et al. Reinfection with *Schistosoma haematobium* following school-based chemotherapy with praziquantel in four highly endemic villages in Côte d'Ivoire. *Trop Med Int Health*. 2001;6:817-25
11. Landouré A, Dembélé R, Goita S, Kané M, Tuinisma M, Sacko M, et al. Significantly reduced intensity of infection but persistent prevalence of schistosomiasis in a highly endemic region in Mali after repeated treatment. *PLoS Negl Trop Dis*. 2012;6:e1774.
12. Garba A, Lamine MS, Barkire N, Djibo A, Sofu B, Gouvras AN, et al. Efficacy and safety of two closely spaced doses of praziquantel against *Schistosoma haematobium* and *S. mansoni* and re-infection patterns in school-aged children in Niger. *Acta Trop*. 2013;128 2:334-44.
13. Kittur N, King CH., Campbell CHJ., Kinung'hi S, Mwinzi PMN, Karanja DMS, et al. Persistent hot spots in Schistosomiasis Consortium for Operational Research and Evaluation studies for gaining and sustaining control of schistosomiasis after four years of mass drug administration of praziquantel. *Am J Trop Med Hyg*. 2019;101:617-27.
14. Laamrani H, Mahjour J, Madsen H, Khallaayoune K, Gryseels B. *Schistosoma haematobium* in Morocco: moving from control to elimination. *Parasitol Today*. 2000;16 6:257-60.
15. Sang HC, Muchiri G, Ombok M, Odiere MR, Mwinzi PN. *Schistosoma haematobium* hotspots in south Nyanza, western Kenya: prevalence, distribution and co-endemicity with *Schistosoma mansoni* and soil-transmitted helminths. *Parasit Vectors*. 2014;7:125.
16. Zou L, Ruan S. Schistosomiasis transmission and control in China. *Acta Trop*. 2015;143:51-7.
17. Pennance T, Person B, Muhsin MA, Khamis AN, Muhsin J, Khamis IS, et al. Urogenital schistosomiasis transmission on Unguja Island, Zanzibar: characterisation of persistent hot-spots. *Parasit Vectors*. 2016;9:646.
18. Steinmann P, Keiser J, Bos R, Tanner M, Utzinger J. Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infect Dis*. 2006;6 7:411-25.
19. Dabo A, Diarra AZ, Machault V, Touré O, Niambélé DS, Kanté A, et al. Urban schistosomiasis and associated determinant factors among school children in Bamako, Mali, West Africa. *Infect Dis Poverty*. 2015;4:1-12.
20. Grimes J, Croll D, Harrison W, Utzinger J, Freeman M, Templeton M. The roles of water, sanitation and hygiene in reducing schistosomiasis: a review. *Parasites Vectors*. 2015;8:156.
21. Huang YX, Manderson L. The social and economic context and determinants of schistosomiasis japonica. *Acta Trop*. 2005;96:223-31.
22. Muhumuza S, Kitimbo G, Oryema-Lalobo M, Nuwaha F. Association between socio economic status and schistosomiasis infection in Jinja District, Uganda. *Trop Med Int Health*. 2009;14:612-9.
23. Utzinger J, N'Goran EK, Ossey YA, Booth M, Traoré M, Lohourignon KL, et al. Rapid screening for *Schistosoma mansoni* in western Côte d'Ivoire using a simple school questionnaire. *Bull World Health Organ*. 2000;78 3:389-98.
24. Raso G, Matthys B, N'Goran EK, Tanner M, Vounatsou P, Utzinger J. Spatial risk prediction and mapping of *Schistosoma mansoni* infections among schoolchildren living in western Côte d'Ivoire. *Parasitology*. 2005;131:97-108.
25. Assaré RK, Lai Y-S, Yapi A, Tian-Bi Y-NT, Ouattara M, Yao PK, et al. The spatial distribution of *Schistosoma mansoni* infection in four regions of western Côte d'Ivoire. *Geospatial Health*. 2015;10:345.
26. Yapi YG, Touré M, Boka OM, Tia E, Boby OA-M, Kadjo KA. Dynamique des populations de *Biomphalaria pfeifferi* et de *Bulinus globosus* en zone d'endemie schistosomienne en Côte d'Ivoire. *Eur Sci J*. 2014;17:339-63.

27. Assaré RK, Hürlimann E, Ouattara M, N'Guessan NA, Tian-Bi YNT, Yapi A, et al. Sustaining the control of *Schistosoma mansoni* in western Côte d'Ivoire: baseline findings before the implementation of a randomized trial. *Am J Trop Med Hyg.* 2016;94:352-60.
28. Montresor A, Crompton DWT, Hall A, Bundy D.A.P., Savioli L. Guidelines for the evaluation of soil-transmitted helminthiasis and schistosomiasis at community level. World Health Organization, Geneva. 1998:1-45.
29. WHO. Helminth control in school-age children, A guide for managers of control programmes, Second edition. Geneva, World Health Organization; 2011.
30. Yap P, Fürst T, Müller I, Kriemler S, Utzinger J, Steinmann P. Determining soil-transmitted helminth infection status and physical fitness of school-aged children. *J Vis Exp.* 2012;66:e3966.
31. Plouvier S, Leroy J.-C., Colette J. A propos d'une technique simple de filtration des urines dans le diagnostic de la bilharziose urinaire en enquête de masse. *Med Trop.* 1975;35:229-30.
32. Knopp S, Person B, Ame SM, Ali SM, Muhsin J, Juma S, et al. Praziquantel coverage in schools and communities targeted for the elimination of urogenital schistosomiasis in Zanzibar: a cross-sectional survey. *Parasites Vectors.* 2016;9.
33. Frandsen F, Christensen NO. An introductory guide to the identification of cercariae from African fresh-water snails with special reference to cercariae of Trematode species of medical and Veterinary Importance. *Acta Trop.* 1984;41:181-202.
34. WHO. Prevention and control of schistosomiasis and soil-transmitted helminthiasis: report of a WHO expert committee. WHO Tech Rep Ser. 2002;912:1-57.
35. Ouattara M, Diakité NR, Yao PK, Saric J, Coulibaly JT, Assaré RK, et al. Effectiveness of different school-based preventive chemotherapy strategies for sustained schistosomiasis control in Côte d'Ivoire: a 5-year cluster randomised trial. *Lancet Glob. Health.* 2019;(Submitted).
36. Garba A, Lamine MS, Barkire N, Djibo A, Sofo B, Gouvras AN, et al. Efficacy and safety of two closely spaced doses of praziquantel against *Schistosoma haematobium* and *S. mansoni* and re-infection patterns in school-aged children in Niger. *Acta Tropica.* 2013;128:334-44.
37. McManus DP, Dunne DW, Sacko M, Utzinger J, Vennervald BJ, Zhou XN. Schistosomiasis. *Nat Rev Dis Primers.* 2018;4:13.
38. Grimes JE, Croll D, Harrison WE, Utzinger J, Freeman MC, Templeton MR. The relationship between water, sanitation and schistosomiasis: a systematic review and meta-analysis. *PLOS Negl Trop Dis.* 2014;8 12:e3296.
39. Schmidlin T, Hürlimann E, Silué KD, Yapi RB, Hougbedji C, Kouadio BA, et al. Effects of hygiene and defecation behavior on helminths and intestinal protozoa infections in Taabo, Côte d'Ivoire. *PLoS One.* 2013;8:e65722.
40. Esrey SA, Potash JB, Roberts L, Shiff C. Effects of improved water supply and sanitation on ascariasis, diarrhoea, dracunculiasis, hookworm infection, schistosomiasis, and trachoma. *Bull World Health Organ.* 1991;69:609-21.
41. Tanser F, Azongo DK, Vandormael A, Bärnighausen T, Appleton C. Impact of the scale-up of piped water on urogenital schistosomiasis infection in rural South Africa. *eLife.* 2018;7:1-24.
42. Krauth SJ, Musard C, Traoré SI, Zinsstag J, Achi LY, N'Goran EK, et al. Access to, and use of, water by populations living in a schistosomiasis and fascioliasis co-endemic area of northern Côte d'Ivoire. *Acta Trop.* 2015;149:179-85.
43. Dzik AJ. Snails, schistosomiasis, and irrigation in the tropics. *Public Health.* 1983;97 4:214-7.
44. Tukahebwa EM, Vennervald BJ, Nuwaha F, Kabatereine NB, Magnussen P. Comparative efficacy of one versus two doses of praziquantel on cure rate of *Schistosoma mansoni* infection and re-infection in Mayuge District, Uganda. *Trans R Soc Trop Med Hyg.* 2013;107:397-404.
45. Sow S, de Vlas SJ, Mbaye A, Polman K, Gryseels B. Low awareness of intestinal schistosomiasis in northern Senegal after 7 years of health education as part of intense control and research activities. *Trop Med Int Health.* 2003;8:744-9
46. Diakité NR, Winkler MS, Coulibaly JT, Guindo-Coulibaly N, Utzinger J, N'Goran EK. Dynamics of freshwater snails and *Schistosoma* infection prevalence in schoolchildren during the construction and operation of a multipurpose dam in central Côte d'Ivoire. *Infect Dis Poverty.* 2017;6:93:1-9.
47. Yapi YG, Touré M, Boka OM, Tia E, Boby OA-M. Dynamique de transmission des schistosomes par *Biomphalaria pfeifferi* dans la région de Man, Côte d'Ivoire. *Bull Soc Pathol Exot.* 2014;107:317-22.
48. Kittur N, Binder S, Campbell CH, King CH, Kinung'hi S, Olsen A, et al. Defining persistent hotspots: areas that fail to decrease meaningfully in prevalence after multiple years of mass drug administration with praziquantel for control of schistosomiasis. *Am J Trop Med Hyg.* 2017;97 6:1810-7.
49. Utzinger J, Becker SL, van Lieshout L, van Dam GJ, Knopp S. New diagnostic tools in schistosomiasis. *Clin Microbiol Infect.* 2015;21:529-42.

## Tables

**Table 1** Prevalence of *Schistosoma* infection in children aged 9-12 years in the six schools of a large SCORE study in western Côte d'Ivoire in October 2016

Variable		Children examined			<i>S. mansoni</i>		<i>S. haematobium</i>	
		N	%	95% CI	%	95% CI	%	95% CI
Sex	Boys	150	10.1	5.5-17.10	1.3	0.2-4.8		
	Girls	124	8.8	2.2-21.4	4.0	1.3-9.1		
Age group <sup>a</sup>	9-10 years	166	9.0	4.3-16.6	0.0	0.0-1.8		
	11-12 years	108	10.2	2.4-23.2	6.5	1.0-19.2		
Schools								
Persistent hotspot school	Gbadrou	44	13.6	5.2-27.4	0.0	0.0-6.6		
	Mona	46	4.3	0.5-14.8	0.0	0.0-6.3		
	Ziondrou	44	22.7	11.5-37.8	4.5	0.6-15.5		
	Zoukougbeu	49	4.1	0.5-14.0	8.2	0.2-16.1		
	Total	183	10.9	6.4-15.5	3.3	1.2-7.0		
Low-prevalence school	Tobly Bangolo	46	8.7	2.4-20.8	0.0	-		
	Zê	45	4.4	0.5-15.1	2.2	0.1-11.8		
	Total	91	6.6	1.4-11.8	1.1	0.0-6.0		

<sup>a</sup> Significant difference between the two age groups in the prevalence of *S. haematobium* infection (p <0.001)

**Table 2** Prevalence and intensity of *S. mansoni* infection in children aged 9-12 years at baseline (2012) and endline (2016) in six selected schools in western Côte d'Ivoire

School	Study arm	Number of treatment rounds	Baseline (2012)			Endline (2016)			Relative change (comparing endline with baseline)		School category
			Prevalence	Arithmetic mean egg counts (EPG)	Treat-ment coverage (%)	Prevalence	Arithmetic mean egg counts (EPG)	Treat-ment coverage (%)	Prevalence (%)	Intensity (%)	
Ziondrou	C	2	19.0	114.9	77	12.0	234.7	89	-37	104	PHS
Mona	A	4	14.0	65.8	109	4.1	56.0	101	-71	-15	PHS
Ziondrou	B	2	32.7	81.0	79	22.0	76.4	85	-33	-6	PHS
Zoukougbeu	B	2	15.2	22.6	74	6.0	49.3	69	-60	118	PHS
Tobly Bangolo	A	4	26.0	33.0	71	8.0	10.0	93	-69	-70	Low-prevalence
	A	4	27.0	71.4	96	4.0	36.0	71	-85	-50	Low-prevalence

PHS, persistent hotspot

**Table 3** Characteristics of study population in six selected schools from a large SCORE study in western Côte d'Ivoire in November 2016

Variable	Overall		Persistent hotspot schools						Low-prevalence schools				
			Gbadrou		Mona		Ziondrou		Zoukougbeu		Tobly		Z
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	Bangolo	è	
Sex													
Male	124	45.3	12	27.3	20	43.5	23	52.3	28	57.1	18	39	25
Female	150	54.7	32	72.7	26	56.5	21	47.7	21	42.9	28	60	44
Age													
9-10 years	166	60.6	29	65.9	31	67.4	19	43.2	30	61.2	35	76	54
11-12 years	108	39.4	15	34.1	15	32.6	25	56.8	19	38.8	11	23	17
Mean number of persons per house	8.3		8.7		7.2		8.8		8.5		7.9		8.7
Presence of persons	267	97.5	44	100.0	45	97.8	44	100.0	47	96.0	43	93	97

tr  
e  
a  
t  
m  
e  
n  
t

N	6	2.2	0	0.0	1	2.2	0	0.0	1	2.0	3	6	1	2
o												5		2
Don't	1	0.3	0	0.0	0	0.0	0	0.0	1	2.0	0	0	0	0
reme														
mber												0		0

Recei  
ved  
prazi  
quant  
el

Y	261	95.3	43	97.7	44	95.6	44	100.0	49	100.0	37	8	4	9
e												0	4	7
s														
												4		8
N	13	4.7	1	2.3	2	4.4	0	0.0	0	0.0	9	1	1	2
o												9		
														2
												6		

Received and  
took all  
praziquantel  
tablets

Y	256	93.4	42	95.5	41	89.1	44	100.0	49	100.0	36	7	4	9
e												8	4	7
s														
												3		8
No or	18	6.6	2	4.5	5	10.9	0	0.0	0	0.0	10	2	1	2
don't														
know												1		
														2
												7		

Don't  
know  
what  
praziquan  
tel treats

Y	179	65.3	34	77.3	23	50.0	23	52.3	38	77.5	38	8	2	5
---	-----	------	----	------	----	------	----	------	----	------	----	---	---	---

e												2	3	1
s												.	.	.
												6		1
N	95	34.7	10	22.7	23	50.0	21	47.7	11	22.5	8	1	2	4
o												7	2	8
												.	.	.
												4		9

**Table 4** Demographic, socioeconomic and environmental factors stratified by two categories of school

Factor	Persistent hotspot schools		Low-prevalence schools		p-value*
	(%)	95% CI	(%)	95% CI	
Age group (11-12 years)	37.4	15.8-65.5	40.4	28.2-54.0	0.805
Sex	54.9	43.0-66.4	54.6	38.8-69.6	0.971
Received praziquantel	11.0	2.1-41.9	1.6	0.3-7.6	0.055
Drinking water is given once at school	4.4	0.6-26.5	16.4	5.6-39.5	0.005
Drinking water is given twice at school	4.4	1.6-11.8	39.3	10.6-78.1	
Houses covered by metal roofs	82.4	51.5-95.4	77.6	35.5-95.6	0.755
Walls made of bricks, stones or cement	84.6	67.3-93.6	74.3	45.8-90.8	0.338
Floors done with cement, tiles or linoleum	83.5	80.8-85.9	82.5	51.6-95.4	0.908
Praziquantel treats worm disease or stomach ache	20.9	5.6-53.8	25.7	11.3-48.4	0.714
Latrine at home	78.0	73.9-81.7	72.1	54.0-85.1	0.367
Open defecation outside school	31.9	3.1-87.3	29.0	5.6-73.6	0.919
Mobile phone	94.5	68.2-99.3	95.6	84.4-98.9	0.815
Received and took all praziquantel tablets	38.5	37.1-39.8	41.5	26.3-58.5	0.654
Open defecation	11.0	4.9-22.8	10.4	5.2-19.7	0.893
Laundry in stagnant water	11.0	3.2-31.6	14.2	3.4-43.9	0.726
Latrine	89.0	47-98.7	85.8	57.1-96.5	0.789
Safe water sources	76.9	64.9-85.7	72.1	58.8-82.4	0.472
Playing in stagnant water	8.8	1.9-32.6	13.1	4.3-33.7	0.591
Electricity	69.2	26.8-93.3	77.6	59.7-89.0	0.603
Bathing in stagnant water**	3.3	0.4-20.7	11.5	2.5-39.3	0.217
Open surface water for drinking and cooking	3.3	0.4-21.5	9.8	6.0-15.8	0.204

\* of the respective difference in prevalence between PHS- and low prevalence schools.

Confidence intervals and p-values were calculated taking into account the clustered data structure with schools as primary sampling units

\*\*Bathing in stagnant water source such as pond, lake or backwater

**Table 5** Association between demographic and environmental factors and *S. mansoni* prevalence

Variable	Odds ratio	p-value	95% CI
Bathing in stagnant water	3.67	0.047	1.02-13.18
Latrine at home	0.25	0.013	0.10-0.64
Latrine at home or at school	1.92	0.044	1.02-3.61
Floors done with cement, tiles or linoleum	3.55	0.139	0.56-22.60

Odds ratios were obtained from a logistic regression model including all four variables together and taking into account the clustered data structure with schools as primary sampling units.

**Table 6** WASH indicators in six study schools from a large SCORE study in western Côte d'Ivoire in November 2016



Overa		Persistent hotspot schools								Low-prevalence schools			
II		Gbadr		Mona		Ziond		Zoukougbeu		Tobly Bangolo		Zè	
		ou				rou							
n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
67	24.5	8	18.2	13	28.3	1	2.3	15	30.6	18	39.1	12	26.7
207	75.5	36	81.8	33	71.7	43	97.7	34	69.4	28	60.9	33	73.3
93	33.9	23	52.3	5	10.9	29	65.9	13	26.5	4	8.7	19	42.2
181	66.1	21	47.7	41	89.1	15	34.1	36	73.5	42	91.3	26	57.8
42	15.3	7	15.9	12	26.1	2	4.5	4	8.2	11	23.9	6	13.3
232	84.7	37	84.1	34	73.9	42	95.5	45	91.8	35	76.1	39	86.7
202	73.7	38	86.4	30	65.2	32	72.7	32	65.3	33	71.7	37	82.2
72	26.3	6	13.6	16	34.8	12	27.3	17	34.7	13	28.3	8	17.8
203	74.1	27	61.4	32	69.6	29	65.9	44	89.8	35	76.1	36	80.0
71	25.9	17	38.6	14	30.4	15	34.1	5	10.2	11	23.9	9	20.0
140	51.1	2	4.5	35	76.1	36	81.8	17	34.7	9	19.6	41	91.1

134	48.9	42	95.5	11	23.9	8	18.2	32	65.3	37	80.4	4	9.0
238	86.9	28	63.6	40	87	44	100	45	91.8	36	78.3	45	100

36	13.1	16	36.4	6	13	0	0	4	8.2	10	21.7	0	0.0
111	40.5	40	90.9	9	19.6	3	6.8	20	40.8	35	76.1	4	8.9

163	59.5	4	9.9	37	80.4	41	93.2	29	59.2	11	23.9	41	91.1
-----	------	---	-----	----	------	----	------	----	------	----	------	----	------

---

**Table 7** Water-related activities in the six schools of a large SCORE study in western Côte d'Ivoire in November 2016

	Over all		Persistent hotspot schools								Low-prevalence schools				p-value
			Gba drou		Mon a		Zion drou		Zoukougbeu		Tobly Bangolo		Zè		
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	
es	169	61.7	35	79.6	23	50	32	72.7	22	44.9	21	45.6	36	80	
lo	105	38.3	9	20.4	23	50	12	27.3	27	55.1	25	54.4	9	20	0.028
ies	138	50.4	30	68.2	17	36.9	27	61.4	14	28.6	17	36.9	33	73.3	

lo	136	49.6	14	31.8	29	63.1	17	38.4	35	71.4	29	63.1	12	26.7	0.40
es	5	1.8	0	0	0	0	0	0	1	2	4	8.7	0	0	2
lo	269	98.2	44	100	46	100	44	100	48	98	42	91.3	45	100	0.02
es	7	2.5	2	4.5	0	0	5	11.4	0	0	0	0	0	0	5
lo	267	97.5	42	95.5	46	100	39	88.6	49	100	46	100	45	100	0.05
es	21	7.7	4	9.1	3	6.5	9	20.5	1	2	3	6.5	1	97.8	9
lo	253	92.3	40	90.9	43	93.5	35	79.5	48	98	43	93.5	44	2.3	0.15
es	94	34.3	23	52.3	10	21.7	9	20.5	10	79.6	11	23.9	31	68.9	2
lo	180	65.7	21	47.7	36	78.3	35	79.5	39	20.4	35	76.1	14	31.1	0.00
es	5	1.8	0	0	0	0	0	0	1	98	4	8.7	0	0	4

lo	269	98.2	44	100	46	100	44	100	48	2	42	91.3	45	100	0.02
															5
es	7	2.6	2	4.5	0	0	5	11.4	0	0	0	0	0	0	0

lo	267	97.4	42	95.5	46	100	39	88.6	49	100	46	100	45	100	0.05
															9

---

\*Activities: to do the laundry, playing, bathing and collecting water for domestic or agriculture use.

**Table 7** Water-related activities in the six schools of a large SCORE study in western Côte d'Ivoire in November 2016 (continued)

	Overall		Persistent hot-spot schools								Low-prevalence schools				p-value
			Gbadrou		Mona		Ziondrou		Zoukougbe		Tobly Bangolo		Zê		
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	
es	25	9.2	5	11.4	2	4.4	12	27.3	0	0	4	8.7	2	4.4	0.305
Jo	249	90.8	39	88.6	44	95.6	32	72.7	49	100	42	91.3	43	95.6	
es	97	35.4	27	61.4	14	69.6	9	20.5	11	22.5	4	8.7	32	71.1	0.310
Jo	177	64.6	17	38.6	32	30.4	35	79.5	38	77.5	42	91.3	13	28.9	
es	3	1.1	0	0	0	0	0	0	1	2	2	4.4	0	0	0.216
Jo	271	98.9	44	100	46	100	44	100	48	98	44	95.6	45	100	
es	6	2.2	2	4.5	0	0	4	9.1	0	0	0	0	0	0	0.081
Jo	268	97.8	42	95.5	46	100	40	90.9	49	100	46	100	45	100	
es	16	5.8	2	4.5	1	2.2	11	25	1	2	1	2.2	0	0	0.018
Jo	258	94.2	42	95.5	45	97.8	33	75	48	98	45	97.8	45	100	
es	69	25.2	20	45.5	7	15.2	9	20.5	8	16.3	3	6.5	22	48.9	0.538
Jo	205	74.8	24	54.5	39	84.8	35	79.5	41	83.7	43	93.5	23	51.1	
es	3	1.1	0	0	0	0	0	0	1	2	2	4.4	0	0	0.212
Jo	270	98.9	44	100	46	100	44	100	48	97.9	44	95.6	44	100	
es	5	1.8	3	6.8	0	0	2	4.5	0	0	0	0	0	0	0.112

---

40	269	98.2	41	93.2	46	100	42	95.5	49	100	46	100	45	100	10
															0

---

**Table 7** Water-related activities in the six schools of a large SCORE study in western Côte d'Ivoire in November 2016 (continued)

Variable	Overall	Persistent hot-spot schools								Low-prevalence schools				p-value		
		Gbadrou		Mona		Ziondrou		Zoukougbeu		Tobly Bangolo		Zê				
		n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)			
Overall	Yes	28	10.2	5	11.4	2	4.4	15	34.1	3	6.1	2	4.4	1	2.2	0.008
	No	246	89.8	39	88.6	44	95.6	29	65.9	46	93.9	44	95.6	44	97.8	
Overall	Yes	43	15.7	14	31.8	6	13.0	9	20.5	4	8.2	2	4.4	8	17.8	0.131
	No	231	84.3	30	68.8	40	87.0	35	79.5	45	91.8	44	95.6	37	82.2	
Overall	Yes	32	11.7	6	13.6	3	6.5	13	29.5	2	4.1	7	15.2	1	2.2	0.294
	No	242	88.3	38	86.4	43	93.5	31	70.5	47	95.9	39	84.8	44	97.8	
Overall	Yes	36	13.1	7	15.9	2	4.4	16	36.4	1	2.0	8	17.4	2	4.4	0.458
	No	238	86.8	37	84.1	44	95.6	28	63.6	48	98.0	38	82.6	43	95.6	
Overall	Yes	34	12.4	7	15.9	2	4.4	16	36.4	4	8.2	4	8.7	1	2.2	0.014
	No	238	86.8	37	84.1	44	95.6	28	63.6	48	98.0	38	82.6	43	95.6	




at	No	240	87.6	37	84.1	42	95.6	28	63.6	45	91.8	42	91.3	44	97.8	0.0
	Yes	24	8.8	4	9.1	1	2.2	14	31.8	2	4.1	3	6.5	0	0.0	
s*	No	250	91.2	40	90.9	45	97.8	30	68.2	47	95.9	43	93.5	45	100.0	24
	Yes	24	8.8	4	9.1	1	2.2	14	31.8	2	4.1	3	6.5	0	0.0	


\*use stagnant water source such as pond, lake, or backwater for bathing

**Table 8** Snail species in the six schools of a large SCORE study in western Côte d'Ivoire in December 2016

	Persistent hot-spot schools				Low-prevalence schools		Total
	Gbadrou	Mona	Ziondrou	Zoukougbeu	Tobly	Zê	
	19	10	12	12	9	14	76
<i>Galari</i>	0	77	12	0	3	0	92
<i>eri</i>	2	6	3	6	4	4	25
<i>ii</i>	0	6	0	7	3	0	16
<i>us</i>	0	0	0	0	1	0	1
<i>us</i>	0	0	0	0	0	4	4
<i>nsis</i>	3	13	1	6	0	0	23
<i>tus</i>	0	0	0	4	0	0	4
<i>æa</i>	7	54	1	21	20	0	103
<i>sis</i>	0	333	0	67	0	0	400
<i>lata</i>	0	5	0	0	0	0	5
<i>a</i>	0	7	0	8	0	0	15



**Figure 1**  
 Characteristics of house building materials, access to the power grid and possession of mobile phones in six villages in western Côte d'Ivoire in 2016



**Figure 2**  
 Latrine (at home and/or school) coverage and open defecation behaviour among school-aged children in six villages of western Côte d'Ivoire in 2016

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterial2.docx](#)
- [Supplementarymaterial1.docx](#)