

Mortality at the Pediatric Emergency Unit of the Mohammed VI Teaching Hospital of Marrakech

mounir bourrous (✉ mounirbourrous@yahoo.fr)

CADI AYYAD UNIVERSITY <https://orcid.org/0000-0003-2044-3514>

Widad Lahmini

cadi Ayyad University

Research article

Keywords: Pediatric mortality, pediatric emergencies, etiologies, prevention

Posted Date: January 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.21789/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 23rd, 2020. See the published version at <https://doi.org/10.1186/s12873-020-00352-9>.

Abstract

Background : Pediatric mortality is still one of the development indicators of a country. In our country the published data concerning the causes and the factors related to pediatric mortality especially in the pediatric emergency wards is very rare. This study aimed to determine the global rate of the pediatric mortality in the pediatric emergency department (PED), to study its epidemiology and to identify its most frequent causes. **Methods :** It is a retrospective and descriptive study, over five years (1st january 2012 and 31st december 2016) including all children aged from 0 to 15 years old who died at the PED in the Mohamed VI Hospital in Marrakech. **Results :** During the period of the study a total of 172.691 patients were admitted, among which 628 died in the PED (global pediatric mortality rate : 3,63%). The masculine gender was predominant with a gender ratio of 1.59. Two-thirds of the patients died in the first 24 hours. The median of time from admission to death was around 12 hours. Majority of the deceased children (75%) were from a low socioeconomic status. The most frequent cause of admissions was respiratory distress (47%) followed by neurological disorders (11%). Neonatal mortality was predominant with (75,1%) followed by postnatal mortality (11,6%). No autopsy was performed during the study period. **Conclusion :** Our data once again underline the crucial importance of prevention. This requires correct follow-up of the pregnancies, an adequate assistance of births, and perfecting healthcare provision to newborns in order to attain proper assistance.

Introduction

The death of a child at the emergency ward is a real challenge and one of the most difficult problems that the clinicians of these wards have to deal with [1]. Pediatric mortality is still one of the development indicators of a country. At the international level, relevant progress has been made in reducing infantile mortality over the last decades. The mortality rate of children under five years old has seen a global decline between 2000 and 2016 going from 69,4 to 38,4 for 1000 live births [2]. In our country the published data concerning the causes and the factors related to pediatric mortality especially in the pediatric emergency wards is very rare. Therefore, we aim through this study, to determine the global rate of the pediatric mortality in the pediatric emergency department (PED), to study its epidemiology and to identify its most frequent causes.

Patients And Methods

It is a retrospective and descriptive study, over five years, of files that belonged to the deceased children of the PED in the Mohamed VI Hospital in Marrakech, exploited with a pre established document. The study's population consisted of children aged from 0 to 15 years old who died at the PED between 1st january 2012 and 31st december 2016. We divided the deaths into 4 groups on the basis of age at death :

- Neonatal mortality: death between birth and the 28th day after birth.
- Postnatal mortality: death between the 29th day and the 1st year of life.

-Juvenile mortality: death between the 1st and the 5th year of life.

-The mortality of children aged between 5 and 15 years old.

Only children aged between 0 and 15 years old and who died in the PED were included in this study. Children that were dead on arrival at the emergency unit were excluded from this study. We mainly exploited the socio-demographic and clinical characteristics of the deceased patients ; the causes of deaths, the date and the time of death. The statistical data were analyzed by the SPSS software. Conditions of confidentiality and anonymous status were thoroughly respected.

Results

During the period of the study (2011-2016) a total of 172.691 patients were admitted, among which 628 died in the PED, leading to a global pediatric mortality rate estimated at 3,63% over 5 years. The masculine gender at any age was predominant with a rate of 61% (gender ratio: 1.59). Neonatal mortality was predominant with a global rate of 75,1% followed by postnatal mortality with 11,6%. Juvenile mortality was at 7,3 % and the one for children aged between 5 and 15 years old was at 5,8 %. The most frequent cause of admissions was respiratory distress (47%) followed by neurological disorders (11%), prematurity (7%) and lastly fever (4%).

The most frequent causes of pediatric mortality, whatever the age range, were dominated by neonatal pathologies (62,3%), followed by infectious causes bronchopulmonary infections included (7,7%), and last but not least birth deformities (7,3%) while traumas were merely (0,9%) (Table I). Majority of the deceased children (75%) were from a low socioeconomic status. Their geographic origin was mostly rural (51%). More than two thirds (67%) consulted directly in the PED, while 30% were referred from an external health institution. Most of the deaths (40%) occurred in the winter season. Majority of the deaths (68%) were recorded between the time range of 4pm to 8pm. The deceased children at the PED spent a median of time around 12 hours. Two-thirds of the patients died in the first 24 hours, among which 20% died in the first hour after their admission (Table II). The most frequent causes of neonatal mortality were : prematurity (24,4%), neonatal infections (32,2%) and perinatal asphyxia (15%) (Table III). The causes of the post natal mortality were dominated by pneumonia (27,1%) followed by birth defects (18,5%). The causes of the juvenile mortality were dominated equally by pneumonia, meningitis, birth defects and neoplasia with a rate of 11,1% each. The mortality causes of children aged between 5 and 15 years old were mainly neoplastic pathologies (23,6%) (Table IV). No autopsy was performed during the study period.

Discussion

The national data on pediatric mortality is poorly handled. Our study is the first to lead an audit on pediatric mortality in a third level ward of the south region. Our ward was open in July 2008 and serves not only the entire region of Marrakech but also the south of Morocco especially when it comes to third level pathologies. Its activity didn't stop going from 21.289 in 2008 to 70.176 admissions 2018. During

our study period, 628 deaths were noted leading to a global mortality rate of 3,63‰. While comparing our results to similar available series, we found that our global rate (3,63‰) was way underneath that of the Bassey et al [3] study with 27‰, the Robinson et al [4] study (37,9‰), the Ndu et al [5] study (58‰), the Joffiro et al [6] study 41‰ and the Santhanam study 122‰ [7]. This global rate though is underneath that of the Zhu et al [8] study with 0.5‰. All these studies were conducted in developing countries. In developed countries, the death of children in pediatric emergency is rare and even has a tendency of dropping [9,10]. Therefore our global rate is largely superior to the 1.7 per 10000 admissions recorded in the Maniktala et al study in the USA [11], and to the rate of 1.5 deaths per 10000 admissions recorded in the Lopez et al study in Spain [12]. Parents earning low income and residing in rural areas were the main risk factors of pediatric mortality around the world [13]. In our study, we had a high rate of children coming from disadvantaged areas. In fact, the mortality level was higher in low income households (52%) and within those living in rural areas (51%). Financial and geographic inaccessibility on top of a parent's low level of education may explain the high rate of child mortality in this group. Some studies found links between the mother's social and economical factors and the mortality, morbidity and even the children's congenital abnormalities. Understanding these associations may contribute to interventions that aim at better equity in the healthcare system [14,15]. More than one third of the deaths in our ward occurred in the winter season coinciding with the peak of bronchopulmonary infections. This observation is reported in other studies especially from Africa [3,5]. Two-thirds of our patients (67%) died in the first 24 hours compared to 32% for Joffiro et al [6]. Our median of the period of stay in the PED was 12 hours largely exceeding that of Zhu which was around 1,5 hours [8]. Ideally, the period of stay of patients shouldn't exceed 48 hours. On the other hand, it was noted that the delay in the patients' transfer to emergency units was associated to a higher mortality level. Therefore shortening the period of stay of patients in emergency units and avoiding any delay in their transfers to the intensive care unit may reduce the mortality rate [8,16,17]. In our study, neonatal mortality was dominant with a rate of 75.1 %. The causes of this neonatal mortality were dominated by neonatal infections (32,2%), prematurity (24,4%) and perinatal asphyxia (15%). Although, this rate is high, it joins the data of other studies [7,8,18-20]. Neonatal mortality is a real tragedy of the public health especially in developing countries, even with its drop in all of the regions of the world. Between 1990 and 2017, the global rate of neonatal mortality dropped by 51% going from 36,6 deaths in 1990 to 18 per 1000 deaths in 2017 [20]. Although, a lot of substantial progress has been made to reduce the neonatal mortality in 1990, more efforts should be made to achieve the sustainable development goal by 2030 [20]. A global approach containing all of the mothers' and the newborns' aspects of health, with an improvement of the environmental and socioeconomic conditions of the mom, follow-ups of the pregnancy, medicalisation of the births and an improvement of the care given to the newborn seem necessary knowing that prematurity and perinatal asphyxia are still the main causes of mortality. On the other hand, increasing the number of adapted units and neonatology wards for a better approach of the healthcare provision seems to be a priority [18,21]. Our study noted a high level of mortality (93,3%) in children under 5 years old in our hospital, which rhymes with other studies in developing countries [5,6,8]. The main causes of this mortality were dominated by pneumonia and meningitis there by correlating with results from other countries [6,8,18]. With regards to children between 5 and 15 years old, we found that neoplasms were the main causes, similar to other studies

made in China and Ethiopia [6,8]. In our context, even though we have a pediatric oncology hospital, the lack of places for hospitalization and oncology-specific care explains this high mortality rate. Scorpions stings occur in an endemic state in the region of Marrakech. But seeing the elaboration of a national strategy to provide proper care and faster admission for severe cases in the pediatric intensive care unit, the mortality in our study does not exceed 0,3% (2 cases). The same conclusion was reported for traumas and burns, which are frequent in the emergency units but their mortality rate remain low (0,9 and 0,4% respectively) due to the urgent care in the pediatric intensive care unit. The infant sudden death syndrome was also reported as a cause of death in the Lopez et al and Maniktala et al studies with the following rates: 20,7% and 31% respectively [11,12]. We couldn't explore more about this cause due to the criteria of our study that excluded children that were dead on arrival. In our context, the parents' hesitation towards the practice of the autopsy make it hard to determine the causes of the infant's sudden death syndrome. The same report was given by the Ntuli et al study [18]. Zhu et al reported a rate of 45,2% of children that were dead on arrival for whom no autopsy was done too [8]. On the other hand, other studies showed that more than the half of the deaths had one or multiple problems at the same time implying mainly severe asphyxia at birth, congenital defects, little weight at birth and prematurity [8,11]. Bad nutrition and diarrhea were also comorbidities in Africa [6,22]. This wasn't found in our study. One of the limits of our study, seeing its retrospective nature, was the intrication of multiple factors in certain patients making the search for the main cause of death complicated. On another note, the indeterminated mortality rate in our work (15,2%) was lower than the Zhu et al study (20%) [8] and higher than the Ntuli et al study (6%) [18].

Conclusion

The analysis of the causes of death in the pediatric emergency units may help in the development of practical care and prevention. Our study keeps all its interest in being the first epidemiologic study of its type made in the biggest Pediatric Emergency Unit in the south of Morocco. It showed a relatively high mortality rate in children under 5 years old dominated by the neonatal mortality. Its causes were dominated by perinatal accidents particularly that of neonatal infection and prematurity. Our data once again underline the crucial importance of prevention. This requires correct follow-up of the pregnancies, an adequate assistance of births, increasing the number of the neonatal units and perfecting healthcare provision to newborns in order to attain proper assistance.

Abbreviations

PED: pediatric emergency departement.

Declarations

Acknowledgements: the authors want to acknowledge warmly Dr. Anass Errahoui, Physician in the Pediatric Emergency Departement of Mohammed VI University Hospital of Marrakesh, for her close cooperation and all the efforts made for data collection.

Funding: Not applicable.

Availability of data and material: The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Authors' contributions:

MB and **WL** analyzed and corrected all the details of the article. **WL** contributed to study conception and design; data collection, analysis, and interpretation; and manuscript preparation. She approved the final manuscript.

MB conceived the study, reviewed the article, analysis and approved the submission of the manuscript.

Ethics approval and consent to participate: This project was approved by the ethics committee of the Faculty of Medicine of the University of Cadi Ayyad in Marrakech. Given the retrospective nature of this study, the project was deemed to be of very low risk for the patients and the anonymity and confidentiality of the patients was respected throughout the study.

Consent for publication: Not applicable.

Competing interests: The authors declare that they have no competing interests.

Publisher's Note : Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors' details :

Paediatric Emergency Department, UHC Mohamed VI, Marrakech. Department of Paediatric, Cadi Ayyad University, Faculty of medicine and pharmacy; PO Box: 7010; Sidi Abbad Street; 40000; Marrakech; Morocco.

References

- 1- O'Malley PJ, Barata IA, Snow SK. Death of a Child in the Emergency Department. *Pediatrics*. 2014 Jul; 134(1):198-201.
- 2- Global, regional, and national under-5 mortality, adult mortality, age-specific mortality, and life expectancy, 1970–2016: a systematic analysis for the Global Burden of Disease Study 2016. GBD 2016 Mortality Collaborators. *Lancet* 2017; 390: 1084–1150.
- 3-Bassey EU, Ijezie E. Pediatric Emergencies Seen in a Tertiary Hospital in Uyo, Akwa Ibom State of Nigeria: A two Year Review. *Int J Sci Stud* 2016;4(4):42-45.

- 4-Robison JA, Ahmad ZP, Nosek CA, Durand C, Namathanga A, Milazi R, et al. Decreased pediatric hospital mortality after an intervention to improve emergency care in Lilongwe, Malawi. *Pediatrics* 2012; 130: e676–682.
- 5-Ndu IK et al. Pattern of Morbidity and Mortality at the Children Emergency Unit of Enugu State Teaching Hospital, Park lane, Enugu. *Niger. Journal of Experimental Research* 2016; 4 (1): 48-54.
- 6- Jofiro G, Jemal K, Beza L, Bacha Heye T. Prevalence and associated factors of pediatric emergency mortality at Tikur Anbessa specialized tertiary hospital: a 5 year retrospective case review study. *BMC Pediatrics* (2018); 18:316.
- 7-Santhanam I, Pai M, Kasturi K, Radhamani MP. Mortality after admission in the pediatric emergency department: a prospective study from a referral children's hospital in southern India. *Pediatr Crit Care Med*. 2002 Oct; 3(4):358-63.
- 8- Zhu CP, Wu XH, Liang YT, Ma WC, Ren L. The mortality of patients in a pediatric emergency department at a tertiary medical center in China : An observational study. *World J Emerg Med*, Vol 6, No 3, 2015.
- 9-Whitfill T, Auerbach M, Scherzer DJ, Shi J, Xiang H, Stanley RM. Emergency Care for Children in the United States : Epidemiology and Trends Over Time. *J Emerg Med*. 2018 Sep ; 55(3) :423-434.
- 10- Matthew O'Meara and Susan Trethewie. Managing paediatric death in the emergency department. *J Paediatr Child Health*. 2016 Feb;52(2):164-7.
- 11- Maniktala A, Pruitt C, Poirier M. Deaths in a non-trauma center pediatric emergency department: a ten-year experience. *J Emerg Med* 2002; 1(2).
- 12- Lopez E, Udaondo J, Olabarri M, Martinez-Indart L, Benito J, Mintegi S; Mortality Group of RISEUP-SPERG. Mortality in Spanish pediatric emergency departments: a 5-year multicenter survey. *Eur J Emerg Med*. 2017 Dec;24(6):392-397.
- 13-Chao F, You D, Pedersen J, Hug L, Alkema L. National and regional under-5 mortality rate by economic status for low-income and middle-income countries: a systematic assessment. *Lancet Glob Health*. 2018 May;6(5):e535-e547.
- 14- Hajizadeh M, Nandi A, Heymann J. Social inequality in infant mortality : what explains variation across low and middle income countries? *Soc Sci Med*. 2014 Jan ;101:36-46.
- 15- Kosowan L, Mignone J, Chartier M, Piotrowski C. Maternal Social and Economic Factors and Infant Morbidity, Mortality, and Congenital Anomaly: Are There Associations? *Fam Community Health*. 2019 Jan/Mar;42(1):54-61.
- 16- Chalfin DB, Trzeciak S, Likourazos A, Baumann BM, Dellinger RP; DELAY-ED Study Group. Impact of delayed transfer of critically ill patients from the emergency department to the intensive care unit. *Crit*

Care Med 2007;35: 1477–1483.

17- McKelvie B, McNally JD, Chan J, Momoli F, Ramsay C, Lobos AT. Increased Mortality and Length of Stay Associated With Medical Emergency Team Review in Hospitalized Pediatric Patients : A Retrospective Cohort Study. *Pediatr Crit Care Med*. 2017 Jun ; 18(6):571-579.

18-Sam Thembelihle Ntuli, Ntambwe Malangu & Marianne Alberts. Causes of Deaths in Children under-Five Years Old at a Tertiary Hospital in Limpopo Province of South Africa. *Global Journal of Health Science* ; Vol. 5, No. 3 ; 2013.

19-Nadir Ijaz, Matthew Strehlow, N. Ewen Wang, Elizabeth Pirrotta, Areeba Tariq, Naseeruddin Mahmood and Swaminatha Mahadevan. Epidemiology of patients presenting to a pediatric emergency department in Karachi, Pakistan. *BMC Emergency Medicine* (2018) 18:22.

20-Mansoor Ahmed, Youngjoon Won. Cross-National Systematic Review of Neonatal Mortality and Postnatal Newborn Care : Special Focus on Pakistan. *Int J Environ Res Public Health*. 2017 Dec; 14(12): 1442.

21-Yalçın SS, Tezel B, Köse MR, Tugay D, Mollahaliloğlu S, Erkoç Y. Changes and determinants in under-five mortality rate in Turkey since 1988. *Cent Eur J Public Health*. 2013 Jun; 21(2):80-7.

22-Tette EMA, Nyarko MY, Nartey ET, Neizer ML, Egbeforme A, Akosa F et al. Under-five mortality pattern and associated risk factors: a case-control study at the Princess Marie Louise Children's Hospital in Accra, Ghana. *BMC Pediatr*. 2016 Aug 31;16(1):148.

Tables

Table I : Main causes of pediatric mortality for all ages

Causes of death	Number of deaths	Rate
Neonatal infections	152	24,2%
Prematurity	115	18,3%
Perinatal asphyxia	71	11,3%
Neonatal respiratory distress	57	9%
Infectious causes	24	3,8%
Pulmonary infections	25	3,9%
Malformations	46	7,3%
Heart Diseases	13	2%
Neoplasia	16	2,5%
Uncertain	96	15,2%
Poisoning	9	1,4%
Traumatic causes	6	0,9%
Various	18	0,4%

Table II : Demographic characteristics and chronological distribution of deaths

VARIABLES	NUMBER	RATE
Patients admitted to the ED	172 691	
Gender (ratio : 1,59)		
Girl	245	61 %
Boy	383	39 %
Age (0 days to 15 years)		
≤ 1 month	472	75,1%
1 month-1 year	73	11,6%
1-5 years	46	7,3%
5-15 years	37	5,8 %

Socio-economic level

Low	471	75%
Average	138	22%
High	19	03%

Geographic origin

Rural	320	51%
Suburban	94	15%
Urban	214	34%

Time of death after admission (median : 12 hours)

≤ 01 hours	63	10%
≤ 12 hours	138	22 %
≤ 24 hours	220	35%
≤ 48 hours	138	22%
> 48 hours	69	11%

Reference mode

Self-reference	433	69%
Referral from another health center	195	31%

Table III : Etiologies of neonatal mortality

Causes of death	Number of deaths	Percentage
Neonatal infection	152	32,2%
Prematurity	115	24,4%
Perinatal asphyxia	71	15%
Neonatal respiratory distress	57	12,1%
Malformations	28	5,9%
Others	48	10,1%
TOTAL	471	100%

Table IV : Causes of mortality in children over 1 month

5 to 15 years	1 month to 1 year	1 to 5 years
Pneumonia -	19 (27,1%)	5 (11,1%)
Meningitis -	7 (10%)	5 (11,1%)
Diarrheal disease -	7 (10%)	-
Sepsis 5 (13,1%)	-	-
Malformations -	13 (18,5%)	5 (11,1%)
Heart diseases 3 (7,8%)	7 (10%)	3 (6,6%)
Neoplasia 9 (23,6%)	2 (2,8%)	5 (11,1%)
Poisoning -	3 (4,2%)	-
Traumatic causes 3 (7,8%)	-	3 (6,6%)
Scorpions stings -	-	2 (4,4%)
Burns -	-	3 (6,6%)
Unclear 18 (47,3%)	12 (17,1%)	14 (31,1%)
Total 38 (100%)	70 (100%)	45 (100%)