

Effect of Cognitive-behavioral Therapy on Domestic Violence and Its Consequences in Transgender Youth: a Randomized Clinical Trial, Parallel Group Study

Mahdiah Damanpak-Rizi

Tehran University of Medical Sciences

Farnaz Farnam (✉ Fz.farnam@gmail.com)

Tehran University of Medical Sciences

Parisa Khodakhah

Support center for Iranian Transgender

Research Article

Keywords: Transgender youths, Family violence, Cognitive-behavioral therapy, Mental health, Randomized controlled trial, Iran

DOI: <https://doi.org/10.21203/rs.3.rs-128510/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Family violence against transgender people is a common issue and affects their mental health. Very few if any interventions have been designed to reduce family violence against transgender youths. This RCT will evaluate the efficacy of cognitive-behavioral therapy for transgender parents on their violent behavior towards their children and transgender youth's mental health.

Methods: This study is a randomized controlled trial conducted on 50 transgender youths with selected inclusion criteria in Iran. The intervention will be undertaken on the parents or guardians of these transgender youths in eight 1-hour online sessions for the intervention group to increase their knowledge of gender dysphoria, to help control their anger regarding their offspring's gender dysphoria and learn to manage stressful situations. This study's primary outcomes include frequency of family violence towards transgender youths, depression, anxiety, stress, self-esteem, suicidal thoughts, and suicide attempts in transgender youth and the secondary outcome is parental conflict resolution tactics before and after the intervention.

Discussion: To the best of our knowledge, this is one of the first RCT on family violence against transgender youth in the world. Findings will help to provide better education and intervention for transgender parents to reduce violence against their children.

Results: N/A

Conclusion: N/A

Trial registration: IRCT20120609009975N7 (08/03/2020).

Background

Gender identity is defined as being male or female, which corresponds to one's gender(1). There is usually a harmony between sex and gender identity, but sometimes these two items are incompatible(2). According to the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5), "the incongruity between a person's experienced gender and the gender assigned to him or her at birth" leads to "Gender Dysphoria," a term that replaced the previous "Gender Identity Disorder" in order to avoid stigma(1, 3). The prevalence of gender dysphoria in trans women ranges from 0.005–0.014% and trans men from 0.002–0.003%. Since not all adults seek hormonal and surgical treatment at specialized clinics, this rate is probably underestimated (3).

Transgender people face many problems include lack of social support, community rejection, physical and family violence. Among these problems, family violence is a common issue (1, 4–7). Family violence is defined as any act or omission by cohabitating or living together as a family that results in serious injury to other family members. Family violence has several distinct subgroupings, such as child abuse, intimate partner abuse, and elder abuse(8).

Some families do not accept transgender youth because of the religious atmosphere and fear of disgrace and harassment of neighbors and other members of society, and they continuously treat their children with violence(9). A 2010 study found that 60% of transgender people are subject to domestic violence (10). According to a report in Iran, parents support their transgender child in only 30% of cases; in 10% of cases, parents do not care about this issue and choose to ignore it, but in 70% of cases, parents deal with anger and repression or are saddened to hear their child's demand for gender reassignment. These parents usually do not accept their child's problems or respond with inhumanity, anger, and resentment (2). In qualitative studies conducted by the research team (under submission), many of the problems faced by these individuals in society stem from violence and a lack of family support (11).

Although transgender people are a minority of the population, this population is increasing, and these individuals, like other individuals in the community, have the right to reproductive health (3). Several studies reported a high level of family violence against transgender youth, which can lead to suicidal thoughts, drug abuse, running away from home, and high-risk behaviors; but very few if any interventions have been conducted to reduce violence against these youths by working with potential perpetrators of violence (10, 12). This study is one of the first RCT on family violence against transgender youth globally and in Iran to the best of our knowledge.

Method/design

Aim:

Research Hypothesis

1. CBT can reduce family violence perpetrated by parents or guardians towards transgender youths.

Primary Objective:

1. The main primary objective is the frequency of family violence towards transgender youth before and after the intervention.
2. Other primary objectives are depression, anxiety, stress, self-esteem, suicidal thoughts, and suicide attempts changes in transgender youths before and after the intervention.

Secondary Objectives:

1. The secondary objective is comparing the parents' conflict resolution tactics about transgender youths before and after the intervention.

Study design:

This trial is a parallel randomized, controlled trial with block randomization and a 1:1 allocation ratio. The project has been reviewed and approved by the Tehran University of Medical Sciences under the ethical

approval code IR.TUMS.FNM.REC.1398.201 on 2020/02/18 and scientific and funding grant number 98-3-100-45678.

Study setting

The exact prevalence of gender dysphoria in Iran is not determined. Based on a survey that recorded the subjects referred to the Tehran Psychiatric Institute (TPI) from 2002 to 2009 with a diagnosis of gender dysphoria by two independent psychiatrists, the prevalence of Gender dysphoria in Iran considered 1:141000 (referral population considered people aged between 15 and 44) (13). Iran is one of the first Islamic countries where gender reassignment has been legalized since 1964 and was second only to Thailand for the most gender reassignment surgeries in 2008 (14). However, despite the legality of gender reassignment, transgender youth are still not sufficiently accepted in society due to cultural and religious pressures; and society, family, friends, school, and co-workers perpetrate violence and discrimination against transgender people. The steps for obtaining allowance of gender reassignment surgery for transgender people in Iran are as follows: Request to Family Court; refer the person to the Legal Medicine Organization (LMO) by the court; evaluate hormonal and karyotype tests and at least 12 counseling sessions with an expert psychiatrist in LMO; approve gender dysphoria diagnosis by a psychiatrist and then reapproved in LMO Commission with attendance of an expert group and referred accepted the request to the court again for the final decision. Only after this approval will the transgender youth change their identity card, have the desired sex clothes, and undergo gender reassignment surgery (although many have started hormone therapy many years sooner by a private physician or my own).

Participants

Our participants will randomly select from the list of transgender youth in the “support center for transgender Iranians (Mahtaa)”. An announcement about this study will first post on the mahtaa pages on Instagram and Twitter. Inclusion criteria for transgender youths are the definitive diagnosis of Gender Dysphoria; age older than 12 years; living in a home with parents or guardians; transgender youth's and their parents' willingness to participate in the study and being a victim of family violence. This later criterion assesses through one question, "Have you experienced any physical, emotional, verbal, or sexual violence from your family in the past four months?" The only exclusion criterion is transgender parents' unwillingness to attend classes during the intervention process. The eligible participant who intends to participate in the study will apprise of the nature of the trial. Verbal and written informed consent will obtain before randomization. Our requisite sample size would be 50 transgender youth that will randomize into intervention and control groups using a randomized block design with four block sizes with a 1:1 allocation ratio. Study outcomes will measure through self-report questionnaires (Fig. 1).

Participant timeline

Randomization

The sample size of this study was calculated based on the most similar available article titled: “individual cognitive-behavioral treatment and family therapy for physically abused children and their offending parents: A comparison of clinical outcomes” (15). Based on this article's mean and standard deviation and with $\alpha = 0.05$ and power $1-\beta 0.90$ and considering a 20% drop, the total sample size required for each group is 25 people.

$$n_1 = n_2 = \frac{(S_1^2 + S_2^2)(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta})^2}{(\bar{X}_1 - \bar{X}_2)^2} \quad N1 = N2 = \frac{(62+4.52)(1.96+1.28)^2}{(9-3.6)^2} = 21$$

For concealment and implementation mechanism, FF will generate allocation sequence and opaque sealed envelopes of participants in each group. All people who give consent for participation and who fulfill the inclusion criteria will be randomized. Intervention sessions will be conducted by one of the research team members (MD) who is not involved in randomization and not assessing the study's outcomes.

Complete blinding is not possible due to the nature of the intervention. Neither participants nor staff can be blinded to allocation. An employee outside the research team will analyze data. Therefore, in this study, the outcome assessor will be blinded.

Intervention:

Fifty eligible transgender youths will assign to the intervention and control group. For assessing the main primary objective (frequency of family violence), all transgender youths will complete the family violence self-report checklist daily for one month as a pre-test tool and again for one month as a post-test (2 months after ending intervention). For other primary outcomes, DASS-21 and Rosenberg self-esteem questionnaires will be sent via an internet website for transgender youth as a pre-test and post-test. After completing the pre-tests by transgender youths, their parents will be invited to the CBT intervention group. For assessing secondary outcome, the parents will complete Conflict Tactics Scales; Parent-child version (CTSPC) as the pre-test and at the end of intervention as post-test. After receiving pre-test questionnaires, our intervention (CBT) will begin with the parents of the transgender youths. Parents in the intervention group will attend eight 1-hour online group therapy sessions. Furthermore, for better modification, these sessions will be held in four 2-hour sessions if parents could not attend eight sessions. Two months after the last session of intervention, post-test questionnaires will send to transgender youths.

The first session of intervention contains information for increasing the parent's/guardian's awareness of the nature of gender dysphoria, with talking about the history of gender dysphoria in Iran and the world, the theoretical causes and risk factors, different reactions of parents, expressing the research team's sympathy with the parents because of their sadness and concerns. The second session contains the problems faced by transgender youth, consequences of lack of family and community support, and examples of successful transgender youth. The third session includes an overview of existing counseling and treatments and health care needs before and after gender reassignment. The fourth session will be

about the definition of violence, family violence, and anger symptoms. The fifth and sixth sessions contain anger management strategies, relaxation, and home exercises, and the seventh and eighth sessions include speech and discussion skills training, empathy skills, conflict resolution skills, and stressful conditions management.

For ethical consideration, the control group will receive all meeting content after ending intervention.

Outcomes measurement

Primary Outcome Measures:

1. Family violence against transgender youth: This study's main primary outcome is the family violence against transgender youth that will be assessed by a research-made daily self-report checklist completed by transgender youths for one month prior and two months after the intervention. In our research, family violence means the frequency of physical, psychological, verbal, and sexual violence perpetrated by a transgender parent or guardian⁽¹⁶⁾

2. Depression, anxiety, and stress: The standard questionnaire of Depression Anxiety Stress Scale (DASS – 21) will use to determine these outcomes, It has 21 items, and each item has a 4-point Likert scale with a score ranging from 0 ("did not apply to me at all") to 3 ("applied to me very much").⁽¹⁷⁻¹⁹⁾

2. Self-esteem: Self-esteem will be determined by the standard questionnaire of Rosenberg Self-Esteem scale (RSES)⁽²⁰⁾, which has a score range between 0–30, where a score of less than 15 may indicate problematic low self-esteem. It is a ten-item Likert-type scale with items answered on a four-point scale—from strongly agree to disagree strongly. The scale measures global self-worth by measuring both positive and negative feelings about the self.⁽²¹⁾

3. Suicide thought and attempt: This outcome will be determined by two questions "How many times have you seriously considered suicide in the last **4** months? Never/once/ 2–3 times/4 times or more" and "How many times have you committed for suicide in the last four months? Never/once/ 2–3 times/4 times or more".

Secondary outcome measures:

1. Parental conflict resolution tactics: The Conflict Tactics Scales will assess this outcome; Parent-child version (CTSPC). The basic form of CTSPC consists of 22 items divided into three scales: non-violent discipline, psychological aggression, and physical assault. The latter is further split into three subscales according to severity, which is respectively called corporal punishment, physical maltreatment, and severe physical maltreatment. The questionnaire also contains supplementary scales on disciplinary methods, neglect, and sexual abuse in the preceding week.⁽²²⁾

Adherence:

1. Researchers will call the transgender youth once a week to remind them to fill out the family violence self-report checklist in the pre-test and post-test period.
2. Researchers will call the transgender youth's parents once a week to review any issues from the previous session and remind them of the next session.
3. Researchers will answer parents' sub-questions about contraception, sexual disorders, and reproductive health at the end of the sessions to encourage them to continue meetings.
4. Researchers help participants to use internet websites for data collection.

Concomitant care:

Parents or guardians who enroll in the study will be warned not to attend any family counseling from sampling and two months after the intervention.

Statistical methods

Outcomes:

The intervention group will compare against the control group, and also, both groups will compare before and after the intervention. Data analysis will perform with SPSS 21. The first description of the data (in percentage and frequency, mean and standard deviation) will provide. And then by using Independent T-test (examining the questionnaire scores in terms of binary variables such as the study group and gender), paired t-test (Study before and after intervention in one group), Pearson correlation test (for correlating questionnaire scores and quantitative variables such as age), ANOVA and Tukey post hoc tests (Analysis questionnaire scores and categorical variables such as job and economic status), Chi-square test (Relationship of two qualitative variables such as the relationship between levels of mental health and family violence with gender, marital status, ...) will analyze.

Additional analyses:

For qualitative variables and quantitative variables with a cut-off point, logistic regression, and quantitative numerical variables, linear regression will use.

Analysis population and missing data:

Outcome data obtained from all participants are included in the data analysis, regardless of protocol adherence. If there is missing data in an internet-based questionnaire, the questionnaire will be returned to the person to complete. Finally, if there is missing data, statistical methods of single imputation and multiple imputations will be used.

Access to data

MD and FF will be given access to the cleaned data sets. For confidentiality, data dispersed to project team members will be blinded to any identifying participant information.

Data management

Electronic files of participants will be stored in numerical order and stored in a secure and accessible place and manner. Participant files will maintain in storage for two years after completion of the study.

Formal committee and Interim analysis:

This trial does not have a data monitoring committee and any stopping guidelines.

Harms:

It seems that our RCT does not have any adverse effects on participants.

Auditing:

The Tehran University of Medical Sciences will do auditing.

Protocol amendments:

Any modifications to the protocol which may impact on the conduct of the study, the potential benefit of the participants or may affect participants' safety, including changes of study objectives, study design, participant population, sample sizes, study procedures, or significant administrative aspects will be notified to the Ethics Committee of Tehran University of Medical Sciences.

Consent or assent:

Comprehensible information about the research will provide in writing to the participants to ensure that they understand the research's purpose and methodology and are fully satisfied with the study. Participants can discuss with the researcher if they have any questions to be left vague to them. Written informed consent will maintain until the end of the study.

Ancillary studies:

Participants complete a consent form for using their data (without the name) in future research under the supervision of FF.

Confidentiality:

All study-related information will be stored electronically. Each participant will have his or her code to fill out forms and questionnaires; therefore, all reports, data collection, process, and administrative forms will be identified by a coded number only to maintain participant confidentiality. All records that contain names or other personal identifiers will be stored separately from study records identified by code

number. All files containing data will protect by password, and no one except the researcher will have access to it.

Abbreviations

Cognitive-behavioral therapy (CBT)

Tehran Psychiatric Institute (TPI)

Legal Medicine Organization (LMO)

Conflict Tactics Scales; Parent-child version (CTSPC)

Depression Anxiety Stress Scale (DASS -21)

Rosenberg Self-Esteem scale (RSES)

Declarations

Ethics approval:

- The study was approved by the research ethics committee of Tehran University of Medical Sciences on the 18.02.2020 (IR.TUMS.FNM.REC.1398.201).
- All methods were carried out in accordance with relevant guidelines and regulations.
- Informed consent obtain from all participants or, if participants are under 18, from a parent and/or legal guardian

Consent for publication:

Not applicable

Availability of data and materials:

The data will be available from the corresponding author on reasonable request.

Competing interests:

We have no conflict of interest to declare.

Funding:

Funding for this study was provided by the Vice-Chancellor for Research and Technology of Tehran University of Medical Sciences (Grant NO. 98-3-100-45678). Email: resdeputy@tums.ac.ir

Authors' contributions:

FF (Farnaz Farnam): she is a grant holder and contributes to conception and design of the study, design, and conduct of intervention, resolve the problems during the project, coordination with related organization, final analysis, interpretation of data, revise and approve final report and papers

MD (Mahdiah Damanpak): she contributes to the design of the study and conduct of the intervention, coordination with related organization, writing the first draft of the proposal and papers, data interpretation

PK (Parisa Khodakhah): she contributes to the sampling and designs the content of the intervention.

Acknowledgments:

The authors thank Vice-Chancellor for Research and Technology of the Tehran University of Medical Science for financial assistance. We thank Dr. Farshad Mamudi for his valuable help in improving the English used in the manuscript and Dr. Maryam Rad for assistance with the study's Statistical sections.

References

1. BJ S, VA S, p R. Kaplan and Sadock's Comprehensive Textbook of Psychiatry. 50 ed2017.
2. Momeni Javid M, Shoa kazemi M. Comparison of personality traits of people with Gender identity disorder and normal people.
3. Diagnostic and statistical manual of mental disorders : DSM-5. 5th ed2013.
4. Budge SL, Adelson JL, Howard KA. Anxiety and depression in transgender individuals: The roles of transition status, loss, social support, and coping. *Journal of consulting and clinical psychology.* 2013;81(3):545.
5. Zucker KJ, Bradley SJ. Gender identity and psychosexual disorders. *Focus.* 2005;3(4):598-617.
6. Roch A, Ritchie G, Morton J. Transgender people's experiences of domestic abuse. Scotland: LGBT Youth Scotland and the Equality Network. 2010.
7. Rezapour fardian R, Rasouli R, Mostafa V. Comparison of Perceived Social Support and Life Satisfaction in People with Gender Dysphoria.
8. Harvey Wallace P, Roberson C. Family violence, legal, medical and social perspectives. Eighth ed2016.
9. Javaheri F, Kouchakian Z. Gender identity disorder and its social dimensions: A study of the phenomenon of Gender Identity in Iran. 2001.
10. Dempsey B. Trans People's Experience of Domestic Abuse. *SCOLAG Legal Journal.* 2010(396):208-12.
11. Farnam F. Investigation of Discrimination, Violence and Suicide in Transgender Women in Iran.

12. Kenagy GP. Transgender health: Findings from two needs assessment studies in Philadelphia. *Health & social work*. 2005;30(1):19-26.
13. Ahmadzad-Asl M, Jalali A-H, Alavi K, Naserbakht M, Taban M, Mohseninia-Omrani K, et al. The epidemiology of transsexualism in Iran. *Journal of Gay & Lesbian Mental Health*. 2010;15(1):83-93.
14. Tait R. Sex change funding undermines no gays claim. *The Guardian*. 2007;26.
15. Kolko DJ. Individual cognitive behavioral treatment and family therapy for physically abused children and their offending parents: A comparison of clinical outcomes. *Child Maltreatment*. 1996;1(4):322-42.
16. Webster J, Holt V. Screening for partner violence: direct questioning or self-report? *Obstetrics & Gynecology*. 2004;103(2):299-303.
17. Lee E-H, Moon SH, Cho MS, Park ES, Kim SY, Han JS, et al. The 21-item and 12-item versions of the Depression Anxiety Stress Scales: Psychometric evaluation in a Korean population. *Asian nursing research*. 2019;13(1):30-7.
18. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): Construct validity and normative data in a large non-clinical sample. *British journal of clinical psychology*. 2005;44(2):227-39.
19. Gomez R. Depression Anxiety Stress Scales: Factor structure and differential item functioning across women and men. *Personality and Individual Differences*. 2013;54(6):687-91.
20. Rosenberg M. *Society and the adolescent self-image*: Princeton university press; 2015.
21. Blascovich J, Tomaka J. Measures of self-esteem. *Measures of personality and social psychological attitudes*. 1991;1:115-60.
22. Reichenheim ME, Moraes CL. Psychometric properties of the Portuguese version of the Conflict Tactics Scales: Parent-child Version (CTSPC) used to identify child abuse. *Cadernos de Saúde Pública*. 2006;22:503-15.