

Glycogen synthase kinase-3 β inhibitor lithium chloride protects against inflammation-mediated skeletal muscle wasting


Ji-Hyung Lee¹, Seon-Wook Kim¹, Jun-Hyeong Kim¹, Hyun-Jun Kim¹, JungIn Um¹, Da-Woon Jung^{1*} and Darren R. Williams^{1*}

¹New Drug Targets Laboratory, ²Laboratory of Drug Discovery, School of Life Sciences, Gwangju Institute of Science and Technology, 1 Oryong-Dong, Buk-Gu, Gwangju 61005, Republic of Korea

*To whom correspondence should be addressed: 1) Da-Woon Jung, School of Life Sciences, Gwangju Institute of Science and Technology, 1 Oryong-Dong, Buk-Gu, Gwangju 61005, Republic of Korea. Fax: +82 62-715-2484; Tel: +82-62-715-3554; email: jung@gist.ac.kr 2) Darren R. Williams, School of Life Sciences, Gwangju Institute of Science and Technology, 1 Oryong-Dong, Buk-Gu, Gwangju 61005, Republic of Korea. Fax: +82 62-715-2484; Tel: +82-62-715-2509; email: darren@gist.ac.kr

Supplementary figure 1

For figure 1B Uncropped Western blot



For figure 1F Uncropped Western blot




Figure S1 related to figure 1B and 1F. Uncropped western blot images for figure 1B and 1F. Red boxes were cropped images for designated figures.

Supplementary figure 2

For figure 2E Uncropped Western blot




Figure S2 related to figure 2E. Uncropped western blot images for figure 2E. Red boxes were cropped images for designated figures.