

Molecular characterization of *Leishmania* species from stray dogs and patients in Saudi Arabia

Abdullah D Alanazi (✉ aalanazi@su.edu.sa)

Shaqra University <https://orcid.org/0000-0002-4862-7668>

Abdulazi S Alouffi

King Abdulaziz City for Science And Technology

Mohamed S Alyousif

King Saud University

Abdulsadah A Rahi

University of Wasit

Magda A Ali

University of Wasit

Hend H.A.M. Abdullah

National Research Centre

Fabio A Brayner

Universidade Federal do Pernambuco, Recife

Jairo Alfonso Mendoza-Roldan

Universita degli Studi di Bari

Domenico Otranto

Universita degli Studi di Bari

Research

Keywords: Leishmania tropica, Leishmania major, Dogs, Patients, kDNA, nPCR, Saudi Arabia

Posted Date: September 11th, 2020

DOI: <https://doi.org/10.21203/rs.2.22961/v5>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: *Leishmania major* and *Leishmania tropica* cause cutaneous leishmaniasis in humans and dogs in several parts of the world, with a large number of cases recorded in the Middle East. However, when occurring in sympatry, the role of each species of *Leishmania* in the epidemiology of cutaneous leishmaniasis (CL) is not clear.

Methods: To determine the prevalence and to identify the species of *Leishmania* that infects humans and stray dogs in Riyadh and Al-Qassim (Saudi Arabia), 311 stray dogs and 27 human patients, suspected for *Leishmania*, were examined for CL by a nested PCR (nPCR).

Results: Nested PCR (nPCR) detected seven patients (25.9%) positive for cutaneous leishmaniasis. Five patients from Riyadh were infected by *L. major* and two from Al-Qassim by *L. tropica*. In addition, five dogs (1.6%) were infected by *L. tropica*.

Conclusions: This is one of the first molecular studies of leishmaniasis from Saudi Arabia. The relationship between the sand fly vectors and the reservoirs of both *Leishmania* spp. is still scarcely known in this region, and further epidemiological investigations of domestic and wild canids infected with *L. major* and *L. tropica* are needed towards a control and prevention of the infection in canine and human populations.

Background

Leishmaniasis are a complex of sand fly transmitted protozoa diseases, listed amongst the neglected tropical diseases affecting millions of people, mainly the world's most vulnerable populations [1]. These diseases are transmitted by phlebotomine sand flies of the genus *Phlebotomus* in the Old World and *Lutzomyia* in the New World. Leishmaniasis include cutaneous (CL), visceral (VL) and mucocutaneous leishmaniasis (MCL) forms, all of which have been reported in Saudi Arabia [2-4]. Moreover, in that country CL by *Leishmania major* has the highest prevalence mainly in Riyadh, Qassim, Al-Madinah, Al-Hassa, Hail and Asir [5-7] with an estimated number of more than 26,300 cases [2] over the past 10 years (2006–2016). In addition, in Saudi Arabia there are several reports of leishmaniasis by *Leishmania infantum*, *Leishmania major* and *Leishmania tropica* among humans and wild animals [8-12].

In spite of the number of molecular studies available to diagnose and identify *Leishmania* species worldwide [13-16], there is a lack of information on human CL patients as well as dog populations from endemic areas of Saudi Arabia. El-Beshbishy et al., [17] reported the molecular characterization of CL in patients from Al-Madinah Al-Munawarah province, western Saudi Arabia by internal transcribed spacer 1 (ITS-1) PCR-restriction fragment length polymorphism (RFLP) and kinetoplast DNA (kDNA) PCR established *L. major* and *L. tropica* as the causative organisms, with a kDNA PCR sensitivity of 90.7%, whereas ITS-1 PCR had a sensitivity of 70.1%. Rasheed et al., [18] reported the prevalence of *Leishmania* species among patients with CL in Qassim province, central Saudi Arabia. They recorded that out of total 206 CL biopsies, 49.5% biopsies were found to be positive for *L. major*, 28.6% biopsies were positive

for *L. tropica*, 3.9% were found to be positive for *L. infantum/donovani*. Therefore, the aims of the current study were to use molecular tools to detect and identify the *Leishmania* species infecting humans and stray dogs in Al-Qassim province and Riyadh city, Saudi Arabia in order to better understand the epidemiology of the infection.

Methods

Ethical approval

This study was reviewed and approved by the Ethics Committee of the Department of Biological Science at Shaqra University, according to the ethical principles of animal research (protocol SH 2-2017).

Study areas

The investigation was conducted from January 2018 to May 2019 in Al-Qassim province and Riyadh city, Saudi Arabia. Al-Qassim province is located at the central part of Saudi Arabia (latitude 25°–23° N and longitude 42°–24° E). It has an area of about 58.046 km² with a population of 1,423.000 people in 2017 [19]. Al-Qassim province is known as an agricultural region and it has a typical [desert climate](http://www.pme.gov.sa), with an average temperature of 13°C and hot summer (an average temperature of 35°C), with few annual rainfalls (214 mm) and low humidity ranging from 25% to 76% (<http://www.pme.gov.sa>). Conversely, Riyadh city is the capital of Saudi Arabia (latitude 24°–08°N and longitude 47°–18° E), with an area of about 1798 km² and inhabited by approximately seven million people in 2017 [19]. Riyadh is characterized by very hot summers with an average temperature of 45°C in July, whereas winters are cold. The overall climate is arid, with scarce annual precipitations (21.4 mm), with a relative humidity ranging from 10% to 47% throughout the year. Riyadh is also known to have many dust storms (<http://www.pme.gov.sa>) (Fig. 1).

Patients Biopsy tissue collections and DNA extraction

A total of 27 suspected patients were attended in both King Saud Medical City in Riyadh city (n =16) and *Buraydah Central Hospital* (n =11) in Al-Qassim province. All samples were diagnosed after clinical and microscopy examination [20]. Briefly, skin biopsies (i.e., 5-10 mm of diameter) were taken under sterile conditions from the border of the ulcer and cutaneous lesions and DNA samples were extracted from all biopsies by MagNaA pure DNA extraction Pure LC DNA Isolation Kit (Roche Applied Science, Mannheim, Germany) according to the manufacturer's instructions and the extracted DNA was quantified by Nanodrop spectrophotometer (Thermo, USA), DNA concentration differed from sample to another, but it ranged from 18ng/ul to 33ng/ul, and an aliquot (100 µl of DNA from each sample) was stored at -20 °C prior to nPCR amplification and analysis.

Sampling of stray dogs

From January 2018 to May 2019, 311 stray dogs were trapped in Al-Qassim province by bait traps (Havahart®) (Fig. 1). Dogs were examined physically for canine leishmaniasis skin lesions in the field. Seven dogs were suspected for canine leishmaniasis due to the presence of cutaneous nodules or ulcerated lesions on the skin (Table 1). Skin biopsies (5 mm in diameter) were taken under sterile conditions from the border of the ulcer and inoculated into medium M199 supplemented (Gibco, Life technologies, Germany) with 25 mmol/L HEPES (pH:7.5) and 20% fetal bovine serum (Gibco, Life technologies, Germany) followed by incubation at 24 °C. Ten days later parasites were harvested and washed with ice-cold phosphate buffered saline (PBS, pH: 7.4) and stored in -20 °C before DNA isolation. DNA from parasites cultures was isolated by using the ReliaPrep™ gDNA Tissue Miniprep System Kit (Promega, Madison, United States), following the manufacturer's instructions.

Table 1: Data of suspect dogs by canine leishmaniasis that trapped in Al-Qassim province.

Dog ID	Site of trapped dogs	Gender	Age (years)	Location of cutaneous lesions
Dog No.1	Unayzah	Male	2	left ear
Dog No.2	Al-Bukkayriah	Male	1	nose
Dog No.3	<i>Buraydah</i>	Male	4	abdomen
Dog No.4	Al-Bukkayriah	Female	2	under mouth lips
Dog No.5	<i>Buraydah</i>	Male	3	muzzle
Dog No.6	Ar Rass	Female	3	nose
Dog No.7	<i>Buraydah</i>	Male	2	upper right leg

Leishmania Nested PCR (nPCR)

The specific external CSB2XF primers (5'-ATTTTTCGCGATTTTCGCAGAAACG-3') and CSB1XR (5'-CGAGTAGCAGAACTCCCGTTCA-3') were used initially. In the second step, specific internal 13Z primers (5'-ACTGGGGGTTGGTGTAATAAG-3') and LiR (5'-TCGCAGAACGCCCT-3') were applied [21]. The specificity and sensitivity of this primers is reported to be 92% and 100%, respectively [21]. In addition, these primers were able to track and multiply the variable part of all forms of the *Leishmania* kDNA. Amplified fragments of *L. infantum* were 680 bp in length and fragments of *L. tropica* and *L. major* were 750 and 560 bp in length, respectively [21].

The first step of PCR master mix that included CSB2XF and CSB1XR were applied using AccuPower® PCR PreMix kit (Bioneer, Daejeon, Korea). The prepared PCR pre-mix volumes contained KCl 30 mM, MgCl₂ 1.5mM, Tris-HCL (pH 9.0) 10mM, Taq DNA polymerase, and dNTP were adjusted to 2 µl. In addition, 1 µl of the first step of each initial CSB1XR and CSB2XF primers at concentrations of 10 pmol/µl (Bioneer, Daejeon, Korea) and 3 µl of DNA were added to the complex. Finally, 13 µl of deionized water (ddH₂O) were added for a total volume of 20 µl for reaction. In this study, no negative control was included in the final nPCR, but all the necessary measures were taken in vitro to nullify unexpected results (i.e; each sample performed separately and individually in isolated PCR room). The reaction was performed in a thermal cycler (Techne TC-3000, USA) by set up the following conditions; initial denaturation temperature

of 94 °C for 5 min; followed by 30 cycles at denaturation 94 °C for 30 s, annealing 55 °C for 60 s, extension 72 °C for 60 s, final extension at 72 °C for 7 min and then the reaction was held at 4 °C. The second step of PCR included 13Z and LiR primers and the same PCR master mix except 3µL of template PCR product.. After that, PCR products were electrophoresed on a 1.5% agarose gel containing 1 µL Syber safe (Thermo Scientific™, Nalgene, UK) in Tris-acetate–EDTA buffer at 100 V for 45 min and visualized under UV imaging system (ImageQuant Laz4000, GE Healthcare Life Science, Hammersmith, UK). The size of each sample was estimated by comparison with a 100 bp DNA Ladder Marker (Solis BioDyne OU, Estonia).

***Leishmania* kDNA sequencing and BLAST analysis**

Positive amplified products of *Leishmania* species were sent to **Macrogen (South Korea)** for sequencing, and the results were compared with the sequences available in GenBank database using BLAST (<http://blast.ncbi.nlm.nih.gov/>). The obtained sequences were alignment with other reference sequences in GenBank using CLASTAL W in MEGA software version 7.0 [22]. The phylogenetic tree was constructed using the Maximum-Likelihood method and Tamura-Nei model with 500 bootstrap replicates in MEGA 7.0 software [22, 23].

Results

Of the 27 human patients examined, five from Riyadh and two from Al-Qassim were positive to *L. major* and *L. tropica* respectively (Fig. 2 and Fig. 3). Of 311 dogs, seven (2.3%) presented with thick cutaneous lesions (i.e., 1.5 × 5 centimeters) in different anatomical sites (e.g., nose, muzzle, abdomen and between fingers) and five of them were positive to *L. tropica*. Sequencing analysis of *Leishmania* kDNA confirmed that the five positive samples (nos. H1-H5) of the human patients from Riyadh were all *L. major* with a size ranging from 441 bp to 451 bp yielding a nucleotide identity ranging from 99.34% to 100% with query cover 100% with previous *L. major* sequences from Iraq (MN313423). *Leishmania* sequences from two human patients (no. H1 and H3) from Al-Qassim were most similar 99.66% to 100% with query cover 100%, respectively to *L. tropica* from Iraq (MF166799) (Fig.4). Sequences of *Leishmania* kDNA from stray dogs (no. D5, D6 and D7) were closely related (i.e., 99.33% to 99.80%) to kDNA of *L. tropica* from Iraq (MF166799), whereas two sequences (no. D2 and D4) displayed a close nucleotide identity (i.e., 99.35% and 99.80) to *L. tropica* kDNA from Iraq (MN334665) and UK (AF308689), respectively (Fig. 4). Phylogenetic tree clustered *L. major* (no. H1-H5) and *L. tropica* (no H1 and H3) sequences from human samples to those from Iraq (accession numbers MN313423 and MF166799, respectively), and the phylogenetic tree clustered *L. tropica* (no. D2 and D4) from Iraq and UK (accession numbers MN334665 and AF308689, respectively) , while (no. D5, D6 and D7) sequences from dog's samples to those from Iraq isolate (accession number MF166799) Fig. 4. In addition, the isolates of *L. tropica* from human and dogs in present study were closely related (i.e., 98.60% to 99.65% with query cover ranged 98.20% - 99.50%) to kDNA of *L. tropica* (Saudi strain, MHOM/SA/91/WR1063) that recorded on GenBank (accession number X84845.1). Representative sequences of *L. major* and *L. tropica* from this study were deposited in the GenBank database under the accession numbers MT787488 - MT787499.

Discussion

This study provides molecular evidence of the circulation of *L. major* and *L. tropica* in human and dog populations from the investigated areas. The above *Leishmania* spp. have already been recorded as agents of cutaneous leishmaniases in Saudi Arabia and Middle Eastern countries [23-26]. However, *L. tropica* infection has been herein molecularly diagnosed for the first time in humans and dogs in the central part of the Saudi Arabia since it was previously reported in the west and southern west of Saudi Arabia in association with the distribution of the sand fly species (i.e., *Phlebotomus sergenti*), which is a proper vector for that species [27]. Conversely, *L. major* is more prevalent throughout the country and can be found in the open deserts regions of Saudi Arabia [2, 28]. Previous studies in Saudi Arabia have reported the natural infection of *L. major* in dogs using enzymatic biochemical methods [29, 30], though no clinical information was available, nor serology or molecular confirmation were performed. The high nucleotide identity of human *L. major* and *L. tropica* isolates with those of Iraq (accession number MN313423 and MF166799) as well as of dog *L. tropica* isolates with Iraq and UK (accession number MN334665, MF166799 and AF308689, respectively) was also confirmed by the phylogenetic dendrogram herein presented. Moreover, this study showed that the isolates of *L. tropica* from human and dogs were closely related to kDNA of *L. tropica* (Saudi strain, MHOM/SA/91/WR1063) that recorded on GenBank (accession number X84845.1). This might be due to the distribution of similar sand flies species in the different parts of Saudi Arabia and Middle East, which may act as proper vectors of both *Leishmania* spp. [24, 31]. However, the phylogenetic analysis in this study was limited and performed based on the available sequences on GenBank based on the sequences of *L. tropica* and *L. major* in the Middle East countries particularly those from Iraq. The phylogenetic tree revealed that both *L. tropic* and *L. major* were clustered in separate clades. Moreover, *L. tropica* genotypes were clustered in a separate clade far from other *Leishmania* spp. (*L. infantum*, *L. chagasi* and *L. donovani*), this might be related to variations in geographical and environmental conditions [32].

Of the 25 species of *Phlebotomus* (*Ph.*) reported in Saudi Arabia only five (i.e., *Ph. papatasi*, *Ph. sergenti*, *Ph. bergeroti*, *Ph. kazeruni*, and *Ph. arabicus*) have been incriminated as vectors of CL [27, 28, 33, 34]. Of these, *Ph. papatasi* is the main and most predominant vector species for *L. major* [28, 33], while *Ph. sergenti* is the natural vector species for *L. tropica* [27]. Presence of *Ph. papatasi* and *Ph. sergenti* in Al-Qassim province suggests that they could have a potential role in the transmission of human and canine leishmaniasis. Nonetheless, more studies are required to elucidate the role of *Phlebotomus* spp. in CL disease transmission in Saudi Arabia.

Stray dogs have been often diagnosed in Saudi Arabia, with clinical disease associated with *Leishmania* species, however previous studies focused on the epidemiology, clinical, histopathological and biochemical aspects [11, 29, 30]. Conversely, molecular studies have reported the occurrence of dog infection by *Leishmania* spp. in Qatar [35], *L. tropica* in Iran [26, 36] and Israel [37, 38] and by *L. major* in Iraq [23] and Israel [25], which are in agreement with the current study. Though CL is endemic in many parts of Saudi Arabia, the paucity of data concerning the relationship between the disease, the vectors and reservoirs is a major hindrance to understand the transmission cycles in endemic areas, considering

that the distribution patterns can easily change through the years in specific geographic areas [39]. Data herein provided contribute to fill existing gaps in order to increase the alert by the Ministry of Health in Saudi Arabia in preventing outbreaks and the spread of CL.

Conclusion

This was the first study that detected and identified the causative agent of CL in stray dogs and patients from Saudi Arabia thus confirming that *L. major* and *L. tropica* are endemic in Al-Qassim province and Riyadh City. However, it is still unclear the relationship between the sand fly vectors and reservoirs and their specific role in the transmission cycles in endemic areas of Saudi Arabia. Further epidemiological and molecular studies focusing on CL these areas are advocated also in order to elaborate better strategic control plans and assess the risk for human health.

Declarations

Acknowledgements

This work was kindly supported by Researcher Supporting Project (RSP-2020-192) King Saud University, Riyadh, Saudi Arabia.

Ethics approval and consent to participate

Blood and tissue sampling for this study was approved by the Ethical Research Committee, Shaqra University and complied with relevant guidelines for animal handling and welfare. (Approval no. SH 02-2017)

Consent for publication

Not applicable.

Availability of data and material

All data generated or analyzed during this study are included in this published article.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

ADA, ASA and AAR conceived the study. MSA, ASA performed field works. ASA collected patient's and dog's samples. ASA, AAR and MAA carried out the biopsy tissue collections and DNA extraction. MAA, AAR and FAB performed laboratory works and analyzed data. HHAMA and ADA performed phylogenetic analysis. ADA, MSA and FAB wrote the first draft of the manuscript. JM-R, DO and ADA reviewed and wrote the final draft of the manuscript. All authors read and approved the final manuscript.

References

1. WHO: Report on the Interregional meeting on leishmaniasis among neighbouring endemic countries in the Eastern Mediterranean, African and Europe, Amman, 23–25 September 2018.
2. Abuzaid AA, Abdoon AM, Aldahan MA, Alzahrani AG, Alhakeem RF, Asiri AM, et al. Cutaneous Leishmaniasis in Saudi Arabia: A Comprehensive Overview. *Vector Borne Zoonotic Dis.* 2017, 17;10:673-684.
3. Sirdar MK, Al-Zahrani MH, Dahlan AA, Sahli AA, Mohamed WS, Hejri YM, et al. Epidemiology and incidence of leishmaniasis in Jazan region, Saudi Arabia (2007-2015): An overview. *J Entomol Zool Stud.* 2018; 6: 859-864.
4. Hawash YA, Ismail KA, Abdel-Wahab MM, Khalifa M. Diagnosis, treatment and clinical features of cutaneous Leishmaniasis in Saudi Arabia. *Korean J Parasitol.* 2018; 56 :229-236.
5. Al-Tawfiq JA, AbuKhamis A: Cutaneous leishmaniasis: a 46-year study of the epidemiology and clinical features in Saudi Arabia (1956–2002). *Int J Infect Dis.* 2004; 8:244-250.
6. Amin TT, Al-Mohammed HI, Kaliyadan F, Mohammed BS: Cutaneous leishmaniasis in Al Hassa, Saudi Arabia: epidemiological trends from 2000 to 2010. *Asian Pac J Trop Biomed.* 2013; 6 :667-672.
7. Alanazi AD, Alyousif MS, Saifi M, Alanazi IO. Epidemiological studies on cutaneous leishmaniasis in Ad-Dawadimi District, Saudi Arabia. *Trop J Pharm Res.* 2016 ; 15: 2709-2712.
8. Alanazi AD, Puschendorf R, Alyousif MS, Al-Khalifa MS, Alharbi S, et al. **Molecular Detection of *Leishmania* spp. in Skin and Blood of Stray Dogs from Endemic Areas of Cutaneous Leishmaniasis in Saudi Arabia.** *Iran J Parasitol.* 2019; 14(2):231-239.
9. Alanazi AD, Rahi AA, Ali AM, Alyousif MS, Alanazi IO, Mahmoud MS, et al. Molecular detection and phylogenetic analysis of *Leishmania major* in stray dogs in Riyadh Province, Saudi Arabia. *Trop Biomed.* 2019; 36(2):315-323.
10. Peters W, Elbihari S, Evans D. Leishmania infecting man and wild animals in Saudi Arabia. 2. *Leishmania arabica* n. sp. *Trans R Soc Trop Med Hyg.* 1986; 80(4):497-502.
11. Elbihari S, Cheema A, El-Hassan A. Leishmania infecting man and wild animals in Saudi Arabia. 4. Canine cutaneous leishmaniasis in the Eastern Province. *Trans R Soc Trop Med Hyg.* 1987; 81(6):925-927.
12. Al-Zahrani M, Peters W, Evans D: Visceral leishmaniasis in man and dogs in south-west Saudi Arabia. *Trans R Soc Trop Med Hyg.* 1988; **82**(6):857.

13. Akhavan AA, Mirhendi H, Khamesipour A, Alimohammadian MH, Rassi Y, Bet al. Leishmania species: detection and identification by nested PCR assay from skin samples of rodent reservoirs. *Exp Parasitol*. 2010;126(4):552-556.
14. Toz SO, Culha G, Zeyrek FY, Ertabaklar H, Alkan MZ, et al. A real-time ITS1-PCR based method in the diagnosis and species identification of *Leishmania* parasite from human and dog clinical samples in Turkey. *PLoS Negl Trop Dis*. 2013; 7(5):e2205.
15. Silva RC, Richini-Pereira VB, Kikuti M, Marson PM, Langoni H. Detection of *Leishmania (L.) infantum* in stray dogs by molecular techniques with sensitive species-specific primers. *Vet Q*. 2017; 37(1):23-30.
16. Ferreira SA, Ituassu LT, Melo MN, Andrade AS. Evaluation of the conjunctival swab for canine visceral leishmaniasis diagnosis by PCR-hybridization in Minas Gerais State. *Brazil Vet Parasitol*. 152; 257–263.
17. El-Beshbishy HA, Al-Ali KH, El-Badry AA. Molecular characterization of cutaneous leishmaniasis in Al-Madinah Al-Munawarah province, western Saudi Arabia. *Int J Infect Dis*. 2013; 17(5):e334-e338.
18. Rasheed Z, Ahmed AA, Salem T, Al-Dhubaibi MS, Al Robaee AA, et al. Prevalence of *Leishmania* species among patients with cutaneous leishmaniasis in Qassim province of Saudi Arabia. *BMC Public Health*. 2019; 19(1):384.
19. General Authority for Statistic. Population Characteristics surveys, Saudi Arabia, pp. 25. https://www.stats.gov.sa/sites/default/files/population_characteristics_surveysar.pdf. 2017.
20. Akilov OE, Khachemoune A, Hasan T. Clinical manifestations and classification of Old World cutaneous leishmaniasis. *Intl J Dermatol*. 2007;46(2):132-142.
21. Noyes HA, Reyburn H, Bailey JW, Smith D. A nested-PCR-based schizodeme method for identifying *Leishmania* kinetoplast minicircle classes directly from clinical samples and its application to the study of the epidemiology of *Leishmania tropica* in Pakistan. *J Clin Microbiol* 1998;36(10):2877-2881.
22. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 2016; 33(7):1870-1874.
23. Al-Bajalan MM, Niranji SS, Al-Jaf SM, Kato H. First identification of *L. major* in a dog in an endemic area of human cutaneous leishmaniasis in Iraq: molecular and phylogenetic studies. *Parasitol Res*. 2018;1-6.
24. Al-Salem WS, Pigott DM, Subramaniam K, Haines LR, Kelly-Hope L, et al. Cutaneous Leishmaniasis and Conflict in Syria. *Emerg Infect Dis* 2016; 22(5):931-933.
25. Baneth G, Nachum-Biala Y, Simon MS, Brenner O, Gaier S, et al. *Leishmania major* infection in a dog with cutaneous manifestations. *Parasites & Vectors*. 2016; 9(1):246.
26. Bamorovat M, Sharifi I, Dabiri S, Mohammadi MA, Harandi MF, et al. *Leishmania tropica* in stray dogs in southeast Iran. *Iran J Public Health*. 2015, 44(10):1359-1366.
27. Al-Zahrani M, Peters W, Evans D, Chin C, Smith V, Lane R. *Phlebotomus sergenti*, a vector of *Leishmania tropica* in Saudi Arabia. *Trans R Soc Trop Med Hyg*. 1988;82(3):416.

28. Haouas N, Amer O, Alshammri FF, Al-Shammari S, Remadi L, et al. Cutaneous leishmaniasis in northwestern Saudi Arabia: identification of sand fly fauna and parasites. *Parasit Vectors*. 2017; 10(1):544.
29. Peters W, Elbihari S, Liu C, Le Blancq S, Evans D, et al. Leishmania infecting man and wild animals in Saudi Arabia 1. General survey. *Trans R Soc Trop Med Hyg*. 1985; 79(6):831-839.
30. Elbihari S, Kawasmeh Z, Al Naiem A. Possible reservoir host (s) of zoonotic cutaneous leishmaniasis in Al-Hassa oasis, Saudi Arabia. *Ann Trop Med Parasit*. 1984; 78(5):543-545.
31. Du R, Hotez PJ, Al-Salem WS, Acosta-Serrano A. Old world cutaneous leishmaniasis and refugee crises in the Middle East and North Africa. *Public Library of Science*; 2016.
32. Medkour H, Laidoudi Y, Athias E, Bouam A, Dizoé S, et al.. Molecular and serological detection of animal and human vector-borne pathogens in the blood of dogs from Côte d'Ivoire. *Comp Immunol Microbiol Infect Dis*. 2020. 69:101412.
33. Mustafa M, Hussein S, Ibrahim E, Al-Seghayer S, Al Amri S, Gradoni L: *Phlebotomus papatasi* (Scopoli), vector of zoonotic cutaneous leishmaniasis in Riyadh province, Saudi Arabia. *Trans R Soc Trop Med Hyg*. 1994; 88(1):40.
34. El-Beshbishy HA, Al-Ali KH, El-Badry AA. Molecular characterization of *Leishmania* infection in sand flies from Al-madinah Al-munawarah province, western Saudi Arabia. *Exp Parasitol*. 2013; 134(2):211-215.
35. Lima C, Colella V, Latrofa MS, Cardoso L, Otranto D, Alho AM. Molecular detection of *Leishmania* spp. in dogs and a cat from Doha, Qatar. *Parasit Vectors*. 2019; 12(1):125.
36. Mohebbali M, Malmasi A, Hajjarian H, Jamshidi S, Akhoundi B, Rezaei M, Janitabar S, Zarei H, Charehdar S: Disseminated leishmaniasis caused by *Leishmania tropica* in a puppy from Karaj, Central Iran. *Iran J Parasitol*. 2011; 6(2):69.
37. Baneth G, Zivotofsky D, Nachum-Biala Y, Yasur-Landau D, Botero A-M. Mucocutaneous *Leishmania tropica* infection in a dog from a human cutaneous leishmaniasis focus. *Parasit Vectors*. 2014; 7(1):118.
38. Baneth G, Yasur-Landau D, Gilad M, Nachum-Biala Y. Canine leishmaniosis caused by *Leishmania major* and *Leishmania tropica*: comparative findings and serology. *Parasit Vectors*. 2017; 10(1):113.
39. Mendoza-Roldan J, Benelli G, Panarese R, Iatta R, Furlanello T, Beugnet F, Zatelli A, Otranto D. *Leishmania infantum* and *Dirofilaria immitis* infections in Italy, 2009-2019: changing distribution patterns. *Parasit Vectors*. 2020;13(1):193.

Figures




Figure 1

Map showing the location of the study areas in Saudi Arabia.




Figure 2

Agarose gel electrophoresis (1.7%) image that show the Nested PCR product analysis of kDNA in Cutaneous *L. major* from human skin lesion samples from Riyadh city. Where M: marker (3000-100bp) and lane (H1 – H5) positive Human *L. major* at (560bp) nPCR product.




Figure 3

Agarose gel electrophoresis (1.7%) image that show the Nested PCR product analysis of kDNA in Cutaneous *L. tropica* from dogs and human skin lesion samples from Al-Qassim province. Where M: marker (3000-100bp), lane (D2 and D4-D7) positive dogs *L. tropica* and lane (H1 and H3) positive Human *L. tropica* at (750bp) nPCR product.




Figure 4

Kinetoplast DNA based phylogenetic analysis of genotypes identified in this study. Phylogenetic tree highlighting the position of *Leishmania* sp. in the present study (Bold) related to other *Leishmania* sp. available in GenBank. The sequences of KDNA were aligned using CLUSTAL W and phylogenetic inferences were constructed in MEGA 7.0 using Maximum Likelihood based on Tamura-Nei Model for nucleotide sequences with 500 bootstrap replicates. There were a total of 587 positions in the final dataset. The scale bar represents a 2% nucleotide sequence discrepancy.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.pdf](#)