

Comparative Role of Compost, Press Mud and Moringa Leaf Extract to Eliminate the Stress and Growth of Maize in Cadmium Polluted Soil

Saqib Bashir

Department of Soil and Environmental Sciences, Ghazi University, Dera Ghazi Khan

Mohamed Soliman Elshikh

Department of botany and microbiology college of science, King Saud University, Riyadh11451,

Mona S Alwahibi

Department of botany and microbiology college of science, King Saud University, Riyadh11451,

Allah Gulshan

Department of Botany, Ghazi University, Dera Ghazi Khan

Javaid Iqbal

Department of Agronomy, Ghazi University, Dera Ghazi Khan

Arif Hussain

Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Punjab

Ali Bakhsh

Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Punjab

Niaz Ahmed

Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Punjab

Muhammad Jamal Khan

School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, Dookie Campus, The University of Melbourne, Victoria 3647,

Shah Fahad

Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresource, College of Tropical Crops, Hainan University, Haikou Hainan 570228

Rahul Datta

Department of Geology and Pedology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemedelska1, 61300 Brno

Subhan Danish (✉ sd96850@gmail.com)

Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan, Punjab

Research Article

Keywords: Maize, Compost, Press mud, Cadmium, wastewater irrigation

Posted Date: January 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-142836/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Cadmium contamination in croplands is recognized as one of the major threats, seriously affecting soil health and sustainable agriculture around the globe. Cd mobility in wastewater irrigated soils can be curtailed through eco-friendly and cost effective organic soil amendments compost (CP), press mud (PM) and moringa leaf extract (ME) at 3% rate that eventually reduces its translocation from polluted soil to plant. This study explored the possible effects of various types of organic soil amendments on cadmium (Cd) phytoavailability in wastewater degraded soil and its subsequent accumulation in maize tissues. Maize plant was grown in Ghazi University as a test plant and Cd accumulation was recorded in its tissues, translocation from root to shoot, chlorophyll contents, plant biomass, yield and soil properties (pH, NPK, OM and Soluble Cd) were also examined. Results revealed that the addition of amendments significantly minimized Cd mobility in soil by 45.8%, 23% and 19.4% when CP, PM and ME were added at 3% over control. Comparing the control soil, Cd uptake effectively reduced via plants shoots by 33.3%, 27.7% and 19.4% when CP, PM and ME. In addition, NPK were significantly increased among all the added treatments in the soil-plant system as well as improved chlorophyll contents relative to non-treated soil. The Current study suggested that among all the amendments, compost at 3% rate performed well and might be considered a suitable approach for maize growth in polluted soil.

Introduction

Cadmium (Cd) is naturally present in earth crust as a trace element, while its concentration varies because of its solubility and mobility¹⁻⁴. Human beings and animals are greatly exposed to heavy metals because of the contaminated food chain⁵. Cd is recognized as a serious global concern and its concentration in industrial and sewage wastewater, and drinking water exceeds the permissible limits (0.003 mg L⁻¹)¹. Wastewater irrigation to the farmlands has become a serious environmental threats because of food chain deterioration⁶. Cd is highly reactive and mobile, which has the potential to influence the plants growth.

The increase of heavy metal accumulation in cultivated soils leads in deteriorating the soil health and enhancing the plant Cd uptake, which affects food safety and quality^{7,8}. Nowadays, various soil restoration strategies (physical, chemical and biological) are adopted in order to improve soil health and minimize food security risks⁹. Additionally, the use of organic materials as a soil amendment could use as a practical and cost effective strategy to restore polluted soils through changing the heavy metals mobility and bioavailability^{10,11}. However, several research studies have shown that organic based soil amendments (compost, biofertilizers, biochar and manure) are more effective to ameliorate the soil health and minimize the use of chemical fertilizer due to their less production cost^{8,12-15}.

Moringa (*Moringa oleifera* L.) that is also known as miraculous tree (evergreen), which is widely grown in tropical areas of Pakistan, having enough tolerance to drought and is easily available at low cost. Its tree contains most important constituents including proteins, carbohydrates, fatty acids, vitamins, flavonoids

and phenolic acids. It has a number of applications in the medical field, for example, anti-inflammatory, antihypertensive activity and analgesic, weight losses, etc. It has attained much attention by the researchers because of its environmental friendly usage. To some extent, it acts as plants growth activator and reduces drought and heavy metals stress on plants¹⁶. Previous study proposed by¹⁷ studied that Moringa seeds were used for wastewater treatment and effectively removed Cd, Cu, Cr and Pb from drinking water. Similarly, its seed ash was also used in the study conducted by¹⁸ for the removal of fluoride from aqueous solutions. Moringa seeds acts as a natural organic biosorbent for the stabilization of heavy metals¹⁹. It can be demonstrated that addition of moringa husks, pods, and seeds showed the noticeable removal of Pb(II) from contaminated water²⁰.

Compost is organic carbon and nutrient rich organic material that can be used as a soil amendment to substitute the chemical fertilizers in agricultural production. The incorporation of compost prominently improves the physical structure and fertility status, microbial communities in degraded soils, improves crop biomass and promote crop growth^{12,21}. In a similar manner, it has the ability to act as biosorbent for the sorption of heavy metals from polluted soils and water³. It has been demonstrated that the compost is a suitable candidate to minimize the risk of crop growth, yield and economic losses through ameliorating the soil properties and thereby, ensure the human health^{11,22}. Abbas et al.¹² found that incorporation of compost effectively reduced the toxic elements from polluted water by 85–89% through chemical absorption.

Press mud is the by-product of sugarcane obtained from the sugar industry after the clarification process of sugarcane juice. Press mud is a dark brown, spongy and amorphous material containing sugar, fiber and coagulated colloids including soil particles, and mainly comprises of CaO, organic matter, nitrogen, phosphorus, potassium, calcium, sulphur in varying amounts²¹.¹⁶ demonstrated that addition of soil amendments such as press mud, poultry manure farmyard manure (FYM) and fish manure to Cr polluted soil. Results revealed that these amendments have potential to stabilize Cr in soil. It can be demonstrated that press mud had been recognized as the filter cake which can contribute to enhance the soil fertility status and can create extra binding sites for the toxic metals adsorption and complexation²³. Press mud has the ability to buffer the soil pH and reduces the uptake of heavy metals. It contains functional groups like hydroxyl ions that play an important role in the adsorption of heavy metals^{24,25}, and is also considered a key source of NPK and carbon contents. Press mud is considered a key source of NPK and carbon contents²³. Sabir et al.²⁶ confirmed that press mud incorporation as a soil amendment could help to improve soil health, structure and thereby, reduces the metals accumulation in plant tissues.

Maize (*Zea mays* L.) belongs to Gramineae family and is recognized as one of the most important nutritious cereal crop that is also consumed by animals as a forage¹⁰. It is widely grown all around the globe and has greater potential to uptake heavy metals. Particularly, this crop being the highest yielding cereal crop in the world has a significant importance in Pakistan. There is only little scientific knowledge is present about the effect of ME as a biofertilizer, compost and press mud on the growth, metabolic and antioxidant enzymes of plants. Thus, the comparative efficiency of compost, press mud and *M. oleifera*

leaf extract was assessed for wastewater irrigated soil restoration as well as reducing the Cd stress on maize plant in calcareous soil condition. The prime objectives of the current study were to evaluate the effect of compost, press mud and *M. oleifera* leaf extract on Cd solubility and maize growth in polluted soil. Additionally, to examine the efficacy of these soil amendments on soil and plant nutrients status.

Materials And Methods

Soil and amendments collection

Soil samples were taken from productive land in the suburbs of Dera Ghazi Khan city, Pakistan that was irrigated with the wastewater of manka canal. The polluted soil samples were gathered from the upper layer at 0–30 cm depth in 2019. The obtained soil was then shifted to green house for air drying about three days. After drying, soil samples were ground, stones and pebbles were removed and then the soil passed through a 2-mm sieve for pot study. Approximately 0.5 kg of the prepared soil was further ground and passed through 0.25-mm sieve for the basic physico-chemical characteristics of the studied soil (Table 1). In our previous study Bashir et al. ⁷ explained that the studied soil of this area has more than 5 mg kg⁻¹ Cd which is greater than the proposed value of Cd (0.2 mg kg⁻¹) in Agricultural soils by Chinese environmental and quality standard for soils (GB15618-1995). Similarly, press mud was purchased from Fatima sugar mill for the current study. While the vegetables waste product derived compost was prepared in Ghazi university and used in current study (Table 1).

Additionally, Moringa leaf extract was prepared according to the method proposed by Mónica et al. ²⁷. For this purpose, moringa (*Moringa oleifera* L.) tree leaves were gathered from the local area of Dera Ghazi Khan, Pakistan and rinsed with tap water to remove dust particles from leaves and dried under shade. After drying plant samples were preserved in freezer, then plant leaves were ground and the collected was filtered through Whatman filter paper. After filtration, the extract was stored at 4 °C for further chemical analysis. There are the followings selected properties were measured: moisture (82 mg); Protein (5.82 mg); Vitamin C (906 mg) and others were presented in (Table 1).

Pot Experiment

The maize growth experiment was performed to examine the influence of various kinds of organic soil amendments (compost, press mud and moringa leaf extract) at 3% application rates on maize growth and Cd accumulation in its tissues. The pot study was organized with the following treatments (1) Control (CK); (2) Compost (CP); (3) Press mud (PM); (4) Moringa leaf extract (ME) and each treatment had three replicates. Each pot (16 cm diameter and 18 cm height) had 4 kg soil and homogeneously mixed with their respective treatments. All the experimental units were irrigated with distilled water and kept their water contents at 70% water holding capacity and left them for 2 weeks for equilibrium. After 15 days, the sterilized Pioneer-1543 seeds were placed in each experimental unit. Each pot was fertilized with the basal dose of N (180 kg ha⁻¹ soil), P (120 kg ha⁻¹ soil) and K (90 kg ha⁻¹ soil) in the form of urea, Ca(H₂PO₄)₂, and KCl respectively. After seed germination, 3 seedlings in each pot were maintained. After

4 months, the mature maize plants were harvested and the further plants and soil analysis were performed.

Soil Analysis

After the crop harvest, soil samples were taken from each unit and measured soil pH and EC using pH and EC meter according to the described method in our previous study ⁸. Bioavailable Cd contents from each potted soil were estimated using the CaCl_2 extractable technique ^{6,7}. Soil organic matter contents were calculated by using the wet oxidation method ²⁸. Similarly, soil available nitrogen, phosphorus and potassium contents were estimated as described by Bashir et al. ⁸

Plant Analysis

Chlorophyll contents were measured using the SPAD meter. However, the harvested portion of maize shoots and roots were oven dried and digested with di-acid $\text{HNO}_3:\text{HClO}_4$ mixture for the estimation of Cd contents and nutrients (NPK) as described in our previous study ⁶⁻⁸.

Statistical analysis

One-way analysis of variance (ANOVA) followed by the LSD test were postulated at ($P < 0.05$) to analyze the data using Statistic 8.1 (Analytical software, USA). In addition, the means and standard deviations (SD) clearly indicated the statistically significant differences among all the treatments for each examined parameters ²⁹.

Results

Effect of amendments on soil pH and Cd bioavailability

The incorporation of various kinds of soil amendments showed the significant alteration in soil pH and soil bioavailable Cd, relative to without soil amendments. The greater increase in soil pH was observed in press mud by 0.4 units relative to control (Fig. 1). However, the addition of compost at 3% rate showed the prominent decline in soil pH by 0.2 units, relative to control soil.

While, the mixing of soil amendments in sewage wastewater irrigated polluted soil showed the significant decline in the soil bioavailable Cd contents against without treated soil (Fig. 2). The maximum reduction was recorded in compost by 49%. Compared to the control soil, the addition of PM and ME at 3% rate also showed the prominent reduction by 28% and 25.3% respectively.

Effect of amendments on soil organic matter and Nutrients availability

The addition of soil amendments in polluted soil led to enhance soil organic carbon contents, over without treated soil. The significant improvement in soil carbon was recorded by 42.7%, 20.2% and 4.5%, when compost, press mud and moringa leaf extract were incorporated at 3% rate relative to non-amended soil. Similarly, the bioavailable status of NPK nutrients efficiently improved in the treated soil relative to non-amended soil (control). The prominent increase in available-N was recorded by 26.7 mg kg⁻¹ when compost was mixed with polluted soil at 3% rate. Whereas, addition of press mud and moringa leaf extract at 3% dose level also showed the prominent increment in soil available-N by 24.2 and 23.2 mg kg⁻¹ relative to non-amended soil (21.8 mg kg⁻¹) (control), respectively. Likewise, addition of compost, press mud and moringa leaf extract into wastewater polluted soil efficiently improved soil available-P by 9.5, 6.7 and 6.6 mg kg⁻¹ respectively, over control soil (5.7 mg kg⁻¹) Available-K contents in wastewater irrigated polluted soils was prominently improved when compost, press mud and moringa leaf extract at 3% rate by 90.7, 81.7 and 73.3 mg kg⁻¹ respectively, over control (72.3 mg kg⁻¹).

Effect Of Amendments On Cd Uptake And Chlorophyll Contents

Results showed that the addition of soil amendments in waste water irrigated soil improved the soil health and plant growth. The chlorophyll contents prominently improved in treated soil relative to control soil (Fig. 3). The maximum increment in chlorophyll contents was recorded in CP, PM and ME amended soils by 27.7%, 16.3% and 13.3% at 3% application rate, respectively over control. The results revealed that Cd accumulation in maize shoots (Fig. 4) and roots (Fig. 5) tissues was increased in wastewater polluted soil relative to soil treated with organic amendments. However, addition of various kind of amendments has ability to reduce Cd accumulation in plant tissues. The greater reduction in Cd accumulation in plant shoots was recorded by 50%, 36.8% and 23.7% when compost (CP), press mud (PM) and moringa leaf extract (ME) were added at 3% rate into wastewater polluted soil respectively, over control. Likewise, incorporation of CP, PM and ME also significantly reduced the Cd accumulation in maize roots. The highest reduction was recorded by 39%, 25% and 17.2% after CP, PM and ME were incorporated at 3% application level. Our current study results revealed that the addition of soil amendments showed a significant difference in Cd translocation from root to shoot. The maximum reduction was recorded in the compost treated soil by 30.3%, while the addition of press mud and moringa leaf extract showed the reduction by 22.7% and 16.6% respectively at 3% application rate, relative to control.

Effect Of Amendments On Maize Biomass And Nutrients Accumulation

Maize fresh and dry shoot root biomass was also improved after the addition of compost (CP), press mud (PM) and moringa leaf extract (ME) at 3% dose level. The maximum fresh shoot and root biomass

was recorded when soil was amended with soil amendments relative to control (Table 2). While, the maximum dry biomass was recorded in CP amended soil. Likewise, all other growth and yield parameters were improved among all the incorporated treatments (Table 2). Similarly, nutrients contents in maize shoots were improved when CP, PM and ME were added at 3% application rate. Among all the soil amendments, CP showed the greater NPK contents in plant shoot tissues relative to other treatments indeed control (Table 3).

Discussion

Restoration of wastewater polluted soil and wastewater treatment is an emerging challenge for all around the globe and scientist to protect food chain. Recently, use of various organic amendments, especially compost (CP), press mud (PM) and moringa leaf extract (ME) is suitable approach for polluted soil restoration. The current investigation confirmed that the incorporation of various organic materials has exhibited their potential to alleviate Cd mobility and phytoavailability to maize in wastewater irrigated farmlands³⁰. As described in Table 1, all materials have sufficient amount of C, H, N and O. The greater amount of carbon contents and C:N in compost and press mud attributed the presence of highest amount of lignin and cellulose. Incorporation of organic amendments has the ability to improve soil organic carbon and nutrients status which ultimately promote crop growth. The addition of compost significantly ameliorates the soil alkalinity and reduced soil pH because of sufficient amount of humic acid in compost, which is helpful for the mobility of soil nutrients in calcareous soil. Similar findings were reported in the recent study described by²² indicated that the addition of compost could reduce the soil pH due to the humification nature of studied compost. Zhou et al.³¹ described that the vegetable derived compost significantly reduced the soil pH, which might be due to the abundance of humic acid and organic compounds in the compost. Similar findings were reported by³² confirmed that organic soil amendments especially compost could release humic substances which could contribute to lessen soil pH.

The recent studies reported by^{33,34} suggested that the incorporation of soil organic amendments (biochar, compost, and biogas slurry) prominently influenced the soil pH, EC and CEC as well as provided the essential nutrients (C, N, P, K, and S) to plant. This enhancement in soil fertility leads to an improvement in plant growth through the uptake of essential elements provided by the organic soils amendments¹². However, the use of press mud significantly enhances the soil pH relative to without treated soil (control)³⁵. Present study confirmed that the significant increase in soil pH and essential soil nutrients (NPK) might be due to the presence of base cations and CaO in press mud. Due to the alkalinity of press mud, OH⁻ ions released in soil solution and prominently increase soil pH, which could create the bindings sites for heavy metals complexation and adsorption³⁶. It can be demonstrated that the increase in soil pH could enhance the sorption cavities and bindings sites on soil collides and as well as increased the net electronegative charges on soil constituents^{6,7}, which are thus responsible for the Cd immobilization through adsorption and precipitation⁸.

The findings of current study confirmed that the addition of organic soil amendments (compost, press mud and moringa leaf extract) effectively minimized Cd mobility in polluted soil, which might be possible due to the presence of organic substance and functional groups that have great contribution to stabilize Cd through complexation. In the similar manner⁹ examined that municipal solid waste compost has the potential to remediate Cu polluted soil by reducing the exchangeable portion of Cu. Addition of compost in degraded soil significantly improve soil health by reducing the metals mobility because of minerals, organic acids and microbes that have much contribution to promote the metals stabilization^{37,38}. Anastopoulos et al.³⁹ and Soares et al.⁴⁰ found that compost has a potential role as a biosorbent for adsorbing heavy metals. It can be recognized that the prominent decline in Cd mobility might be ensured because of all the applied soil amendments properties such as physical (porous structure) and chemical (high pH, greater CEC and surface functional groups) properties could cause the greatest reduction in the solubility of Cd through complexation and adsorption on soil surface. Another possibility was that the addition of press mud would be able to release the greater amount of carbonates, oxides and hydroxides into the soil solution, which increase soil pH that would be able to enhance Cd precipitation as $\text{Cd}(\text{OH})_2$ and CdCO_3 .^{7,8} This could influence the reduction of Cd contents in maize tissues. It can be attributed that press mud contains functional groups like hydroxyl ions that play an important role in adsorption of heavy metals^{24,25}. Press mud reverse the toxic effect of heavy metals. Sabir et al.²⁶ demonstrated that the addition of press mud significantly decreased the Cr (VI) accumulation in root and shoot by 33% and 20%, respectively relative to untreated soil.

Present study confirmed that the addition of green waste derived organic compost has the ability to overcome the Cd solubility and mobility issue in polluted soil due to its organic nature. Additionally, compost incorporation improved the contents of soil carbon by 6–7 times relative to the untreated soil. It has sufficient quantity of humic substances, which held various kinds of functional groups e.g., carboxyl and hydroxyls. Some of these functional groups could contribute to bind the heavy metals with their organic complexes and chelates⁴¹. Previous study explained⁴² that the compost addition could transform soluble portion of Cd by 47.8–69.8% to the most stable form, and therefore, reduced Cd accumulation in Pakchoi tissue by 56.2–62.5% relative to un-amended soil. The mitigation of Cd solubility by compost was recognized primarily to complexation of Cd by organic substance that was present in compost⁴². Karami et al.⁴³ confirmed the effectiveness of green waste induced compost on the phytoavailability of metals such as Cu and lead Pb in plants tissues. Moreover, the incorporation of compost expressively reduced the Cu and Pb toxicity by hindering their uptake in plant tissues.

Our results revealed that moringa leaf extract prominently reduced Cd mobility and enhanced maize growth. The reduction in Cd accumulation in maize plants is might be due to the hormonal behavior of moringa that lead to mitigate the toxicity of Cd in plants parts. Similarly, the use of moringa parts such as seeds, leaf and bark have the potential to hinder the heavy metals mobility and retain them in their surface through surface functional groups present on moringa parts^{44,45}. Current findings demonstrated that ME has ability to alter bioavailable pool of Cd to most stable residual form, which could contribute to reduce the stress on maize plant in polluted soil. Addition of *Moringa oleifera* parts including seeds,

leaves and fruits are useful for polluted water purification and improvement of water quality that make it possible for the drinking purpose^{17,46}. The modification of the moringa as a biosorbent was used to enhance the adsorption of Ni from aqueous solution. It can be demonstrated that *M. oleifera* seed were used as a good approach for the removal of Co, Cu, Pb and Cd⁴⁴. In addition, they studied that Cd has greater ability to adsorb on the moringa bark and leaf as well as enhance crop growth.

After the treatment of polluted soil with organic amendments the maize biomass significantly improved compared to control. This improvement might be attributed to the nutritious significance of compost, press mud and moringa leaf extract that improves the soil productivity. Other possible reason might be considered because of their potential to boost the organic matter mineralization and enhanced crop yield and growth^{12,30}. It is important to discuss that compost and press mud acts as buffer and contains sufficient quantity of vital plant nutrients which pointedly rise crop yield³⁰. The establishment of these results might occurred due to the obvious reduction in the phytoavailable Cd in treated soil, thus alleviating Cd uptake by maize. From the findings of current study, it has been established that use of organic materials as a soil amendment has capability to ameliorate soil alkalinity. In addition, they could release basic nutrients and exchange metals from soil solution, which might be able to tackle metals mobility in polluted soils and improved crop growth^{12,30,44}.

Conclusion

This study established the efficiency of compost (CP), press mud (PM) and moringa leaf extract (ME) as a soil amendment to sorb Cd and its translocation from soil to plant tissues and nutrients status that was investigated in pot study. The findings revealed that CP, PM and ME addition at 3% rate significantly altered soil pH, nutrients availability in soil-plant system, reduced the mobility and uptake of Cd in maize plant as well as minimize its toxicological and physiological stress in polluted soil relative to control. Hence, it has been established that the use of organic waste induced materials to remediate the Cd-polluted soil is considered an emerging approach to restore the polluted soil health and improve the better crop growth. Additionally, the findings of current study need to be further confirmed on large scale field experiments under the presence of multiple potential toxic elements. Organic materials production from different feedstock's could be systematically arranged on their comprehensive soil restoration performance in the following order CP > PM > ME.

Declarations

Author Contributions: Conceptualization, S.B.; M.S.E.; M.S.A.; methodology, A.B.G.; software, J.I.; A.H.; validation, A.B.; N.A.; formal analysis, S.B.; investigation, S.B.; resources, J.I.; A.H.; data curation, J.I.; A.H.; writing—original draft preparation, A.B.; N.A.; S.B.; S.D.; S.F.; writing—review and editing, M.J.K.; R.D.; S.D.; S.F.; M.S.E.; M.S.A.; funding acquisition, S.D.; M.S.E.; M.S.A.;

Acknowledgment

The authors extend their appreciation to the researchers supporting project number (RSP-2020/173) King Saud university, Riyadh, Saudi Arabia

Funding: The authors extend their appreciation to the researchers supporting project number (RSP-2020/173) King Saud university, Riyadh, Saudi Arabia

Institutional Review Board Statement: Not applicable

Informed Consent Statement: Not applicable

Data Availability Statement: Not applicable

Acknowledgments: Not applicable

Conflicts of Interest: The authors declare no conflict of interest

References

1. Shahid, M. *et al.* Arsenic Level and Risk Assessment of Groundwater in Vehari, Punjab Province, Pakistan. *Expo. Heal.* **10**, 229–239 (2018).
2. Younis, U., Shah, M. H. R., Danish, S., Malik, S. A. & Ameer, A. Biochar role in improving biometric and growth attributes of *S. oleracea* and *T. corniculata* under cadmium stress. *Int. J. Biosci.* **5**, 84–90 (2014).
3. Abid, M. *et al.* Biochar increased photosynthetic and accessory pigments in tomato (*Solanum lycopersicum* L.) plants by reducing cadmium concentration under various irrigation waters. *Environ. Sci. Pollut. Res.* **24**, 22111–22118 (2017).
4. Zafar-Ul-Hye, M., Shahjahan, A., Danish, S., Abid, M. & Qayyum, M. F. Mitigation of cadmium toxicity induced stress in wheat by ACC-deaminase containing PGPR isolated from cadmium polluted wheat rhizosphere. *Pakistan J. Bot.* **50**, 1727–1734 (2018).
5. Xiong, T. T. *et al.* Measurement of metal bioaccessibility in vegetables to improve human exposure assessments: field study of soil–plant–atmosphere transfers in urban areas, South China. *Environ. Geochem. Health.* **38**, 1283–1301 (2016).
6. Bashir, S. *et al.* Role of sepiolite for cadmium (Cd) polluted soil restoration and spinach growth in wastewater irrigated agricultural soil. *J. Environ. Manage.* **258**, 110020 (2020).
7. BASHIR, S. *et al.* Efficiency of KOH-modified rice straw-derived biochar for reducing cadmium mobility, bioaccessibility and bioavailability risk index in red soil. *Pedosphere.* **30**, 874–882 (2020).
8. Bashir, S. *et al.* Comparative efficiency of wheat straw and sugarcane bagasse biochar reduces the cadmium bioavailability to spinach and enhances the microbial activity in contaminated soil. *Int. J. Phytoremediation.* **21**, 1098–1103 (2019).
9. Liu, L., Wang, S., Guo, X. & Wang, H. Comparison of the effects of different maturity composts on soil nutrient, plant growth and heavy metal mobility in the contaminated soil. *J. Environ. Manage.* **250**,

- 109525 (2019).
10. Bashir, M. A. *et al.* Combined application of biochar and sulfur regulated growth, physiological, antioxidant responses and Cr removal capacity of maize (*Zea mays* L.) in tannery polluted soils. *J. Environ. Manage.* **259**, 110051 (2020).
 11. Bolan, N. *et al.* Remediation of heavy metal(loid)s contaminated soils - To mobilize or to immobilize?. *Journal of Hazardous Materials.* vol. **266**, 141–166 (2014).
 12. Abbas, A. *et al.* Synergistic use of biochar and acidified manure for improving growth of maize in chromium contaminated soil. *Int. J. Phytoremediation.* **22**, 52–61 (2020).
 13. Fiaz, K. *et al.* Drought impact on Pb/Cd toxicity remediated by biochar in *Brassica campestris*. *J. soil Sci. plant Nutr.* **14**, 845–854 (2014).
 14. Younis, U. *et al.* Growth, survival, and heavy metal (Cd and Ni) uptake of spinach (*Spinacia oleracea*) and fenugreek (*Trigonella corniculata*) in a biochar-amended sewage-irrigated contaminated soil. *J. Plant Nutr. Soil Sci.* **178**, 209–217 (2015).
 15. Danish, S., Younis, U., Nasreen, S., Akhtar, N. & Iqbal, M. T. Biochar consequences on cations and anions of sandy soil. *J. Biodivers. Environ. Sci.* **6**, 121–131 (2015).
 16. Nouman, W., Siddiqui, M. T. & Basra, S. M. A. *Moringa oleifera* leaf extract: An innovative priming tool for rangeland grasses. *Turkish J. Agric. For.* **36**, 65–75 (2012).
 17. Reddy, D. H. K., Seshiah, K., Reddy, A. V. R. & Lee, S. M. Optimization of Cd(II), Cu(II) and Ni(II) biosorption by chemically modified *Moringa oleifera* leaves powder. *Carbohydr. Polym.* **88**, 1077–1086 (2012).
 18. Dobaradaran, S. *et al.* Fluoride removal from aqueous solutions using *moringa oleifera* seed ash as an environmental friendly and cheap biosorbent. *Fresenius Environ. Bull.* **24**, 1269–1274 (2015).
 19. Araujo, C. S. T. *et al.* Bioremediation of Waters Contaminated with Heavy Metals Using *Moringa oleifera* Seeds as Biosorbent. in *Applied Bioremediation - Active and Passive Approaches* (eds. Patil, Y. & Rao, P.) 225–253 (Croacia: InTech Open Science, 2013). doi:10.5772/56157.
 20. Tavares, F. O. *et al.* Environmentally friendly biosorbents (husks, pods and seeds) from *Moringa oleifera* for Pb(II) removal from contaminated water. *Environ. Technol. (United Kingdom).* **38**, 3145–3155 (2017).
 21. Calleja-Cervantes, M. E. *et al.* Thirteen years of continued application of composted organic wastes in a vineyard modify soil quality characteristics. *Soil Biol. Biochem.* **90**, 241–254 (2015).
 22. Li, M. *et al.* Population characteristics and influential factors of nitrogen cycling functional genes in heavy metal contaminated soil remediated by biochar and compost. *Sci. Total Environ.* **651**, 2166–2174 (2019).
 23. Khan, S., Qureshi, I., Shifa, M. S. & Waziri, A. H. An efficient calixarene-based functional material for the sorption and recovery of Cr(VI) from water. *Int. J. Environ. Anal. Chem.* **99**, 1123–1134 (2019).
 24. Lee, T. M., Lai, H. Y. & Chen, Z. S. Effect of chemical amendments on the concentration of cadmium and lead in long-term contaminated soils. *Chemosphere.* **57**, 1459–1471 (2004).

25. Mahmood, T. Phytoextraction of heavy metals - the process and scope for remediation of contaminated soils. *Soil and Environment*. **vol. 29**, 91–109 (2010).
26. Sabir, M. *et al.* Comparative effect of activated carbon, pressmud and poultry manure on immobilization and concentration of metals in maize (*Zea mays*) grown on contaminated soil. *Int. J. Agric. Biol.* **15**, 559–564 (2013).
27. Valdez-Solana, M. A. *et al.* Nutritional content and elemental and phytochemical analyses of moringa oleifera grown in Mexico. *J. Chem.* 2015, <https://doi.org/10.1155/2015/860381>(2015) (2015).
28. Walkley, A. An Examination of Methods for Determining Organic Carbon and Nitrogen in Soils. *J. Agric. Sci.* **25**, 598 (1935).
29. Steel, R. G., Torrie, J. H. & Dickey, D. A. Principles and Procedures of Statistics: A Biometrical Approach(McGraw Hill Book International Co., 1997).
30. Ali, S. *et al.* Comparative evaluation of wheat straw and press mud biochars for Cr(VI) elimination from contaminated aqueous solution. *Environ. Technol. Innov.* **19**, 101017 (2020).
31. Zhou, R. *et al.* Remediation of Cu, Pb, Zn and Cd-contaminated agricultural soil using a combined red mud and compost amendment. *Int. Biodeterior. Biodegrad.* **118**, 73–81 (2017).
32. Tang, J. *et al.* Physicochemical features, metal availability and enzyme activity in heavy metal-polluted soil remediated by biochar and compost. *Sci. Total Environ.* **701**, 134751 (2020).
33. Yuan, Y. *et al.* Is biochar-manure co-compost a better solution for soil health improvement and N₂O emissions mitigation?. *Soil Biol. Biochem.* **113**, 14–25 (2017).
34. Kizito, S. *et al.* Role of nutrient-enriched biochar as a soil amendment during maize growth: Exploring practical alternatives to recycle agricultural residuals and to reduce chemical fertilizer demand. *Sustain.* **11**, 3211 (2019).
35. Borchard, N., Spokas, K., Prost, K. & Siemens, J. Greenhouse Gas Production in Mixtures of Soil with Composted and Noncomposted Biochars Is Governed by Char-Associated Organic Compounds. *J. Environ. Qual.* **43**, 971–979 (2014).
36. Karwal, M. & Kaushik, A. Co-composting and vermicomposting of coal fly-ash with press mud: Changes in nutrients, micro-nutrients and enzyme activities. *Environ. Technol. Innov.* **18**, 100708 (2020).
37. Białobrzewski, I. *et al.* Model of the sewage sludge-straw composting process integrating different heat generation capacities of mesophilic and thermophilic microorganisms. *Waste Manag.* **43**, 72–83 (2015).
38. Awasthi, M. K., Pandey, A. K., Bundela, P. S. & Khan, J. Co-composting of organic fraction of municipal solid waste mixed with different bulking waste: Characterization of physicochemical parameters and microbial enzymatic dynamic. *Bioresour. Technol.* **182**, 200–207 (2015).
39. Anastopoulos, I., Massas, I. & Ehaliotis, C. Composting improves biosorption of Pb²⁺ and Ni²⁺ by renewable lignocellulosic materials. Characteristics and mechanisms involved. *Chem. Eng. J.* **231**, 245–254 (2013).

40. Soares, M. A. R., Marto, S., Quina, M. J., Gando-Ferreira, L. & Quinta-Ferreira, R. Evaluation of Eggshell-Rich Compost as Biosorbent for Removal of Pb(II) from Aqueous Solutions. *Water. Air. Soil Pollut.* **227**, 150 (2016).
41. Chang Chien, S. W., Wang, M. C. & Huang, C. C. Reactions of compost-derived humic substances with lead, copper, cadmium, and zinc. *Chemosphere.* **64**, 1353–1361 (2006).
42. CHEN, H. S. *et al.* Poultry Manure Compost Alleviates the Phytotoxicity of Soil Cadmium: Influence on Growth of Pakchoi (*Brassica chinensis* L.). *Pedosphere.* **20**, 63–70 (2010).
43. Karami, N., Clemente, R., Moreno-Jiménez, E., Lepp, N. W. & Beesley, L. Efficiency of green waste compost and biochar soil amendments for reducing lead and copper mobility and uptake to ryegrass. *J. Hazard. Mater.* **191**, 41–48 (2011).
44. Maina, I. W., Obuseng, V. & Nareetsile, F. Use of *Moringa oleifera* (Moringa) Seed Pods and *Sclerocarya birrea* (Morula) Nut Shells for Removal of Heavy Metals from Wastewater and Borehole Water. *J. Chem.* 2016, [\(https://doi.org/10.1155/2016/9312952\(2016\)\)](https://doi.org/10.1155/2016/9312952(2016)) (2016).
45. George, K. S. *et al.* A Study on the Potential of Moringa Leaf and Bark Extract in Bioremediation of Heavy Metals from Water Collected from Various Lakes in Bangalore. *Procedia Environ. Sci.* **35**, 869–880 (2016).
46. Reddy, D. H. K., Ramana, D. K. V., Sessaiah, K. & Reddy, A. V. R. Biosorption of Ni(II) from aqueous phase by *Moringa oleifera* bark, a low cost biosorbent. *Desalination.* **268**, 150–157 (2011).

Tables

Table 1: Selected properties of soil, press mud (PM); Compost (CP) and Moringa leaf extract (ML)

Properties	Soil	Press mud	Compost	Properties	Moringa Leaf extract
pH	7.8	8.4	7.3	pH	6.8
EC	0.6	2.7	2.4	EC	0.2
OM (g kg ⁻¹)	0.08	0.3	14	Ca (mg/100g)	5.32
Texture	silty clay			Mg (mg/100g)	0.75
Total N (g kg ⁻¹)	0.2	3.6	9.4	P (mg/100g)	0.84
Avail. P (g kg ⁻¹)	0.005	0.74	3.2	Na (mg/100g)	0.21
Extractable K (g kg ⁻¹)	0.07	1.8	11.5	K (mg/100g)	0.32
				Fe (mg/100g)	0.03

Table 2: Change in plant growth parameters after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (ME) at 3% application level Error bars are the SD of the means (n=3) and

different letters indicate that values are significantly different $p < 0.05$. Plant height (PH); Shoot length (SL); Root length (RL); Fresh shoot biomass (FS); Dry shoot biomass (DS); Fresh root biomass (FR); Dry root (DR); Number of leaves (NL).

Treatment	PH	SL	RL	FS	DS	FR	DR	NL
CK	55.3c	45.3c	8.5c	15.4c	6.5c	5.5c	3.1c	14.0c
PM	60.7bc	50.0bc	9.5b	17.8b	8.4b	7.3b	4.1b	16.7b
CP	75.3a	63.0a	12.5a	22.3a	10.3bc	9.5a	4.8a	19.0a
ME	64.3b	52.7b	9.1b	18.4b	7.0a	7.9b	3.5bc	15.3bc

Table 3: Change in soil and plant nutrients after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (ML) at 3% application level. Different letters indicate that values are significantly different $p < 0.05$.

Treatment	Soil (mg kg^{-1})			Plant (g kg^{-1})			
	N	P	K	N	P	K	OM (g kg^{-1})
CK	21.8c	5.0c	72.3c	11.8c	1.5c	12.7c	0.63c
PM	24.2b	7.0b	81.7b	13.9ab	2.1b	15.5b	0.79b
CP	26.7a	9.5a	90.7a	14.7a	3.9a	17.7a	1.10a
ME	23.2bc	6.0bc	73.3c	13.4b	1.9b	14.8b	0.66c

Figures




Figure 1

Change in soil pH after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (ME) at 3% application level Error bars are the SD of the means (n=3) and different letters indicate that values are significantly different $p < 0.05$.




Figure 2

Change in soil Cd bioavailability after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (MLE) at 3% application level Error bars are the SD of the means (n=3) and different letters indicate that values are significantly different p <0.05.




Figure 3

Change in Chlorophyll contents after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (MLE) at 3% application level Error bars are the SD of the means (n=3) and different letters indicate that values are significantly different p <0.05.




Figure 4

Change in Cd contents in maize shoot after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (MLE) at 3% application level Error bars are the SD of the means (n=3) and different letters indicate that values are significantly different $p < 0.05$.




Figure 5

Change in Cd contents in maize root after treated with control (CK), Press mud (PM), Compost (CP) and Moringa Leaf Extract (MLE) at 3% application level Error bars are the SD of the means (n=3) and different letters indicate that values are significantly different $p < 0.05$.