

Comparison of three different prophylactic treatments for postoperative nausea and vomiting after total joint arthroplasty under general anesthesia: a randomized clinical trial

Jinwei Xie

Sichuan University West China School of Public Health

Yingcun Cai

Sichuan University West China Hospital

Jun Ma

Sichuan University West China Hospital

Qiang Huang

Sichuan University West China Hospital

Zongke Zhou

Sichuan University West China Hospital

Fuxing Pei (✉ peifuxinghuaxi@126.com)

Sichuan University West China Hospital <https://orcid.org/0000-0002-5284-178X>

Research article

Keywords: Postoperative nausea and vomiting; Arthroplasty; Enhanced recovery after surgery

Posted Date: May 4th, 2020

DOI: <https://doi.org/10.21203/rs.2.24233/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Postoperative nausea and vomiting (PONV) after total joint arthroplasty is common and associated with delayed recovery. This study was performed to evaluate the efficacy of three different prophylactic regimens for PONV after total joint arthroplasty under general anesthesia.

Methods: Patients undergoing primary total hip or knee arthroplasty were randomized to Group A (8 mg ondansetron, n= 108), Group B (10 mg dexamethasone plus mosapride, n= 108), or Group C (three doses of 10 mg dexamethasone plus mosapride, n=116). The primary outcome was the total incidence of PONV during postoperative 48 hours. The secondary outcomes were complete response, rescue antiemetic treatment, opioid consumption, time until first defecation, postoperative appetite score, satisfaction score, length of hospital stay, blood glucose level, and complications.

Results: Patients in Group C experienced a lower incidence of total PONV (29.3%, p= 0.001), severe nausea (1.72%, p<0.001) and a higher incidence of complete response (70.69%, p=0.001) than did patients in Groups A and B. Moreover, less rescue antiemetic treatment and postoperative opioid consumption was needed in Group C (p<0.05). Additionally, a shorter time until first defecation, shorter length of stay, and better postoperative appetite scores and satisfaction scores were detected in patients in Group C. A slight increase in the fasting blood glucose level was observed in Group C, and the complications were comparable among the groups.

Conclusion: Prophylactic use of mosapride and three doses of dexamethasone can provide better antiemetic effectiveness, postoperative appetite, bowel function recovery, and pain relief than a single dose or ondansetron only.

Background

Postoperative nausea and vomiting (PONV) are one of the most common and distressing complications after surgery. This is especially true in patients undergoing general anesthesia, among whom the incidence of PONV may reach 25%- 30% and even rise to 80% among patients without prophylactic intervention(1, 2). Optimal management of PONV is important for rapid recovery after joint arthroplasty because effective treatments that limit post-operative nausea allow patients to mobilize earlier, take food earlier, and improve overall patient satisfaction(3).

Numerous precautionary measures have been taken in an attempt to prevent PONV, such as administration of 5HT₃ receptor antagonists, neurokinin 1 receptor antagonists, corticosteroids, butyrophenones, and antihistamines(4). Although the efficacy of these prophylactic drugs has been confirmed by high-level evidence, PONV remains a persistent problem(5, 6). One reason for this phenomenon is the huge gap between implementation and the “PONV-free” goal. Another reason is the side effects associated with anti-PONV treatments, and no ideal regimen has been established. For example, use of the 5-HT₃ receptor antagonist ondansetron can aggravate the postoperative constipation which occurs in up to 57% of patients who undergo total joint arthroplasty(7). Dexamethasone, another

common antiemetic agent, has effects on reducing the incidence of PONV in total joint surgery (8-10). Meantime, concerns about infection safety has aroused because of its suppressive effect on immune system and increase in blood glucose level, although the current evidence doesn't support this(11). Furthermore, the most appropriate dosage and timing of dexamethasone have not been determined. A single dose was routine practice against PONV, and extended prevention may provide additional analgesic effect.

The previous study showed that intravenous use of opioids or nonsteroidal anti-inflammatory drugs were risk factors for PONV following primary total hip or knee arthroplasty (odds ratio=2.052, $p=0.008$; published in Chinese) (12) because these drugs inhibit gastrointestinal peristalsis. In contrast, selective 5-HT₄ agonists can stimulate the gastrointestinal tract and promote motility(13), and mosapride, one type of 5-HT₄ agonist, is reportedly effective in reducing the incidence of vomiting caused by chemotherapy(14). To the best of our knowledge, however, very few studies have evaluated and compared the antiemetic efficacy of mosapride with that of other types of PONV prophylaxis.

Thus, in the present study, we compared the anti-emetic efficacy of concurrent use of different doses of dexamethasone and mosapride with ondansetron. We hypothesized that combined application of three doses of dexamethasone and mosapride would provide a greater clinical benefit.

Patients And Methods

Study design

This prospective randomized clinical trial was conducted and reported in accordance with CONSORT statement. Patients receiving primary total hip or knee arthroplasty from November 2017 to December 2018 were included. Institutional review board approval (2012-268) was obtained before patient enrollment. Written informed consent and research authorization were obtained from all patients before surgery. The study was conducted in compliance with the tenets of the Declaration of Helsinki, and the original protocol was registered at the Chinese Clinical Trial Registry (ChiCTR1800015896, April 27, 2018). The study was a little different from the registered protocol, regarding the exclusion criteria, samples size and intervention measures in Group B. Firstly, a 50% reduction in PONV was expected, which was the calculation basis of sample size in registered protocol. However, during the pilot study, patients in group B and C also experienced a high incidence of PONV, then we adjusted the expected reduction to 45% and the interventions in group B (oral mosapride instead of single dose of ondansetron) with the approval of monitoring committee.

Inclusion and exclusion criteria

Eligible patients included those at least 18 years old who were at risk of PONV (at least 1 score of Apfel), and scheduled for primary total hip or knee arthroplasty for end-stage joint diseases such as osteoarthritis, development dysplasia of hip, and osteonecrosis of the femoral head. Exclusion criteria included a history of intolerance of any drugs used in the current study, administration of another anti-

emetic drug or systemic steroid within 24 h before surgery, allergy to experimental drugs or history of adverse reactions, diabetes with poor blood glucose control, history of steroid or immunosuppressive drug use within the previous 6 months, history of cardiac disease such as heart failure, heart block, ventricular arrhythmia or severe impairment of bowel motility, renal function or hepatic function.

Randomization and treatment

According to the inclusion and exclusion criteria, the researchers would enroll appropriate patients before surgery and obtain written informed consents. All included patients in this triple-blinded study were randomly allocated to three groups of prophylactic treatment for PONV using a computer-generated randomization list in a 1:1:1 ratio. The group assignments were performed by a research assistant and kept in opaque sealed envelopes that were only opened immediately before surgery.

- Group A received 8 mg ondansetron (4 ml) at the end of surgery; three doses of 2 ml normal saline at anesthetic induction and 6 and 24 hours later; and oral placebo 3 hours before surgery and three times per day after the operation.
- Group B received one dose of 10 mg dexamethasone (2 ml) at anesthetic induction, two doses of normal saline 6 and 24 hours later, 5 mg oral mosapride 3 hours before surgery and three times per day after the operation, and a dose of 4 ml normal saline at the end of surgery.
- Group C received three doses of 10 mg dexamethasone (2 ml) at anesthetic induction and 6 and 24 hours later, 5 mg oral mosapride 3 hours before surgery and three times per day after the operation, and a dose of 4 ml normal saline at the end of surgery.

The intraoperative dexamethasone and ondansetron were administered by an anesthesiologist, and the postoperative oral drugs were administered by nurses who were not involved in the study. The patients, surgeons, data collectors, and analysts were all blinded to the group assignments.

Anesthesia and perioperative pain management

All surgeries were performed under general anesthesia by the same team (F.X.P. and Z.K.Z.) using the standard medial parapatellar arthrotomy for total knee arthroplasty and posterolateral approach for total hip arthroplasty. A cemented posteriorly stabilized prosthesis was implanted in all patients undergoing total knee arthroplasty, and cementless acetabular and femoral components were used in all patients undergoing total hip arthroplasty.

All patients received the same anesthetic regimen and multimodal pain management protocol. Cefuroxime (1.5 g) was given intravenously as a prophylactic antibiotic prior to the first incision. Sufentanil (0.2 µg/kg), propofol (2 mg/kg), atracurium (1 mg/kg), and midazolam (2 mg) were used for anesthetic induction. Sevoflurane (1%–3%) and single bolus of sufentanil (0.1 µg/kg), and atracurium (0.5 mg/kg) were then used to maintain anesthesia during surgery. After prosthesis insertion, propofol (4 mg/kg. h) and remifentanil (0.1 µg/kg. min) were used to maintain anesthesia. The anesthetic drugs were

discontinued before wound closure. All patients received preoperative oral hydration for up to 2 hours prior to surgery and adequate intravenous fluids intraoperatively.

After prosthesis insertion, a periarticular infiltration of 200 mg ropivacaine (100 mg/10 ml) in 60 ml normal saline was injected around the capsule before closure. One dose of 40 mg parecoxib was injected intravenously to manage pain. Postoperative pain control consisted of 50 mg of oral diclofenac sodium (Voltaren sustained-release tablets; Novartis Pharmaceuticals, Basel, Switzerland) every 12 hours. Breakthrough pain was recorded using a Numeric Rating Scale (NRS) ranging from 0 points (no pain) to 10 points (the most severe pain). Once the patients provided feedback that the pain was at a NRS score of >4 points, 10 mg oral oxycodone was administered. If the VAS score was >6 points, 50 mg of pethidine was given by intramuscular injection as required up to every 6 hours. No nerve block or intravenous patient-controlled analgesia was utilized perioperatively.

Outcome measurements

The primary outcome was the incidence of total PONV, and the secondary outcomes were complete response, times until first defecation and ambulation, postoperative appetite score, satisfaction score, length of hospital stay, blood glucose level, and complications. A blinded clinical investigator (Y.C.C.) recorded all episodes of nausea and vomiting, the severity of nausea, the antiemetic rescue requirement, and complete response to evaluate the antiemetic efficacy. Nausea was defined as a subjective unpleasant sensation associated with awareness of the urge to vomit, and vomiting was defined as the forceful expulsion of gastric contents from the mouth [12]. The incidence of total PONV was determined during the 48-hour study period by calculating the proportion of patients who experienced PONV at least once. Furthermore, newly developed PONV in each of the four periods (0–6, 6–12, 12–24, and 24–48 hours) was also calculated. Following institutional guidelines, 10 mg of intramuscular metoclopramide was used as a first-line antiemetic rescue treatment when patients experienced two or more episodes of PONV within 2 hours. This was followed by administration of 4 mg ondansetron intravenously when two consecutive boluses of metoclopramide alone administered at a 30-minute interval were ineffective.

We used a standardized scoring algorithm to classify the severity of nausea during the 48-hour observation period into four degrees. Complete response was defined as no PONV or no requirement for a rescue antiemetic. Mild nausea was defined as the occurrence of nausea caused by an exogenous stimulus (drinking or movement). Moderate nausea was defined as the experience of nausea that required only one dose of a rescue antiemetic. Finally, severe nausea was defined as more than two emetic episodes or the need for more than one dose of a rescue antiemetic(15).

The blinded investigator (Y.C.C.) also recorded the time of first defecation, time of first ambulation, length of hospital stays, appetite score on postoperative days 0 to 2, and patients' satisfaction score. *The patients' appetite was scored by comparison with the preoperative level (the morning on the day 1 before surgery); 1 point represented a worse appetite, 2 points represented no change in appetite from the preoperative state, and 3 points represented a better appetite.* Patients' satisfaction before discharge was estimated using a NRS that ranged from 0 points (extremely dissatisfied) to 10 points (very satisfied).

The fasting blood glucose level was measured in all patients on postoperative days 1 and 2, while the 2-hour postprandial blood glucose level was measured in patients with diabetes mellitus on postoperative day 1. All patients were followed up for 3 months postoperatively, and any complications such as prolonged QT syndrome, wound discharge, surgical site infection, or readmission were recorded.

Statistical analysis

We performed an *a priori* power analysis based on our preliminary results showing that the incidence of PONV was 49% in patients receiving ondansetron prophylaxis alone after total joint arthroplasty(12). We calculated that 303 patients (101 in each arm) were required to detect a 45% reduction in the incidence of PONV in Group C at an alpha level of 0.05 and power of 0.9 using a two-sided test. To allow for exclusions and dropouts, we enrolled 348 patients in the current trial.

The chi-square test or Fisher's exact test was used to determine the statistical significance of differences in the categorical variables, such as the incidence of PONV and proportion of patients with complete response; if significant, multiple comparisons between groups were performed by Bonferroni's corrected *post hoc* test (Z test). Continuous variables were analyzed with one-way analysis of variance if normality test was affirmative (body mass index, length of hospital stay, and times until first defecation or ambulation) or the Wilcoxon signed-rank test (patients' appetite and satisfaction scores); if significant, multiple comparisons between groups were performed by the *post hoc* Tukey test. A p value of <0.05 was considered statistically significant. Statistical analysis was conducted using SPSS 21.0 (IBM Corp., Armonk, NY, USA).

Results

In total, 348 patients were initially allocated to Groups A, B, and C (116 patients in each group). We excluded eight patients each from Groups A and B according to the defined exclusion criteria. Thus, 332 patients (108 in Group A, 108 in Group B, and 116 in Group C) were included in the final analysis (**Figure 1**). We found no differences in demographic characteristics or clinical data among the groups (**Table 1**).

During the whole evaluation period, the patients in Group C experienced lowest incidence of total PONV (29.3%), nausea (15.5%) and vomiting (13.8%, **Table 2**). Similarly, more patients who received three doses of dexamethasone and mosapride in Group C achieved a complete response (70.69%) compared with Groups A (p=0.001) and B (p=0.219). In addition, fewer patients in Group C had severe PONV (1.72%) compared with Group B (p=0.001) and Group A (p<0.001).

According to the timing and duration of PONV, concurrent use of mosapride and three doses of dexamethasone mainly reduced the incidence of PONV during the first 12 hours, especially from 0 to 6 hours (**Tables 3, 4**). 94.4% of all PONV events occurred from 0 to 6 hours postoperatively in Group C, which was higher than that in Group B (p=0.363) and Group A (p=0.001). Of all patients with episodes of PONV in Group C, 88.89% of patients had episodes that lasted <6 hours, which was a higher proportion

than in Group B ($p=0.001$) and Group A ($p<0.001$). The mean time of PONV in Group C was shorter than that in Groups A ($p=0.002$) and B ($p=0.001$).

With the use of multiple doses of dexamethasone, fewer patients needed antiemetic rescue treatment in Group C; however, the total consumption of rescue treatment was not significantly different. Moreover, the consumption of oxycodone and pethidine in Group C was less than that in Groups B and A (**Table 5**).

The time until first defecation was shortest in Group C, and all intergroup differences were statistically significant ($p<0.05$ for all) (**Table 6**). Moreover, the patients in Group C had a better appetite on POD 1, were more satisfied with their healthcare, and had a shorter length of hospitalization than the patients in Groups B and A ($p<0.05$ for all).

Among patients with diabetes mellitus, the fasting blood glucose level was higher in Group C than in Group A on POD 1 (**Table 7**). Among patients without diabetes mellitus, the fasting blood glucose level was higher in Group C than Groups A and B on POD 1 and 2, although the blood glucose level was acceptable in all groups. During the study and follow-up period, the incidence of complications was comparable among the three groups.

Discussion

As an important part of an enhanced recovery after surgery program, PONV management remains a challenge during the postoperative period, especially in the setting of general anesthesia. Therefore, we should pay more attention to PONV prevention and treatment because PONV can cause anxiety and loss of appetite and can increase the risk of complications and prolong length of hospital stay. Although some clinical guidelines and suggestions have been recommended, the incidence of PONV remains a “little big” problem because of the lack of an ideal PONV prophylaxis regimen.

In the present study, we compared three different PONV prophylactic protocols and found that with the use of oral mosapride and three doses of dexamethasone, patients experienced a lower incidence and severity of PONV from 0 to 6 hours postoperatively, a better appetite, and a higher satisfaction score. More importantly, the combined use of three doses of dexamethasone and mosapride reduced the consumption of opioids and promoted postoperative bowel function recovery, thus reducing the incidence of postoperative constipation.

Effective treatments that limit PONV allow patients to mobilize earlier, decrease the length of stay, and improve patients' overall satisfaction. Many factors are attributable to the incidence of PONV, including female sex, a history of PONV or motion sickness, and opioid use(16). 5-HT₃ receptor antagonists and glucocorticoids are the most common treatment options for PONV prevention. Ondansetron, one type of 5-HT₃ receptor antagonist, has been recommended as the first-line agent for PONV prevention by some clinical guidelines(4). Additionally, a study by Apfel et al.(17) showed comparable efficacy in reducing the incidence of PONV between 4 mg ondansetron and 4 mg dexamethasone. For this reason, our control group comprised patients treated with ondansetron only.

Dexamethasone is a high-potency, long-acting glucocorticoid with good bioavailability and few corticoid-related adverse effects(18). Glucocorticoids reduce PONV through a central antiemetic effect by inhibiting prostaglandin synthesis and release of endogenous opioids(19). Another reason for the antiemetic effect of dexamethasone is the anti-inflammatory response that inhibits the release and reduces the levels of inflammatory factors such as C-reactive protein and interleukin-6(8-10). Our research team systematically explored the potential clinical benefit of multiple low doses of dexamethasone, including two or three doses in the setting of total hip and knee arthroplasty(8-10, 20). In studies performed by Xu et al.(8)and Lei et al.(9), two doses of 10 mg dexamethasone were intravenously administered upon anesthetic induction and return to the inpatient unit. Two other studies by the same research teams evaluated the efficacy and safety of three doses of dexamethasone (10 or 20 mg at anesthetic induction, 10 mg after returning to the inpatient unit, and 10 mg at 24 hours after the first dose) following primary total hip and knee arthroplasty(10, 20). In all of the aforementioned studies, the authors focused on the effect of dexamethasone on the postoperative inflammatory response, pain relief, and joint function. All studies also indicated promising effects of three doses of dexamethasone. In contrast to these previous studies, the current study mainly focused on the outcomes of PONV, appetite, bowel function, and safety. Moreover, the prophylactic regimen in the current study was different from that in the previous studies; i.e., the current regimen combined dexamethasone with a 5-HT₄ agonist at different time intervals instead of a 5-HT₃ antagonist. Additionally, we compared the new regimen with other common regimens (ondansetron only and a single dose of dexamethasone). This is a major strength of our study.

Another advantage of dexamethasone for PONV prevention is the additional anti-inflammatory effect. A previous study demonstrated that PONV is specific to the patients' perioperative inflammatory control [20], which would peak at postoperative 24 – 48 hours, and last for at least 72 hours. Systematic reviews and meta-analyses have also shown that a high dose of dexamethasone (10mg or 0.1mg/kg) more effectively reduces the incidence of PONV and provides better postoperative pain control(21, 22). In addition, the inflammation was closely related with fibrinolysis, which would peak at postoperative 6 hours and last for 24 hours. And similar to another study(23), our preliminary results indicated that most of the PONV episodes (87%) occurred during the first 12 hours postoperatively. Therefore, we hypothesized that repeated dose at postoperative 6 hours and 24 hours would be more effective although the half- life of dexamethasone is 36 hours. Our results also revealed a lower incidence of severe PONV (1.72%) and a shorter duration (<6 h) in the patients treated with three doses of dexamethasone. Moreover, the patients in this group (Group C) experienced better pain control with respect to less opioid consumption (**Table 5**), and this also contributed to the lower incidence of PONV. However, we can't conclude whether the lower incidence of PONV was owed to the first or second or third dose of dexamethasone.

Mosapride, a 5-HT₄ receptor agonist, is a gastroprokinetic agent indicated for the treatment of gastrointestinal symptoms such as heartburn, nausea/vomiting associated with chronic gastritis, and functional dyspepsia(13). The *in vivo* study by Mine et al.(24) showed that mosapride was helpful to improve the selective serotonin reuptake inhibitor (SSRI)-induced delay in gastric emptying and the

incidence of SSRI-induced emesis, which was in accordance with another clinical study(25). Moreover, another study also showed the potential anti-emetic effect of mosapride in patients undergoing chemotherapy(14). Interestingly, in the present study, the patients in Group B (oral mosapride in addition to a single dose of dexamethasone) had a lower incidence of PONV (although not significantly lower) during the first 6 hours. Additionally, a shorter duration of time passed until the first defecation. This raises the possibility that oral mosapride is helpful to reduce the incidence of PONV by promoting gastrointestinal motility. Although our results indicated no adverse events such as prolonged QT syndrome associated with mosapride, in view of the fact that another 5-HT₄ receptor agonist, cisapride, was withdrawn because of severe cardiotoxicity, further studies are warranted to investigate the potential antiemetic efficacy and safety profile of mosapride in the setting of joint arthroplasty before routine recommendation.

In addition to the fewer episodes and decreased severity of PONV in Group C, these patients had better bowel function, higher appetite scores, shorter hospital stays, and higher satisfaction scores. This might have been partially due to the application of mosapride, although multiple doses of dexamethasone could have improved the pain control, joint function, early ambulation, and length of hospital stay.

The main obstacle to the widespread application of dexamethasone is the concern about its safety, especially in terms of the risk of infection and hyperglycemia. Although the literature has provided some evidence that the use of low-dose dexamethasone does not increase the risk of adverse events(26, 27), the current results were inconclusive. In our study, the fasting blood glucose level in patients both with and without diabetes receiving dexamethasone significantly increased, but it remained at an acceptable level. Additionally, the incidence of wound complications during the follow-up period was comparable among the three groups. Nevertheless, we cannot conclude that the use of dexamethasone is safe because our study sample was not large enough to detect statistical significance. A previous power analysis showed that >3500 patients would be required to meaningfully evaluate an increase in surgical infection(21). Thus, we must acknowledge that this study lacks the power to fully assess the low incidence of events because of the relatively small sample size. The rate of such negative outcomes could be much more severe than what we expected, and the results should be interpreted with caution. Although large-scale prospective studies are still needed, our study might provide some insight into the safety of dexamethasone.

We acknowledge that our study has some limitations. First and foremost, the generalizability of this regimen was limited because of same postoperative pain management and anesthesia protocol. Different from most western countries, general anesthesia was preferable by the anesthesia team in our institution because of better feasibility of early ambulation and anticoagulation. Second, we did not evaluate the effect of this regimen on the postoperative inflammatory response and other functional outcomes because previous studies have confirmed its efficacy in this regard(8-10). Thirdly, the follow-up period was too short to detect the possible infectious complications associated with dexamethasone. The last but not the least, the interventions between the three groups were not definitely comparable and consistent, especially the lack of isolation of mosapride, which this made us difficult to conclude the

prophylactic role of mosapride in the prevention of PONV. Furthermore, because of confounding bias, we couldn't conclude how much PONV prevention effect was related to mosapride or dexamethasone.

Conclusion

Prophylactic use of mosapride and three doses of dexamethasone effectively reduced the incidence of PONV. Additionally, this regimen resulted in a better postoperative appetite, bowel function recovery, and pain relief when compared with a single dose or ondansetron only in patients undergoing primary total joint arthroplasty under general anesthesia.

List Of Abbreviations

PONV, postoperative nausea and vomiting; THA, total hip arthroplasty; TKA, total knee arthroplasty; ASA, American Society of Anesthesiologists; PACU, post-anesthesia care unit; BMI, body mass index.

Declarations

Ethics approval and consent to participate

The study protocol was approved by the Institutional Review Board of West China Hospital, Sichuan University (2012-268), and all procedures were performed according to the Declaration of Helsinki. Written informed consent was obtained prospectively from all patients prior to surgery.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The research was supported by the Post-Doctor Research Project, West China Hospital, Sichuan University (2018HXBH073); the National Clinical Research Center for Geriatrics, West China Hospital, Sichuan University (Z2018B11); and the National Health and Family Planning Commission of the People's Republic of China program (201302007).

Authors' contributions

Conceptualization, Z. Zhou and F. Pei; Methodology, J. Xie and Y. Cai; Formal analysis, J. Ma and Q. Huang; Investigation, Y. Cai and J. Xie; Resources, Z. Zhou and F. Pei; Data curation and Writing, Y. Cai and J. Xie; Project administration and Funding Acquisition, F. Pei and J. Xie.

Acknowledgements

We wish to express our sincere appreciation to all the patients and staff involved in this study. We also thank Xiaowei Feng from Chengdu First People's Hospital for the language editing and writing assistance. Finally, we thank Angela Morben, DVM, ELS, from Liwen Bianji, Edanz Editing China (www.liwenbianji.cn/ac), for editing the English text of a draft of this manuscript.

References

1. Lin CJ, Williams BA. Postoperative nausea and vomiting in ambulatory regional anesthesia. *Int Anesthesiol Clin*. 2011;49(4):134-43.
2. Sansonnens J, Taffe P, Burnand B, group ADSs. Higher occurrence of nausea and vomiting after total hip arthroplasty using general versus spinal anesthesia: an observational study. *BMC Anesthesiol*. 2016;16(1):44.
3. Sculco PK, Pagnano MW. Perioperative solutions for rapid recovery joint arthroplasty: get ahead and stay ahead. *J Arthroplasty*. 2015;30(4):518-20.
4. Gan TJ, Diemunsch P, Habib AS, Kovac A, Kranke P, Meyer TA, et al. Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg*. 2014;118(1):85-113.
5. Kranke P, Diemunsch P. The 2014 consensus guidelines for the management of postoperative nausea and vomiting: a leapfrog towards a postoperative nausea and vomiting-free hospital. *Eur J Anaesthesiol*. 2014;31(12):651-3.
6. Eberhart LH, Kranke P. Postoperative nausea and vomiting: is everything now solved or still more questions than answers? *Eur J Anaesthesiol*. 2016;33(12):878-80.
7. Ross-Adjie GM, Monterosso L, Bulsara M. Bowel management post major joint arthroplasty: results from a randomised controlled trial. *Int J Orthop Trauma Nurs*. 2015;19(2):92-101.
8. Xu B, Ma J, Huang Q, Huang ZY, Zhang SY, Pei FX. Two doses of low-dose perioperative dexamethasone improve the clinical outcome after total knee arthroplasty: a randomized controlled study. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(5):1549-56.
9. Lei YT, Xu B, Xie XW, Xie JW, Huang Q, Pei FX. The efficacy and safety of two low-dose peri-operative dexamethasone on pain and recovery following total hip arthroplasty: a randomized controlled trial. *Int Orthop*. 2018;42(3):499-505.
10. Lei Y, Huang Q, Xu B, Zhang S, Cao G, Pei F. Multiple Low-Dose Dexamethasone Further Improves Clinical Outcomes Following Total Hip Arthroplasty. *J Arthroplasty*. 2018;33(5):1426-31.
11. Vuorinen MA, Palanne RA, Makinen TJ, Leskinen JT, Huhtala H, Huotari KA. Infection safety of dexamethasone in total hip and total knee arthroplasty: a study of eighteen thousand, eight hundred

- and seventy two operations. *Int Orthop*. 2019;43(8):1787-92.
12. Cai Y XJ, Ma J, Pei F. Risk factors of postoperative nausea and vomiting following primary total hip or knee arthroplasty under general anesthesia:a retrospective study. . *Chinese Journal of Bone and Joint Surgery*. 2016;9(1):5.
 13. Mizuta Y, Shikuwa S, Isomoto H, Mishima R, Akazawa Y, Masuda J, et al. Recent insights into digestive motility in functional dyspepsia. *J Gastroenterol*. 2006;41(11):1025-40.
 14. Ganesan P, Sharma A, Mohanti BK, Gogia A. Protracted cisplatin-induced vomiting responding to mosapride. *Indian J Cancer*. 2010;47(1):73-4.
 15. Reibaldi M, Fallico M, Longo A, Avitabile T, Astuto M, Murabito P, et al. Efficacy of Three Different Prophylactic Treatments for Postoperative Nausea and Vomiting after Vitrectomy: A Randomized Clinical Trial. *J Clin Med*. 2019;8(3).
 16. Gan TJ. Risk factors for postoperative nausea and vomiting. *Anesth Analg*. 2006;102(6):1884-98.
 17. Apfel CC, Korttila K, Abdalla M, Kerger H, Turan A, Vedder I, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med*. 2004;350(24):2441-51.
 18. McKee MD. Efficacy and safety of steroid use for postoperative pain relief. *J Bone Joint Surg Am*. 2007;89(5):1134; author reply
 19. Holte K, Kehlet H. Perioperative single-dose glucocorticoid administration: pathophysiologic effects and clinical implications. *J Am Coll Surg*. 2002;195(5):694-712.
 20. Xu H, Zhang S, Xie J, Lei Y, Cao G, Pei F. Multiple Doses of Perioperative Dexamethasone Further Improve Clinical Outcomes After Total Knee Arthroplasty: A Prospective, Randomized, Controlled Study. *J Arthroplasty*. 2018;33(11):3448-54.
 21. Lunn TH, Kehlet H. Perioperative glucocorticoids in hip and knee surgery - benefit vs. harm? A review of randomized clinical trials. *Acta Anaesthesiol Scand*. 2013;57(7):823-34.
 22. Yue C, Wei R, Liu Y. Perioperative systemic steroid for rapid recovery in total knee and hip arthroplasty: a systematic review and meta-analysis of randomized trials. *J Orthop Surg Res*. 2017;12(1):100.
 23. Koh IJ, Chang CB, Lee JH, Jeon YT, Kim TK. Preemptive low-dose dexamethasone reduces postoperative emesis and pain after TKA: a randomized controlled study. *Clin Orthop Relat Res*. 2013;471(9):3010-20.
 24. Mine Y, Oku S, Yoshida N. Anti-emetic effect of mosapride citrate hydrate, a 5-HT₄ receptor agonist, on selective serotonin reuptake inhibitors (SSRIs)-induced emesis in experimental animals. *J Pharmacol Sci*. 2013;121(1):58-66.
 25. Bergeron R, Blier P. Cisapride for the treatment of nausea produced by selective serotonin reuptake inhibitors. *Am J Psychiatry*. 1994;151(7):1084-6.
 26. Polderman JA, Farhang-Razi V, Van Dieren S, Kranke P, DeVries JH, Hollmann MW, et al. Adverse side effects of dexamethasone in surgical patients. *Cochrane Database Syst Rev*. 2018;11:CD011940.

27. Dissanayake R, Du HN, Robertson IK, Ogden K, Wiltshire K, Mulford JS. Does Dexamethasone Reduce Hospital Readiness for Discharge, Pain, Nausea, and Early Patient Satisfaction in Hip and Knee Arthroplasty? A Randomized, Controlled Trial. *J Arthroplasty*. 2018;33(11):3429-36.

Tables

Table 1. Baseline characteristics of all patients

	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p value
Age (years)	57.4 ± 14.2	62.2 ± 12.7	61.2 ± 13.3	.087
BMI (kg/m ²)	23.9 ± 4.1	24.1 ± 3.2	24.1 ± 3.9	.899
Female / Male	74 / 34	75 / 33	80 / 36	.993
Smoking (n, %)	34 (31.5%)	35 (32.4%)	22 (18.2%)	.164
History of kinetia				.506
YES	44	42	36	
NO	64	66	80	
ASA Score				.813
ASA=2	76	75	86	
ASA=3	32	33	30	
Type of surgery				
THA	52	45	64	.245
TKA	56	63	52	
Number of comorbidities				.518
1	42	52	50	
≥2	8	12	14	
Operation time (min)	77.4 ± 20.9	77.7 ± 26.2	78.3 ± 19.3	.973
Anesthesia time (min)	126.7 ± 28.7	123.9 ± 31.1	122.1 ± 22.0	.688
PACU time (min)	72.7 ± 47.4	78.0 ± 38.9	75.0 ± 36.0	.72
Sufentanil (mg)	24.3 ± 3.4	24.6 ± 3.3	24.6 ± 5.6	.888
Remifentanil (mg)	0.6 ± 0.1	0.6 ± 0.2	0.6 ± 0.3	.530
Propofol (mg)	202.8 ± 109.4	206.0 ± 129.9	225.0 ± 122.1	.565
Midazolam (mg)	2.1 ± 0.5	2.1 ± 0.5	2.3 ± 0.8	.059
Atracurium (mg)	13.6 ± 2.6	13.9 ± 3.0	13.3 ± 3.8	.511
Sevoflurane (ml)	30.8 ± 6.6	30.2 ± 9.5	28.6 ± 7.2	.341

ASA, American Society of Anesthesiologists; BMI, body mass index; THA, total hip arthroplasty; TKA, total knee arthroplasty; PACU, post-anesthesia care unit.

Table 2. Incidence of PONV during the first postoperative 48 hours

	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	P*	p1 [†]	p2 [†]	p3 [†]
Total PONV	56(51.9%)	40(37.0%)	34(29.3%)	.002	.028	.001	.219
Vomiting	22 (20.4%)	17 (15.7%)	16 (13.8%)	.401	-	-	-
Nausea	34 (31.5%)	23 (21.3%)	18 (15.5%)	.016	.086	.005	.264
Complete response	52 (48.2%)	68 (63.0%)	82 (70.7%)	.002	.028	.001	.219
Mild Nausea	4 (3.7%)	2 (1.9%)	8 (6.9%)	.176	-	-	-
Moderate Nausea	10 (9.3%)	7 (6.5%)	8 (6.9%)	.763	-	-	-
Severe Nausea	20 (18.5%)	14 (13.0%)	2 (1.7%)	<.001	.262	<.001	.001
Moderate + severe Nausea	30 (27.8%)	21 (19.4%)	10 (8.6%)	.001	.149	<.001	.019

Data are presented as number of patients (percentage).

PONV, postoperative nausea and vomiting

*Uncorrected p values (for the three-way comparison)

[†]Bonferroni-corrected p values: p1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C. The corrected significance threshold was 0.016.

Table 3. Timing of PONV events during the postoperative period

Time	Total (n= 332)	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p*	p1 [†]	p2 [†]	p3 [†]
0 - 6 h	52 (69.3%)	16 (47.1%)	19 (82.6%)	17 (94.4%)	.001	.007	.001	.363
6 - 12 h	9 (12.0%)	6 (17.7%)	2 (8.70%)	1(0.6%)	.106	-	-	-
12 - 24 h	7 (9.3%)	6 (17.7%)	1 (4.4%)	0	.006	.119	.012	.482
24 - 48 h	7 (9.3%)	6 (17.7%)	1 (4.4%)	0	.006	.119	.012	.482

Data are presented as number of patients (percentage).

PONV, postoperative nausea and vomiting

*Uncorrected p values (for the three-way comparison)

[†]Bonferroni-corrected p values: p1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C. The corrected significance threshold was 0.016.

Table 4. Duration of PONV during the first postoperative 48 hours

Duration	Total (n= 332)	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p*	p1 [†]	p2 [†]	p3 [†]
Mean time (h)	13.6 ± 5.4	16.2 ± 5.0	18.2 ± 8.2	2.9 ± 1.7	.002	.606	.002	.001
< 6 h	37 (49.3%)	12 (35.3%)	9 (39.1%)	16 (88.9%)	.001	.768	<.001	.001
6 -12 h	14 (18.7%)	8 (23.5%)	4 (17.4%)	2 (11.1%)	.610	-	-	-
12-24 h	8 (10.7%)	4 (11.8%)	4 (17.4%)	0	.189	-	-	-
> 24 h	16 (21.3%)	10 (29.4%)	6 (26.1%)	0	.039	.784	.010	.027

Data are presented as number of patients (percentage) or mean \pm standard deviation.

PONV, postoperative nausea and vomiting

*Uncorrected p values (for the three-way comparison)

[†] Bonferroni-corrected p values: p1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C. The corrected significance threshold was 0.016.

Table 5. Requirement of rescue treatment for PONV and pain during the postoperative 48h

	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p*	p1 [†]	p2 [†]	p3 [†]
Metoclopramide (mg)	6.5 \pm 1.1	2.9 \pm 1.2	1.4 \pm 0.5	.047	.055	.017	.415
Ondansetron (mg)	1.7 \pm 0.5	1.7 \pm 0.5	0.6 \pm 0.1	.238	-	-	-
Oxycodone (mg)	33.2 \pm 8.7	17.9 \pm 9.7	1.8 \pm 0.3	<.001	.001	<.001	<.001
Pethidine (mg)	20.4 \pm 8.3	20.6 \pm 9.8	6.0 \pm 1.0	.002	.959	.005	.001

Data are presented as mean \pm standard deviation.

PONV, postoperative nausea and vomiting

*p values with one-way analysis of variance; [†]p values with Tukey's post hoc test.

P1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C.

-

Table 6. Other clinical outcomes

	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p	p1	p2	p3
Time to first defecation (h)	52.4 \pm 16.5	46.7 \pm 11.8	37.2 \pm 9.0	<.001	.006	<.001	<.001
Appetite score on POD 0	2.2 \pm 0.8	2.4 \pm 0.7	2.6 \pm 0.6	.009	.106	.002	.054
Appetite score on POD 1	2.3 \pm 0.7	2.4 \pm 0.7	2.7 \pm 0.5	.001	.298	<.001	.002
Appetite score on POD 2	2.6 \pm 0.5	2.6 \pm 0.5	2.7 \pm 0.4	.121	-	-	-
Time to first ambulation (h)	27.1 \pm 8.2	28.5 \pm 10.8	25.6 \pm 6.3	.298	-	-	-
Satisfaction score	9.3 \pm 0.7	9.3 \pm 0.9	9.7 \pm 0.4	.001	.661	.007	<.001
LOH (d)	5.4 \pm 2.1	5.0 \pm 2.2	4.1 \pm 2.2	.02	.321	.001	.049

Data are presented as mean \pm standard deviation.

POD, postoperative day; LOH, length of hospitalization.

*p values with one-way analysis of variance; [†]p values with Tukey's post hoc test.

P1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C.

Table 7. Incidence of complications and level of blood glucose on POD 1 and 2

	Group A (n= 108)	Group B (n= 108)	Group C (n= 116)	p*	p1 [†]	p2 [†]	p3 [†]
Diabetes Mellitus patients							
FBG on POD1	9.5 ± 2.2	10.7 ± 2.4	11.0 ± 2.9	.001	.001	.001	.525
2-h PBG after breakfast	11.6 ± 1.5	12.3 ± 3.1	14.9 ± 3.9	.002	.177	.001	.001
2-h PBG after lunch	11.5 ± 1.3	13.2 ± 3.7	14.8 ± 2.6	.096	-	-	-
2-h PBG after dinner	11.0 ± 2.3	12.7 ± 3.2	10.1 ± 2.7	.173	-	-	-
FBG on POD2	9.4 ± 1.0	9.8 ± 2.1	7.7 ± 3.1	.127	-	-	-
Non- Diabetes Mellitus patients							
FBG on POD1	6.8 ± 1.2	6.9 ± 1.6	7.8 ± 1.3	.021	.212	.017	.041
FBG on POD2	5.6 ± 1.0	5.7 ± 1.8	7.1 ± 1.2	.031	.76	.039	.048
Prolonged QT syndrome	0	0	0	-			
Wound site discharge	2 (1.9%)	3 (2.8%)	4 (3.5%)	.913	-	-	-
Surgical site infection	0	0	0	-			
Pulmonary infection	2 (1.9%)	6 (5. 6%)	1 (0.9%)	.104	-	-	-
Re-admission	1 (0.9%)	1 (0.9%)	1 (0.9%)	1.000	-	-	-

Data are presented as mean ± standard deviation or number (percentage).

POD, postoperative day; FBG, fasting blood glucose; PBG, 2-hour postprandial blood glucose.

*p values with one-way analysis of variance; [†]p values with Tukey's post hoc test.

p1, Group A vs. B; p2, Group A vs. C; p3, Group B vs. C.

Figures




Figure 1

A flow diagram shows the patients recruitment.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.pdf](#)