

Using Pranayama and Deep Breathing Exercises to Reduce Cancer-Related Fatigue and Insomnia During Radiotherapy: A Randomized Controlled Study

Fatma Gündođdu (✉ fatma.gundogdu@karatay.edu.tr)

KTO Karatay Üniversitesi <https://orcid.org/0000-0001-8147-220X>

Sema Koçaşlı

Ankara Yıldırım Beyazıt Üniversitesi: Ankara Yıldırım Beyazıt Üniversitesi

Research Article

Keywords: Breathing exercises, pranayama, breast cancer, cancer-related fatigue, insomnia, radiotherapy

DOI: <https://doi.org/10.21203/rs.3.rs-160848/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Patients receiving radiotherapy often experience cancer-related fatigue and insomnia. These experiences negatively affect patients' physiological and psychological well-being. The study aims to evaluate the effectiveness of pranayama and deep breathing exercise in reducing fatigue and insomnia in patients receiving radiotherapy due to breast cancer.

Materials and methods: This randomized controlled interventional study was carried out with 60 patients divided into the pranayama (20), deep breathing exercise (20) and control group (20). Data were collected with the Piper Fatigue Scale (PFS), the Pittsburgh Sleep Quality Index (PSQI) and the Visual Analog Scale (VAS).

Results: While patient fatigue in the control group significantly increased during the treatment process ($p < 0.05$), the deep breathing exercise group ($p < 0.05$) did not experience the same increase in fatigue. The post-treatment insomnia VAS score of the pranayama and deep breathing exercise groups decreased compared to the pre-treatment scores ($p < 0.05$). In the group comparison of PSQI scale scores, the subdimensions *subjective sleep quality* and *daytime dysfunction* were negatively affected at the end of the treatment in the control group ($p < 0.05$) and the subdimension *subjective sleep quality* was positively affected in the deep breathing exercise group ($p < 0.05$).

Conclusion: Pranayama and deep breathing exercises are effective in reducing fatigue and insomnia in patients receiving radiotherapy due to breast cancer.

Introduction

Cancer-related fatigue is defined as "a disturbing, persistent, subjective physical, emotional and/or cognitive exhaustion or burnout feeling associated with cancer and its treatment unrelated to recent daily activities". Unlike the fatigue experienced by healthy individuals, cancer-related fatigue is not known to resolve by rest and gives the individual more trouble than a normal feeling of fatigue [1]. The prevalence of fatigue related to cancer or cancer 33treatment varies between 41% to 88% [2-7]. In addition, studies have shown that together with fatigue; depression, anxiety disorders and a decrease in life and sleep quality occur in cancer patients [3, 8-12].

Not only pharmacological but also complementary/nonpharmacological methods are used to prevent, reduce and treat cancer-related fatigue. Some of these methods include exercise, acupuncture, massage, acupressure, cognitive behavior therapy, nutritional counseling, awareness-based stress reduction and yoga [1,13-18]. Yoga is a mind-body focus and consists of eight limbs (yama, niyama, asana, pranayama, pratyahara dharana, dhyana, samadhi-trans) developing harmony with all aspects of consciousness [13,19,20]. Studies have shown that yoga is effective in decreasing fatigue, sleep disorders, depression and anxiety and in increasing quality of life in breast cancer patients [13, 19, 21-24].

Pranayama however, one of the eight aspects of yoga, consists of breathing techniques that aim to regulate one or more respiratory functions directly and consciously [25-30]. The calm, soothing breathing techniques of pranayama (nadi shodana, bhramari pranayama etc.) reduce sympathetic stimulus and increase parasympathetic activity [24-31]. These breathing exercise methods carried out with slow breaths, ensure relaxation and is effective in relieving stress and related fatigue. Besides pranayama there are also other slow breathing exercises (diaphragmatic breathing, pursed lip breathing, etc.). These breathing exercises constitute an effective integrative body-mind technique to deal with stress and psychosomatic conditions, which decreases fatigue by increasing parasympathetic activity [32-35].

Chakrabarty et al. (2015), Dhruva et al. (2012), and Mustian et al. (2013) reported in their studies that pranayama reduces cancer-related fatigue, increases sleep quality and reduces the use of sleeping medication [14, 25, 36]. Pranayama also lowers cortisol, which is a factor in the development of fatigue, increases functional capacity and quality of life in breast cancer patients and reduces sleep disorders [22, 38, 39]. Therefore, both pranayama (yogic breathing exercises) and slow breathing exercises can be easily used with oncology patients such as breast cancer patients to reduce fatigue. These exercises are easy-to-learn, can be done individually or as a group anytime and anywhere and do not cause harm to the person [25, 40, 41].

The oncology nurse plays an important role in the evaluation, prevention, reduction and follow-up of cancer-related fatigue, developing a care plan related to fatigue management. Including nonpharmacological methods such as yoga, an effective exercise, together with pharmacological methods in the management of fatigue increases the quality of care. The nurse is in a significant position to manage fatigue using evidence-based practice in observing the patient, teaching the patient non-pharmacological practices that ensures the control of fatigue and by encouraging the patient to use these techniques. Well-designed clinical trials are needed to evaluate the contribution of breathing exercises to the recovery process of cancer patients. Therefore, this research was conducted to evaluate the effect of pranayama and deep breathing exercise on fatigue and sleep quality in women receiving radiotherapy due to breast cancer.

Methods

2.1 Study Design

Randomized controlled intervention study.

2.2 Study sample

The study population consisted of patients diagnosed with breast cancer, who received radiotherapy in the radiation oncology unit of a hospital in Ankara, between June 01, 2017 and January 31, 2019.

The sample size was calculated with the G*Power (G*Power, Ver. 3.0.10, University Kiel Germany software program (<http://www.psych.uni-duesseldorf.de/aap/projects/gpower/>)) using preliminary study data

since there are no similar studies in the literature. A preliminary study was conducted for each of the three groups containing eight, eight and seven participants, respectively. In the variables, the lowest effect width was determined as 0.25 and the highest effect width was 0.62. As a result of the analysis on fatigue and insomnia, which was the main variable in the study, the number of groups were determined as three, the repetitions to be carried out as two, type I error rate as 0.05, the power of the study as 0.95. The minimum number of volunteers required to determine the difference between the three groups with an effect width of $f=0.25$ was determined as 54 individuals (18 in each group). At least three additional volunteers (16%) for each group were added, considering monitoring and data loss during the follow-up. Thus, the number of subjects was determined as 63 individuals (21 for each group).

Of the 63 patients, one wanted to interrupt the treatment and two did not want to continue in the study, hence the study was concluded with 20 patients in each group (60 patients) (Figure 1).

Literate patients over the age of 18 years, who underwent breast resection or modified radical mastectomy, were planned for radiotherapy for the chest wall/breast and lymphatic area, had Stage II-III breast cancer according to the TNM (Tumor diameter, Node, Metastasis) classification system, had a hemoglobin (Hgb) level of 10 gm/dL and above, scored 0 or 1 on the ECOG (Eastern Cooperative Oncology Group) Performance Scale and who agreed to participate in the study were included in the sample. Exclusion criteria included; having any problems that would prevent communication, having stage IV breast cancer according to the TNM classification system, using opioids or sedating drugs, scoring above 1 on the ECOG Performance Scale, having psychiatric illnesses, taking yoga/pranayama/deep breathing exercise lessons before or having done these exercises prior to diagnosis, having recurrent breast cancer and being unable to perform the exercises due to physical inability/respiratory distress. Interruption of the treatment regimen due to any complications, participation in less than 20 of the 25 sessions and the desire to leave the study were determined as termination criteria.

Individuals, who met the inclusion criteria, were assigned to the groups using a simple randomization method with a closed envelope. Individuals were asked to complete the data collection forms themselves. Those, who were not able to do so received support from a nurse within the unit.

2.3. Instruments

Descriptive characteristics questionnaire: This form contained seven questions regarding the patients' descriptive characteristics (age, educational status, type of surgery etc.) and was created by the researchers in line with the literature [14, 23, 32, 42].

Fatigue Visual Analog Scale (VAS) Score: This scale which evaluates the severity of fatigue, was scored as follows: no fatigue= 0 points, mild fatigue= 1-3 points, moderate fatigue= 4-6 points, severe fatigue= 7-9 points and unbearable fatigue= 10 points.

Piper Fatigue Scale (PFS): This scale consisted of 22 items and evaluated the patient's subjective fatigue perception with four subdimensions. Each item was scored between 0 and 10. A high score obtained from the scale indicated a high level of perceived fatigue. The Turkish validity-reliability study of the scale was carried out by Can [43,44]. In this study, the Cronbach's alpha internal consistency coefficient was found as 0.98.

Insomnia Visual Analog Scale (VAS) Score: The scoring system of this scale was as follows: no insomnia= 0 points, mild insomnia= 1-3 points, moderate insomnia= 4-6 points, severe insomnia= 7-9 points and unbearable insomnia= 10 points.

Pittsburgh Sleep Quality Index (PSQI): PSQI is a self-report-based screening and evaluation tool that provides detailed information on sleep quality and type and severity of sleep disorder in the past month. The scale consists of 24 questions and seven components to evaluate subjective sleep quality, time it takes to fall asleep, sleep time, habitual sleep efficiency, sleep disturbance, use of sleeping pills and daytime dysfunction. The total PSQI score could vary from 0 to 21. A PSQI score over 5 points indicated that the person suffered serious problems in at least two areas related to sleep or that they suffered mild to moderate problems in more than three areas [45]. The validity and reliability study of the index was carried out by Ağargün et al.[46]. In this study, the Cronbach's alpha internal consistency coefficient was found as 0.80.

Questionnaire on Patients' Opinions of the Exercises: A form consisting of ten questions prepared by the researcher to determine the opinions of the patients on the feasibility and results of the exercises (difficulties in learning and doing the exercises, satisfaction with doing the exercises, etc.). The opinions of patients in the pranayama and deep breathing exercises groups were taken at the end of radiotherapy treatment.

Patients in the pranayama, deep breathing exercise and control group were evaluated six times in terms of fatigue and insomnia VAS scores. One time prior to the start of the 25-day session of radiotherapy treatment, four times during the radiotherapy treatment process (at the 1st through the 4th week) and one time at the end of the treatment. PFS and PSQI were evaluated once prior to and after the 25-day session of radiotherapy treatment.

2.4 Interventions

Protocols for pranayama and deep breathing exercises were prepared in line with the literature and the intervention was carried out (Figure 2). Prior to develop the protocol, certified training was received by the researcher for breathing exercises. Written opinions were received from seven experts regarding the content validity in order to evaluate the protocols in terms of expression, clarity, coverage and whether scientific errors are present. Following the opinions, the protocol content was edited and finalized.

Pranayama and Deep Breathing Exercise Groups: The patients were taught pranayama and deep breathing exercises by the researcher before the radiotherapy started. The patients were observed

performing these exercises for 10 minutes after each radiotherapy session (25 days) accompanied by the researcher in a quiet room within the radiotherapy unit containing two couches and a chair.

Control Group: The patients in this group did not receive any intervention.

2.5 Statistical analysis

Statistical analyses were made using IBM SPSS for Windows Version 21.0 software program. The compatibility of numerical variables to normal distribution was examined by using Kolmogorov-Smirnov and Shapiro-Wilk tests. Numerical variables that conform to normal distribution were reported using mean \pm standard deviation and numerical variables that do not conform to normal distribution were reported using the median [minimum- maximum] values. In terms of numerical variables that conform to normal distribution, comparisons between groups were analyzed with one-way analysis of variance (ANOVA), taking into account the number of groups to be compared. Kruskal-Wallis test was used for comparison of numerical measurements that do not conform with normal distribution. Wilcoxon test was used for comparisons in dependent groups. The comparison of categorical variables between groups was examined with the Chi-Square test. Bonferroni-corrected Post-Hoc tests were used to identify the groups that made the difference in comparisons between groups with significant differences. The statistical significance level for all hypothesis tests was set at 0.05 and below.

2.6 Ethical Considerations

Ankara Yıldırım Beyazıt University Ethics Committee gave permission (study code: 486, approval date: 31.03.2017, number: 28) and the scientific committee of the hospital where the study was performed gave permission for the study. Patients gave written informed consents.

Results

Table 1. Patients' sociodemographic characteristics (n=60)

| Characteristics | Control (n=20) | Pranayama (n=20) | Deep Breathing (n=20) | p |
|---|----------------|------------------|-----------------------|------|
| Age (mean±SD) | 50.3±12.07 | 53±12.20 | 50.5±10.07 | 0.33 |
| Marital Status | | | | |
| Married | 14 (70) | 18 (90) | 12 (60) | 0.09 |
| Single | 6 (30) | 2 (10) | 8 (40) | |
| Education | | | | |
| Primary School | 5 (25) | 2 (10) | 5 (25) | 0.62 |
| High School | 3 (15) | 2 (10) | 4 (20) | |
| Bachelor and above | 12 (60) | 16 (80) | 11 (55) | |
| Employment status (During treatment) | | | | |
| Employed | 2 (10) | 3 (15) | 4 (20) | 0.67 |
| Unemployed | 18 (90) | 17 (85) | 16 (80) | |
| Lives alone (During treatment) | | | | |
| Yes | 1 (5) | 2 (10) | 3 (15) | 0.57 |
| No | 19 (95) | 18 (90) | 17 (85) | |
| Surgical Treatment | | | | |
| Modified Radical Mastectomy(MRM) | 12 (60) | 10 (50) | 16 (80) | 0.13 |
| Breast Conserving Surgery (BCS) | 8 (40) | 10 (50) | 4 (20) | |
| Hormone treatment | | | | |
| Yes | 8 (40) | 5 (25) | 9 (45) | 0.39 |
| No | 12 (60) | 15 (75) | 11 (55) | |

There was no statistically significant difference between the patients in the control, pranayama and deep breathing exercise groups in terms of sociodemographic characteristics, surgical treatment and hormone treatment. As specified in the table, all patients received 25 workdays of radiotherapy and prior to that all patients received chemotherapy (Table 1).

According to the results, the fatigue VAS score of the patients in the control group significantly increased after radiotherapy compared to before radiotherapy ($p=0.01$). The VAS scores decreased after radiotherapy as compared to before radiotherapy but there was no significant difference ($p>0.05$) in the patients in the pranayama and deep breathing exercise groups. In the fourth week, the VAS score of the pranayama (3.0) and deep breathing exercise (3.0) groups decreased in comparison to the VAS scores of the control group (5.0) and this decrease was more significant in the deep breathing group compared to the control group ($p=0.04$). In the fifth week of radiotherapy, although the VAS score of the control group (4.5) was higher in comparison to the pranayama (3.0) and deep breathing exercise (3.0) groups, there was no statistically significant difference ($p>0.05$) (Figure 3).

Table 2. Comparison of the patients' intra-group and inter-group total PFS and subdimension scores before and after radiotherapy (n=60)

| Piper Fatigue Scale Subdimension | Control Group | | Pranayama | | Deep Breathing | | Inter-group | |
|------------------------------------|-------------------|---------|-------------------|-----------|-------------------|-----------|-----------------|------|
| | Median | | Median | | Median | | KW ^a | P |
| | (minimum-maximum) | | (minimum-maximum) | | (minimum-maximum) | | | |
| Behavior Violence | | | | | | | | |
| Before radiotherapy | 3.0 | (0-6.1) | 2.6 | (0.1-6.1) | 2.5 | (0-5.1) | 0.31 | 0.85 |
| After radiotherapy | 4.2 | (0-8.8) | 2.7 | (0.1-8.4) | 3.4 | (0-7) | 2.39 | 0.30 |
| Intra-group Z^b p | -1.867 | 0.06 | -0.411 | 0.68 | -1.09 | 0.27 | | |
| Affectivity | | | | | | | | |
| Before radiotherapy | 3.7 | (0-7.6) | 4.4 | (0.4-7.6) | 2.9 | (0-8.8) | 2.58 | 0.27 |
| After radiotherapy | 4.2 | (0-8.6) | 3.5 | (0.2-8.2) | 3.4 | (0-7) | 4.11 | 0.12 |
| Intra-group Z p | -1.37 | 0.17 | -0.8 | 0.42 | -0.26 | 0.79 | | |
| Sensorial | | | | | | | | |
| Before radiotherapy | 3.3 | (0-6) | 4.4 | (0.2-8.2) | 3.2 | (0-5.6) | 3.79 | 0.15 |
| After radiotherapy | 4.2 | (0-8.6) | 4.0 | (0.6-9) | 3.7 | (0-6.4) | 1.21 | 0.54 |
| Intra-group Z p | -1.67 | 0.09 | -0.32 | 0.74 | -1.15 | 0.24 | | |
| Cognitive/Spiritual | | | | | | | | |
| Before radiotherapy | 3.1 | (0-7.8) | 3.4 | (0.1-6.1) | 2.1 | (0.7-7.1) | 0.92 | 0.63 |
| After radiotherapy | 4.2 | (0-8.6) | 3.4 | (0.8-8.3) | 2.9 | (0-6) | 2.84 | 0.24 |
| Intra-group Z p | -1.02 | 0.30 | -1.38 | 0.16 | -0.45 | 0.65 | | |
| Total | | | | | | | | |
| Before radiotherapy | 3.3 | (0-6.3) | 3.7 | (0.4-6.5) | 3.1 | (0-6.5) | 1.69 | 0.42 |
| After radiotherapy | 4.0 | (0-8.3) | 3.1 | (0.4-8) | 3.1 | (0-6.1) | 2.37 | 0.30 |
| Intra-group Z p | -1.71 | 0.08 | -0.40 | 0.68 | -0.84 | 0.39 | | |

^aKW Kruskal Wallis

^bZ Wilcoxon Test

In both the intra-group and inter-group comparison of all three groups no statistically significant difference in PFS total and subdimension scores before and after radiotherapy was found ($p > 0.05$) (Table 2).

There was no statistically significant difference in terms of insomnia VAS score of the pranayama and deep breathing exercises group before and after radiotherapy (respectively $p = 0.007$, $p = 0.027$). However, in the third week of radiotherapy, the insomnia VAS scores of the pranayama and deep breathing exercise groups showed a statistically significant decrease when compared to the control group ($p = 0.03$). In the fourth week of radiotherapy, the pranayama group showed a statistically significant decrease in VAS scores compared to the control and deep breathing exercise groups ($p = 0.009$). (Figure 4).

Table 3. Comparison of the patients' intra-group and inter-group total PSQI and subdimension scores before and after radiotherapy (n=60)

| PSQI subdimensions | Control Group | | Pranayama | | Deep Breathing | | Inter-group | |
|------------------------------------|-----------------------------|-------|-----------------------------|------|-----------------------------|-------|-----------------|-------|
| | Median (minimum-maximum) | | Median (minimum-maximum) | | Median (minimum-maximum) | | KW ^a | P |
| Subjective Sleep Quality | | | | | | | | |
| Before radiotherapy | 1.0 (0-2) | | 1.0 (0-2) | | 1.0 (0-3) | | 0,60 | 0,73 |
| After radiotherapy | 1.0 (0-3) | | 1.0 (0-2) | | 1.0 (0-2) | | 4.01 | 0.13 |
| Intra-group Z^b p | -2.00 | 0.04 | -1.001 | 0.31 | - 2.12 | 0.003 | | |
| Sleep Latency | | | | | | | | |
| Before radiotherapy | 1.0 (0-3) | | 1.0 (0-3) | | 2.0 (0-3) | | 1.96 | 0.37 |
| After radiotherapy | 1.0 (0-3) | | 1.0 (0-3) | | 1.5 (0-3) | | 5.88 | 0.053 |
| Intra-group Z p | -1.06 | 0.28 | -1.66 | 0.09 | -0.57 | 0.56 | | |
| Conventional Sleep Activity | | | | | | | | |
| Before radiotherapy | 1.0 (0-3) | | 3.0 (0-3) | | 1.5 (0-3) | | 2.44 | 0.29 |
| After radiotherapy | 1.0 (0-3) | | 2.0 (0-3) | | 1.5 (0-3) | | 0.06 | 0.96 |
| Intra-group Z p | -0.41 | 0.67 | -1.44 | 0.15 | -0.1 | 0.91 | | |
| Sleep disorder | | | | | | | | |
| Before radiotherapy | 1.0 (0-3) | | 1.0 (0-2) | | 1.0 (0-2) | | 0.14 | 0.93 |
| After radiotherapy | 1.0 (0-2) | | 1.0 (0-2) | | 1.0 (0-2) | | 1.39 | 0.49 |
| Intra-group Z p | -1.41 | 0.15 | -0.70 | 0.48 | -1.41 | 0.15 | | |
| Use of sleeping medication | | | | | | | | |
| Before radiotherapy | 0.0 (0-3) | | 0.0 (0-3) | | 0.0 (0-3) | | 1.06 | 0.58 |
| After radiotherapy | 0.0 (0-3) | | 0.0 (0-3) | | 0.0 (0-3) | | 0.47 | 0.78 |
| Intra-group Z p | -1.00 | 0.31 | -1.00 | 0.31 | -1.34 | 0.18 | | |
| Daytime Dysfunction | | | | | | | | |
| Before radiotherapy | 0.0 (0-2) | | 1.0 (0-3) | | 1.0 (0-2) | | 4.34 | 0.11 |
| After radiotherapy | 1.0 (0-3) | | 1.0 (0-2) | | 1.0 (0-3) | | 4.72 | 0.09 |
| Intra-group Z p | -2.94 | 0.003 | -0.08 | 0.93 | -0.83 | 0.40 | | |
| Total | | | | | | | | |
| Before radiotherapy | 6 (1-16) | | 9 (2-16) | | 7 (1-15) | | 1.62 | 0.44 |
| After radiotherapy | 8 (2-16) | | 7.5 (1-12) | | 8.5 (2-15) | | 0.88 | 0.64 |
| Intra-group Z p | -1.86 | 0.06 | -1.76 | 0.07 | -0.115 | 0.90 | | |

^aKruskal Wallis

^bWilcoxon Test

The comparison of the intra-group mean scores before and after radiotherapy showed that the PSQI subjective sleep quality negatively increased in the control group after radiotherapy when compared to before ($p=0.04$). In the deep breathing exercise group, subjective sleep quality positively increased after radiotherapy when compared to before ($p=0.03$). The study showed that daytime dysfunction negatively increased after radiotherapy in the control group ($p=0.003$) but the intra-group and inter-group

comparison showed no statistically significant difference in mean PSQI total scores and mean subdimension scores before and after radiotherapy (Table 3).

Discussion

The medical treatment of cancer-related fatigue, which leads to negative effects on quality of life and daily life activities, is very limited. Previous studies have shown that nonpharmacological methods can be effective in the management of fatigue [13-18]. There are studies showing that breathing exercises are effective in reducing fatigue in cancer patients [14, 25, 34, 47, 48]. Devi et al. determined that deep breathing exercises reduced fatigue in patients receiving radiotherapy [42].

In this research the degree of fatigue has increased in the controlled group thorough the end of the treatment (5th week) compared to the start of the radiotherapy. In both the trial groups, the degree of fatigue decreased at the end of the treatment compared to the start of the radiotherapy but there was no statistical significancy. The lack of significancy is thought to be caused by the fact that in the paranayama and breathing exercises group there were both patients with fatigue and without fatigue at the beginning of the study even though the degree of fatigue decreased. If only with patients with fatigue had been included I, the study would have been more effective therefore it is suggested that only the patients with fatigue must be included in the upcoming studies.

However, what distinguishes this study from other studies is that the breathing exercises were carried out individually accompanied by an observer. It is thought that the patients were more willing to do the breathing exercises and to perform these more efficiently because they were only 10 minutes and because the sessions were accompanied by an observer.

Disturbed sleep patterns and sleep problems can also be observed with fatigue in patients undergoing radiotherapy. Sleep disorders may cause patients to experience fatigue during the day, which negatively affects quality of life and creates difficulties in performing activities of daily life and maintaining social roles [9, 12, 49]. In the current study, both pranayama and deep breathing exercises significantly reduced insomnia at the end of treatment in the group comparison of insomnia severity. Mustian et al. determined that yoga practice involving pranayama, was effective in relieving insomnia in breast cancer patients who had completed treatment [36].

There have been no studies showing that only the pranayama or the breathing exercises increase the quality of sleep in the patients receiving radiotherapy even though it has been stated that the pranayama breathing exercises help decrease sleep disorders.

Both pranayama and deep breathing exercises are easy to learn. They can be used individually or as a group at anytime and anywhere and are exercises that can be done without creating stress on the person and without consuming too much energy [14, 42, 47, 48].

It is thought that the mind relaxes and parasympathetic activity is stimulated during the deep breathing exercises by focusing on breathing and therefore these exercises are effective in improving the quality of sleep. In this study it has been found that each of the different deep breathing exercises (pranayama and deep breathing techniques) have been effective in decreasing the fatigue and improving the quality of sleep but none predominates the other.

In this context, the patients participating in the study stated that they easily learned the exercises and that they were willing to continue the exercises in the future. In the current study, a quiet environment and no special equipment other than a chair were required for the application, and no side effects developed during or after the application.

The limitations of the research: the study has been conducted in one hospital for a duration of 10 minutes under observation after 25 sessions of radiotherapy. The fact that the exercises were done under observation and one to one is the strong point of the study. However the fact that whether patients have continued with the exercises at home is unknown constitutes a weak point. The study has been conducted on the patients receiving radiotherapy for breast cancer. So the results obtained are only related to the patients with breast cancer. In this study, feedback related to fatigue and sleep quality is only based on self-feedback and that is another limitation of the study.

Conclusion

The study determined that fatigue and sleep problems exist in breast cancer patients before radiotherapy and that these symptoms increased with radiotherapy treatment. In this study, both pranayama and deep breathing exercises were found to be effective in reducing fatigue and reducing insomnia, even when performed for a short time (10 minutes a day). In line with these results, we suggest conducting studies with widespread pranayama and deep breathing exercises, which are easy to learn, do not require a lot of energy/effort and can be done easily anytime and anywhere in oncology clinics and to conduct further studies on the use of breathing exercises of different lengths and durations in the prevention and treatment of fatigue and insomnia in breast cancer and other types of cancer.

Declarations

Funding: No funds, grants, or other support was received.

Conflicts of interest The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethics approval: Procedures performed in this study were in accordance with the 1964 Helsinki Declaration and its later amendments and in accordance with the ethical standards of the institutional review board and independent ethics committee.

Consent to participate for publication: All participants gave written informed consent prior to enrolment and randomisation.

Availability of data and material and code availability: The authors have full control of all primary data, material and code availability and agree to allow the journal to review their data if requested.

Authors' contributions: **Fatma Gündođdu:** Conceptualization, Methodology, Formal analysis, Data curation, Investigation, Writing - original draft, Writing - review & editing, Visualization. **Sema Koçařlı:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - review & editing

Acknowledgments: The authors would like to thank all who participated in this study.

References

1. Banerjee C, Breitbart WS, Carpenter KM, et al (2018) NCCN Guidelines Version 2.2018 Panel Members Cancer-Related Fatigue NCCN.org <https://oncolife.com.ua/doc/nccn/fatigue.pdf>. 8 Aralık 2018
2. Biswal BM, Kumaraswamy N, Mukhtar F (2004) Prevalence of fatigue among cancer patients undergoing external radiotherapy. *Southeast Asian J Trop Med Public Health* 35(2):463-7
3. Janaki MG, Kadam RA, Mukesh S, Nirmala S, Ponni A, Ramesh BS, RajeevAG (2010) Magnitude of fatigue in cancer patients receiving radiotherapy and its short term effect on quality of life. *Journal of Cancer Research and Therapeutics* 6:22-26. <https://doi.org/10.4103/0973>
4. Bower JE, Garet D, Sternlieb B, Ganz PA, Irwin MR, Olmstead R, Greendale G (2012) Yoga for persistent fatigue in breast cancer survivors: A randomized controlled trial. *Cancer* 118(15): 3766–3775. <https://doi.org/10.1002/cncr.26702>
5. Horneber M, Fischer I, Dimeo F, Ruffer JU, Weis J (2012) Cancer-Related Fatigue, Epidemiology, Pathogenesis, Diagnosis, and Treatment. *Dtsch Arztebl Int* 109 (9): 161-172. <https://doi:10.3238/arztebl.2012.0161>
6. Manir KS, Bhadra K, Kumar G, Manna A, Patra NB, Sarkar SK (2012) Fatigue in breast cancer patients on adjuvant treatment: Course and prevalence. *Indian Journal of Palliative Care* 18(2):109-116. <https://doi:10.4103/0973-1075.100826>
7. Bower JE, Bak K, Berger A, et al (2014) Screening, assesment, and management of fatigue in adult survivors of cancer: an American Society of Clinical Oncology clinical practice guideline adaptation. *J Clin Oncol* 32(17):1840-50
8. Geinitz H, Zimmermann FB, Stoll P, et al (2001) Fatigue, serum cytokine levels, and blood cell counts during radiotherapy of patients with breast cancer. *Int J Radiat Oncol Biol Phys* 51(3):691-698
9. Roscoe JA, Kaufman ME, Rusby MSE, et al (2007) Cancer-Related Fatigue and Sleep Disorders. *The Oncologist* 12(1):35–42
10. Dirksen SR, Epstein DR (2008) Efficacy of an insomnia intervention on fatigue, mood and quality of life in breast cancer survivors. *J Adv Nurs* 61(6):664-75. doi:10.1111/j.1365-2648.2007.04560.x

11. Jim HS, Small B, Faul LA, Franzen J, Apte S, Jacobsen PB (2011) Fatigue, depression, sleep, and activity during chemotherapy: daily and intraday variation and relationships among symptom changes. *Annals of behavioral medicine: a publication of the Society of Behavioral Medicine* 42(3):321–333. doi:10.1007/s12160-011-9294-9
12. [Holliday EB](#), [Dieckmann NF](#), [McDonald TL](#), Hung AY, Thomas CR, Wood LJ (2016) Relationship between fatigue, sleep quality and inflammatory cytokines during external beam radiation therapy for prostate cancer: A prospective study. *Radiother Oncol* 118 (1):105-111. doi:10.1016/j.radonc.2015.12.015
13. Greenlee H, Du-pont-Reyes MJ, Balneaves LG, et al(2017) Clinical practice guidelines on the evidence-based use of integrative therapies during and following breast cancer treatment. *CA Cancer J Clin* 67(3):194–232 doi:10.3322/caac.21397
14. Chakrabarty J, Vidyasagar M, Fernandes D, Joisa G, Varghese P, Mayya S (2015) Effectiveness of pranayama on cancer-related fatigue in breast cancer patients undergoing radiation therapy: A randomized controlled trial. *Int J Yoga*. 8(1):47-53 doi:10.4103 / 0973-6131.146062
15. Chandwani KD, Perkins G, Nagendra HR et al Randomized controlled trial of yoga in women with breast cancer undergoing radiotherapy. *J Clin Oncol* 32:1058-1065 doi:10.1200/JCO.2012.48.2752
16. [Bennett S](#), [Pigott A](#), [Beller EM](#), [Haines T](#), [Meredith P](#), [Delaney C](#) (2016) Educational interventions for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev* 11:1-62 doi: 10.1002/14651858.CD008144.pub2
17. [Hilfiker R](#), [Meichtry A](#), [Eicher M](#) (2018) Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: a systematic review incorporating an indirect-comparisons meta-analysis. *Br J Sports Med* 52(10): 651–658 doi: 10.1136/bjsports-2016-096422
18. [Barsevick AM](#), [Dudley W](#), [Beck S](#), [Sweeney C](#), [Whitmer K](#), [Nail L](#) (2004) A randomized clinical trial of energy conservation for patients with cancer-related fatigue. *Cancer* 100(6):1302-10 doi:10.1002/cncr.20111
19. [Sadja J](#), [Mills PJ](#) (2013) Effects of Yoga Interventions on Fatigue in Cancer Patients and Survivors: A Systematic Review of Randomized Controlled Trials [Explore: The Journal of Science and Healing](#) 9(4): 232–243 doi: 10.1016/j.explore.2013.04.005
20. [Manaf A](#) (2006) *Yoga, Pranayama, Breath and Energy Control*. first ed., Im Publication 33–70. ISBN: 975-7270-34-2
21. [Cramer H](#), [Lauche R](#), [Klose P](#), [Lange S](#), [Langhorst J](#), [Dobos GJ](#) (2017) Yoga for improving health related quality of life, mental health and cancer related symptoms in women diagnosed with breast cancer. *Cochrane Database Syst Rev* 3:1-201 doi:10.1002/14651858.CD010802.pub2
22. [Banasik J](#), [Williams H](#), [Haberman M](#), [Blank SE](#), [Bendel R](#) (2011) Effect of Iyengar yoga practice on fatigue and diurnal salivary cortisol concentration in breast cancer survivors. *J Am Acad Nurse Pract* 23(3):135-42 doi:10.1111/j.1745-7599.2010.00573.x

23. Vadiraja HS, Rao MR, Nagarthna R et al (2009) Effects of yoga program on quality of life and affect in early breast cancer patients undergoing adjuvant radiotherapy: A randomized controlled trial. *Complementary Therapies in Medicine* 17:274-280 doi:10.1016/j.ctim.2009.06.004
24. Zeichner SB, Zeichner RL, Gogineni K, Shatil S, Ioachimescu O (2017) Cognitive Behavioral Therapy for Insomnia, Mindfulness, and Yoga in Patients With Breast Cancer with Sleep Disturbance: A Literature Review. *Breast Cancer: Basic and Clinical Research* 11:1-11 doi:10.1177/1178223417745564
25. Dhruva A, Miaskowski C, Abrams D et al (2012) Yoga Breathing for Cancer Chemotherapy–Associated Symptoms and Quality of Life: Results of a Pilot Randomized Controlled Trial. *The Journal Of Alternative And Complementary Medicine* 18(5):473–479 doi: 10.1089/acm.2011.0555
26. Zaccaro A, Piarulli A, Laurino M et al (2018) How Breath-Control Can Change Your Life: A Systematic Review on Psycho-Physiological Correlates of Slow Breathing, *Front Hum Neurosci* 12:1-15 doi : 10.3389/fnhum.2018.00353
27. Manjunatha U, Bhat JS, Radish KB et al (2018) Effect of Bhramari Pranayama on the Acoustic and Aerodynamic Parameters of Voice in Normophonic Females. *Evid Based Complement Alternat Med e-CAM* 1-7 doi: 10.1155/2018/4176803
28. Lin PJ, Peppone LJ, Janelins MC et al (2018) Yoga for the Management of Cancer Treatment-Related Toxicities. *Curr Oncol Rep* 20(1):5 doi: 10.1007/s11912-018-0657-2
29. Peterson CT, Bauer SM, Chopra D (2017) Effects of Shambhavi Mahamudra Kriya, a Multicomponent Breath-Based Yogic Practice (Pranayama), on Perceived Stress and General Well-Being. *J Evid Based Complementary Altern Med*. 22(4): 788–797 doi: 10.1177/2156587217730934
30. Şinik E (2017) Nose breathing guide for daily life: life is hidden in a breath. first ed. LİBROS publication 147-195 ISBN: 978-605-9151-16-0
31. Tian L, Lin L, Li HL et al (2016) Prevalence and Associated Factors of Cancer-Related Fatigue Among Cancer Patients in Eastern China. *Oncologist* 21(11): 1349–1354 doi: 10.1634/theoncologist.2015-0537
32. Ma X, Yue ZQ, Gong ZQ et al (2017) The Effect of Diaphragmatic Breathing on Attention, Negative Affect and Stress in Healthy Adults. *Front Psychol* 8:874 doi:10.3389/fpsyg.2017.00874
33. Mulhaeriah, Afiyanti Y, Achmad EK, Sangkala MS (2018) Effectiveness of Relaxation Breathing Exercise on fatigue in gynecological cancer patients undergoing chemotherapy. *International Journal of Nursing Sciences*5(4):331-335 doi:10.1016/j.ijnss.2018.09.004
34. Yokogawa M, Kurebayashi T, Ichimura T, Nishino M, Miaki H, Nakagawa T (2018) Comparison of two instructions for deep breathing exercise: non-specific and diaphragmatic breathing. *J Phys Ther Sci* 30(4): 614–618
35. Jerath R, Beveridge C, Barnes V (2019) Self-Regulation of Breathing as an Adjunctive Treatment of Insomnia. *Front Psychiatry* 9:780 doi:10.3389/fpsyg.2018.00780
36. Mustian KM, Sprod LK, Janelins M et al (2013) Multicenter randomized controlled trial of yoga for sleep quality among cancer survivors. *Clin Oncol* 31(26):3233-41 doi:10.1200/JCO.2012.43.7707

37. Bower JE, Ganz PA, Aziz N (2005) Altered cortisol response to psychologic stress in breast cancer survivors with persistent fatigue. *Psychosom Med* 67(2):277-80
doi:10.1097/01.psy.0000155666.55034.c6
38. Yağlı NV, Şener G, Arıkan H et al (2015) Do yoga and aerobic exercise training have impact on functional capacity, fatigue, peripheral muscle strength, and quality of life in breast cancer survivors? *Integr Cancer Ther* 14(2):125-32 doi:10.1177/1534735414565699
39. Pal GK, Agarwal A, Karthik S, Pal P, Nanda N (2014) Slow yogic breathing through right and left nostril influences sympathovagal balance, heart rate variability, and cardiovascular risks in young adults. *N Am J Med Sci* 6(3):145-151. doi:10.4103/1947-2714.128477
40. Agarwal RP, Maroko AA (2018) Yoga into Cancer Care: A Review of the Evidence-based Research. *Int J Yoga* 11(1):3-29. doi:10.4103/ijoy.IJOY_42_17
41. Saoji AA, Raghavendra BR, Manjunath NK (2019) Effects of yogic breath regulation: A narrative review of scientific evidence. *Journal of Ayurveda and Integrative Medicine* 10:50-58
doi:10.1016/j.jaim.2017.07.008
42. Devi R, Saharia HK (2016) Effectiveness of programmed deep breathing exercises on reducing the level of fatigue during external radiation therapy in cancer patients. *J. Evolution Med. Dent Sci* 5(38):2340-2344 doi:10.14260/jemds/2016/543
43. Piper BF, Dibble SL, Dodd MJ et al (1998) The revised piper fatigue scale: Psychometric evaluation in women with breast cancer, *Oncol Nurs Forum* 25:677-84
44. Can G, Durna Z (2001) Assessment of fatigue and care needs in patients with breast cancer. Phd Thesis Istanbul University Graduate School of Health Sciences 1-130 doi: 108382
45. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ (1989) The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 28(2):193-213
46. Ağargün MY, Kara H, Anlar O (1996) The Validity and Reliability of the Pittsburgh Sleep Quality Index. *Turkish Journal of Psychiatry* 7:107-115
47. Kim SD, Kim HS (2005) Effects of a relaxation breathing exercise on fatigue in haemopoietic stem cell transplantation patients. *Journal of Clinical Nursing* 14:51-55
48. Hayama Y, Inoue T (2012) The effects of deep breathing on 'tension-anxiety' and fatigue in cancer patients undergoing adjuvant chemotherapy. *Complement Ther Clin Pract* 18(2):94-98
doi:10.1016/j.ctcp.2011.10.001
49. Rocha SR, Santos MCL, Lopes MVO et al (2018) Accuracy of the defining characteristics of the nursing diagnosis for fatigue in women under radiotherapy. *Rev Bras Enferm.* 71(3):1445-52
doi:10.1590/0034-7167-2017-0549vv