

# Overexpression of I $\kappa$ B $\alpha$ in cardiomyocytes alleviates hydrogen peroxide-induced apoptosis and autophagy by inhibiting NF- $\kappa$ B activation

**Min Han**

Xinjiang Medical University Affiliated First Hospital

**Xiao-Cui Chen**

Xinjiang Medical University Affiliated First Hospital

**Ming-Hui Sun**

Xinjiang Medical University

**Min-Tao Gai**

Xinjiang Medical University affiliated first hospital

**Yi-Ning Yang**

Xinjiang Medical University Affiliated First Hospital

**Xiao-Ming Gao**

Xinjiang Medical University affiliated first hospital

**Xiang Ma**

Xinjiang Medical University affiliated first hospital

**Bang-Dang Chen**

Xinjiang Medical University Affiliated First Hospital

**Yi-Tong Ma** (✉ [myt\\_xj@sina.com](mailto:myt_xj@sina.com))

---

## Research

**Keywords:** nuclear factor kappa B, inhibitor of kappa B alpha, apoptosis, autophagy, adeno-associated virus serotype 9, oxidative stress, cardiomyocytes

**DOI:** <https://doi.org/10.21203/rs.3.rs-16137/v3>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** Inflammation and oxidative stress play predominant roles in the initiation and progression of ischaemia/reperfusion (I/R) injury, with nuclear factor kappa B (NF- $\kappa$ B) serving as a crucial mediator. Overexpression of the inhibitor of  $\kappa$ B alpha (I $\kappa$ B $\alpha$ ) gene is hypothesized to have protective effects against apoptosis and autophagy in cardiomyocytes subjected to hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by inhibiting the NF- $\kappa$ B pathway.

**Methods:** The I $\kappa$ B $\alpha$ <sup>S32A, S36A</sup> gene was transfected via adeno-associated virus serotype 9 (AAV9) delivery into neonatal rat ventricular cardiomyocytes (NRVMs) prior to H<sub>2</sub>O<sub>2</sub> treatment. NRVMs were divided into control, H<sub>2</sub>O<sub>2</sub>, GFP +H<sub>2</sub>O<sub>2</sub>, I $\kappa$ B $\alpha$ +H<sub>2</sub>O<sub>2</sub>, and pyrrolidine dithiocarbamate (PDTC)+H<sub>2</sub>O<sub>2</sub> groups. Nuclear translocation of the NF- $\kappa$ B p65 subunit was evaluated by immunofluorescence and Western blotting. Cell viability was assessed by Cell Counting Kit-8 assay. Supernatant lactate dehydrogenase (LDH) and intracellular malondialdehyde (MDA) were measured to identify H<sub>2</sub>O<sub>2</sub>-stimulated cytotoxicity. Apoptosis was determined by Annexin V-PE/7-AAD staining, and the mitochondrial membrane potential ( $\Delta\Psi$ m) was detected by JC-1 staining. Western blotting was used to detect apoptosis- and autophagy-related proteins.

**Results:** I $\kappa$ B $\alpha$  transfection significantly increased cell viability and  $\Delta\Psi$ m but decreased the supernatant LDH and cellular MDA levels in cardiomyocytes exposed to H<sub>2</sub>O<sub>2</sub>. Meanwhile, I $\kappa$ B $\alpha$  overexpression decreased H<sub>2</sub>O<sub>2</sub>-induced apoptosis by upregulating the Bcl-2/Bax ratio and reduced autophagy by downregulating the expression of Beclin-1 and the LC3-II/LC3-I ratio. These effects partly accounted for the ability of I $\kappa$ B $\alpha$  to inhibit the NF- $\kappa$ B signalling pathway, as evidenced by decreases in p65 phosphorylation and nuclear translocation. Indeed, the effects of inactivation of NF- $\kappa$ B signalling with the specific inhibitor PDTC resembled the cardioprotective effects of I $\kappa$ B $\alpha$  during H<sub>2</sub>O<sub>2</sub> stimulation.

**Conclusion:** I $\kappa$ B $\alpha$  overexpression can ameliorate H<sub>2</sub>O<sub>2</sub>-induced apoptosis, autophagy, oxidative injury, and  $\Delta\Psi$ m loss through inhibition of the NF- $\kappa$ B signalling pathway. These findings suggest that I $\kappa$ B $\alpha$  transfection can result in successful resistance to oxidative stress-induced damage by inhibiting NF- $\kappa$ B activation, which may provide a potential therapeutic target for the prevention of myocardial I/R injury.

## Introduction

Acute myocardial infarction (AMI) is the leading cause of death worldwide, and reperfusion therapy is the most effective treatment for AMI [1]. Paradoxically, the process of myocardial reperfusion also induces a series of adverse cardiac events such as inflammation, necrosis, apoptosis and autophagy, ultimately leading to myocardial ischaemia/reperfusion (I/R) injury [2]. Recent evidence has suggested that excessive inflammation and oxidative stress play predominant roles in the initiation and progression of I/R injury [3, 4].

Nuclear factor kappa B (NF- $\kappa$ B) is an inflammatory inducer and redox-sensitive transcription factor in most cell types [5]. The p65/50 heterodimer, the most common pattern of NF- $\kappa$ B dimer, normally exists as a component of inactive cytoplasmic complexes bound to the inhibitor of  $\kappa$ B alpha (I $\kappa$ B $\alpha$ ). Upon stimulation, I $\kappa$ B $\alpha$  is phosphorylated and undergoes ubiquitylation and proteasomal degradation, subsequently leading to phosphorylation and nuclear translocation of the NF- $\kappa$ B p65 subunit [6]. Activated NF- $\kappa$ B then initiates the expression of corresponding target genes, many of which may regulate apoptosis, inflammation and autophagy [7].

However, whether NF- $\kappa$ B is protective or detrimental for cardiomyocyte apoptosis remains controversial [8]. Notably, our previous study indicated that the p65 ribozyme could prevent cell apoptosis in H9C2 cardiomyocytes exposed to hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) [9]. Autophagy, an evolutionarily conserved form of “self-digestion”, plays dual roles in the heart [10]. Recent studies on autophagy have shown both the protective [11] and deleterious [12] effects of autophagy in cardiomyocytes against oxidative stress. Evidence has revealed a strong correlation between modulation of NF- $\kappa$ B and the autophagic response [13, 14]. In addition, cross-talk between autophagy and apoptosis has been noted [15], and NF- $\kappa$ B is known to mediate the balance between autophagy and apoptosis [16].

Therefore, NF- $\kappa$ B activation is thought to be the key point of I/R injury; thus, inhibiting NF- $\kappa$ B may be a targeted therapy for I/R injury. Phosphorylation of I $\kappa$ B $\alpha$ , the key inhibitor of the canonical NF- $\kappa$ B pathway, at Ser 32 and Ser 36 is necessary for its degradation, and any mutation of these two serine residues blocks I $\kappa$ B $\alpha$  degradation [6]. Recently, adeno-associated virus serotype 9 (AAV9) was demonstrated to be the best gene carrier due to its high efficiency in the heart [17]. H<sub>2</sub>O<sub>2</sub>, a common reactive oxygen species (ROS), is generally utilized to mimic I/R injury *in vitro* [12]. Thus, the I $\kappa$ B $\alpha$ <sup>S32A, S36A</sup> gene was transfected into cardiomyocytes via AAV9-mediated delivery to investigate the role of inhibition of the NF- $\kappa$ B pathway in H<sub>2</sub>O<sub>2</sub>-induced apoptosis and autophagy. Pyrrolidine dithiocarbamate (PDTC), a specific inhibitor of NF- $\kappa$ B, was used as a positive control in this study.

## Materials And Methods

### Ethics statement

The experimental protocol was approved by the Ethics Committee of the First Affiliated Hospital of Xinjiang Medical University (No. IACUC-20180223-69). One- to three-day-old neonatal Sprague-Dawley (SD) rats were purchased from the Experimental Animal Center of Xinjiang Medical University and handled in accordance with the recommendations in the Guidelines for the Care and Use of Laboratory Animals of the National Institutes of Health.

### Reagents

Briefly, rabbit anti-Bax polyclonal antibody (#2772) and rabbit anti-p65 (#8242), anti-p-p65 (#3033), anti-I $\kappa$ B $\alpha$  (#4812), anti-GFP (#2956), anti-Becn-1 (#3495), and anti-LC3  $\alpha/\beta$  (#12741) monoclonal antibodies were all obtained from Cell Signalling Technology (Danvers, MA, USA). Rabbit anti-Bcl-2 (ab196495), anti-

Histone H3 (ab1791) and anti- $\beta$ -actin (ab8227) polyclonal antibodies and horseradish peroxidase (HRP)-conjugated anti-rabbit secondary antibody (ab205718) were obtained from Abcam (Cambridge, UK). RIPA buffer and Halt™ Protease and Phosphatase Inhibitor Cocktail were obtained from Thermo Fisher Scientific (Waltham, MA, USA). Enhanced chemiluminescence (ECL) reagent and JC-1 were obtained from Millipore (Bedford, MA, USA). Trypsin, PDTC and bromodeoxyuridine (BrdU) were obtained from Sigma (St. Louis, MO, USA). Dulbecco's modified Eagle's medium (DMEM), foetal bovine serum (FBS), and penicillin-streptomycin solution were obtained from Gibco (Grand Island, NY, USA). Collagenase II was obtained from Worthington (Minnesota, USA). H<sub>2</sub>O<sub>2</sub> was obtained from Sangon (Shanghai, China).

## Vector design

Recombinant AAV-9 vectors generated by a recombinant baculovirus (rBac)-based system in SF9 cells as previously described were purchased from Virovek (Hayward, CA, USA) [18]. The recombinant AAV9 vectors were packaged as double-stranded DNA and contained the enhanced green fluorescent protein (eGFP) gene (dsAAV9-GFP) or the I $\kappa$ B $\alpha$ <sup>S32A, S36A</sup> gene (dsAAV9-I $\kappa$ B $\alpha$ ) driven by the human cytomegalovirus (CMV) promoter.

## Isolation and culture of rat cardiomyocytes

The protocol for the isolation and purification of neonatal rat ventricular cardiomyocytes (NRVMs) was reported in our previous study [19]. Briefly, the hearts of 1- to 3-day-old neonatal SD rats were dissected and digested with 0.1% trypsin and 0.08% collagenase II. Following differential adhesion twice for 50 min each time, nonadherent cells were resuspended and cultivated in high-glucose DMEM containing 10% FBS, 1% penicillin-streptomycin, and 0.1 mM BrdU for 48 h. The medium was replaced every 48 h.

## AAV9 transfection of cardiomyocytes

After 48 h of culture, NRVMs were transfected with dsAAV9-GFP or dsAAV9-I $\kappa$ B $\alpha$  as previously described [12]. Briefly, cells were first transfected with dsAAV9 (multiplicity of infection, MOI = 5 $\times$ 10<sup>6</sup> vg/cell) in serum-free medium, and then DMEM at an equal volume containing 20% FBS, 2% penicillin-streptomycin and 0.2 mM BrdU was added to every dish 3 h later. Images showing GFP were captured using a fluorescence inverted microscope (Leica DMI4000B, Wetzlar, Germany), and the green fluorescence intensities were analysed using ImageJ software (National Institutes of Health, NY, USA).

## Experimental design and cell grouping

The experiment was designed to explore whether AAV9-delivered I $\kappa$ B $\alpha$ <sup>S32A, S36A</sup> gene transfection could protect cardiomyocytes against H<sub>2</sub>O<sub>2</sub>-induced apoptosis and autophagy via inhibition of NF- $\kappa$ B activation. Cardiomyocytes were starved with serum-free DMEM for 12 h to ensure cell synchronization before H<sub>2</sub>O<sub>2</sub> stimulation. The experimental cardiomyocytes were randomly divided into 5 groups as follows: (1) the control group, which contained primary cardiomyocytes cultivated under normal conditions; (2) the H<sub>2</sub>O<sub>2</sub> control group (H<sub>2</sub>O<sub>2</sub>): the model control group, which contained primary

cardiomyocytes subjected to 100  $\mu\text{M}$   $\text{H}_2\text{O}_2$  alone [12]; (3) the GFP control group (GFP): the vector control group, which contained primary cardiomyocytes transfected with dsAAV9-GFP virus for 5 days before being subjected to 100  $\mu\text{M}$   $\text{H}_2\text{O}_2$ ; (4) the I $\kappa$ B $\alpha$  treatment group (I $\kappa$ B $\alpha$ ): the treatment group, which contained primary cardiomyocytes transfected with dsAAV9-I $\kappa$ B $\alpha$  virus for 5 days before being subjected to 100  $\mu\text{M}$   $\text{H}_2\text{O}_2$ ; and (5) the PDTC treatment group (PDTC): the positive control group, which contained primary cardiomyocytes pretreated with 100  $\mu\text{M}$  PDTC for 60 min before being subjected to 100  $\mu\text{M}$   $\text{H}_2\text{O}_2$ .

### **Measurement of cardiomyocyte viability and cytotoxicity**

The Cell Counting Kit-8 (CCK-8; Dojindo, Japan) assay was used to assess cell viability. In brief,  $2 \times 10^4$  cells were seeded into each well of a 96-well plate and transfected with GFP or I $\kappa$ B $\alpha$  for five days. After the cells were exposed to  $\text{H}_2\text{O}_2$ , 10  $\mu\text{L}$  of CCK-8 stock solution was added to each well, followed by incubation at 37  $^\circ\text{C}$  for 2 h. The absorbance at 450 nm was measured with a GO microplate spectrophotometer (Thermo Fisher Scientific). The extent of cell death was determined by quantifying lactate dehydrogenase (LDH) released into the culture supernatant with an LDH kit (Jiancheng Bioengineering Institute, Nanjing, China). Intracellular malondialdehyde (MDA), an indicator of oxidative injury, was also measured using an MDA assay kit (Jiancheng Bioengineering Institute).

### **Flow cytometry analysis**

Cell apoptosis was measured using PE Annexin V Kit I (BD Biosciences, NJ, USA). Briefly, cells were collected and resuspended in  $1 \times$  binding buffer. Then, the solution ( $1 \times 10^5$  cells) was supplemented with 5  $\mu\text{L}$  of PE Annexin V and 7-AAD and incubated in the dark for 15 min at room temperature. Apoptotic cells were identified by flow cytometry (Beckman Coulter, CA, USA). All the experiments were performed in triplicate.

### **Western blotting analysis**

Nuclear and cytoplasmic proteins were extracted following the instructions of a Nuclear and Cytoplasmic Extraction Kit (Thermo Fisher Scientific, USA). Total proteins were extracted with RIPA buffer containing [Halt™ Protease and Phosphatase Inhibitor Cocktail](#). Phosphorylated p65 in the total lysate and the nuclear p65 to cytosolic p65 ratio were both detected to identify activation of the NF- $\kappa$ B signalling pathway [20]. Equal amounts of protein were loaded and separated on precast SDS-PAGE gels (Invitrogen, Grand Island, NY, USA) and transferred to Millipore PVDF membranes. After blocking with 5% skim milk, the membranes were blotted overnight with specific primary antibodies against p65 (1:1000), p-p65 (1:500), I $\kappa$ B $\alpha$  (1:1000), Bax (1:1000), GFP (1:1000), Beclin-1 (1:1000), LC3  $\square/\square$  (1:1000), Bcl-2 (1:1000), Histone H3 (1:1000), and  $\beta$ -actin (1:1000) at 4  $^\circ\text{C}$ , followed by incubation with anti-rabbit HRP secondary antibody (1:5000) at room temperature for 2 h. ECL solution was added to the membranes to visualize signals.  $\beta$ -actin and Histone H3 were regarded as loading controls. Images were captured and analysed by Image Lab 4.0 software (Bio-Rad Laboratories, Hercules, CA, USA).

## Immunofluorescence

Immunofluorescence was employed to identify H<sub>2</sub>O<sub>2</sub>-induced nuclear translocation of the NF- $\kappa$ B p65 subunit in cardiomyocytes. Briefly, 2×10<sup>5</sup> cells were seeded into confocal dishes. After H<sub>2</sub>O<sub>2</sub> treatment, cardiomyocytes were fixed with 4% paraformaldehyde for 20 min and permeabilized with 0.25% Triton X-100 for 10 min. After blocking with 1% BSA for 1 h, cells were probed overnight with anti-p65 antibody (1:200) at 4 °C and incubated with Alexa Fluor 594-labelled secondary antibody (Invitrogen, 1:200, labelled with red fluorescence) for 2 h at room temperature, followed by 10 min of DAPI staining of nuclei (labelled with blue fluorescence). Signals were detected using a confocal spectral microscope (Leica SP8, Germany).

## Measurement of the mitochondrial membrane potential

JC-1 is an ideal fluorescent probe used to detect the mitochondrial membrane potential ( $\Delta\Psi$ m) in cardiomyocytes. Briefly, a 10 nmol/L JC-1 working solution was prepared prior to use, and cardiomyocytes were stained at 37 °C in the dark for 15 min. Cells doubly stained with JC-1 were visible by either green or red fluorescence. Fluorescence images and intensities were obtained using a fluorescence microscope and ImageJ software. Generally,  $\Delta\Psi$ m is represented by the red to green fluorescence ratio, which decreases in proportion with the severity of cell injury.

## Statistical analysis

All statistical analyses were performed with SPSS 22.0 (SPSS, Inc., Chicago, IL, USA). Data are presented as the mean  $\pm$  SEM. Multiple comparisons were carried out using one-way analysis of variance (ANOVA) followed by Bonferroni's *post hoc* test. A value of  $P < 0.05$  indicated statistical significance.

# Results

## H<sub>2</sub>O<sub>2</sub>-induced activation of NF- $\kappa$ B in NRVMs

The results indicated that H<sub>2</sub>O<sub>2</sub> elicited time-dependent I $\kappa$ B $\alpha$  degradation and p65 translocation after the NRVMs were incubated with 100  $\mu$ M H<sub>2</sub>O<sub>2</sub> for different durations (0, 15, 30, and 60 min). (Fig. 1A-C). The ratio of nuclear p65 to cytosolic p65 peaked at 60 min. Consistent with the nuclear translocation of p65, the level of p-p65/p65 increased following H<sub>2</sub>O<sub>2</sub> stimulation with the incubation time (Fig. 1D and E) and was highest at 60 min. Thus, treatment with 100  $\mu$ M H<sub>2</sub>O<sub>2</sub> for 60 min was used in the following experiments.

## Efficiency of I $\kappa$ B $\alpha$ transfection in NRVMs

As shown in Fig. 1F, the green fluorescence signal was robust, and the dsAAV9-GFP transfection efficiency in NRVMs reached more than 70%. Western blotting analysis showed that the GFP protein was

more highly expressed in the GFP group than in the other groups, while the I $\kappa$ B $\alpha$  protein level was significantly elevated in the I $\kappa$ B $\alpha$  group compared with the control and GFP groups (Fig. 1G-I).

### **I $\kappa$ B $\alpha$ protected cardiomyocytes from H<sub>2</sub>O<sub>2</sub>-induced apoptosis**

The proportion of apoptotic cells in the control group was  $7.0 \pm 1.5\%$ . After treatment with  $100 \mu\text{M}$  H<sub>2</sub>O<sub>2</sub>, the apoptotic rates of cardiomyocytes in the H<sub>2</sub>O<sub>2</sub> group and GFP group increased to  $21.20 \pm 0.95\%$  and  $19.97 \pm 0.97\%$ , respectively, which were decreased by I $\kappa$ B $\alpha$  or PDTC pretreatment (Fig. 2A). Indeed, compared with the levels in the control group, the anti-apoptotic protein Bcl-2 was downregulated, but the pro-apoptotic protein Bax was upregulated in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>, leading to a higher Bax/Bcl-2 ratio, but this effect was completely abolished by pretreatment with I $\kappa$ B $\alpha$  or PDTC (Fig. 2B).

### **I $\kappa$ B $\alpha$ protected cardiomyocytes from H<sub>2</sub>O<sub>2</sub>-induced cell injury**

Compared to that in the control group,  $\Delta\Psi\text{m}$  was significantly decreased in the H<sub>2</sub>O<sub>2</sub> control group, but this decrease was rescued by I $\kappa$ B $\alpha$  or PDTC treatment (Fig. 3A). Additionally, H<sub>2</sub>O<sub>2</sub> treatment significantly decreased cell viability but elevated supernatant LDH and intracellular MDA levels; however, these changes were reversed by I $\kappa$ B $\alpha$  or PDTC treatment (Fig. 3B-D).

### **I $\kappa$ B $\alpha$ suppressed H<sub>2</sub>O<sub>2</sub>-induced NF- $\kappa$ B activation and autophagy in NRVMs**

Compared with the control group, H<sub>2</sub>O<sub>2</sub> treatment significantly elicited p65 translocation, and these changes were successfully reversed by I $\kappa$ B $\alpha$  or PDTC pretreatment (Fig. 4A and B). Consistently, H<sub>2</sub>O<sub>2</sub> increased p-p65/p65 ratio in NRVMs, but I $\kappa$ B $\alpha$  or PDTC dramatically downregulated the H<sub>2</sub>O<sub>2</sub>-induced expression of p-p65. Meanwhile, Beclin-1 and the LC3-II/LC3-I ratio, the autophagy-associated markers, were markedly upregulated in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>, whereas these effects were inhibited by I $\kappa$ B $\alpha$  or PDTC treatment (Fig. 4C).

## **Discussion**

This study shows that I $\kappa$ B $\alpha$  degradation and NF- $\kappa$ B activation occurred in a time-dependent manner in NRVMs subjected to H<sub>2</sub>O<sub>2</sub>. Cells treated with H<sub>2</sub>O<sub>2</sub> showed reductions in cell viability and  $\Delta\Psi\text{m}$  but elevations in LDH and MDA levels, apoptosis and autophagy. I $\kappa$ B $\alpha$  transfection or PDTC pretreatment ameliorated H<sub>2</sub>O<sub>2</sub>-induced cell injury by inhibiting NF- $\kappa$ B activation.

I/R injury severely attenuates the benefit of revascularization after AMI and has therefore become an important focus of cardiovascular research [2]. The inflammatory response induced by AMI is essential for heart repair, but excessive generation of ROS and inflammation following reperfusion therapy exacerbate heart damage [21].

The NF- $\kappa$ B signalling pathway plays key roles in the inflammatory response, oxidative stress, apoptosis, and autophagy in the heart [8]. Phosphorylation and nuclear translocation of the p65 subunit

are signs of NF- $\kappa$ B activation [20]. Previous studies [22-24] identified that H<sub>2</sub>O<sub>2</sub> treatment for different durations (30 min-24 h) elicited significant p65 phosphorylation and nuclear translocation in NRVMs. In line with these studies, p65 was time-dependently phosphorylated and translocated from the cytoplasm to the nucleus with I $\kappa$ B $\alpha$  degradation in NRVMs subjected to H<sub>2</sub>O<sub>2</sub>.

However, whether NF- $\kappa$ B activation protects or damages cardiomyocytes remains debatable. An early study demonstrated that activation of NF- $\kappa$ B reduced cell apoptosis in hypoxic cardiomyocytes [25], whereas most recent studies [26, 27] have shown that NF- $\kappa$ B is a pro-apoptotic transcription factor correlated with myocardial injury and that blockade of NF- $\kappa$ B activity prevents myocardial apoptosis. Gray et al [22] recently reported that ROS generated by ischaemia-reperfusion could rapidly activate calmodulin kinase II (CaMKII), which increased cell injury by inducing I $\kappa$ B $\alpha$  degradation and nuclear p65 accumulation in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>. Importantly, knockout of the CaMKII $\delta$  gene significantly attenuated myocardial infarct size by inhibiting I $\kappa$ B $\alpha$  degradation and NF- $\kappa$ B activation. All these findings show that NF- $\kappa$ B activation deteriorates the heart in I/R injury.

Herein, we hypothesized that direct overexpression of I $\kappa$ B $\alpha$  to prevent NF- $\kappa$ B activation may have a good protective effect in cardiomyocytes. Then, dsAAV9-I $\kappa$ B $\alpha$ <sup>Ser 32A,36A</sup> was designed to prevent I $\kappa$ B $\alpha$  degradation due to its phosphorylation at the Ser 32 and Ser 36 sites and successfully transfected into cardiomyocytes. Western blotting and immunofluorescence demonstrated that I $\kappa$ B $\alpha$  transfection successfully maintained cytoplasmic I $\kappa$ B $\alpha$  levels and suppressed p65 phosphorylation and translocation in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>. Additionally, I $\kappa$ B $\alpha$  elevated cell viability, decreased LDH and MDA levels, and attenuated apoptosis, implying the protective role of I $\kappa$ B $\alpha$  in H<sub>2</sub>O<sub>2</sub>-induced cell injury in NRVMs. The mechanisms may account for the role of NF- $\kappa$ B in mediating the expression of various proteins that promote or inhibit apoptosis. Notably, NF- $\kappa$ B regulates the expression of certain anti-apoptotic genes, such as Bcl-2 [28], and an increased Bcl-2/Bax ratio decreases cell apoptosis. In this study, treatment with I $\kappa$ B $\alpha$  or PDTC significantly elevated the Bcl-2/Bax ratio in NRVMs subjected to H<sub>2</sub>O<sub>2</sub>. These data indicate that I $\kappa$ B $\alpha$  protects NRVMs against H<sub>2</sub>O<sub>2</sub>-induced apoptosis by increasing the Bcl-2/Bax ratio.

Opening of the mitochondrial permeability transition pore (MPTP) in the first few minutes of reperfusion leads to  $\Delta\Psi$ m loss and is responsible for necrotic and apoptotic cell death processes, contributing differentially to myocardial infarct size [29]. Importantly, inhibition of the opening of the MPTP attenuated I/R injury. Thus,  $\Delta\Psi$ m loss reflects mitochondrial dysfunction, indicates early-stage apoptosis and is a critical determinant of I/R injury [30]. A previous study demonstrated that H<sub>2</sub>O<sub>2</sub> induced a significant decrease in  $\Delta\Psi$ m [12]. In this study, H<sub>2</sub>O<sub>2</sub> treatment attenuated  $\Delta\Psi$ m and enhanced Bax expression in NRVMs, and these effects were reversed by pretreatment with I $\kappa$ B $\alpha$  or PDTC. NF- $\kappa$ B is involved in the regulation of mitochondrial dysfunction [31], and Bax antagonizes the anti-apoptotic ability of Bcl-2 and simultaneously promotes permeability of the mitochondrial outer membrane and reduces the level of  $\Delta\Psi$ m [32]. The results herein suggest that I $\kappa$ B $\alpha$  decreases cell injury and apoptosis by inhibiting NF- $\kappa$ B activation and Bax expression, ultimately elevating  $\Delta\Psi$ m after H<sub>2</sub>O<sub>2</sub> stimulation.

Autophagy, a cellular process of lysosome-mediated degradation of cytoplasmic

components or damaged organelles, is thought to be an adaptive response and protective for cell survival [10]. However, autophagy causes a redox effect in cardiomyocytes when exposed to different stimuli. Evidence supports the benefit of autophagy to cardiomyocytes during myocardial ischaemia through its improvement of myocardial energy metabolism and organelle recycling [33], but excessive autophagy causes lethal damage to cells during cardiac I/R injury [21], which is mediated in part by upregulation of Beclin-1 expression [34].

However, the communication between autophagy and NF- $\kappa$ B is bidirectional. Autophagy is required for activation of NF- $\kappa$ B [13]; in turn, NF- $\kappa$ B further increases autophagosome maturation by upregulating Beclin-1 and LC3-II expression in I/R injury [35]. Importantly, PDTC attenuates Beclin-1 expression and the formation of autophagosomes by suppressing I/R injury-induced NF- $\kappa$ B activation [16]. In accordance with these findings, treatment of NRVMs with H<sub>2</sub>O<sub>2</sub> induced p65 phosphorylation and translocation, enhanced Beclin-1 expression, and increased the LC3-II/LC3-I ratio, and these effects were rescued by I $\kappa$ B $\alpha$  transfection or PDTC treatment. These results imply that I $\kappa$ B $\alpha$  can protect cardiomyocytes by inhibiting H<sub>2</sub>O<sub>2</sub>-induced autophagy. In addition, Bcl-2 can bind Beclin-1 to inhibit autophagy [36]. This study also demonstrates that I $\kappa$ B $\alpha$  transfection elevated the expression of Bcl-2, which may disturb the function of Beclin-1 and thus further inhibit H<sub>2</sub>O<sub>2</sub>-induced autophagy, implying cross-talk between apoptosis and autophagy.

### **Study strengths and limitations**

The present study has several strengths. To the best of our knowledge, this study is the first to report the cardioprotective effects of I $\kappa$ B $\alpha$  transfection through inhibition of H<sub>2</sub>O<sub>2</sub>-induced apoptosis and autophagy. Moreover, AAV9 vectors, which can be successfully transfected into cardiomyocytes, may be used as an effective tool for gene therapy for AMI. This study also has some limitations. The present findings were derived from neonatal cardiomyocytes *in vitro* and may differ from findings in animal experiments due to the complicated features of the *in vivo* environment. Further animal studies should be conducted to confirm the cardioprotective effects of I $\kappa$ B $\alpha$  against I/R injury.

## **Conclusions**

The findings of this study show that pretreatment with dsAAV9-I $\kappa$ B $\alpha$  or PDTC protected NRVMs from H<sub>2</sub>O<sub>2</sub>-induced apoptosis, autophagy, mitochondrial dysfunction, and oxidative damage by restraining the NF- $\kappa$ B signalling pathway, suggesting that I $\kappa$ B $\alpha$  transfection can protect cardiomyocytes against cardiac oxidative damage. Thus, AAV9 vectors, as high-efficiency gene carriers to heart, may be used to carry the I $\kappa$ B $\alpha$  gene to protect the heart through targeted inhibition of myocardial NF- $\kappa$ B in future preclinical or clinical studies, which may provide a promising gene therapy for preventing cardiac I/R injury.

## **Declarations**

## **Ethics approval and consent to participate**

All animal experimental protocols were approved by the Ethics Committee of the First Affiliated Hospital of Xinjiang Medical University (No. IACUC-20180223-69).

## **Funding**

This work was supported by National Natural Science Foundation of China (No. 81873490, No. 81860072), the Xinjiang Uygur Autonomous Region Graduate Research and Innovation Project (No. XJ2019G183), the Xinjiang Medical University Graduate Innovation and Entrepreneurship Project (No. CXCX2018020), and the Tianshan Innovative Research Team Plan (No. 2018D14006).

## **Authors' contributions**

MH, XCC, MHS, and MTG performed the research. YNY, XMG, and XM provided guidance throughout the whole study. MH and XCC analysed the data and drafted the paper. YTM and BDC designed and provided final approval of the version to be published.

## **Acknowledgements**

The authors thank Xue Zhang and Ning Yang for their technical assistance in this work.

## **Consent for publication**

Not applicable.

## **Availability of data and materials**

All data generated or analysed during this study are included in this article.

## **Competing interests**

The authors declare no competing interests.

## **Authors' information**

<sup>a</sup> Xinjiang Key Laboratory of Cardiovascular Disease Research, Clinical Medical Research Institute, the First Affiliated Hospital of Xinjiang Medical University, Urumqi, 830054, PR China; <sup>b</sup> Department of Cardiology, the First Affiliated Hospital of Xinjiang Medical University, Urumqi, 830054, PR China; <sup>c</sup> Department of Nephrology, Fifth Affiliated Hospital of Xinjiang Medical University, Urumqi, 830000, PR China.

## **Abbreviations**

NF- $\kappa$ B: nuclear factor kappa B; I $\kappa$ B $\alpha$ : inhibitor of kappa B alpha; NRVMS: neonatal rat ventricular cardiomyocytes; I/R: ischaemia/reperfusion; AMI: acute myocardial infarction; H<sub>2</sub>O<sub>2</sub>: hydrogen peroxide; GFP: green fluorescent protein; PDTC: pyrrolidine dithiocarbamate; CCK-8: Cell Counting Kit-8; LDH: lactate dehydrogenase; MDA: malondialdehyde; MPTP: mitochondrial permeability transition pore;  $\Delta\Psi_m$ : mitochondrial membrane potential; ROS: reactive oxygen species.


## References

1. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2018;39(2):119-177.
2. Del Re DP, Amgalan D, Linkermann A, Liu Q, Kitsis RN. Fundamental Mechanisms of Regulated Cell Death and Implications for Heart Disease. *Physiol Rev*. 2019;99(4):1765-1817.
3. Verma VK, Malik S, Mutneja E, Sahu AK, Bhatia J, Arya DS. Attenuation of ROS-mediated myocardial ischemia-reperfusion injury by morin via regulation of RISK/SAPK pathways. *Pharmacol Rep*. 2020. doi: 10.1007/s43440-019-00011-2.
4. Wallert M, Ziegler M, Wang X, Maluenda A, Xu X, Yap ML, et al.  $\alpha$ -Tocopherol preserves cardiac function by reducing oxidative stress and inflammation in ischemia/reperfusion injury. *Redox Biol* 2019;26:101292.
5. Sies H, Berndt C, Jones DP. Oxidative Stress. *Annu Rev Biochem*. 2017;86:715-748.
6. Zhang Q, Lenardo MJ, Baltimore D. 30 Years of NF- $\kappa$ B: A Blossoming of Relevance to Human Pathobiology. *Cell*. 2017;168(1-2):37-57.
7. Sun SC. The non-canonical NF-kappaB pathway in immunity and inflammation. *Nat Rev Immunol*. 2017;17(9):545-558.
8. Kumar R, Yong QC, Thomas CM. Do multiple nuclear factor kappa B activation mechanisms explain its varied effects in the heart? *Ochsner J*. 2013;13(1):157-65.
9. Sun Z, Ma YT, Chen BD, Liu F. Recombinant adeno-associated virus serotype 9 with p65 ribozyme protects H9c2 cells from oxidative stress through inhibiting NF-kappaB signaling pathway. *J Geriatr Cardiol*. 2014;11(4):311-5.
10. Sciarretta S, Maejima Y, Zablocki D, Sadoshima J. The Role of Autophagy in the Heart. *Annu Rev Physiol*. 2018;80:1-26.
11. Zhu HH, Wang XT, Sun YH, He WK, Liang JB, Mo BH, et al. Pim1 Overexpression Prevents Apoptosis in Cardiomyocytes After Exposure to Hypoxia and Oxidative Stress via Upregulating Cell Autophagy. *Cell Physiol Biochem*. 2018;49(6):2138-2150.
12. Sun MH, Chen XC, Han M, Yang YN, Gao XM, Ma X, et al. Cardioprotective effects of constitutively active MEK1 against H<sub>2</sub>O<sub>2</sub>-induced apoptosis and autophagy in cardiomyocytes via the ERK1/2

- signaling pathway. *Biochem Biophys Res Commun*. 2019;512(1):125-130.
13. Criollo A, Chereau F, Malik SA, Niso-Santano M, Marino G, Galluzzi L, et al. Autophagy is required for the activation of NFκB. *Cell cycle*. 2012;11(1):194-9.
  14. Wang X, Tao YL, Huang YW, Zhan KG, Xue M, Wang Y, et al. Catalase Ameliorates Diabetes-Induced Cardiac Injury Through Reduced p65/RelA- Mediated Transcription of BECN1. *J Cell Mol Med*. 2017;21(12):3420-34.
  15. Dong Y, Chen H, Gao J, Liu Y, Li J, Wang J. Molecular machinery and interplay of apoptosis and autophagy in coronary heart disease. *J Mol Cell Cardiol*. 2019;136:27-41.
  16. Zeng M, Wei X, Wu Z, Li W, Li B, Zhen Y, et al. NF-κB-mediated induction of autophagy in cardiac ischemia/reperfusion injury. *Biochem Biophys Res Commun*. 2013;436(2):180-5.
  17. García-Olloqui P, Rodríguez-Madoz JR, Di Scala M, Abizanda G, Vales Á, Olagüe C, et al. Effect of heart ischemia and administration route on biodistribution and transfection efficiency of AAV9 vectors. *J Tissue Eng Regen Med*. 2020;14(1):123-134.
  18. Chen H. Exploiting the Intron-splicing Mechanism of Insect Cells to Produce Viral Vectors Harboring Toxic Genes for Suicide Gene Mol Ther Nucleic Acids. 2012;1(11):e57.
  19. Tao J, Chen BD, Ma YT, Yang YN, Li XM, Ma X, et al. FrzA gene protects cardiomyocytes from H2O2-induced oxidative stress through restraining the Wnt/Frizzled pathway. *Lipids Health Dis*. 2015;14:90.
  20. Prabhu SD, Frangogiannis NG. The Biological Basis for Cardiac Repair After Myocardial Infarction: From Inflammation to Fibrosis. *Circ Res*. 2016;119(1):91-112.
  21. Huang Z, Han Z, Ye B, Dai Z, Shan P, Lu Z, et al. Berberine alleviates cardiac ischemia/reperfusion injury by inhibiting excessive autophagy in cardiomyocytes. *Eur J Pharmacol*. 2015;762:1-10.
  22. Gray CB, Suetomi T, Xiang S, Mishra S, Blackwood EA, Glembotski CC, et al. [CaMKIIδ subtypes differentially regulate infarct formation following ex vivo myocardial ischemia/reperfusion through NF-κB and TNF-α](#). *J Mol Cell Cardiol*. 2017;103:48-55.
  23. Wei C, Li L, Kim IK, Sun P, Gupta S. NF-κB mediated miR-21 regulation in cardiomyocytes apoptosis under oxidative stress. *Free Radic Res*. 2014;48(3):282-91.
  24. Tang B, Ma J, Ha X, Zhang Y, Xing Y. Tumor necrosis factor-alpha upregulated PHLPP1 through activating nuclear factor-kappa B during myocardial ischemia/reperfusion. *Life Sci*. 2018;207:355-63.
  25. Misra A, Haudek SB, Knuefermann P, Vallejo JG, Chen ZJ, Michael LH, et al. Nuclear factor-kappaB protects the adult cardiac myocyte against ischemia-induced apoptosis in a murine model of acute myocardial infarction. *Circulation*. 2003;108(25):3075-8.
  26. Zou G, Zhong W, Wu F, Wang X, Liu L. Inhibition of lncRNA Neat1 by catalpol via suppressing transcriptional activity of NF-κB attenuates cardiomyocyte apoptosis. *Cell Cycle*. 2019;18(24):3432-3441.


27. Zhang W, Zhang Y, Ding K, Zhang H, Zhao Q, Liu Z, et al. Involvement of JNK1/2-NF- $\kappa$ Bp65 in the regulation of HMGB2 in myocardial ischemia/reperfusion-induced apoptosis in human AC16 cardiomyocytes. *Biomed Pharmacother.* 2018;106:1063-1071.
28. Liu Q, Liu Z, Zhou LJ, Cui YL, Xu JM. The long noncoding RNA NKILA protects against myocardial ischaemic injury by enhancing myocardin expression via suppressing the NF- $\kappa$ B signalling pathway. *Exp Cell Res.* 2020;387(2):111774.
29. Morciano G, Bonora M, Campo G, Aquila G, Rizzo P, Giorgi C, et al. Mechanistic Role of mPTP in Ischemia-Reperfusion Injury. *Adv Exp Med Biol.* 2017;982:169-89.
30. Zhao Y, Guo Y, Chen Y, Liu S, Wu N, Jia D. Curculigoside attenuates myocardial ischemia-reperfusion injury by inhibiting the opening of the mitochondrial permeability transition pore. *Int J Mol Med.* 2020;45(5):1514-1524.
31. Han Y, Cai Y, Lai X, Wang Z, Wei S, Tan K, et al. [lncRNA RMRP Prevents Mitochondrial Dysfunction and Cardiomyocyte Apoptosis via the miR-1-5p/hsp70 Axis in LPS-Induced Sepsis Mice.](#) *Inflammation.* 2020;43(2):605-618.
32. Garcaí-Sáez AJ. The secrets of the Bcl-2 family[J]. *Cell Death Differ.* 2012;19(11):1733-40.
33. Huang L, Dai K, Chen M, Zhou W, Wang X, Chen J, et al. The AMPK Agonist PT1 and mTOR Inhibitor 3HOI-BA-01 Protect Cardiomyocytes After Ischemia Through Induction of Autophagy. *J Cardiovasc Pharmacol Ther.* 2016;21(1):70-81.
34. Zuo Z, Zuo PF, Sheng ZL, Wang X, Ding JD, Ma GS. Tetramethylpazine attenuates myocardial ischemia/reperfusion injury through modulation of autophagy. *Life Sci.* 2019; 239:117016.
35. Haar L, Ren X, Liu Y, Koch SE, Goines J, Tranter M, et al. Acute consumption of a high-fat diet prior to ischemia-reperfusion results in cardioprotection through NF- $\kappa$ B-dependent regulation of autophagic pathways. *Am J Physiol Heart Circ Physiol.* 2014;307(12):H1705-13.
36. Pattingre S, Tassa A, Qu X, Garuti R, Liang XH, Mizushima N, et al. Bcl-2 antiapoptotic proteins inhibit Beclin 1-dependent autophagy. *Cell.* 2005;122(6):927-39.

## Figures




**Figure 1**

Effects of H<sub>2</sub>O<sub>2</sub> and AAV9 vectors on NRVMs. (A-C) Western blot and quantified cytosolic or nuclear protein levels of IkBα and p65 in NRVMs exposed to 100 μM H<sub>2</sub>O<sub>2</sub> for 0, 15, 30, 60 min, respectively (n = 3, \*P < 0.05 and \*\*P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> (0 min)). (D-E) Western blot and quantified total protein levels of p-p65 and p65 in NRVMs subjected to 100 μM H<sub>2</sub>O<sub>2</sub> for 0, 15, 30, 60 min, respectively (n = 3, \*P < 0.05 and \*\*P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> (0 min)). (F) The green fluorescence intensity of GFP in NRVMs (scale bar: 250 μm). (G-I) Western blot to detect the protein expression of IkBα and GFP. (n = 3, \*\* P < 0.01 vs. Control).




**Figure 2**

IκBα reduced cell apoptosis in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>. (A) Flow cytometry indicated that IκBα or PDTC attenuated cell apoptosis in NRVMs exposed to H<sub>2</sub>O<sub>2</sub> (n = 3, \* P < 0.05 and \*\* P < 0.01 vs. Control, # P < 0.05 vs. H<sub>2</sub>O<sub>2</sub> group). (B) Western blot findings indicated that IκBα or PDTC reduced the ratio of Bax/Bcl-2 in NRVMs subjected to H<sub>2</sub>O<sub>2</sub> (n = 3, \*\* P < 0.01 vs. Control, ## P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> group).



**Figure 3**

IkBa reduced cell injury in NRVMs exposed to H<sub>2</sub>O<sub>2</sub>. (A) Representative images of JC-1 staining in different groups (scale bar: 250 μm). The results indicated that IkBa or PDTC elevated the ratio of red to green fluorescence intensity (n = 3, \* P < 0.05 and \*\* P < 0.01 vs. Control, # P < 0.05 vs. H<sub>2</sub>O<sub>2</sub> group). (B) CCK-8 results demonstrated that IkBa or PDTC increased the cell viability in NRVMs stimulated by H<sub>2</sub>O<sub>2</sub> (n = 3, \* P < 0.05 and \*\* P < 0.01 vs. Control, # P < 0.05 vs. H<sub>2</sub>O<sub>2</sub> group). (C) IkBa or PDTC reduced the levels of supernatant LDH in NRVMs subjected to H<sub>2</sub>O<sub>2</sub> (n = 3, \* P < 0.05 and \*\* P < 0.01 vs. Control, # P < 0.05 and ## P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> group). (D) IkBa or PDTC decreased the content of intracellular MDA in H<sub>2</sub>O<sub>2</sub>-treated NRVMs (n = 3, \*\* P < 0.01 vs. Control, # P < 0.05 vs. H<sub>2</sub>O<sub>2</sub> group).



**Figure 4**

IκBα suppressed H<sub>2</sub>O<sub>2</sub>-induced NF-κB activation and autophagy in NRVMs. (A) Distribution of NF-κB p65 (red fluorescence) was detected using immunofluorescence (scale bar: 25 μm). The results showed that IκBα or PDTC inhibited H<sub>2</sub>O<sub>2</sub>-induced p65 nuclear translocation. (B) Western blot and quantified cytosolic or nuclear protein levels of IκBα and p65 in NRVMs exposed to 100 μM H<sub>2</sub>O<sub>2</sub>. The results showed that IκBα transduction maintained the cytosolic IκBα level and suppressed p65 nuclear translocation (n = 3, \*\* P < 0.01 vs. Control, ## P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> group). (C) Western blot and quantified total protein levels of p-p65, p65, Beclin-1 and LC3-II/LC3-I in NRVMs exposed to 100 μM H<sub>2</sub>O<sub>2</sub>. The findings showed that IκBα or PDTC decreased p-p65/p65 ratio, Beclin-1 expression and LC3-II/LC3-I ratio after H<sub>2</sub>O<sub>2</sub> stimulation. (n = 3, \*\* P < 0.01 vs. Control, ## P < 0.01 vs. H<sub>2</sub>O<sub>2</sub> group).