

Effects of tactile-kinesthetic stimulation on feeding and weight of moderate and late preterm neonates

Emily Muthoni Nyaga (✉ nyaga.emily@gmail.com)

Moi University College of Health Sciences <https://orcid.org/0000-0001-5947-5789>

Nehad Sabry Basiouny

Alexandria University Faculty of Nursing

Fabian Omoding Esamai

Moi University School of Medicine

Gamat El-Sayed Mansy

Alexandria University Faculty of Nursing

Research article

Keywords: Preterm neonates, tactile-kinesthetic stimulation, feeding intolerance, and weight.

Posted Date: March 9th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-16398/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Owing to immaturity of their body systems, preterm neonates are susceptible to feeding intolerance, slowed growth and long hospitalization periods among others. Tactile-kinesthetic stimulation, a moderate pressure of the skin and joint movement has been widely researched on among preterm neonates; the studies have predominately focused on weight gain and the potential underlying mechanism. This study focuses on effect of tactile-kinesthetic stimulation on amount of feeds, feeding intolerance and weight of moderate and late preterm neonates. **Methods:** A quasi-experimental study comprising 72 preterm neonates born at 28 to <37 weeks gestation age (GA) was conducted. Subjects were divided into two groups; control and tactile-kinesthetic stimulation (TKS). Neonates in the TKS group received massage for 15 minutes per session; 3 times a day for 10 days while control group had standard nursery care. Neonates' amount of feeds and signs of feeding intolerance were assessed on days 3, 10, 17, and 23 of life while weight was measured on alternate days starting on day 3 up to day 23. Mann Whitney U test, Student t-test, Chi-square test, and Fisher's exact test were used to determine whether there was any difference in feeding parameters and weight between TKS and control groups. **Results:** Although amount of feeds didn't differ between the groups, TKS group neonates had fewer episodes of feeding intolerance compared to those in control group on the 10 th ($p = .03$), 17 th ($p = .00$), and 23 th ($p = .00$) day of life. Moderate preterm neonates didn't differ in weight gain however; late preterm neonates in TKS group had significantly more weight gain than the control. **Conclusion:** Tactile-kinesthetic stimulation reduces feeding intolerance and enhance weight gain in moderate preterm neonates. **Key words:** Preterm neonates, tactile-kinesthetic stimulation, feeding intolerance, and weight. Trial registration: ClinicalTrials.gov NCT04287322, registered on 27/02/2020. Retrospectively registered.

Background

A steady increase in the rate of preterm births has been witnessed in the last decade constituting 11.1% of 135 million live births worldwide, the majority of which occur in sub-Saharan Africa. In Kenya, the prevalence of preterm birth stands at 6.7% as a result of demographic change, infertility treatment, increased maternal age, multiple births, and increased rate of obesity ⁽¹⁻³⁾.

Immature digestive system of preterm neonates induces feeding intolerance resulting in increased risk of infection, inability to maintain enteral feeds, and weight loss ^(4, 5); in addition to necrotizing enterocolitis - a leading cause of morbidity and mortality of preterm neonates ⁽⁶⁾. With priority to promptly achieve full enteral feeds for the preterm neonates, signs of feeding intolerance are a great set back in care of the preterm neonates. Although in high-income countries supplemental parenteral nutrition is commonly given to preterm neonates with difficulty in tolerating adequate enteral feeds ⁽⁷⁾ it is way too costly for a majority of the population in Kenya and therefore not used in Moi Teaching & Referral Hospital (MTRH). The preterm neonates are rather put on maintenance fluid (Ringer's lactate/5% dextrose) despite its lack of optimal nutritional requirements for the neonates. Inadequate nutrition coupled with feeding intolerance increases length of time to reach full enteral feeding; and the latter has been associated with poorer mental outcome in preterm neonates at 24 months corrected age ⁽⁴⁾. Through parasympathetic activity massage accelerates peristalsis, decrease abdominal distension, accelerate bowel transit time, increase frequency of stooling, and decrease frequency of vomiting ^(4, 5).

Optimal physical growth is important for normal growth and development of preterm neonates. Rapid growth during hospitalization allows early discharge, rapid adaptation to home environment, and emotional attachment to the mother. Poor growth indices have been associated with lower neurodevelopmental and growth outcomes later in life ⁽⁴⁾. Enhancing retention and digestion of feeds for growth and development is a core component in care of preterm neonates. Weight being a critical criterion for discharge from neonatal unit is a major problem. In many instances, insufficient weight gain is the main reason for prolonged hospitalization, high cost care, and related problems ⁽⁸⁾. At MTRH Special Care Nursery (SCN) the hospital policy allows neonates to be discharged after attaining 1700g. Strategies that enhance weight gain not only reduce the cost of care but allow mother to bond with her infant in a relaxed home environment. Tactile-kinesthetic stimulation (TKS), a form of massage therapy involves moderate-pressure stroking of the neonate and movements of large joints. Through parasympathetic activity massage and kinesthetic movements (extension and flexion of joints) increases secretion of digestive hormones (insulin, gastrin and gastric fluid), increases mean enteral intake and weight gain ^(9,10). Massage therapy is safe and tolerated by preterm neonates; the neonates are more responsive to tactile stimulation than other sensory modalities due to early maturation of the tactile system and the large surface area of the skin ⁽¹¹⁾.

Although studies have demonstrated more enteral intake, and greater weight gain in preterm neonates as a result of TKS ^(1,12,13) little has been done on feeding intolerance and comparison of weight between moderate and late preterm neonates. The purpose of this study was to determine effects of TKS on feeding and weight of moderate and late preterm neonates.

Methods

Study design

A quasi-experimental design was carried out to assess effects of tactile-kinesthetic stimulation on feeding parameters and physical growth of preterm neonates. The study involved two groups; control and tactile-kinesthetic stimulation (intervention) groups.

Study setting

The study was conducted at a level II Special Care Nursery (American Academy of Pediatrics, 2012) ⁽¹⁴⁾ of Moi Teaching & Referral Hospital (MTRH), an academic hospital in the Western region of Kenya. It has 70-bed capacity, 11 incubators, and 54 cots. It provides care to preterm and moderately ill neonates who do not require mechanical ventilation.

Study population

The study population comprised preterm neonates admitted at SCN of MTRH. The SCN has an average of 90 neonates of whom majority (75%) are preterm (hospital records, 2017).

Sample size

Using Epi-Info software and main output variable as mean weight gain which was compared in the two groups, sample size was estimated to be 36 participants in each group, totaling to 72 participants for the study to detect average weight gain of 27 ± 3 grams per day among the interventional group compared to average weight gain of 25 ± 3 grams per day among the control group ⁽¹²⁾. Thus, change of average weight gain by two grams was detected. The estimated sample size was made at assumption of 95% confidence level and 80% power of study.

Eligibility criteria; neonates who fulfilled the following criteria comprised the study participants; a) on breast milk or formula feeds, b) born 28 to 37 weeks gestational age, c) ≥ 1000 grams. The gestation was limited to ≥ 28 weeks and ≥ 1000 grams based on significant neonatal mortality rate in neonates born before 28 weeks gestation and/or weighing < 1000 grams in MTRH ⁽¹⁵⁾. Neonates were excluded from the study if they; a) were critically ill and those on continuous positive airway pressure (CPAP) or b) had neonatal infections including severe sepsis or necrotizing enterocolitis.

Sampling method

Consecutive sampling was used to recruit preterm neonates for the study on day 3 of the neonate's life. Study participants that met the eligibility criteria were consecutively selected in order of appearance according to their convenient accessibility. The first study participant was recruited in the control group while the subsequent with similar characteristics in the intervention group, this was continued until the desired sample size was reached.

Study tool

A study tool developed by the authors after thorough review of related literature on effects of TKS ^(1, 16) was used to collect data on a) neonates characteristics including gender, gestational age, and clinical risk index for babies (CRIB II) score; b) amount of feeds, signs of feeding intolerance (gastric residual $> 50\%$ of previous feed as per MTRH hospital practices, abdominal distension and vomiting), and weight in grams. Data on feeding characteristics were analyzed on day 3, 10, 17, and 23 while that for weight on alternate days starting from day 3 up to day 23.

Study tool validity & reliability

Tools were reviewed for content validity by 5 experts in the field of pediatric nursing and suggestions made were incorporated into the tools. Medical records were used to gather information on amount of feeds and signs of feeding intolerance. Inter-rater reliability using two observers (research assistant and staff nurse at SCN) were done for weight recording. Notably, there was no variation in weight recording between the two observers. A pilot study was carried out on 12 preterm neonates (four in each group) at MTRH hospital prior to data collection to test the feasibility of the study. Neonates included in the pilot study were excluded from the main study. There were no amendments made to the study tool.

Study procedures

A researcher was trained on TKS by a specialized pediatric nurse to ensure moderate pressure is applied on the body surface in a manner to cause effect and not harm the neonate. The training was conducted in three sessions each lasting two hours. The first and second session were conducted in the skills laboratory while the third session at MTRH hospital. The first session comprised review of massage procedure and watching the video. The second session was return demonstration on a dummy. The third session was practice on preterm neonates to achieve competency. Two research assistants (RA's - Bachelor of Science in nursing intern) were trained on data entry. The RA's were blinded to aim of study, group assignment, and interventions received by the preterm neonates. In addition RA's had no access to study data; completed assessment tools, and notes.

Control group

Neonates in this group didn't receive any specific stimulation rather had standard care of the SCN.

Tactile-Kinesthetic Stimulation (TKS) group

The group received TKS in addition to standard care of SCN.

Tactile-kinesthetic stimulation involved three sessions per day; morning, afternoon, and evening for 10 days starting day 3 of life (initial contact). After thorough hand scrubbing, the researcher placed her warmed hands on the preterm neonate's body. Access ports were used for neonates in the incubator to prevent hypothermia. The preterm neonates were on a cardiorespiratory monitor for physiological monitoring with set alarm to detect deviation from normal that would warrant discontinuation of the TKS. The stimulation was given 1-2 hours after feeding. A small amount of sunflower oil was used for TKS to decrease injurious friction between surfaces (providers' palms and neonate's skin) and was removed with cotton after the stimulation. Stimulation was temporarily stopped if the neonate started crying or passed urine or stool. The stimulation was continued when the neonate regained stability. The 15-minutes stimulation included three standardized 5 minutes phases.

The phases were as follows:

Phase 1: Preterm neonates were placed in prone position. Moderate pressure (sufficient to produce slight skin indentation or slight skin color change from pink to white) was used to provide 12 strokes with palms of the hands, each stroke lasting 5 seconds. The strokes were provided in each area as follows: (a) head - from forehead hairline over scalp down to neck with alternate hands; (b) neck - from midline outwards with both hands simultaneously; (c) shoulders - from midline outwards with both hands simultaneously, and (d) back - from nape of neck down to buttocks, long stroke with alternate hands.

Phase 2: The preterm neonates were placed in supine position. Twelve moderate pressure strokes with palms of the hands, 5 seconds each, were provided in each area as follows: (a) forehead - from midline, outwards with both hands simultaneously; (b) cheeks - from side of nose, with both hands simultaneously in rotating and clockwise direction; (c) chest - 'butterfly' stroking from midline upwards, outwards, downwards and inwards back to initiating point; (d) abdomen - from the appendix, in a clock wise direction around abdomen avoiding the epigastrium and probes, with gentle strokes; (e) upper limbs (each separately) - from shoulders

to wrist using alternate hands for stroking; (f) lower limbs (each separately) - from hips to ankles using alternate hands for stroking; (g) palms - from wrist to finger tips using alternate hands for stroking; and (h) soles - from heel to toe tips using alternate hands for stroking.

Phase 3: Kinesthetic stimulation was done for 5 minutes. The intervention comprised five passive flexion and extension movements of each large joint (shoulder, elbow, wrist, hip, knee, and ankle) for two seconds.

The massage therapy protocol was adopted from Mathai 2001⁽¹⁷⁾ a modification of Field et al. (1986) protocol for medically stable preterm neonates.

Data management and analysis

Data from study tools was coded and entered into Statistical Package for the Social Sciences (SPSS) version 20 database that was created for the study. Study tools were visually checked for completeness and accuracy before data entry. Descriptive statistics were computed for the baseline clinical data of the study participants. Chi-square test and student t-test were used to show relationship between variables (gender, GA, Apgar score, birth weight, and CRIB II score) between TKS and control groups. Means and (% confidence intervals, standard deviations (SDs) and interquartile ranges (IQR) were computed to describe dependent variables; amount of feed, signs of feeding intolerance, duration to attaining full breastfeeding and weight. Mann Whitney U test, Student t-test, Chi-square test, and Fisher's exact test were used to determine whether there was any difference between TKS and control groups in amount of feeds and signs of feeding intolerance. Student t-test and Mann Whitney U test were used to determine whether a difference existed in weight of moderate and late preterm neonates between the two groups.

Results

Seventy two participants (36 in each group) were recruited to the study on day 3 of life. Twelve participants; six in each group did not complete the study; six participants in TKS group and four in control group were discharged during study period while two participants in control group died before completion of study. The sixty participants all singletons (30 each in group) were followed up to completion of the study (21 days). Chi-square test and student t-test were used to analyze baseline clinical characteristics; results showed no statistical difference between TKS and control groups as regards gender, Apgar score, birth weight, length, head circumference and temperature at birth. A statistically significant lower mean gestational age ($p = .03$) and higher mean CRIB II score ($p = .02$) were observed in TKS group compared to control group. Table 1 summarizes the baseline clinical characteristics of the participants.

Mann Whitney U test and Student t-test were used to analyze amount of feeds between TKS and control groups and the results showed the amount of feeds didn't differ between the two groups. Although TKS group attained full breastfeeding (no gavage or cup feeding) a day earlier than the control group, the difference was not statistically significant. Chi-square test and Student t-test were used to analyze feeding intolerance (gastric residual, abdominal distension, and vomiting) and the findings showed no difference at initial contact; occurring among half of the participants in TKS group and 63% in the control group. However, on the 10th ($p = .03$), 17th ($p = .00$), and 23rd ($p = .00$) day of life TKS group had significantly fewer episodes of

feeding intolerance than control group. Table 2 and 3 summarizes the feeding characteristics of the participants.

Analysis of mean weight for moderate preterm neonates using Student t-test and Mann Whitney U test showed at initial contact (3rd day of life) TKS had 1164.0±160.3 while that for control group was 1262.3±131.7 and at the end of the study (23rd day of life) was 1555.4±231.0 and 1581.9±195.2 respectively. Weight didn't differ between the two groups. However, a statistically significant difference was observed in late preterm neonates between the two groups on 15th ($p = .05$), 21st ($p = .04$) and 23rd ($p = .02$) day of life where TKS group neonates had more weight than control (Table 3a).

Discussion

In the recent years the focus on preterm care has shifted from increasing survival to minimizing morbidity and improving neurodevelopmental outcomes⁽¹⁸⁾. Sensory stimulation such as tactile-kinesthetic stimulation (TKS) provides a platform for learning during infancy with potential to improve the outcomes of preterm neonates⁽¹⁹⁻²¹⁾.

Similar to most studies⁽²²⁻²⁶⁾, we found no difference in mean enteral intake between the intervention and control groups. Few studies^(27, 28) have reported more enteral intake with TKS compared to control. For instance, Fontana et al. (2018) demonstrated a higher consumption of breast milk at discharge among preterm neonates receiving massage therapy compared to control⁽²⁷⁾.

In addition to monitoring amount of feeds intake, signs of feeding intolerance are key in nutritional management of preterm neonates. Although the definition of feeding intolerance varies with population and feeding practices, the term has been based historically on presence of increased amount of gastric aspirate, abdominal distention, and vomiting^(16, 29, 30). Preterm neonates in the interventional group had significantly fewer episodes of signs of feeding intolerance compared to the control group. Despite several studies assessing effects of tactile-kinesthetic stimulation on preterm neonates, we found none has explored feeding intolerance. However, studies^(5, 16, 31) on abdominal massage of preterm neonates have reported similar findings demonstrating increased peristalsis (gastric motility) from first day of massage therapy as the underlying mechanism⁽¹⁶⁾.

Moderate preterm neonate's weight didn't differ between interventional and control groups in this study, however, this could be attributed to subnormal growth that occurs in such neonates⁽³¹⁾. Notably, similar to most studies^(1, 8, 11, 12, 17, 22, 23, 32-36), we found greater weight gain in late preterm neonates in the interventional group compared to the control group. The underlying mechanism for weight gain has been suggested to be increased vagal activity leading to heightened gastric motility, and increased secretion of digestive enzymes - insulin, insulin growth factor - 1, gastrin, and ornithine decarboxylase enzyme^(9, 37-39).

Early establishment of breastfeeding helps preterm neonates bonding, warmth, cuddling, and ultimately growth and skill development. Interventional group preterm neonates attained full breastfeeding a day earlier than those in control group although the difference was not statistically significant.

Conclusion

Preterm neonates that receive tactile-kinesthetic stimulation have fewer signs of feeding intolerance. In addition, tactile-kinesthetic stimulation enhances weight gain in late preterm neonates although the same is not observed in moderate preterm neonates. Our findings indicate the need for teaching mothers of stable preterm neonates on tactile-kinesthetic stimulation and its incorporation as a care strategy of preterm neonates in

Abbreviations

CPAP – Continuous positive airway pressure

CRIB II – Clinical risk index for babies

GA – Gestation age

IQR – Interquartile ranges

IREC – Institutional Research & Ethics Committee

MTRH – Moi Teaching & Referral Hospital

RAs – Research assistants

SCN – Special care nursery

SD – Standard deviation

SPSS – Statistical package for the social sciences

TKS – Tactile-kinesthetic stimulation

Declarations

Ethics approval and consent to participate statement; the study was approved by the Institutional Research & Ethics Committee (IREC) of University's College of Health Sciences and Moi Teaching & Referral Hospital – Kenya. Following informed written consent from their mothers, neonates were recruited to the study. In addition the study is registered under ClinicalTrials.gov trial registration number NCT04287322.

Consent for publication; not applicable.

Competing interests; authors have nothing to disclose with respect to any affiliation or financial agreement that would lead to a conflict of interest.

Authors' contributions; EMN, NSB, GEM, and FOE participated in developing the study protocol, monitoring data collection process and analyzing the data, writing the manuscript. All authors read and approved the final manuscript.

Funding; no funding was received.

Acknowledgements; we thank the health care professionals at nursery unit of MTRH– Kenya and research assistants for support during the study. We are also very grateful to mothers and their preterm neonates for their participation in the study.

References

1. Ahmed RG, Suliman GI, Elfakey WA, Salih KM, El-Amin EI, Ahmed WA, et al. Effect of tactile kinesthetic stimulation on preterm infants' weight and length of hospital stay in Khartoum, Sudan. *Saudi medical journal*. 2015;36(2):196.
2. Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller A-B, et al. Born too soon: the global epidemiology of 15 million preterm births. *Reproductive health*. 2013;10(1):S2.
3. Vogel JP, Lee AC, Souza JP. Maternal morbidity and preterm birth in 22 low-and middle-income countries: a secondary analysis of the WHO Global Survey dataset. *BMC pregnancy and childbirth*. 2014;14(1):56.
4. Kim Hy, Bang KS. The effects of enteral feeding improvement massage on premature infants: A randomised controlled trial. *Journal of clinical nursing*. 2018;27(1-2):92-101.
5. Shaeri M, Ghadami A, Valiani M, Armanian A-M, Amini Rarani S. Effects of abdominal massage on feeding tolerance in preterm infants hospitalized in selected hospitals of Isfahan-Iran. *International Journal of Pediatrics*. 2017;5(3):4503-10.
6. Islam MA, Begum S, Yasmeen BN, Ahmed J, Amin MM. Feeding Intolerance in Preterm Neonates in Neonatal Intensive Care Unit, Padma General hospital, Dhaka. *Northern International Medical College Journal*. 2016;8(1):178-80.
7. Webbe J, Longford N, Uthaya S, Modi N, Gale C. Outcomes following early parenteral nutrition use in preterm neonates: protocol for an observational study. *BMJ open*. 2019;9(7):e029065.
8. Johari S, Haghgou HA, Daemi M, Rezaeiyan T, Mosala Nejad Z. The Effect of Massage on Weight Gain of Low-Weight Hospitalized Infants: A Randomized Clinical Trial. *Physical Treatments-Specific Physical Therapy Journal*. 2016;5(4):205-10.
9. Diego MA, Field T, Hernandez-Reif M. Preterm infant weight gain is increased by massage therapy and exercise via different underlying mechanisms. *Early human development*. 2014;90(3):137-40.
10. Dieter JN, Field T, Hernandez-Reif M, Emory EK, Redzepi M. Stable preterm infants gain more weight and sleep less after five days of massage therapy. *Journal of pediatric psychology*. 2003;28(6):403-11.
11. Aliabadi F, Askary RK. effects of Tactile–kinesthetic Stimulation on low birth Weight neonates. *Iranian journal of pediatrics*. 2013;23(3):289.
12. Massaro A, Hammad T, Jazzo B, Aly H. Massage with kinesthetic stimulation improves weight gain in preterm infants. *Journal of perinatology*. 2009;29(5):352.
13. Diego MA, Field T, Hernandez-Reif M. Temperature increases in preterm infants during massage therapy. *Infant Behavior and Development*. 2008;31(1):149-52.
14. Barfield WD, Papile LA, Baley JE, Benitz W, Cummings J, Carlo WA, et al. Levels of neonatal care. *Pediatrics*. 2012;130(3):587-97.

15. Njuguna FM, Kiptoon P, Nyandiko W. An assessment of the overall mortality of low birth weight neonates at the new birth units of the Moi Teaching and Referral hospital in Eldoret, Kenya. *European Journal of Biology and Medical Science Research*. 2014;2(3):63-71.
16. Tekgündüz KŞ, Gürol A, Apay SE, Caner İ. Effect of abdomen massage for prevention of feeding intolerance in preterm infants. *Italian journal of pediatrics*. 2014;40(1):89.
17. Mathai S, Fernandez A, Mondkar J, Kanbur W. Effects of tactile-kinesthetic stimulation in preterms-A controlled trial. *Indian pediatrics*. 2001;38(10):1091-8.
18. Lai MM, D'Acunto G, Guzzetta A, Boyd RN, Rose SE, Fripp J, et al. PREMM: preterm early massage by the mother: protocol of a randomised controlled trial of massage therapy in very preterm infants. *BMC pediatrics*. 2016;16(1):146.
19. Abdallah B, Badr LK, Hawwari M. The efficacy of massage on short and long term outcomes in preterm infants. *Infant Behavior and Development*. 2013;36(4):662-9.
20. Byers JF. Components of developmental care and the evidence for their use in the NICU. *MCN: The American Journal of Maternal/Child Nursing*. 2003;28(3):174-80.
21. Philpott-Robinson K, Lane SJ, Korostenski L, Lane AE. The impact of the Neonatal Intensive Care Unit on sensory and developmental outcomes in infants born preterm: A scoping review. *British Journal of Occupational Therapy*. 2017;80(8):459-69.
22. Field T, Diego M, Hernandez-Reif M, Dieter JN, Kumar AM, Schanberg S, et al. Insulin and Insulin-Like Growth Factor 1 (IGF-1) increased in preterm neonates. *Journal of developmental and behavioral pediatrics: JDBP*. 2008;29(6):463.
23. Ferber SG, Kuint J, Weller A, Feldman R, Dollberg S, Arbel E, et al. Massage therapy by mothers and trained professionals enhances weight gain in preterm infants. *Early human development*. 2002;67(1-2):37-45.
24. Abigail W, Frank A, Tiffany F. Massage effects on cocaine-exposed preterm neonates. *J Dev Behav Pediatr*. 1993;14:318-22.
25. Wheeden A, Scafidi FA, Field T, Ironson G, Valdeon C, Bandstra E. Massage effects on cocaine-exposed preterm neonates. *Journal of Developmental and Behavioral Pediatrics*. 1993.
26. Moyer-Mileur LJ, Haley S, Slater H, Beachy J, Smith SL. Massage improves growth quality by decreasing body fat deposition in male preterm infants. *The Journal of pediatrics*. 2013;162(3):490-5.
27. Fontana C, Menis C, Pesenti N, Passera S, Liotto N, Mosca F, et al. Effects of early intervention on feeding behavior in preterm infants: A randomized controlled trial. *Early human development*. 2018;121:15-20.
28. Scafidi FA, Field T, Schanberg SM. Factors that predict which preterm infants benefit most from massage therapy. *Journal of Developmental and Behavioral Pediatrics*. 1993.
29. Patole S. *Nutrition for the preterm neonate*: Springer; 2013.
30. Li Y-F, Lin H-C, Torrazza RM, Parker L, Talaga E, Neu J. Gastric residual evaluation in preterm neonates: a useful monitoring technique or a hindrance? *Pediatrics & Neonatology*. 2014;55(5):335-40.
31. Karlberg J, Albertsson-Wikland K, Baber F, Low L, Yeung C. Born small for gestational age: consequences for growth. *Acta Pædiatrica*. 1996;85:8-13.

32. Gh R. Effect of Tactile-Kinesthetic Stimulation in weight gaining of pre-term infants hospitalized in intensive care unit. *Tehran University Medical Journal TUMS Publications*. 2009;67(5):347-52.
33. Rad ZA, Haghshenas M, Javadian Y, Hajiahmadi M, Kazemian F. The effect of massage on weight gain in very low birth weight neonates. *Journal of clinical neonatology*. 2016;5(2):96.
34. Saeadi R, Ghorbani Z, Moghaddam AS. The effect of massage with medium-chain triglyceride oil on weight gain in premature neonates. *Acta Medica Iranica*. 2015:134-8.
35. Kumar J, Upadhyay A, Dwivedi AK, Gothwal S, Jaiswal V, Aggarwal S. Effect of oil massage on growth in preterm neonates less than 1800 g: a randomized control trial. *The Indian Journal of Pediatrics*. 2013;80(6):465-9.
36. Diego MA, Field T, Hernandez-Reif M. Vagal activity, gastric motility, and weight gain in massaged preterm neonates. *The Journal of pediatrics*. 2005;147(1):50-5.
37. Field T. Newborn Massage Therapy. *Int J Ped & Neo Heal*. 2017;1:2-54.
38. Diego MA, Field T, Hernandez-Reif M, Deeds O, Ascencio A, Begert G. Preterm infant massage elicits consistent increases in vagal activity and gastric motility that are associated with greater weight gain. *Acta Paediatrica*. 2007;96(11):1588-91.
39. Lee Y-H, Park BNR, Kim SH. The effects of heat and massage application on autonomic nervous system. *Yonsei medical journal*. 2011;52(6):982-9.

Tables

Table 1: Baseline Clinical Characteristics

Baseline Characteristics	TKS Group [n=30]		Control Group [n=30]		Significance
	No.	%	No.	%	
1.1 Gender					
Male	16	53.3	11	36.7	X ² =1.684 p=0.194
Female	14	46.7	19	63.3	
1.2 Gestational age (weeks)					
Min-Max	28-35		28-35		t=2.198 p=0.032*
Mean±SD	30.3±2.6		31.8±2.4		
1.3 Apgar score 1 minute					
0-3	3	10.0	3	10.0	t=0.950 p=0.346
4-6	11	36.7	9	30.0	
7-10	16	53.3	18	60.0	
Min-Max	1-9		3-10		
Mean±SD	6.3±2.1		6.8±1.9		
1.4 Apgar score 5 minutes					
0-3	1	3.3	0	0.0	t=1.498 p=0.140
4-6	9	30.0	6	20.0	
7-10	20	66.7	24	80.0	
Min-Max	2-10		4-10		
Mean±SD	7.2±1.8		7.9±1.6		
1.5 Birth weight (grams)					
Min-Max	1000.0-1500.0		1000.0-1500.0		t=0.703 p=0.485
Mean±SD	1273.7±167.8		1304.2±168.8		
1.6 Length (cm)					
Min-Max	34.5-44.0		32.0-44.5		t=0.704 p=0.484
Mean±SD	39.3±2.5		39.8±3.0		
1.7 Head circumference (cm)					
Min-Max	25.0-30.0		26.0-32.0		t=0.816 p=0.418
Mean±SD	28.1±1.2		28.4±1.4		
1.8 Temperature at birth (°c)					
Min-Max	33.4-37.8		33.4-37.0		t=0.392 p=0.696
Mean±SD	35.8±1.1		35.7±0.8		
1.9 Clinical risk index for babies (CRIB II)					
7-8	9	30.0	14	46.7	t=2.311 p=0.024*
9-10	4	13.3	7	23.3	
11-12	8	26.7	5	16.7	
13-14	7	23.3	4	13.3	
15-16	2	6.7	0	0.0	
Min-Max	7-16		7-14		
Mean±SD	10.9±2.7		9.5±2.1		

X²: Chi-Square test

t: Student t-test

*significant at p≤0.05

Table 2: Amount of feeds

Amount of feed (mls)	TKS Group [n=30]	Control Group [n=30]	Significance
Day 3			
Min-Max	10.0-191.0	0.0-201.0	Z=1.405
Median (Q1-Q3)	71.5(33.8-110.3)	52.0(11.8-87.0)	p = .160
Day 10			
Min-Max	114-332	0-344	Z=0.399
Median (Q1-Q3)	238.0(188.5-264.0)	220.0(170.3-280.0)	p = .690
Day 17			
Min-Max	092-348	30-360	Z=1.281
Median (Q1-Q3)	288.0(265.5-320.0)	256.0(212.8-323.5)	p = .200
Day 23			
Min-Max	192-368	0-400	Z=0.872
Median (Q1-Q3)	332.0(288.0-360.0)	312.0(266.3-360.0)	p = .383
2.2 Duration to attaining full breastfeeding (days)			
Min-Max	24-68	0-51	t=0.241
Mean±SD	30.4±10.7	29.1±9.6	P = .810

Z: Mann Whitney test

t: Student t-test

*significant at $p \leq 0.05$

Table: 3 Signs of feeding intolerance

Signs of feeding intolerance (gastric residual, abdominal distension, vomiting)	TKS [n=30]		Control [n=30]		Significance
	No.	%	No.	%	
Day 3					
Absent	15	50.0	11	36.7	$X^2=1.090$ P=0.297
Present	15	50.0	19	63.3	
Day 10					
Absent	23	76.7	15	50.0	$X^2=4.590$ P=0.032*
Present	7	23.3	15	50.0	
Day 17					
Absent	30	100.0	21	70.0	$FE_P=0.0009^*$
Present	0	0.0	9	30.0	
Day 23					
Absent	29	96.7	21	70.0	$X^2=7.680$ P=0.006*
Present	1	3.3	9	30.0	

X^2 : Chi-Square test

FE_P : Fisher's Exact test

*significant at $P \leq 0.05$

Table 4: Weight of moderate and late preterm neonates

Outcome (n)	Moderate preterm (28-<32weeks) [n=33]				Late preterm (32<37weeks) [n=27]			
	TKS Group [n=20]	Control Group [n=13]	Significance		TKS Group [n=10]	Control Group [n=17]	Significance	
			t	P			Z	P
Weight (kg)	1216.5±158.5	1312.0±144.4	1.750	0.090	1388.0±126.3	1298.2±189.6	1.143	0.253
Day 3	1164.0±160.3	1262.3±131.7	1.841	0.075	1418.5±118.7	1268.8±211.9	1.734	0.083
Day 5	1180.5±168.3	1272.3±145.7	1.611	0.117	1449.1±125.6	1300.3±225.6	1.734	0.083
Day 7	1191.0±191.5	1278.9±127.8	1.453	0.156	1454.5±125.3	1325.3±229.1	1.281	0.200
Day 9	1228.8±185.6	1299.2±139.9	1.168	0.252	1468.5±111.5	1341.2±229.1	1.155	0.248
Day 11	1273.3±204.2	1323.9±159.2	0.755	0.456	1492.1±105.7	1372.7±237.6	1.256	0.209
Day 13	1307.8±213.7	1372.3±182.2	0.896	0.377	1543.5±95.2	1387.1±248.4	1.633	0.103
Day 15	1365.8±214.7	1420.8±170.4	0.777	0.443	1592.5±106.6	1412.1±243.1	1.984	0.047*
Day 17	1408.0±214.5	1466.9±189.3	0.806	0.426	1639.5±92.1	1471.5±248.8	1.507	0.132
Day 19	1451.3±218.2	1493.1±196.8	0.559	0.580	1684.5±85.7	1505.6±238.2	1.935	0.053
Day 21	1497.8±218.2	1547.3±196.5	0.641	0.526	1723.5±85.1	1510.9±266.1	2.034	0.042*
Day 23	1555.4±231.0	1581.9±195.2	0.342	0.735	1767.5±79.3	1565.9±238.6	2.362	0.018*

t: Student t-test

Z: Mann Whitney test

*significant at P≤0.05

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.doc](#)