

Echocardiography-Guided Hemodynamic Management of Severe Sepsis and Septic Shock in Adults: A Randomized Controlled Trial

Walid S. Alhabashy

Tanta University Faculty of Medicine

Osama M. Shalaby

Tanta University Faculty of Medicine

Ahmed S. Elgebaly (✉ elgebaly_13@hotmail.com)

Tanta University Faculty of Medicine <https://orcid.org/0000-0001-9826-6365>

Mohammed S. Abd El Ghafar

Tanta University Faculty of Medicine

Research

Keywords: Echocardiography, EGDT, Severe sepsis, Septic Shock, Adults

Posted Date: October 12th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-17392/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Echocardiography (ECHO) is used to guide septic shock resuscitation, but without evidence for efficacy. Therefore, we compared the outcome of early goal-directed therapy (EGDT) and ECHO-guided management of hemodynamics in severe sepsis and septic shock.

Materials and Methods: This is a single center, randomized controlled trial conducted on 100 adult patients with severe sepsis or septic shock. Patients were assessed and treated with either EGDT protocol (EGDT group) or ECHO-guided resuscitation protocol (ECHO group).

Results: Only 87 patients (45 in group I and 42 in group II) were analyzed. There was a significant increase of mean norepinephrine and dobutamine doses and a significant decrease in total fluids in the first 24 hours, time to normalization, time to weaning of vasopressors, total MV days, MV free days and ICU and hospital stays in ECHO group. At 30 days, the mortality rate in EGDT group was 35.6% which was significantly higher compared to 14.3% in ECHO group. At 90 days, the overall mortality was significantly higher in EGDT group compared to Echo group (40.0% vs 16.7% respectively). Hazardous ratio of mortality was 1.630 (95% confidence interval (CI): 1.123 - 2.366) and 1.653 (95% CI: 1.137 - 2.404) at 30 and 90 days respectively in EGDT group compared to ECHO group.

Conclusions: In severe sepsis and septic shock, ECHO-guided management of hemodynamics resulted in a decrease in mortality, lower total fluid intake, higher vasopressor and inotrope support, earlier weaning of vasopressors and less MV days, ICU and hospital stay.

Introduction

Severe sepsis and septic shock are common in critically ill patients and are on top of the causes of mortality in intensive care unit (ICU) [1]. Vasodilation, increased permeability, hypovolemia, and ventricular dysfunction are the main findings in septic shock [2].

A paradigm of “early goal-directed therapy” (EGDT) has been dominated in sepsis resuscitation for about fifteen years based on the results of a single-center, randomized, “usual care” controlled study [3]. EGDT was not, however, repeated in large, multinational, multicenter studies, even the patients received more intensive fluid in the “usual care” group than the control group in the first study [4-7].

Due to the failure of several negative randomized controlled trials to obtain similar benefits as the initial trial, the ideal approach to resuscitate patients with septic shock is still uncertain. Research is ongoing to find an optimal balance between administering large volumes of intravenous fluids that can have deleterious effects, or small volumes of fluids that fail to support adequate organ perfusion [8].

Previously, the invasive assessment of hemodynamic parameters using central venous catheters and/or pulmonary artery catheters allowed clinicians to define cardiovascular physiology and dominated fluid management for many years [9]. However, utilization of such techniques has significantly decreased due

to risks associated with their use. In addition, their “static” variable outcomes poorly predicted fluid responsiveness, and prospective studies showed no benefit from their routine use [10-15].

Nowadays, critical care physicians are increasingly employing the echocardiography (ECHO) in the non-invasive assessment of hemodynamic parameters during hemodynamic instability. Many clinicians advocate the use of ECHO as a vital tool in the management of the critically ill patient [16, 17].

Despite the adequacy of focused ECHO in the early stages of septic shock, nevertheless, a comprehensive systemic ECHO assessment of cardiac output, left and right ventricular systolic functions, volume status and filling pressure is crucial to provide a comprehensive hemodynamic management. Unfortunately, outcome studies on the utilization of ECHO in septic shock are not adequate and are therefore strongly required [18].

The aim of this study was to compare the outcome between EGDT and ECHO-guided management of hemodynamics in severe sepsis and septic shock in adult patients.

Methods

This randomized controlled clinical trial was conducted from March 2015 to May 2016 at surgical ICU. After ethics committee approval, registration on PACTR (PACTR201902680224481) and obtaining of an informed written consent from each patient either by the patient or his next of kin. All data of patients were confidential with secret codes and private file for each patient. All given data were used for the current medical research only.

The target population included 100 patients aged 18 to 60 years, admitted or planned for admission to ICU for an episode of severe sepsis and septic shock. Severe sepsis and septic shock was defined by the American College of Chest Physicians/Society of Critical Care Medicine consensus criteria ^[19]. Patients met criteria for inclusion if they had (1) a suspected infection, (2) two or more systemic inflammatory response syndrome criteria, and (3) either: (a) had severe sepsis (end-organ dysfunction) or (b) had septic shock (a systolic blood pressure less than 90 mmHg despite an intravenous fluid challenge of at least 20 ml/kg with evidence of organ dysfunction or hyperlactatemia).

Exclusion criteria were patient refusal to be included, known history of any cardiac disease, acute coronary syndrome, cardiac dysrhythmias (as a primary diagnosis), acute pulmonary edema, status asthmaticus, body mass index > 35, severe respiratory disorders or high PEEP requirements on mechanical ventilation (MV), liver insufficiency, multi-organ system failure, contraindication to central venous catheterization, active gastrointestinal hemorrhage, and/or do-not-resuscitate status.

After stabilization of the airway and breathing, standard continuous monitoring of ECG, respiratory rate (RR), oxygen saturation and invasive arterial blood pressure was done. Then venous access and initiation of fluid resuscitation and empirical antibiotics were done.

Patients were randomly allocated into two equal groups (n = 50). Randomization was accomplished by a volunteer, not sharing in the study by computer generated sequence through sealed opaque envelopes.

Group I (EGDT group):

In which the surviving sepsis campaign dependent resuscitation protocol was applied, targeting all of the following as a part of a stepwise treatment protocol:

500-ml bolus of crystalloid was given every 30 minutes to achieve a central venous pressure (CVP) of 8 to 12 mmHg. If mean arterial pressure (MAP) was less than 65 mmHg, vasopressors (noradrenaline 0.05-0.3 µg/kg/min) were given to maintain a MAP of at least 65 mmHg. If the central venous oxygen saturation (ScvO₂) was less than 70 %, red cells were transfused to achieve a hematocrit of at least 30 %. After the CVP, MAP, and hematocrit were thus optimized, if the ScvO₂ was less than 70 %, Dobutamine administration was started at a dose of 2.5 µg/kg/min, a dose that was increased by 2.5 µg/kg/min every 30 minutes until the ScvO₂ was 70 % or higher or until a maximal dose of 20 µg/kg/min was given. Dobutamine was decreased in dose or discontinued if the heart rate (HR) was above 120 beats per minute (Fig. 1) [3].

Group II (ECHO group):

In which a transthoracic bedside focused echocardiographic assessment of the patient was done, as a baseline ECHO in a five-step approach to monitor hemodynamics using Philips (CX50 – Extreme edition) machine with S5-1 ECHO probe.

Step 1: Starting point was to detect potential signs of pre-existing chronic cardiac dysfunction that needs a full formal ECHO study and exclude the patient from this study, as these findings can mislead in interpretation of subsequent findings (i.e. primary cardiogenic cause of shock, instead of sepsis, LV or LA significant dilatation, and LV marked hypertrophy are signs of chronic volume/pressure overload; RA significant dilatation, RV dilatation and hypertrophy have the same meaning for right-side chronic disease (isolated RV dilatation can vice versa be a sign of acute RV dysfunction).

Step 2: LV/RV contractility assessment by eyeballing categorizing into hyperdynamic, good and poor function with Step 3: LVOT VTI assessment by Echo Doppler, a low output state can then be ascribed to sepsis-related LV systolic dysfunction (± RV dysfunction) or isolated RV dysfunction and treated with inotropic support ± vasopressors (depending on MAP after inotropic support).

Low output with evidence of normal biventricular systolic function should prompt investigation of fluid responsiveness (Step 4): Fluid responsiveness assessed by 3 parameters A) IVC Collapsibility (Fig. 2A, Fig. 2B) B) passive leg raise test and C) velocity time integral (VTI) variation on the left ventricular outflow tract (LVOT) (Fig. 2C) if patient is fluid responder he received a fluid infusion of 500 ml of crystalloids.

When inadequacy of global perfusion and/or hypotension is associated with a non-low output state, persistent preload defect should be investigated (again step 4) and if detected corrected.

If this is not the case, an exclusion diagnosis of vasodilatation is made (Step 5), and systemic arterial tone corrected with upward titration of vasopressors. Whenever this is done, LV Systolic/Diastolic functions should subsequently be re-assessed, as normalization of LV afterload can unmask sepsis related myocardial dysfunction and diastolic function was assessed using spectral doppler on mitral inflow and using tissue doppler of the septal mitral annulus in apical four chamber view to measure e' and a' velocities (Fig. 3).

Echocardiographic parameters: were repeated before each change in the management of fluids, vasopressors or inotropes, then just before discharge from the ICU.

IVC Collapsibility diameter < 2 cm and index $> 50\%$ indicates fluid responsiveness in spontaneously breathing patients. IVC distensibility index $> 18\%$ indicates fluid responsiveness in MV patients. Passive leg raising test $> 10\%$ increase in SV indicates fluid responsiveness (Fig. 4).

In both groups:

The source of sepsis was early treated or eradicated according to the situation. The time of the study was the first 24 hours.

Measurements:

Demographic characteristics and clinical parameters (HR, RR, core temperature, invasive MAP, CVP and urine output) were recorded. Acute Physiology and Chronic Health Evaluation Score II (APACHE II) was calculated on 24 hours of study enrolment in both groups. Laboratory parameters (complete blood count, serum urea and creatinine) were done daily for three days to be incorporated in the severity scoring system. Also, central venous saturation, arterial lactate level and arterial blood gases were recorded. Total fluid requirements in first 24 hours from initiation of therapy, vasopressors and inotropic drugs requirements in first 24 hours, time till normalization of the tissue perfusion indicators, time till weaning of vasopressors and inotropic drugs, MV and ventilator-free days, ICU length of stay, mortality at 30 and 90 days of admission were recorded.

The primary outcome was mortality at 30 days and the secondary outcomes were mortality at 90 days, the total fluid intake, MV-free days and duration of ICU and hospital stay.

Statistical analysis:

The sample size calculation was performed using G. power 3.1.9.2. The sample size was calculated on the following considerations: 0.05 α error and 80% power of the study and group ratio 1:1 to demonstrate a 25% decrease (expected) in mortality (the primary outcome) with ECHO-guided compared to EGDT

management of hemodynamics (35% according to a previous study [20]). To overcome dropout, 7 cases were added to each group. Therefore, 50 patients were recruited in each group.

The collected data were organized, tabulated and statistically analyzed using SPSS version 20 (IBM® SPSS® Inc., Chicago, Illinois, USA). All data were assessed for normal distribution using Shapiro-Wilks test. Quantitative parametric data were presented as mean \pm standard deviation and compared by unpaired student (t) test. Quantitative non-parametric data were presented as median and interquartile range and compared by Mann-Whitney (U) test. Qualitative variables were expressed as frequency and percent and were compared by Chi-square (X^2) test. A P value <0.05 was considered statistically significant.

Results

In this study, only 87 patients with severe sepsis/septic shock were analyzed; 45 patients in group I (5 cases were withdrawn) and 42 patients in group II (8 cases dropped out; 1 case with poor window, 4 cases with chronic cardiac dysfunction and 3 cases were withdrawn). (Fig. 5)

Both groups in our study were matched in the baseline characteristics (age, sex, BMI and body surface area (BSA). The source of sepsis, type of organism, need for mechanical ventilation and APACHE II score were comparable between both groups. (Table 1)

MAP showed significant increase in ECHO group compared to EGDT group at 6 and 12 h. In each group, there was significant increase in MAP at 6, 12 and 24 h compared to baseline values. HR showed significant decrease in ECHO group compared to EGDT group at 6, 12 and 24 h. In addition, there was significant progressive decrease in each group at 6, 12, and 24 hours compared baseline values. Core body temperature showed no significant differences at baseline and at 6, 12 and 24 h between both groups. [Table (2)]

CVP was significantly lower in EGDT group compared to ECHO group at baseline. Then, values in EGDT progressively increased with time, while values in ECHO increased at 6 h, then decreased nearly to baseline values at 12 h and progressively decreased at 24 h. The difference between both groups was non-significant at 6 h, while at 12 and 24 h, there was significant decrease in ECHO group compared to EGDT group. [Table (2)]

ScvO₂ showed significant increase in ECHO group compared to EGDT group at 6, 12 and 24 h. Lactate concentration showed significant decrease at 24 h in ECHO group compared to EGDT group. pH was significantly higher in ECHO group at 12 and 24 h. Urine output showed no significant difference between both groups at the first 6 h. However, urine output was significantly increased in Echo group compared to EGDT group at 12 and 24 h. [Table (2)]

There was a significant increase in mean norepinephrine and dobutamine doses in Echo group compared to EGDT group. In addition, there was significant decrease in total fluids in the first 24 hours, time to

normalization, time to weaning of vasopressors, total MV days, MV free days and ICU and hospital stays in Echo group compared to EGDT group. [Table (3)]

At 30 days, mortality rate in EGDT group was 35.6% (16 patients) which was significantly higher compared to 14.3% (6 patients) in Echo group. At 90 days, the overall mortality was significantly higher in EGDT group compared to Echo group (40.0%; 18 patients vs 16.7%; 7 patients respectively). [Table (3)] Hazardous ratio of mortality was 1.630 (95% confidence interval (CI): 1.123 - 2.366) and 1.653 (95% CI: 1.137 - 2.404) at 30 and 90 days respectively in EGDT group compared to ECHO group.

Discussion

In the present work, we proposed that the use of ECHO to guide treatment of sepsis and septic shock will be associated with better outcome. Thus, we designed the present study to examine this hypothesis in the light of available gold standard of sepsis treatment (i.e. EGDT) at the time of the study. We prospectively evaluated a group of patients whose treatment was guided by ECHO and compared the results with a group of patients who received EGDT.

Our results regarding the mortality is comparable to those reported by Chertoff et al. [21] who reported a mortality rate of 29.69%. In addition, the reported incidence of mortality in the present work lies within reported range in literature; the short-term mortality is 20% to 30%, and up to 50% in patients with septic shock [1, 22].

In contrary to Lanspa et al. [8] who reported insignificant difference between ECHO-guided resuscitation compared to EGDT in mortality, ICU stay or lactate clearance. This could be a result of late ECHO assessment after initial resuscitation as no difference in fluids administration between the two groups in contrary to our study results which showed a statistically significant difference between groups in the fluid resuscitation volume.

The present work revealed that the baseline MAP of both groups shows no significant difference. Then the MAP of the ECHO group became higher significantly than that of the EGDT group after 6 and 12 hours of the study. Thereafter, the MAP of both groups shows no significant difference at 24 hours from enrolment. This means that the ECHO group reached the target MAP earlier.

As well, Lin [23] and his colleagues observed improvement of blood pressure of the intervention group when they tested the effect of modified goal-directed protocol (targeting specific MAP, CVP and UOP without targeting ScvO₂ on the clinical outcome of septic shock patients. However, the MAP of the intervention group stayed significantly higher throughout the study [23].

In the current work, there was no statistically significant difference between both groups as regards the HR at the start of the study, then the HR starting from 6 hours till 24 hours after the enrolment differed significantly between both groups.

The significant decrease in the HR in the ECHO group was shown as well when the HR measurement was compared to the baseline HR at enrolment in the study. This decrease in the HR in the ECHO group can be a preceding indicator of the improvement of the hemodynamics which also correlated with higher MAP in the ECHO group.

The fluid therapy in the EGDT group was according to the CVP (targeting CVP between 8 - 12 mm Hg). This CVP guidance made the fluid therapy relatively higher in the EGDT group. Most of recent studies that test the CVP in guiding fluid therapy criticize the CVP as a predictor of fluid responsiveness as it may be falsely high in volume-depleted patient hindering useful fluid resuscitation or it may be falsely low in volume overloaded patient exposing the patient to more overload or even pulmonary congestion and edema. However, it is still used by many clinicians in ICU.

In agreement with our results, a study by Feng et al. [24] who reported that those who had transthoracic ECHO (TTE) had a higher maximum dose of norepinephrine, but surprisingly were weaned of vasopressors earlier compared to the no TTE group. Dobutamine was used more often in the group who received TTE, they concluded that performance of TTE is associated with a 28-day mortality benefit in a general population of septic, critically ill patients [24].

This is also in agreement with Kanji et al. [25] reported more utilization of Dobutamine in the limited ECHO group compared to the standard management.

In the present study, we reported significantly lower total MV days, MV free days and ICU and hospital stays in ECHO group compared to EGDT group due to earlier weaning from vasopressors and MV.

This is in disagreement with Lanspa et al. [26] who reported an insignificant difference between the echo group and non-echo group (median of 28 days versus 25 days; $p=0.51$). Also, Feng et al. [24] reported no significant difference in ventilation-free days between ECHO group and non-ECHO groups.

Overall, results of the present work indicated that, ECHO as a sole monitoring and guiding tool was associated with better outcome than EGDT as evidenced from enhancement of vital data over time, decreased total fluid administration, significant decrease of 30 day and overall mortality. Critical care echocardiography may be considered the fifth pillar of clinical examination especially in critically ill patients [27, 28].

Finally, it may be the first time to compare outcome between EGDT and use of echocardiographic findings to govern the progress of management in sepsis and septic shock. Results of the present work seems to be promising. However, further studies with larger sample size and in multi-centers are needed to generalize the routine use of echocardiography in management of hemodynamics in severe sepsis and septic shock provides. Anyway, present work advocates routine uses of echocardiography as a crucial, non-invasive bedside tool for the management of patients with severe sepsis and septic shock.

Another limitation of our study is the use of old definitions of sepsis, severe sepsis and septic shock and it's preferred to depend on the new definitions for sepsis and septic shock provided by the International Consensus (Sepsis-3) in 2016 [29].

Further studies are needed to reveal the role of early diagnosis by new scores as Quick Sequential Organ Function Assessment (qSOFA) and qSOFA-65 [30] which helps in prediction and thereby early management.

Limitations in ECHO use still exist; low echogenicity at surface examination. Whenever there is strict requirement of continuous monitoring of cardiac output or pulmonary artery pressure, echocardiography is not the right tool. Also, in centers where adequate training on the use of critical care echocardiography does not exist, and when repeated bedside assessments of hemodynamic variables are required echocardiography is of limited use as a single monitoring tool. Patients excluded from the study e.g. morbid obese and atrial fibrillation still represent challenges in echocardiographic assessments and follow up either due to inconsistency of cardiac output on LVOT VTI or limited window for examination respectively.

Conclusion

In severe sepsis and septic shock, ECHO-guided management of hemodynamic provides additional benefits over EGDT. It is a non-invasive, reproducible, readily available tool with a resultant decrease in mortality and favorable outcomes in the term of lower total fluid intake, earlier weaning of vasopressors and less MV days, ICU and hospital stay. ECHO-guided management should be a routine care in the management of hemodynamics in severe sepsis and septic shock patients as early as possible in the first 24 hours.

Declarations

Ethics approval: 2932/12/14 (Tanta University Research Ethics Committee at 12/2014) and retrospectively registered in Pan African Clinical Trials Registry, (Identifier: PACTR201902680224481, date of registration: 6/12/2018, url: <https://pactr.samrc.ac.za/TrialDisplay.aspx?TrialID=5786>)

Consent to participate: After ethics committee approval and obtaining of an informed written consent from each patient either by the patient or his next of kin

Consent for publication: Contributors agreed to publish this work

Availability of data and material: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: There are no conflicts of interest

Funding: No financial support

Authors' contributions: All authors contributed in writing the manuscript. They read and approved the final manuscript.

Acknowledgements: Nil

References

- [1] Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med*. 2001;29:1303-10.
- [2] Russell JA, Rush B, Boyd J. Pathophysiology of Septic Shock. *Crit Care Clin*. 2018;34:43-61.
- [3] Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345:1368-77.
- [4] Mouncey PR, Osborn TM, Power GS, Harrison DA, Sadique MZ, Grieve RD, et al. Trial of early, goal-directed resuscitation for septic shock. *N Engl J Med*. 2015;372:1301-11.
- [5] Peake SL, Delaney A, Bailey M, Bellomo R, Cameron PA, Cooper DJ, et al. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med*. 2014;371:1496-506.
- [6] Rowan KM, Angus DC, Bailey M, Barnato AE, Bellomo R, Canter RR, et al. Early, Goal-Directed Therapy for Septic Shock - A Patient-Level Meta-Analysis. *N Engl J Med*. 2017;376:2223-34.
- [7] Yealy DM, Kellum JA, Huang DT, Barnato AE, Weissfeld LA, Pike F, et al. A randomized trial of protocol-based care for early septic shock. *N Engl J Med*. 2014;370:1683-93.
- [8] Lanspa MJ, Burk RE, Wilson EL, Hirshberg EL, Grissom CK, Brown SM. Echocardiogram-guided resuscitation versus early goal-directed therapy in the treatment of septic shock: a randomized, controlled, feasibility trial. *J Intensive Care*. 2018;6:50.
- [9] Kortgen A, Niederprum P, Bauer M. Implementation of an evidence-based "standard operating procedure" and outcome in septic shock. *Crit Care Med*. 2006;34:943-9.
- [10] Harvey S, Harrison DA, Singer M, Ashcroft J, Jones CM, Elbourne D, et al. Assessment of the clinical effectiveness of pulmonary artery catheters in management of patients in intensive care (PAC-Man): a randomised controlled trial. *Lancet*. 2005;366:472-7.
- [11] Micek ST, Roubinian N, Heuring T, Bode M, Williams J, Harrison C, et al. Before-after study of a standardized hospital order set for the management of septic shock. *Crit Care Med*. 2006;34:2707-13.
- [12] Richard C, Warszawski J, Anguel N, Deye N, Combes A, Barnoud D, et al. Early use of the pulmonary artery catheter and outcomes in patients with shock and acute respiratory distress syndrome: a

randomized controlled trial. JAMA. 2003;290:2713-20.

[13] Shah MR, Hasselblad V, Stevenson LW, Binanay C, O'Connor CM, Sopko G, et al. Impact of the pulmonary artery catheter in critically ill patients: meta-analysis of randomized clinical trials. JAMA. 2005;294:1664-70.

[14] Shapiro NI, Howell MD, Talmor D, Lahey D, Ngo L, Buras J, et al. Implementation and outcomes of the Multiple Urgent Sepsis Therapies (MUST) protocol. Crit Care Med. 2006;34:1025-32.

[15] Swan HJ, Ganz W, Forrester J, Marcus H, Diamond G, Chonette D. Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. N Engl J Med. 1970;283:447-51.

[16] Cholley BP, Vieillard-Baron A, Mebazaa A. Echocardiography in the ICU: time for widespread use! Intensive Care Med. 2006;32:9-10.

[17] Seppelt IM. All intensivists need echocardiography skills in the 21st century. Crit Care Resusc. 2007;9:286-8.

[18] Via G, Price S, Storti E. Echocardiography in the sepsis syndromes. Crit Ultrasound J. 2011;3:71-85.

[19] Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. Chest. 1992;101:1644-55.

[20] Early Goal-Directed Therapy Collaborative Group of Zhejiang Province. [The effect of early goal-directed therapy on treatment of critical patients with severe sepsis/septic shock: a multi-center, prospective, randomized, controlled study]. Zhongguo wei zhong bing ji jiu yi xue = Chinese critical care medicine = Zhongguo weizhongbing jijiuyixue. 2010;22:331-4.

[21] Chertoff J, Chisum M, Simmons L, King B, Walker M, Lascano J. Prognostic utility of plasma lactate measured between 24 and 48 h after initiation of early goal-directed therapy in the management of sepsis, severe sepsis, and septic shock. J Intensive Care. 2016;4:13.

[22] Annane D, Bellissant E, Cavaillon JM. Septic shock. Lancet. 2005;365:63-78.

[23] Lin SM, Huang CD, Lin HC, Liu CY, Wang CH, Kuo HP. A modified goal-directed protocol improves clinical outcomes in intensive care unit patients with septic shock: a randomized controlled trial. Shock. 2006;26:551-7.

[24] Feng M, McSparron JI, Kien DT, Stone DJ, Roberts DH, Schwartzstein RM, et al. Transthoracic echocardiography and mortality in sepsis: analysis of the MIMIC-III database. Intensive Care Med. 2018;44:884-92.

- [25] Kanji HD, McCallum J, Sirounis D, MacRedmond R, Moss R, Boyd JH. Limited echocardiography-guided therapy in subacute shock is associated with change in management and improved outcomes. *J Crit Care.* 2014;29:700-5.
- [26] Lanspa MJ, Pittman JE, Hirshberg EL, Wilson EL, Olsen T, Brown SM, et al. Association of left ventricular longitudinal strain with central venous oxygen saturation and serum lactate in patients with early severe sepsis and septic shock. *Critical Care.* 2015;19:304.
- [27] Narula J, Chandrashekhar Y, Braunwald E. Time to Add a Fifth Pillar to Bedside Physical Examination: Inspection, Palpation, Percussion, Auscultation, and Insonation. *JAMA Cardiol.* 2018;3:346-50.
- [28] Mayo P, Arntfield R, Balik M, Kory P, Mathis G, Schmidt G, et al. The ICM research agenda on critical care ultrasonography. *Intensive Care Med.* 2017;43:1257-69.
- [29] Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *Jama.* 2016;315:801-10.
- [30] Lee J, Song JU. Performance of a quick sofa-65 score as a rapid sepsis screening tool during initial emergency department assessment: A propensity score matching study. *J Crit Care.* 2020;55:1-8.

Tables

Table 1: Patients' characteristic of the studied patients

		Group I (EGDT) (n=45)	Group II (ECHO) (n=42)	Test	P
Age (years)		44.44 ± 7.65	42.33 ± 8.96	T = 0.18	0.24
Sex	Male	23 (51.1%)	24 (57.1%)	χ ² =0.31	0.57
	Female	22 (48.9%)	18 (42.9%)		
BMI (kg/m²)		27.93 ± 3.04	27.90 ± 2.73	T = 0.04	0.96
BSA (m²)		1.90 ± 0.14	1.90 ± 0.13	T = 0.14	0.88
Source of sepsis	Respiratory tract	19 (42.2%)	16 (38.1%)	χ ² =2.98	0.81
	Soft tissue	11 (24.4%)	7 (16.7%)		
	Blood	5 (11.1%)	6 (14.3%)		
	Urinary tract	4 (8.9%)	6 (14.3%)		
	Abdomen	4 (8.9%)	5 (11.9%)		
	Endocarditis	0 (0%)	1 (2.4%)		
	Unknown	2 (4.4%)	1 (2.4%)		
Type of organism	Gram positive	18 (40.0%)	14 (33.3%)	χ ² =0.59	0.74
	Gram negative	16 (35.6%)	15 (35.7%)		
	Mixed	11 (24.4%)	13 (31%)		
Need for mechanical ventilation		41 (91.1%)	38 (90.5%)	FE: —	0.918
APACHE II score		24.76 ± 4.26	23.95 ± 3.45	T = 0.79	0.42

BSA was calculated by Mosteller Formula ($BSA=0.01667 \times W^{0.5} \times H^{0.5}$). EGDT: early goal directed therapy group; Echo: echocardiography group; BMI: body mass index; BSA: body surface area, T: Student's t test, X²: Chi-square, FE: Fisher's Exact test

Table 2: Vital signs, central venous pressure, central venous oxygen saturation, lactate concentration, pH and urine output of the studied patients

	Group I (EGDT) (n=45)	Group II (ECHO) (n=42)	T	P
Mean arterial pressure (mmHg)				
Baseline	58.31 ± 6.29	56.57 ± 7.49	1.17	0.24
At 6 hours	66.31 ± 4.32	74.98 ± 9.24	5.65	<0.001
At 12 hours	67.91 ± 3.30	81.43 ± 12.06	7.23	<0.001
At 24 hours	69.51 ± 2.46	70.05 ± 2.09	1.09	0.27
Heart rate (beats/min)				
Baseline	119.78 ± 13.23	113.62 ± 16.04	1.97	0.52
At 6 hours	116.02 ± 10.96	100.05 ± 9.77	7.15	<0.001
At 12 hours	107.47 ± 10.63	95.31 ± 7.8	6.04	<0.001
At 24 hours	106.82 ± 22.92	84.90 ± 7.85	5.88	<0.001
Core body temperature (°C)				
Baseline	37.98 ± 1.34	38.16 ± 1.25	0.62	0.53
At 6 hours	37.54 ± 1.08	37.43 ± 1.24	0.41	0.67
At 12 hours	37.46 ± 0.58	37.19 ± 0.74	1.92	0.06
At 24 hours	37.47 ± 0.62	37.24 ± 0.65	1.65	0.10
Central venous pressure (mmHg)				
Baseline	6.89 ± 2.87	10.38 ± 4.03	4.67	<0.001
At 6 hours	11.2 ± 2.39	11.19 ± 3.44	0.02	0.98
At 12 hours	11.73 ± 1.94	10.38 ± 3.13	2.43	0.017
At 24 hours	12.07 ± 1.87	9.6 ± 2.84	4.81	<0.001
Central venous oxygen saturation (%)				
Baseline	61.2 ± 7.43	65.9 ± 3.47	1.57	0.508
At 6 hours	65.82 ± 5.82	69.37 ± 5.36	3.81	0.004
At 12 hours	65.82 ± 5.82	70.79 ± 4.00	2.79	0.006
At 24 hours	68.36 ± 4.97	71.24 ± 4.62	2.69	0.006

Lactate concentration (mmol/L)				
Baseline	6.02 ± 1.51	6.38 ± 1.62	1.07	0.28
At 6 hours	5.03 ± 1.27	5.26 ± 1.68	0.71	0.47
At 12 hours	4.54 ± 1.42	4.08 ± 1.24	1.58	0.12
At 24 hours	3.74 ± 2.42	2.32 ± 0.61	3.68	<0.001
pH				
Baseline	7.16 ± 0.12	7.2 ± 0.09	1.95	0.54
At 12 hours	7.23 ± 0.09	7.28 ± 0.07	1.99	0.049
At 24 hours	7.27 ± 0.08	7.36 ± 0.06	5.46	<0.001
Urine output (mL)				
At 6 hours	216.24 ± 140.13	254.33 ± 190.46	1.06	0.28
At 12 hours	414.67 ± 284.33	622.74 ± 189.19	3.98	<0.001
First day	751.31 ± 485.13	1070.55 ± 338.23	3.71	<0.001

Data are presented as mean ± SD, T: Student's t test

Table 3: Outcomes of the studied patients

	Group I (EGDT) (n=45)	Group II (ECHO) (n=42)	Test	p
Fluids in the first 24 hours (mL)	3635.56 ± 973.32	2564.29 ± 927.58	T = 5.24	<0.001
Norepinephrine dose (µg/kg/h)	0.1 (0.05 - 0.2)	0.3 (0.2 - 0.34)	U = 308.5	<0.001
Dobutamine dose (µg/kg/h)	2.50 (0 - 7.5)	7.50 (5 - 10)	U = 538.0	<0.001
Time to normalization	36 (27.2 - 66.8)	12 (7 - 16)	U = 114.5	<0.001
Time to weaning vasopressors (h)	48 (24 - 72)	24 (12 - 24)	U = 231.0	<0.001
Total MV days	10 (6 - 12)	7 (5 - 8)	U = 501.0	<0.001
MV free days	3 (0 - 4)	2 (2 - 3)	U = 1287.5	0.026
ICU stay (d)	13 (8 - 16)	10 (7.25 - 11)	U = 603.5	0.004
Hospital stay (d)	18 (10 - 21)	13 (10 - 14)	U = 595.5	0.003
Mortality at 30 days	16 (35.6%)	6 (14.3%)	χ ² = 5.20	0.023
Mortality at 90 days	18 (40.0%)	7 (16.7%)	χ ² = 5.77	0.016

Data are presented as mean ± SD, median (IQR) or number (%). IQR: Interquartile range, T: Student's t test, U: Mann Whitney test

Figures




Figure 1

Early goal directed therapy (EGDT) protocol in severe sepsis /septic shock




Figure 2

IVC Collapsibility is showing non-fluid responder in (a) compared to potential fluid responder in (b), LVOT VTI variability is showing a fluid responsive patient (c)




Figure 3

Echocardiography-guided algorithm in severe sepsis/septic shock




Figure 4

Stroke volume calculation using the left ventricular outflow tract (LVOT) velocity time integral (VTI) and LVOT cross sectional area (CSA)




Figure 5

Patient flowchart of the study groups