

Sex differences in the association between self-rated health and high-sensitivity C-reactive protein levels in Koreans: A cross-sectional study using data from the Korea National Health and Nutrition Examination Survey

Se-Won Park

Dongguk University Bundang Oriental Hospital

Seong-Sik Park

Dongguk University Bundang Oriental Hospital

Eun-Jung Kim

Dongguk University Bundang Oriental Hospital

Won-Suk Sung

Dongguk University Bundang Oriental Hospital

In-Hyuk Ha

Jaseng Spine and Joint Research Institute

Boyoung Jung (✉ happiness630@hanmail.net)

Hanyang Women's University <https://orcid.org/0000-0001-9812-0745>

Research

Keywords: High-sensitivity C-reactive protein, Korea National Health and Nutrition Examination Survey, Self-rated health, Cross-sectional study

DOI: <https://doi.org/10.21203/rs.3.rs-17466/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: No studies have investigated the association between self-rated health (SRH) and high-sensitivity C-reactive protein (hs-CRP) levels in South Koreans. We explored this association and analyzed differences between sexes. **Methods:** Using cross-sectional data from the 2015–2017 Korea National Health and Nutrition Examination Survey, we analyzed the association between SRH and high hs-CRP levels (>1.0 mg/L) in 14,544 Koreans aged ≥ 19 years who responded to the SRH survey and had hs-CRP test results. Differences in sociodemographic factors were analyzed using Pearson's chi-square test for categorical variables or the Mann-Whitney U test for continuous variables. Multiple logistic regression analysis was used to measure the association between hs-CRP levels and SRH according to sex while adjusting for other possible confounders. **Results:** The percentage of having a very poor to poor SRH was higher in the high hs-CRP level group (22.4%) than in the low hs-CRP level group (17.66%). Among men, the risk of a high hs-CRP level increased with worse SRH (adjusted for confounders; P for trend <0.001). After adjusting for all confounders, including chronic diseases, men with a very poor SRH showed a higher odds ratio (OR) for high hs-CRP levels than those with a very good SRH (fully adjusted OR, 1.74; 95% confidence interval, 1.04–2.90). Significant correlations were absent among women. **Conclusions:** A poor SRH was correlated with low-grade inflammation (high hs-CRP level) among male Korean adults. The findings could be useful for developing health improvement programs and in goal setting at a national scale.

Background

Self-rated health (SRH) is an index utilized worldwide to summarize how patients perceive their overall health status [1]. SRH is an independent predictor of mortality and disease morbidity, even after adjusting for demographic, sociological, and medical risk factors [2]. Despite criticisms that SRH is assessed based on a single question, it is known to be a strong predictor in both healthy and unhealthy individuals. SRH is not only a predictor of previously diagnosed disease but also a predictor of reactions associated with the progression of disease in the premorbid stage; it encapsulates recent or sporadic health issues that may be missed by one-time objective testing, and it also reflects behavioral and emotional factors [2, 3].

C-reactive protein (CRP) is produced by hepatocytes following acute tissue injury or infection. Though CRP levels are generally elevated in cases of severe inflammation, high-sensitivity CRP (hs-CRP) levels increase nonspecifically in the event of inflammation in the body. In particular, hs-CRP is used as an indicator to assess the risk of cardiovascular disease (CVD), and several studies have suggested hs-CRP as a predictor of mortality. In assessing CVD risk, the American Heart Association (AHA) and Centers for Disease Control and Prevention (CDC) defined hs-CRP levels of >3.0 mg/L as indicating high risk, 1.0–3.0 mg/L as indicating average risk, and <1.0 mg/L as indicating low risk [4, 5].

In South Korea, the percentage of individuals who consider their health status to be good (“very good” or “good”) is low, at 29.2% in 2017. From 1998 to 2017, this percentage has remained in the range 29–47%, indicating that less than half of the population consider themselves to be healthy. Moreover, this percentage is one of the lowest among countries of the Organization for Economic Co-operation and Development; even though South Korea maintains an objective health status similar to that of the US and Europe, South Koreans experience more subjective health anxiety [6]. Several prior studies have reported that a poor SRH is associated with a high hs-CRP level [7–9]. A poor SRH is related to mortality and morbidity because low-grade inflammation is associated with the development of CVD, cancer, and diabetes [10, 11]. Moreover, the level of inflammation differs according to sex and genetic variation [12, 13]. A previous study showed that poor SRH and high hs-CRP levels are significantly associated in women but not in men [9]. In addition, several studies have reported differences in the relationship between SRH and low-grade inflammation according to sex [8, 14]; however, to the best of our knowledge, there have been no large-scale studies on this relationship focusing on the Korean population. Therefore, in this study, using data from the 2015–2017 Korea National Health and Nutrition Examination Survey (KNHANES), we aimed to investigate the correlation between SRH and hs-CRP in Koreans aged ≥ 19 years and to analyze whether the correlation showed any differences between male and female subjects.

Methods

Participants

In this study, we used raw data from KNHANES VI and VII (2015-2017). KNHANES is a nationwide cross-sectional survey conducted by the CDC to ascertain the health and nutritional status of the Korean population. The survey extracts a representative sample of households and conducts a household member verification survey, health questionnaire survey, health examination, and nutritional survey for household members aged ≥ 1 year. In the present study, we aimed to investigate Korean adults aged ≥ 19 years. Of the 23,657 survey participants, we excluded individuals with missing measurements of hs-CRP (4,871 people) and SRH (57 people), those aged < 19 years (1,838 people), and those with unclear measurement values (2,655 people). We also excluded individuals with hs-CRP levels ≥ 10 mg/L (866 people) as these levels could be considered to indicate acute infection, systemic inflammation, or tissue injury [15]. A total of 14,544 subjects (6,281 men, 8,263 women) were included in our analysis (Fig. 1).

Instruments and procedure

SRH assessment

SRH was assessed using the question "In general, how would you rate your health?" There were five possible responses: "very good," "good," "fair," "poor," and "very poor."

CRP measurement

hs-CRP was measured by immunoturbidimetry using a Cobas analyzer (Roche, Germany) and a Cardiac C-Reactive Protein High Sensitivity reagent (Roche, Germany). The minimum value in the specimens was 0.1 mg/L, and the maximum value was 20 mg/L. A high hs-CRP level was defined as an hs-CRP level > 1.0 mg/L.

The AHA and CDC recommend an hs-CRP cutoff value of < 1 mg/L to indicate low risk, 1-3 mg/L to indicate average risk, and > 3 mg/L to indicate high risk. However, these criteria were recommended based on studies focused on Western populations [16, 17]. As an alternative, we used an hs-CRP measurement greater than 1.0 mg/L as the cutoff value based on the studies by Jung et al. [18, 19].

Covariates

For the participants' demographic characteristics, we included age; for socioeconomic characteristics, we included educational level, household income level, and marital status; for lifestyle factors, we included smoking status and alcohol consumption; and for disease and health-related factors, we included body mass index (BMI), chronic diseases, and menopausal status (women). Regarding chronic diseases, we included those directly or indirectly associated with hs-CRP: hypertension, diabetes mellitus, coronary artery disease (including myocardial infarction and angina [20]), dyslipidemia [21], stroke [22], and rheumatoid arthritis [23].

Statistical methods

The KNHANES applies stratified cluster sampling and weighted values to a nationally representative sample, which is based on the reciprocal of selection probabilities (primary sampling unit, household), the inverse of response rate (household, subject), and a post-stratification factor that provides age and sex-specific survey result estimates for the Korean population [24].

Therefore, we performed data analyses based on a complex sampling design with elements of stratification variables, clustering variables, and weights [25].

Data are presented as means \pm standard errors for continuous variables or as frequencies and percentages for categorical variables. The following categorical variables were compared using Pearson's chi-square test: educational level, marital status, household income level, smoking status, alcohol consumption, chronic diseases, SRH, and menopause. The following continuous variables were compared using the Mann-Whitney U test: age, weight, height, and BMI. Associations between hs-CRP levels and various sociodemographic categories and SRH were explored. In addition, multiple logistic regression analysis was used to study the relationship between hs-CRP levels and SRH according to sex. Results are presented as odds ratios (ORs) and 95% confidence intervals (CIs). Additionally, this study investigated if there was an underlying trend in the different levels of SRH in each model (Models 1–4) using Scheffe's test after assigning an ordinal score to each SRH group in the model. A P -value <0.05 was considered significant. All data analyses were performed using the statistical software package SAS version 9.4 (SAS Institute Inc., Cary, NC, USA) and IBM SPSS Statistics 25 (SPSS Inc., Chicago, IL, USA).

Results

Characteristics of participants

Table 1 shows the characteristics of the study participants (6,281 men and 8,263 women). To analyze the differences between sexes, men and women were separately classified into groups according to their hs-CRP levels: low hs-CRP (≤ 1.0 mg/L) and high hs-CRP (>1.0 mg/L). Among all participants, the percentage of those with very poor to poor SRH was higher in the high hs-CRP group (22.47%) than in the low hs-CRP group (17.66%). Similarly, for both men and women, the percentages of participants with very poor to poor SRH were higher in the high hs-CRP groups than in the low hs-CRP groups, whereas the percentages of participants with good to very good SRH were lower in the low hs-CRP groups than in the high hs-CRP groups ($P < 0.001$) (Fig. 2).

Correlation between SRH and high hs-CRP level (overall)

Table 2 shows the overall correlations between SRH and high hs-CRP levels along with their respective ORs and 95% CIs. In the crude model (Model 1), the poorer the SRH, the higher was the risk in relation to the reference group (very good SRH). After adjustment for age, educational level, marital status, and household income level (Model 2), the associations between fair to very poor SRH and high hs-CRP levels remained significant. However, after adjustment for BMI, smoking, and alcohol consumption (Model 3) and after additional adjustment for chronic diseases (Model 4), the associations only remained significant for poor and fair SRH.

Correlation between SRH and high hs-CRP level (male vs. female participants)

When we analyzed male participants, those with fair to very poor SRH had a significantly higher risk of being in the high hs-CRP group than those with very good SRH in both the crude model (Model 1) and Model 2 (adjustment for age, educational level, marital status, and household income level). After adjustment for BMI, smoking, and alcohol consumption (Model 3), the associations between fair to very poor SRH and high hs-CRP levels remained significant. However, after adjustment for chronic diseases (Model 4), the association only remained significant for very poor and fair SRH (Table 3).

When we analyzed female participants, those with fair to very poor SRH had a significantly higher risk of being in the high hs-CRP group than those with very good SRH in the crude model (Model 1). After adjustment for age, educational level, marital status, and household income level (Model 2), the association only remained significant for poor SRH. However, in contrast with that observed for male participants, no significant correlations remained after adjustment for BMI, smoking, and alcohol consumption (Model 3) and after additional adjustment for chronic diseases (Model 4) (Table 4).

Discussion

Our findings were similar to those of previous studies that reported a correlation between a poor SRH and a high hs-CRP level [7–9, 26]. In a study of 4,049 respondent older adults without significant cognitive deficit by Szybalska et al. [27], a worse SRH was associated with increased interleukin-6 (IL-6) and CRP levels. Leshem-Rubinow et al. [7] analyzed the correlations between SRH and the inflammation-sensitive biomarkers hs-CRP and fibrinogen in 13,773 healthy individuals and observed higher biomarker levels in the group with the lowest SRH level; hs-CRP showed a correlation in both men and women, but fibrinogen only showed a correlation in men. Shanahan et al. [8] studied 13,236 young adults and reported that on adjusting for acute/chronic diseases, medication history, and health behaviors, a lower SRH level was associated with a higher hs-CRP level; however, on adjusting for BMI, the correlation in female participants was weakened, whereas the correlation in male participants remained significant. However, a study of 16,256 Japanese individuals reported a significant correlation between a poor SRH and a high hs-CRP level only in female participants [9]. Thus, while correlations between a poor SRH and a high hs-CRP level have been reported, the above studies show limitations such as restricted age of participants, including only older adults [26] or only young adults [8], and lack of consideration for diseases that could affect the relationship between SRH and CRP [9]. Moreover, there have been few studies on the relationship between SRH and CRP [7, 28] so far; there have been no such studies in Koreans.

Several studies have reported correlations between a poor SRH and pro-inflammatory cytokines, including IL-6 [17, 29]. CRP is produced by hepatocytes under the regulatory control of IL-6 and other inflammatory cytokines [30], and these pro-inflammatory cytokines cause sickness behaviors, such as weakness, depression, exaggerated pain (hyperalgesia or allodynia), and lack of appetite [31]. In other words, the relationship between a poor SRH and a high hs-CRP level can be explained by differences caused by pro-inflammatory cytokines, and this, in turn, can explain our results.

Inflammatory indices in women are known to be altered by the menstrual cycle, menopause, and hormone therapy [32], and menopause and estrogen replacement therapy have been reported to affect obesity and inflammation in women [33]. CRP levels can be presumed to change depending on the hormonal environment, and this could act as a confounding factor in the relationship between SRH and hs-CRP [34]. In our study, we were unable to investigate whether participants were taking female hormones or their stage in the menstrual cycle. As we only accounted for female menopause, we may not have observed significant results for the association between SRH and hs-CRP levels in women. Moreover, biological sex is known to affect CRP-related genetic variation [35], and according to a study by Kettunen et al. [36], allelic variants in the CRP gene are associated with CRP levels, and men and women show differences depending on the CRP genotype. Hence, the differences in genetic variation between men and women could have affected our results.

Sex differences have been reported in the relationship between CRP levels and mortality [37–40], but it is unclear why a high CRP level is only associated with an increased mortality risk in men. Dong et al analyzed the middle-aged Chinese population; hs-CRP was associated with increased risk of developing CVD [41]. In addition, Lee JH et al. [42] studied 23,233 rural Koreans and reported that a high CRP level was more strongly associated with higher mortality in men than in women. There have been several studies reporting a stronger correlation between SRH and mortality in men than in women [40, 43]. Specifically, men with a poor SRH have been reported to show a higher risk for conditions related to mortality, such as CVD and cancer. When assessing SRH, the subject rates their current overall health; it has been reported that men rate their own health in comparison to that of other men, and male SRH tends to mostly reflect serious and life-threatening disease, whereas female SRH tends to reflect other factors unrelated to mortality and chronic, non-life-threatening disease, resulting in a weaker correlation between SRH and mortality for women [43–45]. Moreover, in a study of Korean adults by Shin et al. [46], women tended to rate their own health more poorly than men, and in a study by Lee SY et al. [47], traditional Korean gender roles had a negative effect on women, and the risk of a poor SRH was higher among Korean women than among women from the US. Similarly, in our study, we only observed a correlation between SRH and hs-CRP among male participants. The discrepancy between men and women could be related to the fact that CRP is more likely to reflect CVD and mortality in men than in women and that SRH is more likely to directly reflect health and mortality in men than in women.

In our study of Korean adults aged ≥ 19 years, when we analyzed all participants, the poor SRH group was more likely to have high hs-CRP levels (>1.0 mg/L) than the very good SRH group. Especially in male participants, as SRH went from very good to very poor, there was a corresponding increase in the risk of high hs-CRP level (>1.0 mg/L). These results can be explained by the fact that immune-related activity is associated with vague symptoms of malaise and interoceptive perception [29]. Such findings are consistent with those of a previous study of healthy adults, in which a poorer SRH was associated with increased serum inflammatory marker levels (IL-6 and CRP) [26].

Even after correcting for all sociodemographic characteristics, health-related factors, and chronic diseases known to be associated with low-level inflammation, among male participants, the very poor SRH group showed 1.74 times higher risk of high hs-CRP level than the very good SRH group, but there was no significant relationship among female participants. This finding could be because SRH is a dynamic evaluation for judging the trajectory of health, which reflects both clinical stage and preclinical stage disease [48]. Therefore, even after correcting for chronic diseases associated with hs-CRP, we still observed a correlation in male participants.

Our study has several limitations. First, because this was a cross-sectional study, it was not possible to infer causal relationships, and we could only investigate the correlation between SRH and hs-CRP. Nevertheless, the value of this study is that we used data from the KNHANES, which is representative of the Korean population, and that it was a large-scale study of Korean adults. Second, SRH assessment was performed at specific times. Future monitoring studies are necessary to ascertain the long-term relationships between SRH and hs-CRP. Third, we only used hs-CRP as an inflammatory marker; further studies will need to investigate the correlations of SRH with other indicators (e.g., IL-6, tumor necrosis factor-alpha). Finally, because the study was based on data from a survey of South Koreans, the results could have been affected by the racial characteristics of Koreans and may thus be difficult to apply to people of other races. Despite these limitations, our study showed a strong correlation between a poor SRH and a high hs-CRP level in male Korean adults and is valuable as the first study to examine the relationship between SRH and hs-CRP in Korean adults.

Conclusions

We investigated the relationship between SRH and hs-CRP through a survey of Korean adults. Even after correcting for factors that could affect low-grade inflammation, such as age, socioeconomic status, BMI, health-related behaviors, and chronic diseases, male adults with a poor SRH were at greater risk of having high hs-CRP. This study showed a close relationship between low-grade inflammation (high hs-CRP level) and SRH, an instrument reflecting one's own assessment of their health. Since we only observed a correlation in male participants, it suggests that there could be factors affecting the relationship between SRH and hs-CRP differently for men and women. Our findings could provide a basis for developing health improvement programs. Future studies will need to be conducted to examine sex differences in the relationship of SRH with other inflammatory markers.

List Of Abbreviations

AHA, American Heart Association; BMI, body mass index; CDC, Centers for Disease Control; CI, confidence interval; CVD, cardiovascular disease; hs-CRP, high-sensitivity C-reactive protein; KNHANES, Korea National Health and Nutrition Examination Survey; OR, odds ratio; SRH, self-rated health.

Declarations

Ethics approval and consent to participate

The study was approved by the Institutional Review Board of Jaseng Hospital of Korean Medicine in Seoul, Korea (JASENG 2019-08-001).

Consent for publication

Not applicable.

Availability of data and materials

All original data are publicly available free of charge from the KNHANES website (<http://knhanes.cdc.go.kr>) for the purposes of academic research.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no external funding.

Authors' contributions

Conceptualization, B.J. and S.W.P.; methodology, B.J.; software, B.J.; validation, I.H.H.; formal analysis, B.J.; investigation, E.J.K.; resources, I.H.H.; data curation, B.J.; writing—original draft preparation, B.J. and S.W.P.; writing—review and editing, S.S.P., E.J.K., W.S.S. I.H.H., and B.J.; supervision, B.J.

Acknowledgements Not applicable

References

1. Joffer J, Jerdén L, Öhman A, Flacking R. Exploring self-rated health among adolescents: a think-aloud study. *BMC Public Health* 2016;16(1):156; doi: 10.1186/s12889-016-2837-z.
2. Vie TL, Hufthammer KO, Meland E, Bredablik HJ. Self-rated health (SRH) in young people and causes of death and mortality in young adulthood. a prospective registry-based Norwegian HUNT-study. *SSM Popul Health* 2019;7: 100364; doi: 10.1016/j.ssmph.2019.100364.
3. Ploubidis GB, Pongiglione B. Self-rated health over the life course: evidence from the 1958 and 1970 British birth cohorts. *Pathways to Health* 2019;79-97; doi: [1007/978-94-024-1707-4_4](https://doi.org/10.1007/978-94-024-1707-4_4).
4. Brás J, Pinto S, Almeida MI, Prata J, von Doellinger O, Coelho R, et al. Peripheral biomarkers of inflammation in depression: evidence from animal models and clinical studies. *Methods Mol Biol* 2019;2011: 467–92); doi: 10.1007/978-1-4939-9554-7_28.
5. Ridker PM. A test in context: high-sensitivity C-reactive protein. *J Am Coll Cardiol* 2016;67:712–23; doi: 10.1016/j.jacc.2015.11.037.
6. Landi S, Ivaldi E, Testi A. Socioeconomic status and waiting times for health services: an international literature review and evidence from the Italian National Health System. *Health Policy* 2018;122(4):334–51; doi: 10.1016/j.healthpol.2018.01.003.
7. Leshem-Rubinow E, Shenhar-Tsarfaty S, Milwidsky A, Toker S, Shapira I, Berliner S, et al. Self-rated health is associated with elevated C-reactive protein even among apparently healthy individuals. *Isr Med Assoc J* 2015;17:213–8.
8. Shanahan L, Bauldry S, Freeman J, Bondy CL. Self-rated health and C-reactive protein in young adults. *Brain Behav Immun* 2014;36:139–46; doi: 10.1016/j.bbi.2013.10.020.
9. Tanno K, Ohsawa M, Onoda T, Itai K, Sakata K, Tanaka F, et al. Poor self-rated health is significantly associated with elevated C-reactive protein levels in women, but not in men, in the Japanese general population. *J Psychosom Res* 2012;73:225–31; doi: 10.1016/j.jpsychores.2012.05.013.
10. Emerging Risk Factors Collaboration; Kaptoge S, Di Angelantonio E, Lowe G, Pepys MB, Thompson SG, et al. C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet* 2010;375(9709):132–40; doi: 10.1016/S0140-6736(09)61717-7.

11. Dehghan A, Kardys I, de Maat MP, Uitterlinden AG, Sijbrands EJ, Bootsma AH, et al. Genetic variation, C-reactive protein levels, and incidence of diabetes. *Diabetes* 2007;56(3):872–8; doi: 10.2337/db06-0922.
12. Jilma B, Dirnberger E, Loscher I, et al. Menstrual cycle-associated changes in blood levels of interleukin-6, alpha1 acid glycoprotein, and C-reactive protein. *J Lab Clin Med*. 1997;130(1):69-75.
13. Ancelin ML, Farré A, Carrière I, et al. C-reactive protein gene variants: independent association with late-life depression and circulating protein levels. *Transl Psychiatry*. 2015;5:e499.
14. Alho M. The association of systemic low-grade inflammation with health-related quality of life in Finnish young men. *Exercise Physiology* Master's thesis 2019:1–188.
15. Al Aseri ZA, Habib SS, Marzouk A. Predictive value of high sensitivity C-reactive protein on progression to heart failure occurring after the first myocardial infarction. *Vasc Health Risk Manag* 2019;15:221–7; doi: 10.2147/VHRM.S198452.
16. Saito I, Sato S, Nakamura M, Kokubo Y, Mannami T, Adachi H, et al. A low level of C-reactive protein in Japanese adults and its association with cardiovascular risk factors: the Japan NCVC-Collaborative Inflammation Cohort (JNIC) study. *Atherosclerosis* 2007;194:238–44; doi: 10.1016/j.atherosclerosis.2006.07.032.
17. Sung KC, Ryu S, Chang Y, Byrne CD, Kim SH. C-reactive protein and risk of cardiovascular and all-cause mortality in 268 803 East Asians. *Eur Heart J* 2014;35:1809–16. doi: 10.1093/eurheartj/ehu059.
18. Jung SW, Lee KJ, Lee JH. Does weekend catch-up sleep affect high-sensitivity C-reactive protein levels among Korean workers?: A cross-sectional study using KNHANES. *J Occup Environ Med* 2019;61(9):e367–73; doi: 10.1097/JOM.0000000000001657.
19. Momiyama Y, Kawaguchi A, Kajiwaru I, Ohmori R, Okada K, Saito I, et al. Prognostic value of plasma high-sensitivity C-reactive protein levels in Japanese patients with stable coronary artery disease: the Japan NCVC-Collaborative Inflammation Cohort (JNIC) Study. *Atherosclerosis* 2009;207: 272–6. doi: 10.1016/j.atherosclerosis.2009.04.015.
20. Mazidi M, Toth PP, Banach M. C-reactive protein is associated with prevalence of the metabolic syndrome, hypertension, and diabetes mellitus in US adults. *Angiology* 2018;69:438–42; doi: 10.1177/0003319717729288.
21. Lin GM, Liu K, Colangelo LA, Lakoski SG, Tracy RP, Greenland P. Low-density lipoprotein cholesterol concentrations and association of high-sensitivity C-reactive protein concentrations with incident coronary heart disease in the multi-ethnic study of atherosclerosis. *Am J Epidemiol* 2015;183:46–52; doi: 10.1093/aje/kwv144.
22. Yu H, Huang Y, Chen X, Nie W, Wang Y, Jiao Y, et al. High-sensitivity C-reactive protein in stroke patients—The importance in consideration of influence of multiple factors in the predictability for disease severity and death. *J Clin Neurosci* 2017;36:12–9; doi: 10.1016/j.jocn.2016.10.020.
23. Wang Y, Liang Y. Clinical significance of serum calprotectin level for the disease activity in active rheumatoid arthritis with normal C-reactive protein. *Int J Clin Exp Pathol* 2019;12:1009–14.
24. Kweon S, Kim Y, Jang M-j, Kim Y, Kim K, Choi S, et al. Data resource profile: the Korea 390 national health and nutrition examination survey (KNHANES). *International journal of 391 epidemiology*. 2014;43(1):69-77.
25. Kim Y. The Korea National Health and nutrition examination survey (KNHANES): current 421 status and challenges. *Epidemiology and health*. 2014;36.
26. Christian LM, Glaser R, Porter K, Malarkey WB, Beversdorf D, Kiecolt-Glaser JK. Poorer self-rated health is associated with elevated inflammatory markers among older adults. *Psychoneuroendocrinology* 2011;36(10):1495–504; doi: 10.1016/j.psyneuen.2011.04.003.
27. Szybalska A, Broczek K, Puzianowska-Kuznicka M, Slusarczyk P, Chudek J, Skalska A, et al. Self-rated health and its association with all-cause mortality of older adults in Poland: The PolSenior project. *Arch Gerontol Geriatr* 2018;79:13–20; doi: 10.1016/j.archger.2018.07.016.
28. Pinillos-Franco S, Kawachi I. The relationship between social capital and self-rated health: A gendered analysis of 17 European countries. *Soc Sci Med* 2018;219:30–5; doi: 10.1016/j.socscimed.2018.10.010.
29. Lekander M, Elofsson S, Neve M, Hansson LO, Undén AL. Self-rated health is related to levels of circulating cytokines. *Psychosom Med* 2004;66:559–63; doi: 10.1097/01.psy.0000130491.95823.94.

30. Ridker PM. C-reactive protein: eighty years from discovery to emergence as a major risk marker for cardiovascular disease. *Clin Chem* 2009;55:209–15; doi: 10.1373/clinchem.2008.119214.
31. Munshi S, Parrilli V, Rosenkranz JA. Peripheral anti-inflammatory cytokine interleukin-10 treatment mitigates interleukin-1 β -induced anxiety and sickness behaviors in adult male rats. *Behav Brain Res* 2019;372:112024; doi: 10.1016/j.bbr.2019.112024.
32. Govorov I, Bremme K, Larsson A, Holmström M, Komlichenko E, Chaireti R, et al. Blood inflammatory and endothelial markers in women with von Willebrand disease. *PloS One* 2019;14(1):e0210544; doi: 10.1371/journal.pone.0210544.
33. Sung ES, Jeong JA, Choi CK, Shin MH. The relationship between body mass index and poor self-rated health in the Korean population. *BioRxiv* 2019:688994; doi: <https://doi.org/10.1101/688994>.
34. Uchino BN, Landvatter J, Cronan S, Scott E, Papadakis M, Smith TW, et al. Self-rated health and inflammation: a test of depression and sleep quality as mediators. *Psychosom Med* 2019;81:328–32; doi: 10.1097/PSY.0000000000000683.
35. Lee S, Oh SS, Jang SI, Park EC. Sex difference in the association between high-sensitivity C-reactive protein and depression: the 2016 Korea National Health and Nutrition Examination Survey. *Sci Rep* 2019;9:19–8. doi: 10.1038/s41598-018-36402-3.
36. Kettunen T, Eklund C, Kähönen M, Jula A, Päivä H, Lyytikäinen LP, et al. Polymorphism in the C-reactive protein (CRP) gene affects CRP levels in plasma and one early marker of atherosclerosis in men: The Health 2000 Survey. *Scand J Clin Lab Invest* 2011;71(5):353–61. doi: 10.3109/00365513.2011.568123.
37. Evans CR, Long DL, Howard G, McClure LA, Zakai NA, Jenny NS, et al. C-reactive protein and stroke risk in blacks and whites: The REasons for Geographic And Racial Differences in Stroke cohort. *Am Heart J* 2019;217:94–100; doi: 10.1016/j.ahj.2019.08.003.
38. Milano SS, Moura Júnior OVD, Bordin AAS, Marques GL. C-reactive protein is a predictor of mortality in ST-segment elevation acute myocardial infarction. *J International Journal of Cardiovascular Sciences* 2019;32:118–24. doi: <http://dx.doi.org/10.5935/2359-4802.20180086>.
39. Norring-Agerskov D, Bathum L, Pedersen OB, Abrahamsen B, Lauritzen JB, Jørgensen NR, et al. Biochemical markers of inflammation are associated with increased mortality in hip fracture patients: the Bispebjerg Hip Fracture Biobank. *Aging Clin Exp Res* 2019;31(12):1727–34; doi: 10.1007/s40520-019-01140-7.
40. Reile R, Stickley A, Leinsalu M. Large variation in predictors of mortality by levels of self-rated health: results from an 18-year follow-up study. *Public Health* 2017;145:59-66; doi: 10.1016/j.puhe.2016.12.034.
41. Dong Y, Wang X, Zhang L, Chen Z, Zheng C, Wang J, et al. High-sensitivity C reactive protein and risk of cardiovascular disease in China-CVD study. *J Epidemiol Community Health* 2019;73(2):188–92; doi: 10.1136/jech-2018-211433.
42. Lee JH, Yeom H, Kim HC, Suh I, Kim MK, Shin MH, et al. C-reactive protein concentration is associated with a higher risk of mortality in a rural Korean Population. *J Prev Med Public Health* 2016;49(5):275–87; doi: 10.3961/jpmph.16.025.
43. Benjamins MR, Hummer RA, Eberstein IW, Nam CB. Self-reported health and adult mortality risk: an analysis of cause-specific mortality. *Soc Sci Med* 2004;59:1297–306; doi: 10.1016/j.socscimed.2003.01.001.
44. Benyamini Y, Leventhal EA, Leventhal H. Gender differences in processing information for making self-assessments of health. *Psychosom Med* 2000;62(3):354–64; doi: 10.1097/00006842-200005000-00009.
45. Jylhä M, Guralnik JM, Ferrucci L, Jokela J, Heikkinen E. Is self-rated health comparable across cultures and genders? *J Gerontol B Psychol Sci Soc Sci* 1998;53(3): S144–52; doi: 10.1093/geronb/53b.3.s144.
46. Shin HY, Shin MH, Rhee JA. Gender differences in the association between self-rated health and hypertension in a Korean adult population. *BMC Public Health* 2012;12:135; doi: 10.1186/1471-2458-12-135.
47. Lee SY, Kim SJ, Yoo KB, Lee SG, Park EC. Gender gap in self-rated health in South Korea compared with the United States. *Int J Clin Health Psychol* 2016;16:11-20; doi: 10.1016/j.ijchp.2015.08.004.
48. Benyamini Y. Why does self-rated health predict mortality? An update on current knowledge and a research agenda for psychologists. *Psychol Health* 2011;26(11):1407–13; doi: 10.1080/08870446.2011.621703.

Tables

Table 1 Baseline characteristics of the study population in the 2015-2017 KNHANES

Characteristic	Men			Women		
	Low hs-CRP (≤ 1.0 mg/L) (n=4,495)	High hs-CRP (> 1.0 mg/L) (n=1,786)	P- value†	Low hs-CRP (≤ 1.0 mg/L) (n=6,306)	High hs-CRP (> 1.0 mg/L) (n=1,957)	P- value†
Age (years), mean (SD)	49.8 (0.249)	53.3(0.391)	.000*	49.9 (0.203)	54.2 (0.373)	.000*
Educational level, n (%)			.000*			.000*
Elementary school or lower	580 (12.9)	333 (18.6)		1,424 (22.6)	653 (33.4)	
Middle school	446 (9.9)	205 (11.5)		651 (10.3)	203 (10.4)	
High school	1,575 (35.0)	568 (31.8)		2,001 (31.7)	547 (28.0)	
College or higher	1,894 (42.1)	680 (38.1)		2,230 (35.4)	554 (28.3)	
Marital status, n (%)			.002*			.000*
Married	3,536 (78.7)	1,487 (83.3)		5,396 (85.6)	1,772 (90.5)	
Unmarried	959 (21.3)	299 (16.7)		910 (14.4)	185 (9.5)	
Household income level, n (%)			.101			.009*
Low	1,034 (23.0)	457 (25.6)		1,444 (22.9)	522 (26.7)	
Lower middle	1,077 (24.0)	464 (26.0)		1,591 (25.2)	519 (26.5)	
Upper middle	1,156 (25.7)	435 (24.4)		1,609 (25.5)	470 (24.0)	
High	1,228 (27.3)	430 (24.1)		1,662 (26.4)	446 (22.8)	
Weight (kg), mean (SD)	70.0 (0.158)	73.4 (0.351)	.000*	56.8 (0.103)	62.1 (0.253)	.000*
Height (cm), mean (SD)	170.5 (0.101)	169.6 (0.161)	.000*	157.5 (0.081)	156.2 (0.149)	.000*
BMI (kg/m ²), mean (SD)	24.0 (0.044)	25.4 (0.087)	.000*	22.9 (0.040)	25.4 (0.091)	.000*
Smoking, n (%)			.123			.483
Nonsmoker	1,130 (25.1)	389 (21.8)		5,710 (90.5)	1,742 (89.0)	
Past smoker	1,926 (42.8)	777 (43.5)		341 (5.4)	118 (6.0)	
Current	1,439 (32.0)	620 (34.7)		255 (4.0)	97 (5.0)	

smoker					
Alcohol consumption, n (%)			.001*		.000*
Never drink	196 (4.4)	85 (4.8)	951 (15.1)	392 (20.0)	
<Once/month	1,004 (22.3)	457 (25.6)	2,634 (41.8)	864 (44.1)	
<Five times/month	1,691 (37.6)	577 (32.3)	2,020 (32.0)	496 (25.3)	
≥Five times/month	1,604 (35.7)	667 (37.3)	701 (11.1)	205 (10.5)	
Chronic diseases**, n (%)			.000*		.000*
None	2,579 (57.4)	789 (44.2)	4,000 (63.4)	966 (49.4)	
One	1,186 (26.4)	590 (33.0)	1,274 (20.2)	536 (27.4)	
Two or more	730 (16.2)	407 (22.8)	1,032 (16.4)	455 (23.2)	
Self-rated health, n (%)			.000*		.000*
Very poor	80 (1.8)	59 (3.3)	270 (4.3)	108 (5.5)	
Poor	532 (11.8)	264 (14.8)	1,026 (16.3)	410 (21.0)	
Fair	2,271 (50.5)	943 (52.8)	3,328 (52.8)	1,009 (51.6)	
Good	1,317 (29.3)	431 (24.1)	1,431 (22.7)	370 (18.9)	
Very good	295 (6.6)	89 (5.0)	251 (4.0)	60 (3.1)	
Menopause, n (%)					.000*
No			3,211 (50.9)	752 (38.4)	
Yes			3,072 (48.7)	1,189 (60.8)	

* $p \leq .05$

** Chronic diseases include hypertension, diabetes, dyslipidemia, coronary heart disease (myocardial infarction or angina pectoris), stroke, and rheumatoid arthritis.

† P-values of t-test or the Mann-Whitney U test for continuous variables and Pearson's chi-squared test for categorical variables to determine differences between groups according to high-sensitivity C-reactive protein (hs-CRP) levels.

KNHANES, Korea National Health and Nutrition Examination Survey; SE, standard error; SD, standard deviation; BMI, body mass index.

Table 2 Association between high hs-CRP level and SRH in all subjects

SRH	Model 1			Model 2			Model 3			Model 4		
	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value
Very poor	2.02	1.47 2.78	.000*	1.54	1.11 2.12	.010*	1.22	0.87 1.72	.245	1.22	0.87 1.71	.259
Poor	1.82	1.44 2.29	.000*	1.68	1.32 2.13	.000*	1.40	1.09 1.79	.008*	1.38	1.08 1.77	.011*
Fair	1.51	1.22 1.87	.000*	1.46	1.18 1.82	.001*	1.34	1.06 1.68	.013	1.32	1.05 1.66	.016*
Good	1.16	0.92 1.46	.211	1.17	0.93 1.47	.183	1.12	0.88 1.42	.375	1.12	0.87 1.42	.377
Very good (ref.)	1.00			1.00			1.00			1.00		
P for trend			<0.001*			<0.001*			<0.001*			<0.001*

Odds ratios with adjustments using logistic regression models.

* $p \leq .05$

Model 1: unadjusted.

Model 2: adjusted for age, educational level, marital status, and household income level.

Model 3: adjusted for Model 2 confounders + body mass index, smoking, and alcohol consumption.

Model 4: adjusted for Model 3 confounders + chronic diseases.

hs-CRP, high-sensitivity C-reactive protein; SRH, self-rated health; OR, odds ratio; CI, confidence interval.

Table 3 Association between high hs-CRP level and SRH in men

SRH	Model 1			Model 2			Model 3			Model 4		
	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value
Very poor	2.69	1.68 4.32	.000*	2.19	1.37 3.51	.001*	1.78	1.07 2.96	.026*	1.74	1.04 2.90	.034*
Poor	1.79	1.30 2.44	.000*	1.70	1.23 2.35	.001*	1.43	1.02 2.00	.041*	1.40	0.99 1.97	.057*
Fair	1.63	1.22 2.16	.001*	1.61	1.20 2.14	.001*	1.44	1.06 1.96	.019*	1.42	1.04 1.93	.025*
Good	1.12	0.83 1.51	.462	1.14	0.84 1.54	.396	1.08	0.78 1.49	.637	1.08	0.78 1.49	.637
Very good (ref.)	1.00			1.00			1.00			1.00		
P for trend			<0.001*			<0.001*			<0.001*			<0.001*

Odds ratios with adjustments using logistic regression models.

* $p \leq .05$

Model 1: unadjusted.

Model 2: adjusted for age, educational level, marital status, and household income level.

Model 3: adjusted for Model 2 confounders + body mass index, smoking, and alcohol consumption.

Model 4: adjusted for Model 3 confounders + chronic diseases.

hs-CRP, high-sensitivity C-reactive protein; SRH, self-rated health; OR, odds ratio; CI, confidence interval.

Table 4 Association between high hs-CRP level and SRH in women

SRH	Model 1			Model 2			Model 3			Model 4		
	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value	OR	95% CI	P-value
Very poor	1.97	1.28 3.01	.002*	1.35	0.87 2.10	.182	0.99	0.62 1.59	.980	1.01	0.63 1.62	.968
Poor	2.03	1.43 2.90	.000*	1.77	1.24 2.53	.002*	1.39	0.95 2.03	.086	1.40	0.96 2.03	.083
Fair	1.50	1.06 2.11	.021*	1.38	0.98 1.95	.063	1.27	0.89 1.80	.186	1.27	0.89 1.80	.187
Good	1.25	0.88 1.78	.213	1.24	0.87 1.76	.233	1.21	0.84 1.76	.301	1.22	0.84 1.76	.297
Very good (ref.)	1.00						1.00			1.00		
P for trend			<0.001*			<0.001*			<0.001*			<0.001*

Odds ratios with adjustments using logistic regression models.

* $p \leq .05$

Model 1: unadjusted.

Model 2: adjusted for age, educational level, marital status, and household income level.

Model 3: adjusted for Model 2 confounders + body mass index, smoking, alcohol consumption, and menopause.

Model 4: adjusted for Model 3 confounders + chronic diseases.

hs-CRP, high-sensitivity C-reactive protein; SRH, self-rated health; OR, odds ratio; CI, confidence interval.

Figures




Figure 1

Flow diagram showing the number of included and excluded participants and the data for analysis KNHANES, Korea National Health and Nutrition Examination Survey; hs-CRP, high-sensitivity C-reactive protein; SRH, self-rated health; BMI, body mass index




Figure 2

Prevalence of self-rated health (SRH) according to sex The percentage of participants with very poor to poor SRH was higher in the group with high hs-CRP level (22.47%) than that in the group with low hs-CRP level (17.66%) (P<0.001)