

Statin prescription among patients with type 2 diabetes in Botswana: findings and implications

Julius Chacha Mwita (✉ mwitajc@ub.ac.bw)

University of Botswana <https://orcid.org/0000-0002-5947-3684>

Brian Godman

Strathclyde Institute of Pharmacy and Biomedical Sciences

Tonya M Esterhuizen

Stellenbosch University Faculty of Medicine and Health Sciences

Research article

Keywords: statin, type 2 diabetes mellitus, prescription and Botswana

Posted Date: December 30th, 2019

DOI: <https://doi.org/10.21203/rs.2.10881/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 10th, 2020. See the published version at <https://doi.org/10.1186/s12902-020-0516-7>.

Abstract

Background There is evidence of statin benefit among patients with diabetes regardless of their cholesterol levels or prior cardiovascular disease history. Despite the evidence, there is under-prescription of statins in clinical practice. This study aimed to assess statin prescriptions and associated factors among patients with type 2 diabetes in Botswana. **Methods** The study was a secondary data analysis of 500 randomly selected type 2 diabetes patients at a specialised diabetes clinic at Gaborone, Botswana. We assessed the proportion of statin-eligible patients who are prescribed statins and evaluated the adjusted associations between various factors and statin prescription. **Results** Overall, 477(95.4%) participants were eligible for statin prescription. Clinicians prescribed statins in 217 (45.5%) of eligible participants, and only one(4.4%) ineligible participant. The probability of statin prescription was high in participants with high baseline low-density lipoprotein cholesterol (risk ratio [RR]: 1.49; 95%CI: 1.17-1.89), increasing duration of diabetes(RR: 1.01; 95%CI 1.00-1.03) and the presence of chronic kidney disease(RR: 1.35; 95%CI: 1.06-1.74). **Conclusion** Most patients with type 2 diabetes in Gaborone are not receiving statins. Clinicians did not consider most guideline-recommended indications for statin prescription. The findings call for improvement in diabetes quality of care by implementing evidence-based guideline recommendations.

Background

Cardiovascular disease(CVD), which includes coronary artery disease(CAD), cerebrovascular accident(CVA), and peripheral arterial disease (PAD), is common and contributes to over two-thirds of mortality among patients with type 2 diabetes mellitus[1-3]. Although the presence of type 2 diabetes alone confers the highest risk for CVD of any single risk factor, the coexistence of other cardiovascular risk factors is a common phenomenon[3, 4]. Consequently, guidelines advise screening and optimal treatment of CVD risk factors in people with diabetes[5, 6]. Besides, prescribing of 3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors(statins) among patients with type 2 diabetes reduces the risk of major CVD events by 23%-33%[7-9]. There is evidence of statin benefit among patients with diabetes regardless of their low-density lipoprotein cholesterol (LDL-C) values or prior CVD history[7, 9-14]. For each mmol/l reduction in LDL-C, there is evidence of a 9 % relative reduction in all-cause mortality in patients with diabetes[15]. Irrespective of LDL-C, guidelines recommend statins for patients with diabetes aged ≥ 40 years without atherosclerotic cardiovascular disease(ASCVD), or those who are younger than 40 years but with existing ASCVD or additional risk factors[5, 6]. While some studies in developed countries have reported high use of statins among patients with type 2 diabetes, there has generally been under-prescription of statins across many countries[16-18]. Statin prescription among patients with type 2 diabetes is as high as 100% in developed countries, and 3-13% in Africa[17-23]. The suboptimal utilisation of statin therapy in Africa is due to many factors, but mainly limited access to standard diabetes care because of the high cost of tests and medications[22]. Affordability is a critical issue in several African countries where there is no universal healthcare, with the cost of medicines accounting for up to 70% of total healthcare expenditure, much of which is out-of-pocket[24, 25]. This is a

concern given the high growth rates of cardiovascular diseases in sub-Saharan African countries and current poor control of cardiovascular diseases[26-30]. The underuse of statins significantly increases the incidence of cardiovascular events and associated mortality[31]. Although healthcare is free in Botswana, factors not related to cost may still affect the uptake of statins in patients with diabetes. This is an issue given current prevalence rates of diabetes in Botswana and the resultant impact on morbidity and mortality[32-34]. Unfortunately, published data on the impact of diabetes mellitus on CVD in Botswana are scant. About a quarter of patients admitted with heart failure had type 2 diabetes in a recent study in Gaborone[33]. Although data on atherosclerotic diseases in Botswana are missing, we can expect that diabetes-related coronary atherosclerosis underlin some cases of heart failure above[33]. Also, the rising burden and suboptimal control of CVD risk factors in patients with type 2 diabetes have become an increasingly severe concern in Botswana[35]. One of the critical steps that will prevent or slow ASCVD in our suboptimally-treated patients is the use of statins[5]. Currently, there is no study assessing statin prescriptions among patients with type 2 diabetes in Botswana. We aimed to address this by evaluating the extent of statin prescriptions among patients with diabetes in Botswana. Our secondary aim was to determine factors associated with statin prescriptions among Type 2 diabetes.

Methods

Study design

We conducted a secondary analysis of data from a previous study among type 2 diabetics at a specialised diabetes clinic in Gaborone, Botswana. Any concerns with the management of diabetic patients in this dedicated leading clinic are likely to be exacerbated in non-specialist centres such as primary healthcare centres.

Participant recruitment and data collection

The original study took place between August 2017 and February 2018[35]. The primary objective of the original study was to assess glycaemic, low-density lipoprotein, and hypertension control in patients with type 2 diabetes. The study included 500 randomly selected patients with type 2 diabetes aged ≥ 18 years attending the clinic for at least three months before data collection. Using systematic random sampling, we e enrolled every eight patient from the list of patients who attended the clinic every study day. The enrolment occurred in the mornings and afternoons until either the daily target of ten patients was achieved or the clinic came to an end. Demographic data (age, sex, occupation, educational attainments, and marital status), duration of diabetes, and the type of diabetes medications were collected. Other information was the history of hypertension, lipid disorders, ischemic heart diseases, stroke or peripheral vascular disease. We also recorded data on the use of medications for hypertension and lipid disorders (including statins), and anthropometric measurements (weight, height, hip and waist circumferences). With participants' in light clothing and without footwear, we measured participants' height and weight to the nearest 0.1 centimetres(cm) and 0.1 kilograms(kg) respectively[36]. We also measured the hip and waist circumferences (to the nearest 0.1cm using a non-stretchable tape measure at a hip and level

midway between the lowest rib and iliac crest[37]. After 10 minutes of rest, three seated blood pressure (BP) measurements were taken from the right arm using a digital automatic BP monitor (Omron)[36, 38]. We documented the average of the three readings and that of each patient's previous visit. Also, we used the electronic medical records to obtain patients' serum creatinine, low-density lipoprotein-cholesterol(LDL-C), and urine dipstick for proteinuria results within six months of enrolment[35]. The diabetes clinic has a satellite laboratory that performs tests per protocol as defined by the ISO-9000 certified Princess Marina Referral Hospital clinical laboratory. Total cholesterol(TC), triglycerides(TG) and High-density lipoprotein cholesterol (HDL-C) are measured enzymatically in plasma using commercial kits made by Roche Diagnostics, Switzerland. LDL-cholesterol is calculated using the Friedwald formula as $TC-HDL-C-TG/2.2$ in mmol/dL[39].

For the present study, we evaluated the extent of statin prescriptions among the participants in the dataset. The primary outcome measure was receiving a statin prescription among statin-eligible participants. We assessed statin eligibility based on the Society for Endocrinology, Metabolism, and Diabetes of South Africa(SEMDSA) guidelines[6]. According to SEMDSA, the eligibility for statin prescribing included any CVD or chronic kidney disease(CKD), participant's age above 40 years, and diabetes duration longer than ten years. Also, the presence of one or more additional cardiovascular risk factors, i.e. hypertension, cigarette smoker, HDL-C level, family history of early CAD, and any albuminuria were the other eligibility criteria[6]. Thus, we assessed the association of the above eligibility criteria for statin prescribing. Other independent variables included baseline serum LDL-C, body mass index(BMI), waist-hip ratio(WHR) and education attainment.

Definition of terms

The diagnosis of hypertension based on the self-reported history of hypertension, the use of hypertension-lowering medications or sustained blood pressure $\geq 140/90$ mmHg in more than one visit[41]. We defined CVD as the history of CAD, CVA (ischemic stroke, transient ischemic attacks), or PAD [6]. CAD was any documented definite or probable myocardial infarction, CAD-related revascularisation (surgery, angioplasty, stenting, or any combination of these), or stable angina in patients' medical records[42]. Data on CVA and PDA were extracted from patients' medical records as defined by the treating physician. Smoking status was a documented self-report of current smoking habits. We estimated glomerular filtration rate (eGFR) using the Modification of Diet in Renal Disease(MDRD), and classified patients with $eGFR < 60 \text{ ml/minute/1.73m}^2$ as having CKD [43]. BMI was categorized into underweight for BMI $< 18.5 \text{ kg/m}^2$; normal for BMI of $25.0-29.9 \text{ kg/m}^2$; or obese for BMI $\geq 30 \text{ kg/m}^2$ [44]. We also calculated the WHR by dividing waist circumference (cm) by hip circumference (cm) defined WHR ≥ 0.85 for women and ≥ 0.90 for men as high[37]. Dipstick proteinuria appeared as negative(-), trace, (+), (++) , or (+++) in the dataset. We classified proteinuria in individuals with $\geq (+)$ dipstick proteinuria results. For patients already on lipid-lowering medications and whose baseline 'untreated' levels of lipid profile were not available, we estimated the LDL-C levels before the initiation of statin treatment as in previous studies[45]. The adjustment was made based on the assumption that most patients received atorvastatin (the only statin available in the public sector in Botswana) at a dosage of

at least 10mg per day. With an estimated adherence of 58.2 %, we calculated the baseline LDL-C levels by assuming that the measured LDL-C is a result of a 25% reduction from baseline [45]. Baseline LDL-C levels above 4.13 mmol/l were considered high[46].

Statistical analysis

Clean data were imported and analysed using Stata Version 14(Stata Corp, College Station, TX). Categorical variables are presented as percentages and continuous variables as a mean (standard deviation[SD]) or median[first–third quartiles]. Comparison of clinical and demographic factors by gender and statin use is by using the Chi-square or Fisher's exact tests for categorical variables, and independent student's t-tests or Wilcoxon rank-sum test for continuous variables as appropriate. A 2-sided p-value<0.05 was considered as statistically significant. As our outcome of interest (statin prescription) is a common event, reporting relative risks(RRs) was more desirable than odds ratios(ORs). We used generalised linear models for the binomial family, and specified log link to assess for independent predictors for statin prescribing. The multivariable model included all factors with $p < 0.2$ on bivariate analysis. We used a backward selection modelling method, with probabilities set at 0.05 and 0.1 for inclusion and exclusion; respectively. We report adjusted risk ratios (RRs), 95% confidence intervals (CIs), and p-values. With a sample size of 477 statin-eligible participants, we were able to estimate the prevalence of statin use of 13% with a margin of error of 3% on a two-sided alpha level of 0.05[22].

Results

Of the 500 patients in the dataset, the mean(SD) age was 58.9(12.2) years, 330(66%) were females. Table 1 summarises baseline patients' characteristics by gender. The majority(96.7%) of participants aged ≥ 40 years and women were significantly older than men. Approximately a third (34.4%) of participants had a diabetes duration of over ten years. Hypertension(84.7%) and obesity(51.6%) were prevalent, especially in female participants. Overall, CKD(11.3%), proteinuria(10.7%), CVD(8.8%), and smoking(3.4%) were uncommon. The mean(SD) baseline LDL-C was 3.1(1.2) mmol/L, significantly higher in female than male participants.

[Table 1: Demographic and clinical characteristics of patients with type 2 diabetes at a specialised Diabetes clinic in Gaborone (N=477)]

Statin eligibility and prescribing rates

Of the 500 participants, 477 (95.4%) were eligible for a statin prescription. Clinicians prescribed statins (exclusively atorvastatin) in 217 (45.5%) of statin-eligible participants, and only one (4.4%) ineligible participant. Seven (1.5%) of all participants received prescriptions of other lipid-lowering medications alone or in combinations with statins. Of those who were eligible for statins, statin-prescribed individuals differed from those without prescriptions in several parameters on the bivariate analysis (Table 2). Relative to the statin-non-prescribed group, the statin-prescribed group had a longer duration of diabetes (8.9 years vs. 6.0 years; $p < 0.001$); were older participants (61.5 years vs 59.2 years ; $p = 0.018$), and a

higher proportion of hypertensive patients (85.7% vs 78.1% ; $p < 0.032$), a higher proportion of participants with CKD (17.2% vs 6.6%; $p = 0.001$), and a higher baseline LDL-C (3.3 vs 2.9 mmol/L; $p < 0.001$). The two groups did not differ significantly in the presence of CVD, proteinuria and gender.

[Table 2: Factors associated with statin prescription among statin-eligible patients with type 2 diabetes at a specialised Diabetes clinic in Gaborone (N= 477)]

Multivariable analysis

The multivariable model examined adjusted associations between statin prescription and various factors. The best fit had the following covariates: age, the duration of diabetes, BMI, hypertension a high baseline LDL-C, CKD, CVD, and proteinuria. Increasing diabetes duration was associated with an increased likelihood (RR: 1.01; 95%CI 1.00-1.03) of receiving a statin prescription (Table 3), as was the presence of CKD (RR: 1.35; 95%CI: 1.06-1.74) and a high baseline LDL-C (RR: 1.49; 95%CI: 1.17-1.89). Patients' age, BMI, history of CVD, and a diagnosis of hypertension were not associated with statin prescribing after adjustment for the other variables in the model.

[Table 3 Adjusted relative risks for associations between various factors and statin prescription among statin eligible patients with diabetes at a specialised diabetes clinic in Botswana]

Discussion

Less than half of the statin-eligible patients with type 2 diabetes at a specialised diabetes clinic in Botswana received a statin prescription. A longer duration of diabetes, a higher baseline LDL-C and the presence of chronic kidney disease were independently associated with the tendency to prescribe statins.

The under-prescription of statins in our patients is a concern since the use of statins appreciably reduces cardiovascular morbidity and mortality in patients with diabetes irrespective of their LDL-C levels[7, 9-13]. Although the proportion of patients with diabetes who are prescribed statins varies substantially worldwide, there is a low prescribing of statins both in developing and developed countries[18, 19, 21-23, 47-50]. The percentage of our patients with diabetes who received statins (45.5%) in our is consistent with findings from developed countries where 25% to 73% of patients with diabetes are prescribed statins [18, 19, 47-49]. The proportion of our participants with statin prescriptions is higher than some patients in developed countries, such as Germany (25%) and Great Britain(33%)[18, 47]. While the finding of a comparatively higher statin prescription in our setting than some other settings in developed countries is encouraging, there is no reason for complacency as more than half of our patients were without CVD protection by statins. Similar to developed countries, one potential explanation for low statin prescribing rates among our patients with diabetes is inadequate adherence to guidelines[22, 40, 50, 51]. Whereas there may be a fear of the association of statin therapy with a slightly increased risk of developing diabetes, the benefits of statins in reducing cardiovascular morbidity and mortality among patients with established diabetes should dispel these concerns[7-9, 52]. Several epidemiological studies have observed a lower proportion of statin prescription in patients with diabetes in Africa than in our patients

[21-23]. In addition to inadequate adherence to guidelines, the main reasons for low statin prescribing in Africa include limited access to these medicines due to their high cost, inaccessibility to lipid testing facilities and unavailability of guidelines[22]. The presence of free consultations, tests and medications in Botswana might explain our higher statin prescribing rates than those in other African settings without universal health access. Irrespective of the reasons, it is imperative that statins are routinely prescribed to reduce the risk of CVD events in patients with type 2 diabetes[7-13, 15, 53].

Our results of increasing statin prescribing with increasing diabetes duration also agree with previous research findings[54]. This finding is reassuring as a longer duration of diabetes leads to an increased risk of CVD. For this reason, guidelines recommend statins for patients with diabetes for more than ten years [5, 6]. Although the median diabetes duration was seven years in our patients, the association of statin prescribing and duration was still apparent. The finding may suggest that clinicians correctly recognise a longer duration of diabetes as an indication for statin therapy.

As it may take time for the transmission of information between clinicians and patients, the acceptance of new medications is likely easy as the diabetes duration increases[55].

Another finding in our study was that the presence of chronic kidney disease increased the likelihood of statin prescribing. This finding is also encouraging as statins reduce mortality by up to 36% in patients with kidney failure[5, 6, 56, 57]. Besides, this finding is consistent with SEMDSA guideline recommendations of a statin for every patient with diabetes and CKD[6]. Although albuminuria is a marker of renal disease, our patients with proteinuria did not receive a statin prescription. We can postulate that clinicians do not recognise proteinuria as a predictor of CVD and an indication for statins in patients with diabetes. We will investigate this further as it contrasts with Berthold et al. who reported increased odds of statin prescribing in type 2 diabetes patients with proteinuria in Germany[18]

Our findings that a high baseline LDL-C increased the likelihood of statin prescribing agreed with those of Berthold et al. that showed an 11% increase in statin prescribing rates for every 0.26 mmol/L increase in LDL-C [18]. Besides, this finding confirms the observation from previous studies that prescribers tend to respond more to the pre-treatment LDL-C value than to the patients' overall CVD risk profile as described in clinical guidelines [9, 40]. Although there is a lack of local guidelines, the clinic adopted the SEMDSA guidelines which recommend statins along with lifestyle changes regardless of cholesterol levels for all patients with diabetes aged > 40 with or without CVD[6]. Our findings that there is approximately a 50% increased likelihood of statin prescription in our patients may suggest a need for deliberate efforts for improving the understanding and implementation of the adopted guidelines, and we will be taking this further.

In most clinical guidelines, the presence of CVD, CKD, patients age, diabetes and presence of CVD risk factors such as hypertension, albuminuria and cigarette smoking are indicators of prescribing statins among patients with type 2 diabetes[5, 6]. The recommendations are based mainly on the rationale that the presence of any of the above factors is associated with an increased risk of CVD. Except for CKD and duration of diabetes, none of the other indications was a predictor of statin prescriptions in our patients.

Given the high prevalence of hypertension and other indications in our patients, most participants would have qualified for statins if guideline recommendations are followed. As our clinic has adopted the SEMDSA guidelines, this finding is a concern and a call for efforts to improve its implementation for the benefit of this high-risk population. We will be following this up.

We are mindfully aware of the limitations of our study. We estimated the baseline LDL-C levels by a 25% adjustment of measured LDL. There was a risk of either overestimation or underestimation of the baseline LDL-C in case of significant errors in our assumptions of the dosage and the adherence of atorvastatin. Although measured LDL-C results were available for most of the included participants, HDL cholesterol results were mostly missing. Guidelines consider HDL as one of the factors for statin prescriptions in patients with diabetes. However, all other indications for statin prescriptions were available in our patients. We did not document the dosage of statin used in our patients; hence, we are unable to determine whether moderate to high-intensity statins were prescribed as recommended by the guidelines. Also, we were unable to identify patients with contraindications to warfarin as the information was not available. The study was also performed in one leading clinic, hence limiting the generalizability of the study findings to other facilities in the country. However, being one of the few specialised diabetes clinics in Botswana, our results likely characterise the 'finest' diabetes care in the country. Therefore, highlighted concerns are likely to be higher in non-specialist healthcare facilities treating patients with type 2 diabetes in Botswana.

Conclusion

In conclusion, we believe this study provides a useful and reliable picture of current statin prescribing behaviour in Botswana despite the limitations mentioned above. There is under-prescribing of statins in this high-risk population. The presence of CKD, high baseline LDL, and an increased duration of diabetes strongly influence statin prescriptions in patients with diabetes. Clinicians did not consider most guideline-recommended indications for statin prescription. By identifying gaps in the prescription of statins to patients with diabetes, the study provides an opportunity for improvement in the quality of care. Furthermore, the study findings suggest a need for further studies to investigate the reasons for statin under-prescription in our setting. We are following this up to provide future guidance for clinicians in Botswana treating patients with type 2 diabetes, with the results likely to be of interest to other sub-Saharan African countries with high rates of type 2 diabetes.

Abbreviations

ASCVD: atherosclerotic cardiovascular disease; BMI: Body mass index; CAD: coronary artery disease; CKD: chronic kidney disease; CVD: Cardiovascular disease; eGFR: estimated glomerular filtration rate; HDL-C: HbA1c: Haemoglobin A1c; HDL-C: High-density lipoprotein cholesterol; HRDC: Health Research Development Committee; LDL-C: Low-density lipoprotein cholesterol; MDRD: Modification of Diet in Renal Disease PAD: peripheral artery disease; SEMDSA: Society for Endocrinology, Metabolism, and Diabetes of South Africa WHR: Waist-Hip ratio.

Declarations

Ethics approval and consent to participate

The Health Research Development Committee(HRDC) of the Botswana Ministry of Health and Wellness(HPDME:13/18/1) and Stellenbosch University Health Research Ethics Committee (X19/01/001) approved the study. The participating patients provided written informed consent in the primary study.

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request and with permission of the HRDC of Botswana Ministry of Health and Wellness.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding was received.

Authors' contributions

JCM, BG and EMT conceptualised the study. JCM analysed data and drafted the initial draft. ETM and BG critically reviewed data and reviewed the manuscripts. All the authors read and approved the final manuscript.

Acknowledgements

This research project has been conducted as part of the academic requirements of the MSc in Clinical Epidemiology www.sun.ac.za/clinepi, Stellenbosch University[58].

References

1. Di Angelantonio E, Kaptoge S, Wormser D, Willeit P, Butterworth AS, Bansal N, O'Keefe LM, Gao P, Wood AM, Burgess S *et al*: **Association of Cardiometabolic Multimorbidity With Mortality**. *JAMA* 2015, **314**(1):52-60.
2. Huxley R, Barzi F, Woodward M: **Excess risk of fatal coronary heart disease associated with diabetes in men and women: meta-analysis of 37 prospective cohort studies**. *BMJ (Clinical research ed)* 2006, **332**(7533):73-78.

3. Lloyd-Jones DM, Leip EP, Larson MG, D'Agostino RB, Beiser A, Wilson PW, Wolf PA, Levy D: **Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age.** *Circulation* 2006, **113**(6):791-798.
4. Reusch JE DB: **Atherosclerosis in diabetes and insulin resistance.** *Diabetes Obes Metab* 2007(9):455-463.
5. American Diabetes Association: **Standards of medical care in diabetes—2017 abridged for primary care providers.** *Clinical Diabetes* 2017, **35**(1):5-26.
6. SEMDSA Type 2 Diabetes Guidelines Expert Committee: **SEMDSA 2017 guidelines for the management of type 2 diabetes mellitus.** *J Endocr Metab Diabetes S Afr* 2017, **22**(1 Suppl 1):S1 - S196.
7. Colhoun HM, Betteridge DJ, Durrington PN, Hitman GA, Neil HA, Livingstone SJ, Thomason MJ, Mackness MI, Charlton-Menys V, Fuller JH: **Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): a multicentre randomised placebo-controlled trial.** *Lancet (London, England)* 2004, **364**(9435):685-696.
8. Collins R, Armitage J, Parish S, Sleight P, Peto R: **MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial.** *Lancet (London, England)* 2003, **361**(9374):2005-2016.
9. Sever PS, Poulter NR, Dahlof B, Wedel H, Collins R, Beevers G, Caulfield M, Kjeldsen SE, Kristinsson A, McInnes GT *et al*: **Reduction in cardiovascular events with atorvastatin in 2,532 patients with type 2 diabetes: Anglo-Scandinavian Cardiac Outcomes Trial–lipid-lowering arm (ASCOT-LLA).** *Diabetes care* 2005, **28**(5):1151-1157.
10. Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhalra N, Peto R, Barnes EH, Keech A, Simes J *et al*: **Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials.** *Lancet (London, England)* 2010, **376**(9753):1670-1681.
11. de Vries FM, Denig P, Pouwels KB, Postma MJ, Hak E: **Primary prevention of major cardiovascular and cerebrovascular events with statins in diabetic patients: a meta-analysis.** *Drugs* 2012, **72**(18):2365-2373.
12. Hayward RA, Hofer TP, Vijan S: **Narrative review: lack of evidence for recommended low-density lipoprotein treatment targets: a solvable problem.** *Annals of internal medicine* 2006, **145**(7):520-530.
13. Perk J, De Backer G, Gohlke H, Graham I, Reiner Z, Verschuren M, Albus C, Benlian P, Boysen G, Cifkova R *et al*: **European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts).** *European heart journal* 2012, **33**(13):1635-1701.
14. Perreault S, Dragomir A, Blais L, Berard A, Lalonde L, White M, Pilon D: **Impact of better adherence to statin agents in the primary prevention of coronary artery disease.** *European journal of clinical pharmacology* 2009, **65**(10):1013-1024.

15. Kearney P, Blackwell L, Collins Ra, Keech A, Simes J, Peto R, Armitage J, Baigent C: **Cholesterol Treatment Trialists'(CTT) Collaborators. Efficacy of cholesterol-lowering therapy in 18,686 people with diabetes in 14 randomised trials of statins: a meta-analysis.** *Lancet (London, England)* 2008, **371**(9607):117-125.
16. Eliasson B, Svensson AM, Miftaraj M, Jonasson JM, Eeg-Olofsson K, Sundell KA, Gudbjornsdottir S: **Clinical use and effectiveness of lipid lowering therapies in diabetes mellitus—an observational study from the Swedish National Diabetes Register.** *PloS one* 2011, **6**(4):e18744.
17. Balder JW, Scholtens S, de Vries JK, van Schie LM, Boekholdt SM, Hovingh GK, Kamphuisen PW, Kuivenhoven JA: **Adherence to guidelines to prevent cardiovascular diseases: The LifeLines cohort study.** *The Netherlands journal of medicine* 2015, **73**(7):316-323.
18. Berthold HK, Gouni-Berthold I, Bohm M, Krone W, Bestehorn KP: **Patterns and predictors of statin prescription in patients with type 2 diabetes.** *Cardiovascular diabetology* 2009, **8**:25.
19. Harrison TN, Scott RD, Cheetham TC, Chang S-C, Hsu J-WY, Wei R, Ling Grant DS, Boklage SH, Romo-LeTourneau V, Reynolds K: **Trends in Statin Use 2009-2015 in a Large Integrated Health System: Pre- and Post-2013 ACC/AHA Guideline on Treatment of Blood Cholesterol.** *Cardiovascular drugs and therapy* 2018, **32**(4):397-404.
20. Simmons RK, Carlsen AH, Griffin SJ, Charles M, Christiansen JS, Borch-Johnsen K, Sandbaek A, Lauritzen T: **Variation in prescribing of lipid-lowering medication in primary care is associated with incidence of cardiovascular disease and all-cause mortality in people with screen-detected diabetes: findings from the ADDITION-Denmark trial.** *Diabetic medicine : a journal of the British Diabetic Association* 2014, **31**(12):1577-1585.
21. Gudina EK, Amade ST, Tesfamichael FA, Ram R: **Assessment of quality of care given to diabetic patients at Jimma University Specialized Hospital diabetes follow-up clinic, Jimma, Ethiopia.** *BMC Endocrine Disorders* 2011, **11**(1):19.
22. Sobngwi E, Ndour-Mbaye M, Boateng KA, Ramaiya KL, Njenga EW, Diop SN, Mbanya JC, Ohwovoriole AE: **Type 2 diabetes control and complications in specialised diabetes care centres of six sub-Saharan African countries: the Diabcare Africa study.** *Diabetes research and clinical practice* 2012, **95**(1):30-36.
23. Uloko AE, Ofoegbu EN, Chinenye S, Fasanmade OA, Fasanmade AA, Ogbera AO, Ogbu OO, Oli JM, Girei BA, Adamu A: **Profile of Nigerians with diabetes mellitus - Diabcare Nigeria study group (2008): Results of a multicenter study.** *Indian journal of endocrinology and metabolism* 2012, **16**(4):558-564.
24. Cameron A, Ewen M, Ross-Degnan D, Ball D, Laing R: **Medicine prices, availability, and affordability in 36 developing and middle-income countries: a secondary analysis.** *Lancet (London, England)* 2009, **373**(9659):240-249.
25. Ofori-Asenso R, Agyeman AA: **Irrational Use of Medicines—A Summary of Key Concepts.** *Pharmacy* 2016, **4**, 35.
26. Kayima J, Wanyenze RK, Katamba A, Leontsini E, Nuwaha F: **Hypertension awareness, treatment and control in Africa: a systematic review.** *BMC cardiovascular disorders* 2013, **13**:54.

27. Lloyd-Sherlock P, Beard J, Minicuci N, Ebrahim S, Chatterji S: **Hypertension among older adults in low- and middle-income countries: prevalence, awareness and control.** *International journal of epidemiology* 2014, **43**(1):116-128.
28. **IDF Diabetes Atlas** [https://diabetesatlas.org/IDF_Diabetes_Atlas_8e_interactive_EN/]
29. Adeniyi OV, Yogeswaran P, Longo-Mbenza B, Ter Goon D: **Uncontrolled Hypertension and Its Determinants in Patients with Concomitant Type 2 Diabetes Mellitus (T2DM) in Rural South Africa.** *PloS one* 2016, **11**(3):e0150033.
30. Hamid S, Groot W, Pavlova M: **Trends in cardiovascular diseases and associated risks in sub-Saharan Africa: a review of the evidence for Ghana, Nigeria, South Africa, Sudan and Tanzania.** *The aging male : the official journal of the International Society for the Study of the Aging Male* 2019:1-8.
31. Rannanheimo PK, Tiittanen P, Hartikainen J, Helin-Salmivaara A, Huupponen R, Vahtera J, Korhonen MJ: **Impact of Statin Adherence on Cardiovascular Morbidity and All-Cause Mortality in the Primary Prevention of Cardiovascular Disease: A Population-Based Cohort Study in Finland.** *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research* 2015, **18**(6):896-905.
32. Blake AM, Munby HN, Katlego PM, Sebuyuyu PM, Nkomozana O, Bangure R, Kerr-Muir M, Ngondi JM: **Characteristics of patients with diabetic retinopathy in Gaborone, Botswana.** *Tanzania Journal of Health Research* 2015, **17**(1).
33. Mwita JC, Magafu M, Omech B, Tsimba B, Dewhurst MJ, Goepamang M, Mashalla Y: **Undiagnosed and diagnosed diabetes mellitus among hospitalised acute heart failure patients in Botswana.** *SAGE open medicine* 2017, **5**:2050312117731473.
34. Omech B, Mwita JC, Tshikuka JG, Tsimba B, Nkomazna O, Amone-P'Olak K: **Validity of the Finnish Diabetes Risk Score for Detecting Undiagnosed Type 2 Diabetes among General Medical Outpatients in Botswana.** *Journal of diabetes research* 2016, **2016**:4968350.
35. Mwita JC, Francis JM, Omech B, Botsile E, Oyewo A, Mokgwathi M, Molefe-Baikai OJ, Godman B, Tshikuka JG: **Glycaemic, blood pressure and low-density lipoprotein-cholesterol control among patients with diabetes mellitus in a specialised clinic in Botswana: a cross-sectional study.** *BMJ Open* 2019, **9**(7):e026807.
36. STEPS W: **Section 3: guide to physical measurements (Step 2).** In.: Geneva; 2008.
37. World Health Organization: **Waist circumference and waist-hip ratio: report of a WHO expert consultation, Geneva, 8-11 December 2008.** In.; 2011.
38. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., Jones DW, Materson BJ, Oparil S, Wright JT, Jr. *et al.*: **The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report.** *JAMA* 2003, **289**(19):2560-2572.
39. Friedewald WT, Levy RI, Fredrickson DS: **Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge.** *Clinical chemistry* 1972, **18**(6):499-502.

40. Barham AH, Goff DC, Jr., Chen H, Balasubramanyam A, Rosenberger E, Bonds DE, Bertoni AG: **Appropriateness of cholesterol management in primary care by sex and level of cardiovascular risk.** *Preventive cardiology* 2009, **12**(2):95-101.
41. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Ogedegbe O: **2014 evidence-based guideline for the management of high blood pressure in adults: a report from the panel members appointed to the Eighth Joint National Committee (JNC 8).** *JAMA* 2014, **311**(5):507-520.
42. White AD, Folsom AR, Chambless LE, Sharret AR, Yang K, Conwill D, Higgins M, Williams OD, Tyroler HA: **Community surveillance of coronary heart disease in the Atherosclerosis Risk in Communities (ARIC) Study: methods and initial two years' experience.** *Journal of clinical epidemiology* 1996, **49**(2):223-233.
43. Levey AS, Greene T, Schluchter MD, Cleary PA, Teschan PE, Lorenz RA, Molitch ME, Mitch WE, Siebert C, Hall PM *et al*: **Glomerular filtration rate measurements in clinical trials. Modification of Diet in Renal Disease Study Group and the Diabetes Control and Complications Trial Research Group.** *Journal of the American Society of Nephrology: JASN* 1993, **4**(5):1159-1171.
44. Expert Panel on the Identification E, and Treatment of Overweight in Adults,: **Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: executive summary. Expert Panel on the Identification, Evaluation, and Treatment of Overweight in Adults.** *The American journal of clinical nutrition* 1998, **68**(4):899-917.
45. Law MR, Wald NJ, Rudnicka AR: **Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis.** *BMJ (Clinical research ed)* 2003, **326**(7404):1423.
46. Expert Panel on Detection E, Adults ToHBCi: **Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III).** *JAMA* 2001, **285**(19):2486-2497.
47. Chang KC, Soljak M, Lee JT, Woringer M, Johnston D, Khunti K, Majeed A, Millett C: **Coverage of a national cardiovascular risk assessment and management programme (NHS Health Check): Retrospective database study.** *Preventive medicine* 2015, **78**:1-8.
48. Ofori-Asenso R, Ilomaki J, Tacey M, Zomer E, Curtis AJ, Bell JS, Zoungas S, Liew D: **Patterns of statin use and long-term adherence and persistence among older adults with diabetes.** *Journal of diabetes* 2018, **10**(9):699-707.
49. Steen DL, Khan I, Becker L, Foody JM, Gorcyca K, Sanchez RJ, Giugliano RP: **Patterns and predictors of lipid-lowering therapy in patients with atherosclerotic cardiovascular disease and/or diabetes mellitus in 2014: Insights from a large US managed-care population.** *Clinical cardiology* 2017, **40**(3):155-162.
50. Lemstra M BD, Crawley A, Fung R.: **Proportion and risk indicators of nonadherence to statin therapy: a meta-analysis.** *Can J Cardiol* 2012, **28**(5):574-580.

51. Bai JW, Boulet G, Halpern EM, Lovblom LE, Eldelekli D, Keenan HA, Brent M, Paul N, Bril V, Cherney DZI *et al*: **Cardiovascular disease guideline adherence and self-reported statin use in longstanding type 1 diabetes: results from the Canadian study of longevity in diabetes cohort.** *Cardiovascular diabetology* 2016, **15**:14-14.
52. Sattar N, Preiss D, Murray HM, Welsh P, Buckley BM, de Craen AJ, Seshasai SR, McMurray JJ, Freeman DJ, Jukema JW *et al*: **Statins and risk of incident diabetes: a collaborative meta-analysis of randomised statin trials.** *Lancet (London, England)* 2010, **375**(9716):735-742.
53. Godman B, Shrank W, Andersen M, Berg C, Bishop I, Burkhardt T, Garuoliene K, Herholz H, Joppi R, Kalaba M *et al*: **Policies to enhance prescribing efficiency in europe: findings and future implications.** *Frontiers in pharmacology* 2010, **1**:141.
54. Casagrande SS, Aviles-Santa L, Corsino L, Daviglius ML, Gallo LC, Espinoza Giacinto RA, Llabre MM, Reina SA, Savage PJ, Schneiderman N *et al*: **HEMOGLOBIN A1C, BLOOD PRESSURE, AND LDL-CHOLESTEROL CONTROL AMONG HISPANIC/LATINO ADULTS WITH DIABETES: RESULTS FROM THE HISPANIC COMMUNITY HEALTH STUDY/STUDY OF LATINOS (HCHS/SOL).** *Endocrine practice : official journal of the American College of Endocrinology and the American Association of Clinical Endocrinologists* 2017, **23**(10):1232-1253.
55. Tarn DM, Paterniti DA, Kravitz RL, Heritage J, Liu H, Kim S, Wenger NS: **How much time does it take to prescribe a new medication?** *Patient education and counseling* 2008, **72**(2):311-319.
56. Seliger SL, Weiss NS, Gillen DL, Kestenbaum B, Ball A, Sherrard DJ, Stehman-Breen CO: **HMG-CoA reductase inhibitors are associated with reduced mortality in ESRD patients.** *Kidney international* 2002, **61**(1):297-304.
57. Tonelli M, Keech A, Shepherd J, Sacks F, Tonkin A, Packard C, Pfeffer M, Simes J, Isles C, Furberg C *et al*: **Effect of Pravastatin in People with Diabetes and Chronic Kidney Disease.** *Journal of the American Society of Nephrology* 2005, **16**(12):3748-3754.
58. Mwita JC: **Statin prescription among patients with type 2 diabetes at a specialised diabetes clinic, Botswana.** 2019.

Tables

Table 1: Demographic and clinical characteristics of patients with type 2 diabetes at a specialised Diabetes clinic in Gaborone (N= 477)

Characteristics	All (N=477)	Males (n= 160)	Females (n= 317)
Age, mean(SD), years	60.3(10.8)	56.8(11.5)	62.0(10.1)
Age > 40 years n(%)	461(96.7)	150(93.8)	311(98.1)
Diabetes duration, median, IQR, years	7(3-13)	8.9(3-14)	7 (3-13)
Diabetes duration > 10 years	164 (34.4)	63(39.4)	101(31.9)
BMI, mean (SD) kg/m²	30.7(6.0)	29.0(5.2)	31.6(6.1)
Normal weight n(%)	85(17.8)	39(24.4)	46(14.5)
Overweight n(%)	146(30.6)	59(36.7)	87(27.4)
Obese n(%)	246(51.6)	62(38.8)	184(58.0)
Marital status			
Living alone n(%)	247(51.8)	47(29.4)	200(63.1)
Living with a partner n(%)	230(48.2)	113(70.6)	117(36.9)
Education status			
≤ Primary education, n(%)	306(64.1)	82(51.3)	224(70.7)
≥ Secondary or tertiary, n(%)	171(35.9)	78(48.7)	93(29.3)
WHR, mean (SD)	0.94(0.10)	0.97(0.08)	0.93(0.09)
Low WHR n (%)	79(16.6)	67(41.9)	12(3.8)
High WHR n (%)	398(83.4)	93(58.1)	305(96.2)
Hypertension n (%)	404(84.7)	120(75.0)	284(89.6)
Use of antihypertensive n (%)	389(81.6)	110(68.8)	279(88.0)
Smoking n (%)	16(3.4)	12(7.5)	4(1.3)
Lipid-lowering medications n (%)	224(47.0)	70(43.8)	154(48.6)
Statins n (%)	217(45.5)	68(42.5)	149(47.0)
Others n (%)	7(1.5)	2(1.6)	5(1.6)
CVD n (%)	42(8.8)	15(9.4)	27(8.5)
PAD n (%)	11(2.3)	3(1.9)	8(2.5)
Coronary artery disease n (%)	12(2.5)	5(3.3)	7(2.2)
Cerebrovascular disease n (%)	22(4.6)	8(5.0)	14(4.4)
CKD n (%)	54(11.3)	24(15.0)	30(9.5)
Proteinuria n (%)	51(10.7)	27(16.9)	24(7.6)
HbA1c mean (SD), %	8.4(2.4)	8.5(2.6)	8.3(2.3)
Baseline LDL-C, mean (SD), mmol/L	3.1(1.2)	2.8(1.1)	3.3(1.2)

Normal	315(66.0)	113(70.6)	202(63.7)
High	59(12.4)	9(5.6)	50(15.8)
Missing	103(21.6)	38(23.8)	65(20.5)

Legend: BMI- Body Mass Index, CKD – chronic kidney disease, CVD – cardiovascular disease, HbA1c – Haemoglobin A1c, IQR – interquartile range; LDL-C – low-density lipoprotein cholesterol, PAD – peripheral artery disease, SD – standard deviation; WHR – waist-hip ratio.

Table 2: Factors associated with statin prescription among statin-eligible patients with type 2 diabetes at a specialised Diabetes clinic in Gaborone (N= 477)

Characteristics	Statin not prescribed (n=260)	Statin prescribed (n= 217)	P-value
Sex			
Males, n (%)	92(35.4)	68(31.3)	0.351
Female, n (%)	168(64.6)	149(68.7)	
Diabetes duration, median, IQR, years	6(2 – 12)	8.9(4 – 15)	<0.001
Duration ≤ 10 years	179(68.9)	134(61.8)	0.104
Diabetes duration > 10 years	81(31.1)	83(38.2)	
Age, mean (SD), years	59.2(10.8)	61.5(10.7)	0.018
Age ≤ 40 years, n (%)	12(4.6)	4(1.8)	<0.094
Age > 40 years n (%)	248(95.4)	213(98.2)	
Marital status			
Living alone n (%)	135(51.9)	112(51.6)	0.946
Living with a partner n (%)	125(48.1)	105 (48.4)	
Education status			
≤ Primary education, n (%)	166(63.9)	140(64.2)	0.879
≥ Secondary or tertiary, n (%)	94(36.1)	77(35.8)	
Hypertension n (%)	212(81.5)	192 (88.5)	0.036
Antihypertensive use n (%)	203(78.1)	186(85.7)	0.032
Smoking	13(5.0)	3(1.4)	0.029
CVD, n (%)	20(7.7)	22(10.1)	0.538
- PAD, n (%)	7(2.7)	4(1.8)	0.625
- Coronary artery disease n (%)	6(2.3)	6(2.8)	0.751
- Cerebrovascular diseases n (%)	9(3.5)	13(6.0)	0.190
BMI, mean (SD) kg/m ²	30.3(5.7)	31.2(6.2)	0.113
Normal weight n (%)	50(19.2)	35(16.1)	0.669
Overweight n (%)	79(30.4)	67(30.9)	
Obese n (%)	135(47.9)	115(52.8)	
WHR, mean (SD) kg/m ²	0.94(0.08)	0.95(0.10)	0.106
Low WHR n (%)	46(17.7)	33(15.2)	0.467

High WHR n (%)	214(82.3)	184(84.8)	
CKD, n (%)	20(7.7)	34 (15.7)	0.006
Proteinuria, n (%)	30 (11.5)	21(9.7)	0.512
HbA1c mean (SD), %	8.4(2.6)	8.4(2.2)	0.948
Baseline LDL-C mean (SD), mmol/L	2.9(0.9)	3.3(1.4)	0.003
Normal	177(68.1)	120(55.3)	<0.001
High	17(6.5)	42(19.4)	
Missing	66(25.4)	55(25.3)	

Legend: BMI- Body Mass Index, CKD – chronic kidney disease, CVD – cardiovascular disease, HbA1c – Haemoglobin A1c, IQR – interquartile range; LDL-C – low-density lipoprotein cholesterol, PAD – peripheral artery disease, SD – standard deviation; WHR – waist-hip ratio.

Table 3 Adjusted relative risks for associations between various factors and statin prescription among statin eligible patients with diabetes at a specialised diabetes clinic in Botswana

Characteristic	Risk ratio	95% Conf. Interval	p-value
Age	1.006	0.994-1.017	0.362
CKD	1.354	1.055-1.738	0.017
Hypertension	1.336	0.846-2.110	0.213
BMI	1.014	0.994-1.034	0.16
High baseline LDL	1.488	1.173-1.887	0.001
Diabetes duration, years	1.014	1.000-1.027	0.048
Proteinuria	0.979	0.644-1.488	0.922
CVD	0.901	0.623-1.303	0.581

Legend: BMI- Body Mass Index, CKD – chronic kidney disease, CVD – cardiovascular disease, LDL-C – low-density lipoprotein cholesterol.