


Mortality rate of infection with COVID-19 in Korea from the perspective of underlying disease

Yun Jung Kang (✉ lvpig@naver.com)

Short report

Keywords: COVID-19, Mortality rate, Over 70, Perspective, Prevention, Underlying diseases

DOI: <https://doi.org/10.21203/rs.3.rs-18975/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

On December 31st, 2019 the China National Health Commission (NHC) reported that an unknown cause of pneumonia had been detected in Wuhan in Hubei province. On February 12th, the disease caused by novel coronavirus (2019-nCoV) has a formal name, COVID-19. On January 20th, 2020, the first case of COVID-19 was confirmed in Korea. Among the deaths, age-specific death rate was the highest among cases over 70's, with underlying diseases in their circulatory system, such as myocardial infraction, cerebral infraction, arrythmia, and hypertension. Patients with underlying disease who are 70 years of age or older should recognize that there is a high possibility of developing a serious disease in case of viral infection and follow strict precautions.

Short Report

On December 31st, 2019 the China National Health Commission (NHC) reported that an unknown cause of pneumonia had been detected in Wuhan in Hubei province. The NHC later confirmed that the infection was a novel coronavirus-infected pneumonia (NCIP). On February 12th, the disease caused by novel coronavirus (2019-nCoV) has a formal name, COVID-19. On January 20th, 2020, the first case of COVID-19 was confirmed in Korea [1]. After the first COVID-19 case was confirmed on January 20th, 2020, the Korea Centers for Disease Control and Prevention has focused on delaying the inflow of the virus into Korea and its spread in local communities, with considerable success. Also, the mortality rate of the COVID-19 outside China was rather low, leading to the expectation that the disease's impact on national health would be minor. However, things turned over rapidly after Case No. 31. The epidemic is spreading rapidly on global level [2]. The Secretary General of World Health Organization, Tedros Adhanom Ghebreyesus, warned that the COVID-19 epidemic has reached its watershed and that every state should prepare for it [3]. Among the symptoms of COVID-19 are fever and minor respiratory symptoms such as dry coughs, which overlap with other respiratory diseases; therefore, it is not easy to confirm a case based only on early symptoms. When the symptoms get worse and lead to serious respiratory symptoms such as dyspnea, lowering of oxygen saturation, and pneumonia, they can cause death [4].

Current situation in Korea regarding COVID-19 occurrences at 00:00, March 16th, 2020 is as follows: the total number of confirmed cases are 8,236, with 1,137 released from quarantine among them. New confirmed cases are counted to be 74, while total number of deaths is 75. Most of the dead has had underlying diseases or been in their old age (Table 1). The mortality rate is known as 1-2% but cannot be confirmed yet. Among the deaths, age-specific death rate was the highest among cases over 70's, with underlying diseases in their circulatory system, such as myocardial infraction, cerebral infraction, arrythmia, and hypertension. By the term "underlying diseases," we refer to chronic diseases of the patient, such as hypertension, diabetes, asthma, renal failure, and tuberculosis [5]. Korea had its first case of swine flu in May 2009, with the first death on August 15th the same year. After that, on October 26th, the government announced children under 59-month old, pregnant women and mothers within two weeks of delivery, citizens over 65 years old, patients with chronic lung diseases, chronic cardiovascular diseases (except for hypertension), diabetes, chronic renal diseases, chronic liver diseases, cancers, people with weakened immunity, and other patients with absorption risks as high risk group to the complications of the swine flu and recommended them treatment in time, according to its 6th version of the Guidelines for Preventing and Managing Swine Flu [6]. Currently, the Korea Centers for Disease Control and Prevention is planning to categorize patients according to their pulse, age, and underlying diseases they had on being found infected; the Centers would transfer critical patients to negative pressure isolation rooms designated by the government for proper treatment. The Centers came up with this new plan in the situation where the number of confirmed cases and death increased rapidly, leading to the importance of judging and categorizing the seriousness of patients' situation [7].

Korean Diabetes Association argued that patients who are suffering underlying diseases with high risk of death should be given access to early diagnosis and treatment of COVID-19. It requested preferential opportunities of examination and hospitalization for people over 70 with diabetes when they had suspected symptoms. According to a recent research on Chinese patients, which was published in The Journal of the American Medical Association, showed the overall death rate of 2.3% among 44,672 patients; however, the mortality rate leaped to 8.0% in people in their 70's and 14.8% in people in their 80's. Patients with diabetes also showed higher mortality rate of 7.3% [8].

Even when they are exposed to the virus on the same conditions, people with underlying diseases should be aware that they are more susceptible to infection than people without them, as they have weaker immunities; they must adhere to the prevention regulation strictly. Especially, citizens over 70 with underlying diseases should be classified as high-risk group and managed carefully. This research would be provided as a basic material for guidelines regarding disease prevention and management of high-risk group among confirmed cases for future infectious diseases.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Consent for publication was included in the consent to participate form.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

FUNDING

None.

Authors' contributions

The author (YJ Kang) wrote the entire manuscript and holds final responsibility for the decision to submit the manuscript for publication. The author read and approved the final manuscript.

ACKNOWLEDGEMENTS

None.

ORCID

Yun Jung Kang <https://orcid.org/0000-0002-1425-3178>

References

1. Hyun JH, Kim JH, Lee HY, Kim JE, Lee EY, Kim TK, Kim HS, Contact Tracing Results of the First Confirmed COVID-19 Case in the Republic of Korea. *Weekly health and illness* 2020; 13(7): 352-358.
2. Kim NS, Coronavirus Infection-19 Status and Challenges. *Health and Welfare Issue&Focus* 2020; 373: 1-13. DOI: <http://repository.kihasa.re.kr/bitstream/201002/34380/8/%ec%9d%b4%ec%8a%88%ec%95%a4%ed%8f%ac%ec%bb%a4%ec%8a%a4.2020.N0373.pdf>
3. Gu GI, Jo YR. "Corona 19, has a decisive move"... WHO's warning. *Dong A*; 2020 Feb 28 [cited 2020 Mar 16]. Available from: <http://www.donga.com/news/List/article/all/20200228/99928838/1>(Korean).
4. Korea Centers for Disease Control and Prevention.; 2020 Mar 7 [cited 2020 Mar 16]. Available from: http://ncov.mohw.go.kr/tcmBoardView.do?brdId=&brdGubun=&dataGubun=&ncvContSeq=353594&contSeq=353594&board_id=140&gubun=BDJ (Korean).
5. Heyland DK, Frank C, Groll D, Pichora D, Dodek P, Rocker G, et al, Understanding cardiopulmonary resuscitation decision making: Perspectives of seriously ill hospitalized patients and family members. *Chest* 2006; 130: 419-428.
6. Lee HG, Kang YA, Kim HS, Shin SY, Kim JH, Kim JS, Hong SB, Epidemiological Characteristics Based on the Underlying Diseases for the Deaths Related to Pandemic H1N1 Influenza in Korea. *Acute and Critical Care* 2010; 25(2): 83-88. DOI: 10.4266/kjccm.2010.25.2.83
7. "Daegu confirmed 570 waiting for admission"...Efficient efficiency due to lack of beds. *Korea Economy*; 2020 Feb 27 [cited 2020 Mar 16]. Available from: <https://www.hankyung.com/society/article/202002270269Y>(Korean).
8. Kim AR. "If you have an underlying disease, you need an early check for corona." *Health News*; 2020 Mar 5 [cited 2020 Mar 16]. Available from: <http://m.bokuennews.com/news/article.html?no=187467>(Korean).

Table 1

Table. 1 Status and characteristics of domestic deaths (00:00, March 16th)

Category		Persons	Rate (%)	Remarks
Deaths	-	75		Mortality rate (out of 8,236 confirmed patients) 0.91%
Mortality rates by age	30s (849 confirmed)	1	0.12%	
	40s (1,147 confirmed)	1	0.09%	
	50s (1,585 confirmed)	6	0.38%	
	60s (1,024 confirmed)	14	1.37%	
	70s (531 confirmed)	28	5.27%	
	80s ≤ (270 confirmed)	25	9.26%	
Underlying disease (Can be duplicated)	Circulatory system disease (*)	47	62.7%	(*) Myocardial infarction, cerebral infarction, arrhythmia, hypertension, etc. (**) Diabetes, hypothyroidism, etc. (***) Dementia, schizophrenia, etc. (****) Asthma, chronic obstructive pulmonary disease, pneumonia, etc.
	Endocrine and metabolic diseases (**)	35	46.7%	
	Mental illness (***)	19	25.3%	
	Respiratory diseases (****)	18	24.0%	
	Urinary and reproductive system diseases	11	14.7%	
	Malignant neoplasm (cancer)	10	13.3%	
	Nervous system diseases etc.	3	4.0%	
	Digestive system diseases	2	2.7%	
	Blood and hematopoietic diseases	1	1.3%	
High risk group	65 years and older	61	81.3%	
	underlying disease(o)	74	98.7%	

[source] [Korea Centers for Disease Control and Prevention](#)