

407 **Additional Information**

408

409 **Supplementary Table 1. Correlation between VNT and sVNT for different COVID-19 sera\***

410

411

Sample	VNT	sVNT			
		90%	70%	50%	30%
COVID-19 Patient-1	<10	<10	10	20	40
COVID-19 Patient-2	40	20	40	80	320
COVID-19 Patient-3	20	<10	20	40	80
COVID-19 Patient-4	10	<10	<10	10	20
COVID-19 Patient-5	20	10	20	40	80
COVID-19 Patient-6	<10	<10	<10	10	20
COVID-19 Patient-7	40	<10	20	40	80
COVID-19 Patient-8	10	<10	20	40	80
COVID-19 Patient-9	20	10	20	40	80
COVID-19 Patient-10	10	10	40	80	160
COVID-19 Patient-11	20	20	80	160	640
COVID-19 Patient-12	80	20	80	160	640
COVID-19 Patient-13	160	160	320	640	1280
Negative Control-1	<10	<10	<10	<10	<10
Negative Control-2	<10	<10	<10	<10	<10

412

413 \*Neutralization titers were obtained from two biological replicates each with two technical

414 replicates. sVNT titers at 90%, 70%, 50% and 30% inhibition, respectively, are shown.

415

416

417 **Supplementary Fig. 1. Titration of COVID-19 sera with different levels of SARS-CoV-2**  
418 **antibodies.** Serum samples were two-fold diluted starting at 1:10. Percentage inhibition was  
419 plotted at each dilution point for two negative controls and 13 COVID-19 sera from PCR  
420 positive patients.

421  
422  
423

