

# Computing Total Edge Irregularity Strength for Heptagonal Snake Graph and Related Graphs

Fatma Salama (✉ [fatma.salama@science.tanta.edu.eg](mailto:fatma.salama@science.tanta.edu.eg))

Tanta University Faculty of Science Tanta Gharbia Governorate EG <https://orcid.org/0000-0002-3555-7349>

---

## Research Article

**Keywords:** Irregular labelling, Total edge irregularity strength, Edge irregular total labeling, Heptagonal

**Posted Date:** May 3rd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-260228/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Computing Total Edge Irregularity Strength for Heptagonal Snake Graph and Related Graphs

F. Salama

Mathematics Department, Faculty of Science, Tanta University, Tanta, Egypt;

Email: fatma.salama@science.tanta.edu.eg

**Abstract:** A labeling of edges and vertices of a simple graph  $G(V, E)$  by a mapping  $\mathbb{T}: V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, \mathbb{T}\}$  provided that any two pair of edges have distinct weights is called an edge irregular total  $\mathbb{T}$ -labeling. If  $\mathbb{T}$  is minimum and  $G$  admits an edge irregular total  $\mathbb{T}$ -labelling, then  $\mathbb{T}$  is called the total edge irregularity strength (TEIS) and denoted by  $tes(G)$ . In this paper, the definitions of the heptagonal snake graph  $HPS_n$ , the double heptagonal snake graph  $D(HPS_n)$  and an  $l$ -multiple heptagonal snake graph  $L(HPS_n)$  have been introduced. The exact value of TEISs for the new family has also been investigated.

**Keywords:** Irregular labelling; Total edge irregularity strength; Edge irregular total labeling; Heptagonal snake graph.

**2010 Mathematics subject classification:** 05C78.

## 1. Introduction

In graph theory, graph labeling is an assignment of labels or weights to the vertices and/or edges of a graph. Graph labeling plays an important role in many fields such as computer science, coding theory, astronomy and physics. For a connected, simple and undirected graph  $G(V, E)$ , an edge I irregular I total  $\mathbb{T}$ -labeling  $\mathbb{T}: V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, \mathbb{T}\}$  has been defined in [1] as a labeling of its vertices and edges such that for any pair of edges  $pq$  and  $p^*q^*$  in a graph  $G$  their weights are distinct, i.e.  $w_{\mathbb{T}}(pq) \neq w_{\mathbb{T}}(p^*q^*)$  where  $w_{\mathbb{T}}(pq) = \mathbb{T}(pq) + \mathbb{T}(p) + \mathbb{T}(q)$ . A minimum  $\mathbb{T}$  is a total edge irregularity strength of a graph  $G$  when  $G$  admits an edge irregular total  $\mathbb{T}$ -labelling. Also, the authors in [1] introduced an inequality that gives bounds of TEIS of a graph  $G$

$$tes(G) \geq \max \left\{ \left\lceil \frac{|E(G)|+2}{3} \right\rceil, \left\lceil \frac{\Delta G+1}{2} \right\rceil \right\} \quad (1).$$

In [2], Ivančo and Jendroř has deduced TEIS for a tree. They also introduced the following conjecture

Conjecture 1. Let  $G$  be a graph different from  $K_5$ , then

$$tes(G) = \max \left\{ \left\lceil \frac{\Delta G + 1}{2} \right\rceil, \left\lceil \frac{|E(G)| + 2}{3} \right\rceil \right\}.$$

Since then, the conjecture has been verified for centralized uniform theta graphs in Putra et al. [3], for a polar grid graph in Salama [4], for hexagonal grid graphs in Al-Mushayt et al. [5], for Disjoint Union of Wheel Graphs in Jeyanthi [6], for strong product of two paths in Ahmad et al.[7], for subdivision of star  $S_n$  in Siddiqui [8], for generalized prism in Bača and Siddiqui [9], for helm and sun graphs in Ahmad et al.[10], for the corona product of paths with some graphs in Salman and Baskoro [11], for a wheel graph, a fan graph, a triangular Book graph and a friendship graph in Tilukay et al. [12], for a categorical product of

two paths in Ahmad and Bača [13], for subdivision of star in Hinding et al. [14], for series parallel graphs in Rajasingh and Arockiamary [15], for categorical product of two cycles in Ahmad et al. [16].

In the current paper, the definitions of the heptagonal snake graph  $HPS_n$ , the double heptagonal snake graph  $D(HPS_n)$  and an  $l$ -multiple heptagonal snake graph  $L(HPS_n)$  have been introduced. Moreover, the exact value of TEISs for a heptagonal snake graph, a double heptagonal snake graph and an  $l$ -multiple heptagonal snake graph has been investigated.

## 2. Main results:

In this section, we introduce the definition of the heptagonal snake graph  $HPS_n$  and also, we deduce the exact value of TEIS for it.

### 2.1 Computing TEIS of heptagonal snake graph $HPS_n$ :

**Definition1.** A heptagonal snake graph  $HPS_n$  is a graph with  $6n + 1$  vertices and  $7n$  edges obtained by interchanging every edge of a path  $P_n$  by a cycle graph  $C_7$ , as shown in Figure (1).




Figure (1) A heptagonal snake graph  $HPS_n$

**Theorem1.** If  $HPS_n$  is a heptagonal snake graph with  $6n + 1$  vertices, then

$$tes(HPS_n) = \left\lceil \frac{7n+2}{3} \right\rceil.$$

**Proof:** Since  $HPS_n$  is a heptagonal snake graph with maximum degree  $\Delta(HPS_n) = 4$  and size  $|E(HPS_n)| = 7n$ , it thus (1) becomes

$$tes(HPS_n) \geq \left\lceil \frac{7n+2}{3} \right\rceil.$$

To prove the equality by showing the existence of an edge irregular total  $\mathbb{T}$ -labeling for  $HPS_n$  where  $\mathbb{T} = \left\lceil \frac{7n+2}{3} \right\rceil$ , we assume that a map  $\mathbb{T}: V(HPS_n) \cup E(HPS_n) \rightarrow \{1, 2, \dots, \mathbb{T}\}$  is a total  $\mathbb{T}$ -labeling and  $\mathbb{T} = \left\lceil \frac{7n+2}{3} \right\rceil$ . The total  $\mathbb{T}$ -labeling  $\mathbb{T}$  is defined for  $HPS_n$  for three cases as follows:

**Case 1:**  $\mathbb{T} \equiv 0 \pmod{3}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$

$\mathbb{T}$  is defined as:

$$\mathbb{T}(m_c) = \begin{cases} 1 & \text{for } c = 1 \\ 3c - 3 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{3} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n + 1 \end{cases},$$

$$\mathbb{T}(p_d^c) = \begin{cases} d & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(q_r^c) = \begin{cases} r + c - 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(m_c p_d^c) = \begin{cases} 1 & \text{for } c = 1 \\ c + 1 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} - 4 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(m_c q_r^c) = \begin{cases} 2 & \text{for } c = 1 \\ c + 2 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} - 3 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(p_{d+2}^c m_{c+1}) = \begin{cases} c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} + 2 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(q_{r+1}^c m_{c+1}) = \begin{cases} c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} + 1 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(p_d^c p_{d+1}^c) = \begin{cases} c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} - 2 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(p_{d+1}^c p_{d+2}^c) = \begin{cases} c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(q_r^c q_{r+1}^c) = \begin{cases} c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2\mathbb{H} - 1 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases}$$

Obviously, all edges and vertices labels are at most  $\mathbb{H} = \lceil \frac{7n+2}{3} \rceil$ . Now we introduce the edges' weights of  $HPS_n$  as follows:

$$w_{\mathbb{T}}(m_c p_d^c) = \begin{cases} 3 & \text{for } c = 1 \\ 4c + d - 2 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 4 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_r^c) = \begin{cases} 4 & \text{for } c = 1 \\ 5c + r - 2 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 3 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{d+2}^c m_{c+1}) = \begin{cases} 4c + d + 4 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c + 2 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(q_{r+1}^c m_{c+1}) = \begin{cases} 5c + r + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c + 1 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_d^c p_{d+1}^c) = \begin{cases} 2d + c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 2 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{d+1}^c p_{d+2}^c) = \begin{cases} 2d + c + 4 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(q_r^c q_{r+1}^c) = \begin{cases} 3c + 2r + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{3} \\ 7c - 1 & \text{for } \frac{\mathbb{H}}{3} + 1 \leq c \leq n \end{cases},$$

**Case 2:**  $\mathbb{H} \equiv 1 \pmod{3}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$

$\mathbb{T}$  is defined as in case 1 but with some modifications in conditions. We put  $\lceil \frac{\mathbb{H}}{3} \rceil$  instead of  $\frac{\mathbb{H}}{3}$  in the definition of  $m_c, m_c p_d^c$  and  $m_c q_r^c$ . In the others we put  $1 \leq c \leq \lceil \frac{\mathbb{H}}{3} \rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{3}$  and  $\lceil \frac{\mathbb{H}}{3} \rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{3} + 1 \leq c \leq n$ . Obviously, all edges and vertices labels are at most  $\mathbb{H} = \lceil \frac{7n+2}{3} \rceil$ . The edges' weights for  $m_c p_d^c$  and  $m_c q_r^c$  are given by:

$$w_{\mathbb{T}}(m_c p_d^c) = \begin{cases} 3 & \text{for } c = 1 \\ 4c + d - 2 & \text{for } 2 \leq c \leq \lceil \frac{\mathbb{H}}{3} \rceil - 1 \\ 4 \lceil \frac{\mathbb{H}}{3} \rceil + \mathbb{H} - 2 & \text{for } c = \lceil \frac{\mathbb{H}}{3} \rceil \\ 7c - 4 & \text{for } \lceil \frac{\mathbb{H}}{3} \rceil + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_r^c) = \begin{cases} 4 & \text{for } c = 1 \\ 5c + r - 2 & \text{for } 2 \leq c \leq \lceil \frac{\mathbb{H}}{3} \rceil - 1 \\ 4 \lceil \frac{\mathbb{H}}{3} \rceil + \mathbb{H} - 1 & \text{for } c = \lceil \frac{\mathbb{H}}{3} \rceil \\ 7c - 3 & \text{for } \lceil \frac{\mathbb{H}}{3} \rceil + 1 \leq c \leq n \end{cases},$$

For other edges the weights are like in case 1, but we put  $1 \leq c \leq \lceil \frac{\mathbb{H}}{3} \rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{3}$  and  $\lceil \frac{\mathbb{H}}{3} \rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{3} + 1 \leq c \leq n$ .

**Case 3:**  $\mathbb{H} \equiv 2 \pmod{3}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$

$\mathbb{T}$  is defined as:

$$\mathbb{T}(p_d^c) = \begin{cases} d & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{3} \right\rceil - 1 \\ d & \text{for } d, d+1 \\ \mathbb{T} & \text{for } d+2 \\ \mathbb{T} & \text{for } \left\lceil \frac{\mathbb{T}}{3} \right\rceil + 1 \leq c \leq n \end{cases} \quad c = \left\lceil \frac{\mathbb{T}}{3} \right\rceil .$$

For  $m_c, q_r^c, m_c p_d^c, m_c q_r^c, p_d^c p_{d+1}^c$  and  $q_r^c q_{r+1}^c$  we find that  $\mathbb{T}$  is defined as in case 1 but we put  $\left\lceil \frac{\mathbb{T}}{3} \right\rceil$  instead of  $\frac{\mathbb{T}}{3}$ , for the others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{T}}{3} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{T}}{3}$  and  $\left\lceil \frac{\mathbb{T}}{3} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{T}}{3} + 1 \leq c \leq n$ . Clearly, all edges and vertices labels are at most  $\mathbb{T} = \left\lceil \frac{7n+2}{3} \right\rceil$ . Also, the weights for all edges are given as in case 1 such that for  $m_c p_d^c, m_c q_r^c, p_d^c p_{d+1}^c$  and  $q_r^c q_{r+1}^c$  but with  $\left\lceil \frac{\mathbb{T}}{3} \right\rceil$  instead of  $\frac{\mathbb{T}}{3}$  and  $\left\lceil \frac{\mathbb{T}}{3} \right\rceil$ . For others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{T}}{3} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{T}}{3}$  and  $\left\lceil \frac{\mathbb{T}}{3} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{T}}{3} + 1 \leq c \leq n$ . Obviously, for all cases the edge's weights are different. Hence,  $\mathbb{T}$  is an edge-irregular total  $\mathbb{T}$ -labeling. Thus

$$tes(HPS_n) = \left\lceil \frac{7n+2}{3} \right\rceil.$$




Figure (2)  $tes(HPS_6) = 15$  for  $n = 6$

Figure (2) is an illustration of Theorem 1, where all edge vertex and vertex labels are at most  $\mathbb{T} = 15$  and any two different edges have distinct weights.

## 2.2 Computing TEIS of heptagonal snake graph $D(HPS_n)$ :

**Definition 2.** A double heptagonal snake graph  $D(HPS_n)$  consists of two heptagonal snake graphs that have a common path  $P_n$ , as shown in Figure (3).




Figure (3) A double heptagonal snake graph  $D(HPS_n)$

**Theorem 2.** Let  $D(HPS_n)$  be a double heptagonal snake graph on  $n+1$  vertices,  $n \geq 2$ . Then

$$tes(D(HPS_n)) = \left\lceil \frac{14n+2}{3} \right\rceil.$$

**Proof:** Since  $|E(D(HPS_n))| = 14n$  and the maximum degree of a double heptagonal snake graph is given by  $\Delta(D(HPS_n)) = 8$ , I then from (1) we have

$$tes(D(HPS_n)) \geq \left\lceil \frac{14n+2}{3} \right\rceil.$$

To verify equality by showing the existence of an edge irregular I total  $\mathbb{H}$ -labeling for  $D(HPS_n)$  with  $\mathbb{H} = \left\lceil \frac{14n+2}{3} \right\rceil$ , we assume that a map  $\mathbb{F}: V(D(HPS_n)) \cup E(D(HPS_n)) \rightarrow \{1, 2, \dots, \mathbb{H}\}$  is a I total  $\mathbb{H}$ -labeling and define the total  $\mathbb{H}$ -labeling  $\mathbb{F}$  for  $D(HPS_n)$  in the following cases as:

**Case 1:**  $\mathbb{H} \equiv 0 \pmod{6}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{F}$  is defined as:

$$\mathbb{F}(m_c) = \begin{cases} 1 & \text{for } c = 1 \\ 6c - 6 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n + 1 \end{cases},$$

$$\mathbb{F}(p_{1,d}^c) = \mathbb{F}(p_{2,d}^c) = \begin{cases} 2d - 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{1,r}^c) = \mathbb{F}(q_{2,r}^c) = \begin{cases} 2r + 2c - 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ \mathbb{H} & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(m_c p_{1,d}^c) = \begin{cases} 1 & \text{for } c = 1 \\ 2c & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 11 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(m_c p_{2,d}^c) = \begin{cases} 2 & \text{for } c = 1 \\ 2c + 1 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 10 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(m_c q_{1,r}^c) = \begin{cases} 2 & \text{for } c = 1 \\ 2c + 1 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 9 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(m_c q_{2,r}^c) = \begin{cases} 3 & \text{for } c = 1 \\ 2c + 2 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 8 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{1,d+2}^c m_{c+1}) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} + 1 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{2,d+2}^c m_{c+1}) = \begin{cases} 2c + 3 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} + 2 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{1,r+1}^c m_{c+1}) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 1 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{2,r+1}^c m_{c+1}) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{1,d}^c p_{1,d+1}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 7 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{1,d+1}^c p_{1,d+2}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 3 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{2,d}^c p_{2,d+1}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 6 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(p_{2,d+1}^c p_{2,d+2}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 2 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{1,r}^c q_{1,r+1}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 5 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{2,r}^c q_{2,r+1}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2\mathbb{H} - 4 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

Obviously, all edges and vertices labels are at most  $\mathbb{H} = \left\lceil \frac{14n+2}{3} \right\rceil$ . Now we introduce the edges' weights of  $D(HPS_n)$  as follows:

$$w_{\mathbb{F}}(m_c p_{1,d}^c) = \begin{cases} 3 & \text{for } c = 1 \\ 8c + 2d - 7 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 11 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$



$$w_{\mathbb{T}}(m_c p_{2,d}^c) = \begin{cases} 4 & \text{for } c = 1 \\ 8c + 2d - 6 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 10 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_{1,r}^c) = \begin{cases} 5 & \text{for } c = 1 \\ 10c + 2r - 7 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 9 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_{2,r}^c) = \begin{cases} 6 & \text{for } c = 1 \\ 10c + 2r - 6 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 8 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{1,d+2}^c m_{c+1}) = \begin{cases} 8c + 2d + 5 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c + 1 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{2,d+2}^c m_{c+1}) = \begin{cases} 8c + 2d + 6 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c + 2 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(q_{1,r+1}^c m_{c+1}) = \begin{cases} 10c + 2r + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 1 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(q_{2,r+1}^c m_{c+1}) = \begin{cases} 10c + 2r + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{1,d}^c p_{1,d+1}^c) = \begin{cases} 4d + 2c + 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 7 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{1,d+1}^c p_{1,d+2}^c) = \begin{cases} 4d + 2c + 5 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 3 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{2,d}^c p_{2,d+1}^c) = \begin{cases} 4d + 2c + 2 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 6 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(p_{2,d+1}^c p_{2,d+2}^c) = \begin{cases} 4d + 2c + 6 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 2 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{F}}(q_{1,r}^c q_{1,r+1}^c) = \begin{cases} 6c + 4r - 1 & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 5 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{F}}(q_{2,r}^c q_{2,r+1}^c) = \begin{cases} 6c + 4r & \text{for } 1 \leq c \leq \frac{\mathbb{H}}{6} \\ 14c - 4 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases}.$$

**Case 2:**  $\mathbb{H} \equiv 1 \pmod{6}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{F}$  is defined as:

$$\mathbb{F}(m_c q_{1,r}^c) = \begin{cases} 2 & \text{for } c = 1 \\ 2c + 1 & \text{for } 2 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 2 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2\mathbb{H} - 9 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(m_c q_{1,r}^c) = \begin{cases} 3 & \text{for } c = 1 \\ 2c + 2 & \text{for } 2 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 3 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2\mathbb{H} - 8 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases}.$$

For  $m_c, m_c p_{1,d}^c$  and  $m_c p_{2,d}^c$  we find that  $\mathbb{F}$  is defined as in case 1 but with  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil$  instead of  $\frac{\mathbb{H}}{6}$ . For others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ . Clearly, all edges and vertices labels are at most  $\mathbb{H} = \left\lceil \frac{14n+2}{3} \right\rceil$ . The edges weights of  $D(HPS_n)$  are given as:

$$w_{\mathbb{F}}(m_c p_{1,d}^c) = \begin{cases} 3 & \text{for } c = 1 \\ 8c + 2d - 7 & \text{for } 2 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 8 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + \mathbb{H} - 6 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 11 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{F}}(m_c p_{2,d}^c) = \begin{cases} 4 & \text{for } c = 1 \\ 8c + 2d - 6 & \text{for } 2 \leq c \leq \frac{\mathbb{H}}{6} \\ 8 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + \mathbb{H} - 5 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 10 & \text{for } \frac{\mathbb{H}}{6} + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_{1,r}^c) = \begin{cases} 5 & \text{for } c = 1 \\ 10C + 2r - 7 & \text{for } 2 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 8 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + \mathbb{T} - 4 & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c - 9 & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(m_c q_{2,r}^c) = \begin{cases} 6 & \text{for } c = 1 \\ 10C + 2r - 6 & \text{for } 2 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 8 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + \mathbb{T} - 3 & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c - 8 & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases}.$$

For others the edges' weights are given as in case 1 but with  $1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{T}}{6}$  and  $\left\lceil \frac{\mathbb{T}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{T}}{6} + 1 \leq c \leq n$ .

**Case 3:**  $\mathbb{T} \equiv 2 \pmod{6}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{T}$  is defined as:

$$\mathbb{T}(p_{1,d}^c) = \mathbb{T}(p_{2,d}^c) = \begin{cases} 2d - 1 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 2d - 1 & \text{for } d \\ \mathbb{T} & \text{for } d + 1, d + 2 \end{cases} \quad c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil,$$

$$\mathbb{T}(p_{1,d}^c p_{1,d+1}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 2 & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c - 2\mathbb{T} - 7 & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

$$\mathbb{T}(p_{2,d}^c p_{2,d+1}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 3 & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c - 2\mathbb{T} - 6 & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases}.$$

For  $m_c, m_c p_{1,d}^c, m_c p_{2,d}^c, m_c q_{1,r}^c$  and  $m_c q_{2,r}^c$  we find that  $\mathbb{T}$  is defined as in case 1 but with  $\left\lceil \frac{\mathbb{T}}{6} \right\rceil$  instead of  $\frac{\mathbb{T}}{6}$ . For others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{T}}{6}$  and  $\left\lceil \frac{\mathbb{T}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{T}}{6} + 1 \leq c \leq n$ .

Clearly, all edges and vertices labels are at most  $\mathbb{T} = \left\lfloor \frac{14n+2}{3} \right\rfloor$ . The edges' weights of  $D(HPS_n)$  are given as:

$$\begin{aligned}
w_{\mathbb{F}}(m_c p_{1,d}^c) &= \left\{ \begin{array}{ll} 3 & \text{for } c = 1 \\ 8c + 2d - 7 & \text{for } 2 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 8 \lceil \frac{\mathbb{H}}{6} \rceil + 2d - 7 & \text{for } d \\ 8 \lceil \frac{\mathbb{H}}{6} \rceil + \mathbb{H} - 6 & \text{for } d + 1, d + 2 \end{array} \right\} c = \lceil \frac{\mathbb{H}}{6} \rceil, \\
w_{\mathbb{F}}(m_c p_{2,d}^c) &= \left\{ \begin{array}{ll} 4 & \text{for } c = 1 \\ 8c + 2d - 6 & \text{for } 2 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 8 \lceil \frac{\mathbb{H}}{6} \rceil + 2d - 6 & \text{for } d \\ 8 \lceil \frac{\mathbb{H}}{6} \rceil + \mathbb{H} - 5 & \text{for } d + 1, d + 2 \end{array} \right\} c = \lceil \frac{\mathbb{H}}{6} \rceil, \\
w_{\mathbb{F}}(p_{1,d}^c p_{1,d+1}^c) &= \left\{ \begin{array}{ll} 4d + 2c + 1 & \text{for } 1 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 2 \lceil \frac{\mathbb{H}}{6} \rceil + \mathbb{H} + 2d + 1 & \text{for } c = \lceil \frac{\mathbb{H}}{6} \rceil \\ 14c - 7 & \text{for } \lceil \frac{\mathbb{H}}{6} \rceil + 1 \leq c \leq n \end{array} \right\}, \\
w_{\mathbb{F}}(p_{2,d}^c p_{2,d+1}^c) &= \left\{ \begin{array}{ll} 4d + 2c + 2 & \text{for } 1 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 2 \lceil \frac{\mathbb{H}}{6} \rceil + \mathbb{H} + 2d + 2 & \text{for } c = \lceil \frac{\mathbb{H}}{6} \rceil \\ 14c - 6 & \text{for } \lceil \frac{\mathbb{H}}{6} \rceil + 1 \leq c \leq n \end{array} \right\}.
\end{aligned}$$

For  $m_c q_{1,r}^c$  and  $m_c q_{2,r}^c$  the edges' weights are given as in case 1 but with  $\lceil \frac{\mathbb{H}}{6} \rceil$  instead of  $\frac{\mathbb{H}}{6}$ . For others we put  $1 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\lceil \frac{\mathbb{H}}{6} \rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ .

**Case 4:**  $\mathbb{H} \equiv 3 \pmod{6}$ ,  $1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{F}$  is defined as:

$$\begin{aligned}
\mathbb{F}(p_{1,d}^c) = \mathbb{F}(p_{2,d}^c) &= \left\{ \begin{array}{ll} 2d - 1 & \text{for } 1 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 2d - 1 & \text{for } d \\ \mathbb{H} & \text{for } d + 1, d + 2 \end{array} \right\} c = \lceil \frac{\mathbb{H}}{6} \rceil, \\
&\quad \left\{ \begin{array}{l} \mathbb{H} \\ \mathbb{H} \end{array} \right\} \text{for } \lceil \frac{\mathbb{H}}{6} \rceil + 1 \leq c \leq n \\
\mathbb{F}(q_{1,r}^c) = \mathbb{F}(q_{2,r}^c) &= \left\{ \begin{array}{ll} 2r + 2d - 2 & \text{for } 1 \leq c \leq \lceil \frac{\mathbb{H}}{6} \rceil - 1 \\ 2r + 2d - 2 & \text{for } r \\ \mathbb{H} & \text{for } r + 1 \end{array} \right\} c = \lceil \frac{\mathbb{H}}{6} \rceil, \\
&\quad \left\{ \begin{array}{l} \mathbb{H} \\ \mathbb{H} \end{array} \right\} \text{for } \lceil \frac{\mathbb{H}}{6} \rceil + 1 \leq c \leq n
\end{aligned}$$

$$\mathbb{F}(q_{1,r}^c q_{1,r+1}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 2 \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 2 & \text{for } c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \\ 14c - 2\mathbb{H} - 5 & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 1 \leq c \leq n \end{cases},$$

$$\mathbb{F}(q_{2,r}^c q_{2,r+1}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 2 \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 3 & \text{for } c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \\ 14c - 2\mathbb{H} - 4 & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 1 \leq c \leq n \end{cases}.$$

For  $m_c, m_c p_{1,d}^c, m_c p_{2,d}^c, m_c q_{1,r}^c, m_c q_{2,r}^c, p_{1,d}^c p_{1,d+1}^c$  and  $p_{2,d}^c p_{2,d+1}^c$  we find that  $\mathbb{F}$  is defined as in case 1 but we put  $\left\lfloor \frac{\mathbb{H}}{6} \right\rfloor$  instead of  $\frac{\mathbb{H}}{6}$ . For the others we put  $1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ . Clearly, all edges and vertices labels are at most  $\mathbb{H} = \left\lfloor \frac{14n+2}{3} \right\rfloor$ . The edges' weights of  $D(HPS_n)$  are given as:

$$w_{\mathbb{F}}(q_{1,r}^c q_{1,r+1}^c) = \begin{cases} 6c + 4r - 1 & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 4 \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + \mathbb{H} + 2r & \text{for } c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \\ 14c - 5 & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{F}}(q_{2,r}^c q_{2,r+1}^c) = \begin{cases} 6c + 4r & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 4 \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + \mathbb{H} + 2r + 1 & \text{for } c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \\ 14c - 4 & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 1 \leq c \leq n \end{cases}.$$

For  $m_c p_{1,d}^c, m_c p_{2,d}^c, m_c q_{1,r}^c, m_c q_{2,r}^c, p_{1,d}^c p_{1,d+1}^c$  and  $p_{2,d}^c p_{2,d+1}^c$  the edges' weights are given as in case 1 but with  $\left\lfloor \frac{\mathbb{H}}{6} \right\rfloor$  instead of  $\frac{\mathbb{H}}{6}$ . For others we put  $1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ .

**Case 5:**  $\mathbb{H} \equiv 4 \pmod{6}, 1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{F}$  is defined as:

$$\mathbb{F}(p_{1,d}^c) = \mathbb{F}(p_{2,d}^c) = \begin{cases} 2d - 1 & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 2d - 1 & \text{for } d \\ \mathbb{H} & \text{for } d + 1, d + 2 \\ \mathbb{H} & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 1 \leq c \leq n \end{cases} \quad c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor,$$

$$\mathbb{F}(p_{1,d+1}^c p_{1,d+2}^c) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor - 1 \\ 2 \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor + 2 & \text{for } c = \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \\ 14c - 2\mathbb{H} - 3 & \text{for } \left\lfloor \frac{\mathbb{H}}{6} \right\rfloor \leq c \leq n \end{cases},$$

$$\mathbb{T}(p_{2,d+1}^c p_{2,d+2}^c) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 3 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2\mathbb{H} - 2 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil \leq c \leq n \end{cases} .$$

For  $m_c, q_{1,r}^c, q_{2,r}^c, m_c p_{1,d}^c, m_c p_{2,d}^c, m_c q_{1,r}^c, m_c q_{2,r}^c, p_{1,d}^c p_{1,d+1}^c, p_{2,d}^c p_{2,d+1}^c, q_{1,r}^c q_{1,r+1}^c$  and  $q_{2,r}^c q_{2,r+1}^c$  we find that  $\mathbb{T}$  is defined as in case 1 but we put  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil$  instead of  $\frac{\mathbb{H}}{6}$ . For the others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ . Clearly, all edges and vertices labels are at most  $\mathbb{H} = \left\lceil \frac{14n+2}{3} \right\rceil$ . The edges weights of  $D(HPS_n)$  are given as:

$$w_{\mathbb{T}}(p_{1,d+1}^c p_{1,d+2}^c) = \begin{cases} 4d + 2c + 5 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + \mathbb{H} + 2d + 3 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 3 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases} ,$$

$$w_{\mathbb{T}}(p_{2,d+1}^c p_{2,d+2}^c) = \begin{cases} 4d + 2c + 6 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + \mathbb{H} + 2d + 4 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases} .$$

For  $m_c p_{1,d}^c, m_c p_{2,d}^c, m_c q_{1,r}^c, m_c q_{2,r}^c, p_{1,d}^c p_{1,d+1}^c, p_{2,d}^c p_{2,d+1}^c, q_{1,r}^c q_{1,r+1}^c$  and  $q_{2,r}^c q_{2,r+1}^c$  the edges' weights are given as in case 1 but we put  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil$  instead of  $\frac{\mathbb{H}}{6}$ . For the others we put  $1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ .

**Case 6:**  $\mathbb{H} \equiv 5 \pmod{6}, 1 \leq d \leq 3n$  and  $1 \leq r \leq 2n$ .

$\mathbb{T}$  is defined as:

$$\mathbb{T}(q_{1,r+1}^c m_{c+1}) = \begin{cases} 2c + 1 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 2 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2\mathbb{H} - 1 & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases} ,$$

$$\mathbb{T}(q_{2,r+1}^c m_{c+1}) = \begin{cases} 2c + 2 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1 \\ 2 \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 3 & \text{for } c = \left\lceil \frac{\mathbb{H}}{6} \right\rceil \\ 14c - 2\mathbb{H} & \text{for } \left\lceil \frac{\mathbb{H}}{6} \right\rceil + 1 \leq c \leq n \end{cases} ,$$

For  $p_{1,d+2}^c m_{c+1}$  and  $p_{2,d+2}^c m_{c+1}$  we find that  $\mathbb{T}$  is defined as in case 1 but we put  $1 \leq c \leq \left\lceil \frac{\mathbb{H}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{H}}{6}$  and  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{H}}{6} + 1 \leq c \leq n$ . For the others we put  $\left\lceil \frac{\mathbb{H}}{6} \right\rceil$  instead of  $\frac{\mathbb{H}}{6}$ . Clearly, all edges and vertices labels are at most  $\mathbb{H} = \left\lceil \frac{14n+2}{3} \right\rceil$ . The edges' weights of  $D(HPS_n)$  are given as:

$$w_{\mathbb{T}}(q_{1,r+1}^c m_{c+1}) = \begin{cases} 10c + 2r + 1 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 4 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + \mathbb{T} + 2r & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c - 1 & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

$$w_{\mathbb{T}}(q_{2,r+1}^c m_{c+1}) = \begin{cases} 10c + 2r + 2 & \text{for } 1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1 \\ 4 \left\lceil \frac{\mathbb{T}}{6} \right\rceil + \mathbb{T} + 2r + 1 & \text{for } c = \left\lceil \frac{\mathbb{T}}{6} \right\rceil \\ 14c & \text{for } \left\lceil \frac{\mathbb{T}}{6} \right\rceil + 1 \leq c \leq n \end{cases},$$

For  $p_{1,d+2}^c m_{c+1}$  and  $p_{2,d+2}^c m_{c+1}$  the edges' weights are given as in case 1 but we put  $1 \leq c \leq \left\lceil \frac{\mathbb{T}}{6} \right\rceil - 1$  instead of  $1 \leq c \leq \frac{\mathbb{T}}{6}$  and  $\left\lceil \frac{\mathbb{T}}{6} \right\rceil \leq c \leq n$  instead of  $\frac{\mathbb{T}}{6} + 1 \leq c \leq n$ . For the others we put  $\left\lceil \frac{\mathbb{T}}{6} \right\rceil$  instead of  $\frac{\mathbb{T}}{6}$ . clearly, for all cases any pairs of edges the weights are different. Hence,  $\mathbb{T}$  is I an edge I irregular itotal  $\mathbb{T}$  -labeling. Thus

$$tes(D(HPS_n)) = \left\lceil \frac{14n+2}{3} \right\rceil.$$




Figure (4)  $tes(D(HPS_n)) = 34$ , for  $n = 7$

Figure (4) is an illustration of Theorem 2, where all edge vertex and vertex labels are at most  $\mathbb{T} = 34$  and any two different edges have distinct weights.

### 2.3 Computing TEIS of heptagonal snake graph $L(HPS_n)$ :

**Definition3.** An  $l$ -multiple heptagonal snake graph  $L(HPS_n)$  is a graph consists of  $l$  heptagonal snake graphs that have a common path  $P_n$ .

**Therom3.** If  $L(HPS_n)$  is an  $l$  -multiple heptagonal snake graph. Then

$$tes(L(HPS_n)) = \left\lceil \frac{7ln + 2}{3} \right\rceil.$$

### Conclusion

In this paper, the definitions of the heptagonal snake graph  $HPS_n$ , the double heptagonal snake graph  $D(HPS_n)$  and an  $l$  -multiple heptagonal snake graph  $L(HPS_n)$  have been introduced. The exact values of TEISs for a heptagonal snake graph, a double heptagonal snake graph and an  $l$  -multiple heptagonal snake graph have also been investigated and given in the forms

$$tes(HPS_n) = \left\lceil \frac{7n+2}{3} \right\rceil,$$

$$tes(D(HPS_n)) = \left\lceil \frac{14n+2}{3} \right\rceil,$$

$$tes(L(HPS_n)) = \left\lceil \frac{7ln+2}{3} \right\rceil.$$

**Ethics approval and consent to participate:** This article does not contain any studies with human participants or animals performed by any of the authors.

**Conflict of interest:** The author declares that there is no conflict of interest for the paper.

**Authors' contributions:** The author contributed equally to this work. The author read and approved the final manuscript.

**Funding:** Not applicable.

### References

1. **Bača, M.; Jendroí, S.; Miller, M.; Ryan, J.** On irregular total labellings, *Discrete Math.* 2007, 307 (11), 1378–1388.
2. **Ivančo, J.; Jendroí, S.** Total edge irregularity strength of trees, *Discussiones Math. Graph Theory*, 2006, 26, 449–456.
3. **Putra, R.W.; Susanti, Y.** On total edge irregularity strength of centralized uniform theta Graphs, *AKCE Inter. J. Graphs and Comb.*, 2018, 15, 7–13.
4. **Salama, F.** On the total irregularity strength of polar grid graph, *J. Taibah Univ. Sci.*, 2019, 13, 912–916.
5. **Al-Mushayt, O.; Ahmad, A.; Siddiqui, M.K.** On the total edge irregularity strength of hexagonal grid graphs. *Australas. J. Comb.* 2012, 53, 263–271.
6. **Jeyanthi, P.** Total Edge Irregularity Strength of Disjoint Union of Wheel Graphs, *Electronic Notes in Discrete Mathematics*, 2015, 48, 175–182.
7. **Ahmad A, Bača M, Bashir Y, et al.** Total edge irregularity strength of strong product of two paths. *Ars Combin.* 2012;106:449–459.
8. **Siddiqui MK.** On edge irregularity strength of subdivision of star  $S_n$ . *Int J Math Soft Comput.* 2012;2:75–82.
9. **Bača, M.; Siddiqui, M. K.** Total edge irregularity strength of generalized prism, *Applied Math. Comput.*, 2014, 235, 168–173.
10. **Ahmad A, Arshad M, Ižaríková G.** Irregular labelings of helm and sun graphs. *AKCE Int J Graphs Combin.* 2015;12:161–168.
11. **Salman, A.N.M., Baskoro, E.T.** The total edge-irregular strengths of the corona product of paths with some graphs. *J Comb Math Comb Comput*, 2008, 65, 163–175.
12. **Tilukay, M. I.; Salman, A.N.M.; Persulesy, E. R.** On the Total Irregularity Strength of Fan, Wheel, Triangular Book, and Friendship Graphs, *Procedia Computer Science*, 2015, 74, 124 – 131.
13. **Ahmad, A.; Bača, M.** Total edge irregularity strength of a categorical product of two paths, *Ars Combin.* 2014, 114, 203–212.
14. **Hinding, N.; Suardi, N.; Basir, H.** Total edge irregularity strength of subdivision of star, *J. Discrete Math. Sci. Cryptogr.*, 2015, 18 (6), 869–875.
15. **Rajasinh, I.; Arockiamary, S.T.** Total edge irregularity strength of series parallel graphs, *Int. J. Pure Appl. Math.*, 2015, 99 (1), 11–21.



16. **Ahmad, A.; Bača, M.; Siddiqui, M.K.** On edge irregular total labeling of categorical product of two cycles. *Theory Comput. Syst.* , 2014, 54(1), 1–12 .

# Figures




Figure 1

A heptagonal snake graph  $\square\square\square\square$




Figure 2

An illustration of Theorem 1, where all edge vertex and vertex labels are at most  $\bar{\Gamma}=15$  and any two different edges have distinct weights.




Figure 3

A double heptagonal snake graph  $\square(\square\square\square)$




Figure 4

$$\chi(\Gamma(\chi)) = 34 \text{ for } \chi = 7$$