

Cepharanthine and Curcumin inhibited mitochondrial apoptosis induced by PCV2

Yinlan Xu

Shanxi Agricultural University

Jiangang Zheng

Shanxi Agricultural University

Panpan Sun

Shanxi Agricultural University

Jianhua Guo

Texas A&M University College Station

Xiaozhong Zheng

Queen's Medical Research Institute

Yaogui Sun

Shanxi Agricultural University

Kuohai Fan

Shanxi Agricultural University

Wei Yin

Shanxi Agricultural University

Hongquan Li (✉ livets@163.com)

Shanxi Agricultural University

Na Sun (✉ snzh060511@126.com)

Shanxi Agricultural University

Research article

Keywords: PCV2, Cap, Paeonol, Cepharanthine, Curcumin, apoptosis

Posted Date: May 19th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-26158/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 18th, 2020. See the published version at <https://doi.org/10.1186/s12917-020-02568-0>.

Abstract

Background

Porcine circovirus type 2 (PCV2) is an immunosuppressive pathogen with high prevalence rate in pig farms. It has caused serious economic losses to the global pig industry. Due to the rapid variation of PCV2 strain, short vaccine protection period and poor cross-protection effect, etc. It is necessary to screen and develop effective new compounds and explore their anti-PCV2 mechanism. The 13 natural compounds were purchased, with a clear plant origin, chemical structure and content and specific biological activities.

Methods

The maximum no-cytotoxic concentration (MNTC) and 50% cytotoxic concentration (CC₅₀) for 13 tested compounds were determined by the cytopathologic effect (CPE) assay and (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) method in PK-15 cells, respectively, qPCR and Western blot were used to detect the anti-PCV2 effect of compounds. Western blot method, AnnexinV-FITC /PI, JC-1, and ROS Kits were respectively used to detect the anti-PCV2 induced apoptosis mechanism of Cepharanthine and Curcumin. In this study, Ribavirin was used as the positive control.

Results

MNTC and CC₅₀ of 13 compounds were obtained. Compared with the PCV2 infected group, the expression of Cap treated with Paeonol (0.4 mg/mL and 0.2 mg/mL), Cepharanthine (0.003 mg/mL, 0.0015 mg/mL and 0.00075 mg/mL) and Curcumin (0.02 mg/mL, 0.001 mg/mL and 0.005 mg/mL) were significantly lower in a dose-dependent manner. The expression of Cleaved-caspase3 and Bax were up-regulated in the treated with Cepharanthine or Curcumin, while the expression of Bcl-2 was down-regulated, ROS value and MMP value were decreased to different degrees and the apoptosis rate was reduced.

Conclusion

Paeonol, Cepharanthine and Curcumin have significant antiviral effect. And the PCV2-induced Mitochondrial apoptosis was mainly remitted by Cepharanthine and Curcumin.

Background

PCV2 is a DNA virus, which is an important pathogen in pig farms and highly infectious [1]. The diseases caused by PCV2 mainly include postweaning multisystemic wasting syndrome (PMWS), Porcine

dermatitis and nephropathy syndrome (PDNS), Porcine proliferative and necrotizing pneumonia (PNP), etc., which have caused serious economic losses to the global pig industry for 57 years [2]. The capsid protein Cap is a major viral structural protein and a major immunogen involving viral replication of PCV2 [3]. PCV2 was shown to induce apoptosis of spleen and liver cells in vivo [4, 5] and also induce apoptosis in vitro by mitochondrial apoptosis, ER (endoplasmic reticulum) apoptosis and ROS regulation [6–9].

It has been reported that PCV2 vaccine has a short protection period and poor efficacy for protecting pigs from PCV2 infection [10, 11]. Therefore, it is needed to control infection. Chinese herbal medicine has the characteristics of high efficiency, low toxicity and low residue. It can improve the immunity, possess anti-stress and anti-virus ability. A variety of natural compounds or their active ingredients have been used to enhance host immune, prevent and treat viral diseases, such as Coronavirus diseases [12], Hepatitis B (HB) [13], Infectious bronchitis [14], African swine fever (ASF) [15], Influenza [16] and other viral diseases. However, the research on the anti-PCV2 action of natural compounds and its mechanism has also become a hotspot.

In our previous studies, dozens of natural compounds were screened for their antiviral activity, such as Chlorogenic Acid [17], ligustrazine hydrochloride [18], Scutellarin [17, 19], dipotassium glycyrrhizinate [18, 20, 21], Sodium tanshinone IIA sulfonate [21, 22], tea seed saponins [23] and Matrine [19, 24, 25]. Among them, only Scutellarin or Matrine had anti-PCV2 effect [19, 25].

Therefore, in this study, from many natural compounds with clear chemical structure, traditional medicine plant source and specific biological activity, 13 compounds with antiviral potential were selected and tested in the model of PCV2 infected PK-15 cell to screen anti-PCV2 compounds in vitro. Furthermore, the antiviral mechanism of compounds was explored and explained from the perspectives of apoptosis, MMP and ROS. It provides a theoretical basis for the development of a safely, efficiently new compound with anti-PCV2 effect.

Results

Cytotoxicity of 13 compounds on the PK-15 cells detected by MTT

Compared with the normal group of normal cells, the results showed that (See Additional File. 1), the degree of cytopathological change were also different after the treatment of different compounds and different concentrations of the same compound. Due to the low solubility, the maximum solubility concentration (MSC) of 0.8mg/mL Formononetin, 0.8mg/mL Icarine and 0.15mg/mL Astragaloside were used to treat the cells, respectively and no change in cell morphology was observed. The MNTC is the same as MSC. The 10 compounds except Formononetin, Icarine and Astragaloside showed different cytopathic changes (see Additional File. 1) and MNTC values (Table 2). Dose-response curves of the tested compounds (Fig.2) and CC_{50} value (Table 2) were generated by GraphPad Prism 5.0 analysis after calculating the CR value. The curves were “S” type, showing a dose-dependent relationship. As the concentration of the compound increased, more cells with change in morphology the was observed.

The anti-PCV2 effect of Paeonol, Cepharanthine and Curcumin

To explore the anti-PCV2 effect of compounds, the expression levels of Cap was detected by qPCR and Western blot. The results of qPCR showed that, compared to cell control group, the PCV2 copy numbers were significantly increased ($P<0.05$); compared to PCV2 infected group, the PCV2 copy numbers were significantly decreased in these compound treated with Formononetin at 0.8, Icariine at 0.8, Paeonol at 0.4, 0.2 and 0.1, Cepharanthine at 0.003, 0.0015 and 0.00075, Curcumin at 0.02, 0.01 and 0.005 mg/mL, Curcumol at 0.024 and Syringin at 0.625 and 0.3125 mg/mL ($P<0.05$) (Fig.3A).

Based on the results by qPCR, dose-dependent Paeonol, Cepharanthine and Curcumin were selected for Western blot. The results showed that the expression level of Cap protein treated with the three compounds respectively were significantly lower than PCV2 infected group ($P<0.05$), except the 0.1 mg/mL of Paeonol group (Fig.3 B) (The original and unedited blots see Additional File. 2). Therefore, the Cepharanthine and Curcumin were selected for the subsequent anti-apoptotic test.

Inhibition of PCV2-induced apoptosis by Cepharanthine and Curcumin

To evaluate the anti-apoptosis effect of Cepharanthine or Curcumin, Samples were analyzed after being treated with Annexin V/PI by Flow cytometer. The results showed that Compared to the PCV2-infected group, the cell apoptosis rates were significantly decreased in the group treated with Cepharanthine, Curcumin or Ribavirin, demonstrating a dose-dependent response except the group of 0.005 mg/mL Paeonol ($P<0.05$) (Fig.4A and B). Our results indicated that Cepharanthine, Curcumin or Ribavirin could significantly inhibit PCV2-induced apoptosis ($P<0.05$).

Inhibition of PCV2-induced apoptosis by Cepharanthine and Curcumin via the mitochondria pathway

To evaluate the anti-apoptosis mechanism of Cepharanthine or Curcumin, Samples were analyzed after being treated with JC-1 kit by Flow cytometer. The results showed that Compared to the PCV2-infected group, the MMP value were significantly decreased in the group treated with Cepharanthine, Curcumin and Ribavirin, demonstrating a dose-dependent response ($P<0.05$) (Fig.5A and B). The results indicated that Cepharanthine, Curcumin or Ribavirin could significantly inhibit PCV2-induced apoptosis via mitochondrial pathway ($P<0.05$).

The mechanism of inhibition of PCV2-induced mitochondrial apoptosis by Cepharanthine and Curcumin

To evaluate the mechanism of inhibition of PCV2-induced mitochondrial apoptosis by Cepharanthine or Curcumin, Samples were analyzed by Western blot and Flow cytometer. The results of Western blot showed that Compared with the PCV2-infected group, the expression of pro-apoptin Cleaved-caspase 3 and Bax was significantly decreased while the expression of anti-apoptotic protein Bcl-2 was significantly increased in the treatment group of Cepharanthine, Curcumin or Ribavirin (Fig.6A-H) (The original and unedited blots see Additional File. 3). The results of Flow cytometer after being stained with ROS kit demonstrated that the value of ROS in the Curcumin-treated group showed strongly downward trend in a dose-dependent manner, compared with the PCV2 control group (Fig.6I and J). However, only the group of

treated with 0.003mg/mL of Cepharanthine could significantly reduce the amount of ROS($P<0.05$). The results demonstrated that Cepharanthine or Curcumin inhibited the PCV2-induced increase in MMP through the caspase family and Bcl-2 family mechanisms. Curcumin inhibited PCV2-induced mitochondrial apoptosis by reducing ROS, However, ROS reduction is not the main pathway for Cepharanthine and Ribavirin to inhibit PCV2-induced mitochondrial apoptosis.

Discussion

PCV2 is the main pathogen causing porcine circovirus associated diseases (PCVAD). In this study, 13 natural compounds with defined chemical structures, useful clinical effect and from medicinal plants were selected. After the primary screening, three compounds with antiviral effect, Paeonol, Cepharanthine and Curcumin, were further studied. The antiviral mechanism of these two compounds was investigated by evaluating the level of key apoptins, MMP and ROS in the cellular mitochondrial pathway. Ribavirin has a broad antiviral activity and is used as the positive control.

Cepharanthine is a natural alkaloid extracted from the roots of the *Stephania japonica*, which has anti-inflammatory [26], anti-Hepatitis B virus [27], herpes simplex type-1 virus (HSV-1) [28] and etc. Curcumin is an active ingredient extracted from the rhizome of some plants of the Zingiberaceae and Araceae. Medical research shows that curcumin has low toxicity, little adverse reactions and multiple therapeutic effects, such as anti-inflammatory [29], anti-oxidative [30], anti-H9N2 influenza virus [31], dengue virus [32], hepatitis C virus [33] and so on.

Lin et al. proved that the PCV2 could up-regulate MMP and induce apoptosis through mitochondrial pathway [34]; Chen et al. proved the accumulation of ROS and interference with mitochondrial function were demonstrated in RAW264.7 cells infected with PCV2 [35, 36]. Mitochondria have a central role in cell apoptosis. The loss of MMP is the marker of mitochondrial apoptosis, the changes of MMP are caused by Bcl-2 family, Caspase family and Mitochondrial permeability transition pore (MPTP) [37]. Cleaved-caspase 3 is the main executive protein of apoptosis. There are many members of the Bcl-2 family, among which Bax is the main pro-apoptotic protein and Bcl-2 is the main anti-apoptotic protein. The opening and closing of MPTP is influenced by ROS, Ca^{2+} , ADP, oxidative stress, high PH value and other apoptotic factors [37]. In this study, at first, the cell apoptosis and change of MMP were detected by flow cytometry after Annexin V/PI double staining and JC-1 assay. The results indicated that Cepharanthine, Curcumin or Ribavirin could significantly inhibit PCV2-induced apoptosis via mitochondrial pathway. To further evaluate the mechanism of Cepharanthine or Curcumin inhibiting mitochondrial apoptosis, the apoptins of Caspase family and Bcl-2 family were analyzed by Western blot; the values of ROS were analyzed by flow cytometry after ROS kit staining. The results showed that Compared with cell control group, the expression levels of cleaved-caspase3 and Bax were up-regulated, and the expression of Bcl-2 was down-regulated, the ROS values are increased. This result is consistent with the previous reports [8, 34, 38]. Compared with PCV2-infected group, the expression levels of cleaved-caspase3 and Bax were down-regulated, and the expression of Bcl-2 was up-regulated in the treated with Cepharanthine or Curcumin. The value of ROS was decreased but ROS did not decrease in treated with the 0.0015 and

0.00075 mg/mL of Cepharanthine. The results demonstrated that Cepharanthine or Curcumin inhibited the PCV2-induced increase in MMP through the caspase family and Bcl-2 family mechanisms. Curcumin could inhibit cell mitochondrial apoptosis by regulating the opening of MPTP through reducing the level of ROS, but Cepharanthine and Ribavirin did not have the similar effect. It is speculated that various antiviral compounds could inhibit the cell apoptosis by governing MMP through affecting mitochondrial apoptosis pathway.

Conclusion

To summarize, our study revealed that Paeonol, Cepharanthine and Curcumin have significant antiviral effect and Cepharanthine and Curcumin could inhibit PCV2-induced mitochondrial apoptosis by through change the MMP (Fig. 7). The results showed that Cepharanthine and Curcumin have great potential to be developed a new anti-PCV2 drugs.

Methods

Cells, viruses, antibiotics and nature compounds

PCV-free pig kidney endothelial cell line (PK-15) was preserved in our laboratory. Cells were cultured and passed in Dulbecco's modified eagle's medium (DMEM, Hyclone, USA) containing 10% fetal calf serum (FBS, BI, Israel) (10% DMEM) and maintenance in DMEM containing 2% FBS (2% DMEM).

The strain of PCV2-SH was supplied by Professor Jiang Ping of Nanjing Agricultural University. Virus was replicated and harvest in PK-15 cells infected with PCV2. The titer of $10^{6.4}$ TCID₅₀/mL was determined by Indirect Immunofluorescence (IFA).

Antibodies against Cap were purchased from Biorbyt LLC. (California, Britain) Cleaved-caspase3, Bcl-2 and Bax were purchased from Abcam (Cambridge, USA); GAPDH and horseradish peroxidase (HRP)-conjugated secondary antibodies were purchased from Wuhan Sanying Biology Technology Co.,Ltd. Wu Han, China.

Ribavirin, the positive control drug, was purchased from Beijing Solarbio technology co., LTD; Syringin was purchased from Nanjing Puyi biotechnology co., LTD; Glycyrrhizin acid was purchased from DOSF biotechnology co., LTD; Other tested compounds, such as Curcumol, Paeonol, Oxymatrine, Caffeic acid, Formononetin, Cepharanthine, Apigenin, Psoralen, Lcariine, Curcumin and Astragaloside were purchased from National Institutes for Food and Drug Control. Compounds information and chemical structure were respectively listed Table 1 and Figure 1.

Cytotoxicity assay

The viability of PK-15 cell was assessed with MTT. 1.0×10^6 cells/mL of cells were seeded into 96-well plates and incubated until 100% confluency was reached. The tested compounds were added with two-

fold serial diluted and cultured for another 60 h and cell control group was established. Four repeated wells were set up for each concentration with 100 μ L per well. The cytopathic condition of cells was observed every day and photographed. The compounds and dissolution methods are shown in Table 1. Then 25 μ L MTT (4 mg/mL in PBS) was added to each well and incubated at 37 °C for 4 h. 150 μ L of DMSO (Solarbio, Beijing) was added and incubated for 30 min after removing the culture medium. The optical density (OD) were measured at 490 nm using a microplate spectrophotometer (Spectra Max M5, Molecular Devices, USA). Cytopathic ratio (CR) was calculated based on $CR = [(OD \text{ control} - OD \text{ test}) / OD] \times 100$. Then MNTC and CC_{50} on PK-15 cells were calculated using GraphPad PrismTM5.0 (Inc. California, USA).

Real time quantitative PCR

The anti-PCV2 effect of compounds in PK-15 cell was determined by real time quantitative PCR (qPCR). When cell confluency rate of 24-well plate reached 80%, $10^{4.4}$ TCID₅₀ of PCV2 was incubated for 2 h and discarded the supernatant. MNTC of every test compound was in two-fold serial dilution with 2% DMEM into 3 gradients and added to 24-well plates (500 μ L/well) (Table 2). Meanwhile, the cell control group, PCV2-infected group and Ribavirin positive control group were applied and incubated for 48 h with 5% CO₂ at 37 °C. DNA was extracted according to the instruction of DNA extraction kit (TianGen, Beijing, China) and the DNA concentration was measured using a nucleic acid concentration analyzer (NanoDrop Technologies, Wilmington, DE, USA). The copy number of the Cap gene was detected by qPCR. The Primers 5' TAC ATT TCC AGC AGT TTG and 5' CTC CCG CCA TAC CAT AA were used for the amplification of 148 bp fragment.

Western blot

The anti-PCV2 effect and anti-apoptotic mechanism of the compound was detected by Western blot. When cell confluency of 6-well plate reached 80%, $10^{4.4}$ TCID₅₀ of PCV2 was incubated for 2 h and discarded the supernatant. Paeonol, Cepharaanthine and Curcumin were added in turn, 2mL/well. The cell control group, PCV2-infected group and Ribavirin positive control group were applied and incubated for 48 h. Total cell protein was extracted and protein concentration was determined using the BCA protein assay kit (Beyotime Biotechnology, Jiangsu, China). The protein samples were separated by SDS-PAGE, then transferred to the PVDF membrane. The levels of PCV2 Cap protein and apoptin were detected using an eECL Western Blot detection kit (Cwbio Inc, Beijing, China) and chemiluminescence imaging system (BIO-RAD, California, USA).

Annexin V/PI staining for apoptosis

The anti-apoptotic effect of Cepharaanthine and Curcumin were detected by Annexin V/PI. Cells in 6-well plate were incubated with $10^{4.4}$ TCID₅₀ PCV2 for 2 h and discarded the supernatant, then Cepharaanthine and Curcumin were added, the cell control group, PCV2-infected group and Ribavirin positive control group were applied and incubated for 48 h. Samples were collected, centrifugated and treated using

Annexin V/PI kit (Keygen Biotech, Nanjing, China), then analyzed by flow cytometry (BD Biosciences, USA).

Detection of MMP by JC-1

Mitochondrial membrane potential (MMP) of cells was detected by JC-1. Cells in the 6-well plate were incubated with $10^{4.4}$ TCID₅₀ PCV2 for 2 h and discarded the supernatant, then Cepharaanthine and Curcumin were added. After 48 h incubation, cells were stained with JC-1 using a commercial kit (Beyotime Biotechnology, Jangsu, China) treated and analyzed by flow cytometry (BD Biosciences, USA).

Detection of ROS by flow cytometer assay

The amount of reactive oxygen species (ROS) was measured by flow cytometry. Cells in the 6-well plate were incubated with $10^{4.4}$ TCID₅₀ of PCV2 for 2 h and discarded the supernatant, Cepharaanthine and Curcumin were added and incubated for 48 h. Cells were treated with ROS assay kit and the changes of ROS were detected by Flow cytometer.

Statistical analysis

CC₅₀ was calculated using the software of GraphPad Prism TM5.0 (GraphPad Software, Inc. California, USA). The gray intensity of protein bolts was analyzed by Image J (National Institutes of Health, USA). Data are expressed as the mean \pm standard error of the mean (SEM). Different letters indicate statistically significant difference to other groups ($p \leq 0.05$).

Abbreviations

PCV2: Porcine circovirus type 2; PK-15: PCV-free pig kidney endothelial cell line; DMEM: Dulbecco's modified eagle's medium; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; CPE: cytopathologic effect; CR: Cytopathic ratio; MNTC: maximum no-cytotoxic concentration; MSC: the maximum solubility concentration; CC₅₀: 50% cytotoxic concentration; TCID₅₀: 50% tissue culture infective dose; RT-qPCR: real time quantitative PCR; ROS: reactive oxygen species; MMP: Mitochondrial membrane potential.

Declarations

Acknowledgements

We thank Prof. Jiang (Nanjing Agricultural University) for his assistance in the PCV2.

Authors' contributions

YX, JZ, NS and HL designed all the experiments. YX and JZ performed the MTT experiments, qPCR and Western blot assays. YS, KF, and WY performed the Flow cytometer assays. YX, PS, NS, XZ and JG wrote

the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by National key research and development program (Grant No. 2017YFD0501500). The funding body had a role in the design of the study, collection, analysis, and interpretation of data or in the writing of this manuscript.

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The experiment was performed in according to the experiment operational guideline of Shanxi province, China.

Consent for publication

Not applicable.

Competing interests

All authors declared no competing conflict of interest.

Author details

¹ College of Animal Science and Veterinary Medicine, Shanxi Agricultural University, Taigu 030801, Shanxi, China

² Laboratory Animal Center, Shanxi Agricultural University, Taigu 030801, Shanxi, China

³ Department of Veterinary Pathobiology, Schubot Exotic Bird Health Center, Texas A&M University, College Station, Texas, TX 77843, USA

⁴ Medical Research Council (MRC) Centre for Inflammation Research, Queen's Medical Research Institute, The University of Edinburgh, Edinburgh, EH16 4TJ, UK

References

1. Ellis J, Clark E, Haines D, West K, Krakowka S, Kennedy S, Allan GM. Porcine circovirus-2 and concurrent infections in the field. *Vet Microbiol.* 2004;98(2):159–63.
2. Meng XJ. Spread like a wildfire-The omnipresence of porcine circovirus type 2 (PCV2) and its ever-expanding association with diseases in pigs. *Virus Res.* 2012;164(1–2):1–3.

3. Nawagitgul P, Morozov I, Bolin SR, Harms FA, Sorden SD, Paul PS. Open reading frame 2 of porcine circovirus type 2 encodes a major capsid protein. *J Gen Virol.* 2000;81(9):2281–7.
4. Kiupel M, Stevenson GW, Galbreath EJ, North A, HogenEsch H, Mittal SK. Porcine circovirus type 2 (PCV2) causes apoptosis in experimentally inoculated BALB/c mice. *BMC Vet Res.* 2005;1(1):7–15.
5. Sinha A, Schalk S, Lager KM, Wang C, Opriessnig T. Singular PCV2a or PCV2b infection results in apoptosis of hepatocytes in clinically affected gnotobiotic pigs. *Res Vet Sci.* 2012;92(1):151–6.
6. Wei L, Zhu Z, Wang J, Liu J. Activation of the phosphatidylinositol 3-kinase/AKT signaling pathway during porcine circovirus type 2 infection facilitates cell survival and viral replication. *J Virol.* 2012;86(24):13589–97.
7. Wei L, Zhu Z, Wang J, Zhang C, Quan R, Yan X. Regulatory role of ASK1 in porcine circovirus type 2-induced apoptosis. *Virology.* 2013;447(1–2):285–91.
8. Zhou YS, Qi BZ, Gu BZ, Xu F, Du HH, Li XL, Fang WH. Porcine circovirus 2 deploys PERK pathway and GRP78 for its enhanced replication in PK-15 cells. *Viruses.* 2016;8(2):56–72.
9. Zhang Y, Sun R, Geng S, Shan Y, Li X. Porcine circovirus type 2 induces ORF3-independent mitochondrial apoptosis via perk activation and elevation of cytosolic calcium. *J Virol.* 2019;93(7):01784–800.
10. Beach NM, Meng XJ. Efficacy and future prospects of commercially available and experimental vaccines against porcine circovirus type 2 (PCV2). *Virus Res.* 2012;164(1–2):33–42.
11. Feng H, Blanco G, Segalés J, Sibila M. Can Porcine circovirus type 2 (PCV2) infection be eradicated by mass vaccination? *Vet Microbiol.* 2014;172(1–2):92–9.
12. Cinatl J, Morgenstern B, Bauer G, Chandra PP, Rabenau H, Doerr HW. Glycyrrhizin, an active component of liquorice roots, and replication of sars-associated coronavirus. *Lancet.* 2003;361(9374):2045–6.
13. Xu WS, Zhao KK, Miao XH, Ni W, Cai X, Zhang RQ, Wang JX. Effect of oxymatrine on the replication cycle of hepatitis B virus in vitro. *World J Gastroenterol.* 2010;16(16):2028–37.
14. Chen C, Zuckerman DM, Brantley S, Sharpe M, Childress K, Hoiczky E, Pendleton AR. Sambucus Nigra extracts inhibit infectious bronchitis virus at an early point during replication. *BMC Vet Res.* 2014;16(1):10–24.
15. Hakobyan A, Arabyan E, Avetisyan A, Abroyan L, Hakobyan L, Zakaryan H. Apigenin inhibits African swine fever virus infection in vitro. *Arch Virol.* 2016;61(12):3445–53.
16. Cheng W, Ren J, Huang Q, Long H, Jin H, Zhang L, Liu H, Van OL, Lin W. Pregnane steroids from a gorgonian coral subergorgia suberosa with anti-flu virus effects. *Steroids.* 2016;108:99–104.
17. Cheng J, Sun N, Zhao X, Niu L, Song M, Sun Y, Jiang J, Guo J, Bai Y, He J, Li H. In vitro screening for compounds derived from traditional Chinese medicines with antiviral activities against porcine reproductive and respiratory syndrome virus. *J Microbiol Biotechnol.* 2013;23(8):1076–83.
18. Sun Y, Song M, Niu L, Bai X, Li H. Antiviral effects of the constituents derived from Chinese herb medicines on infectious bursal disease virus. *Pharm Biol.* 2013;51(9):1137–43.

19. Sun N, Yu T, Zhao JX, Sun YG, Jiang JB, Duan ZB, Wang WK, Hu YL, Lei HM, Li HQ. Antiviral activities of natural compounds derived from traditional Chinese medicines against porcine circovirus type 2 (PCV2). *Biotechnol Bioproc E*. 2015;20(1):180–7.
20. Wang ZW, Sun N, Wu CH, Jiang JB, Bai YS, Li HQ. In vitro antiviral activity and underlying molecular mechanisms of dipotassium glycyrrhetate against porcine reproductive and respiratory syndrome virus. *Antivir Ther*. 2013;18(8):997–1004.
21. Sun Y, Niu L, Song M, Zhao X, Sun N, He JP, Wu CH, Jiang JB, Bai YS, Guo JH, Li HQ. Screening compounds of Chinese medicinal herbs anti-marek's disease virus. *Pharm Biol*. 2014;52(7):841–7.
22. Zhang JQ, Gu YL, Wu CH, Ma HL, He JP, Bai YS, Li HQ. Effects of sodium tanshinone IIA sulfonate against Marek's disease virus in experimentally infected chickens. *Int J Biol Macromol*. 2013;58:258–62.
23. Li E, Sun N, Zhao JX, Sun YG, Huang JG, Lei HM, Guo JH, Hu YL, Wang WK, Li HQ. In vitro evaluation of antiviral activity of tea seed saponins against porcine reproductive and respiratory syndrome virus. *Antivir Ther*. 2015;20(7):743–52.
24. Sun N, Wang ZW, Wu CH, Li E, He JP, Wang SY, Hu YL, Lei HM, Li HQ. Antiviral activity and underlying molecular mechanisms of Matrine against porcine reproductive and respiratory syndrome virus in vitro. *Res Vet Sci*. 2014;96(2):323–7.
25. Sun N, Sun PP, Lv HP, Sun YG, Guo JH, Wang ZR, Luo TT, Wang SY, Li HQ. Matrine displayed antiviral activity in porcine alveolar macrophages co-infected by porcine reproductive and respiratory syndrome virus and porcine circovirus type 2. *Sci Rep*. 2016;6(1):24401–14.
26. Furusawa S, Wu JH. The effects of biscoclaurine alkaloid cepharanthine on mammalian cells: Implications for cancer, shock, and inflammatory diseases. *Life Sci*. 2007;80(12):1073–9.
27. Zhou YB, Wang YF, Zhang Y, Zheng LY, Yang XA, Wang N, Jiang JH, Ma F, Yin DT, Sun CY, Wang QD. In vitro activity of cepharanthine hydrochloride against clinical wild-type and lamivudine-resistant hepatitis B virus isolates. *Eur J Pharmacol*. 2012;683(1–3):10–5.
28. Okamoto M, Okamoto T, Baba M. Inhibition of human immunodeficiency virus type 1 replication by combination of transcription inhibitor K-12 and other antiretroviral agents in acutely and chronically infected cells. *Antimicrob Agents Chemother*. 1999;43(3):492–7.
29. Suresh MV, Wagner MC, Rosania GR, Stringer KA, Reddy RC. Pulmonary administration of a water-soluble curcumin complex reduces severity of acute lung injury. *Am J Respir Cell Mol Biol*. 2012;47(3):280–7.
30. Dai J, Gu L, Su Y, Wang Q, Zhao Y, Chen XX, Deng HX, Li W, Wang GF, Li KS. Inhibition of curcumin on influenza A virus infection and influenzal pneumonia, via, oxidative stress, TLR2/4, p38/JNK MAPK and NF- κ B pathways. *Int Immunopharmacol*. 2018;54:177–87.
31. Umar S, Shah MA, Munir MT, Yaqoob M, Fiaz M, Anjum S, ouzouaia M, Younus M, Nisa Q, Iqbal M, Umar W. Synergistic effects of thymoquinone and curcumin on immune response and anti-viral activity against avian influenza virus (H9N2) in turkeys. *Poult Sci*. 2016;95(7):1513–20.

32. Padilla SL, Rodríguez A, Gonzales MM, Gallego GJ, Castano OJ. Inhibitory effects of curcumin on dengue virus type 2-infected cells in vitro. *Arch Virol.* 2014;59(3):573–9.
33. Kim KJ, Kim KH, Kim HY, Cho HK, Sakamoto N, Cheong JH. Curcumin inhibits hepatitis C virus replication via suppressing the AKT-SREBP-1 pathway. *Febs Lett.* 2010;584(4):707–12.
34. Lin C, Gu J, Wang H, Zhou J, Li J, Wang S, Jin Y, Liu C, Liu J, Yang H, Jiang P, Zhou J. Caspase-dependent apoptosis induction via viral protein ORF4 of porcine circovirus 2 binding to mitochondrial adenine nucleotide translocase 3. *J Virol.* 2018;92(10):e00238–256.
35. Chen HL, Yang J, Fu YF, Meng XN, Hu TJ. Effect of total flavonoids of *Spatholobus Suberectus* Dunn on PCV2 induced oxidative stress in Raw 264.7 cells. *BMC Complement Altern Med.* 2017;17(1):244–53.
36. Chen XX, Ren F, Hesketh J, Shi XL, Li JX, Gan F, Huang KH. Reactive oxygen species regulate the replication of porcine circovirus type 2 via NF- κ B pathway. *Virology.* 2012;426(1):66–72.
37. Adeyemi RO, Landry S, Davis ME, Weitzman MD, Pintel DJ. Parvovirus minute virus of mice induces a DNA damage response that facilitates viral replication. *Plos Pathog.* 2010;6(10):e1001141-1153.
38. Su ZJ, Wei YY, Yin D, Shuai XH, Zeng Y, Hu TJ. Effect of *Sophora Subprostrate* Polysaccharide on oxidative stress induced by PCV2 infection in RAW264.7 cells. *Int J Biol Macromol.* 2013;62:457–64.

Tables

Table 1 The compounds used in the test

No.	Chemical name	Classification	Main plant source	Cosolvent	Purity
1	Formononetin	Phenolic acids	Mango	1%DMSO	98.10%
2	Icariine	Flavonoid glycosides	Herba epimedii	1%DMSO	94.20%
3	Astragaloside	Saponins	Astragalus membranaceus	1%DMSO	96.90%
4	Paeonol	Monoterpenes	Peony	1%DMSO	99.90%
5	Cepharanthine	Alkaloids	Stephania	1%DMSO	100.00%
6	Curcumin	Polyphenols	Turmeric	1%DMSO	98.70%
7	Curcumol	Hemiketal	Curcuma zedoary	1%DMSO	99.90%
8	Syringin	Phenylpropanolides	Lilac	DMEM	95.20%
9	Oxymatrine	Alkaloid	Sophora	DMEM	92.50%
10	Caffeic acid	Organic acid	Caffeic	1%DMSO	99.70%
11	Apigenin	Flavonoid	Apium graveolens	1%DMSO	99.20%
12	Glycyrrhizic acid	triterpenes	Glycyrrhiza	DMEM	95.00%
13	Psoralen	Furancoumarins	Fructus psoraleae	1%DMSO	99.70%

Table 2 The results of cytotoxicity

No.	Chemical name	MNTC[mg/mL]	CC ₅₀ [mg/mL]
1	Formononetin	0.8	0.8
2	Icariine	0.8	0.8
3	Astragaloside	0.15	0.15
4	Paeonol	0.4	1.2696±0.08983
5	Cepharanthine	0.003	0.008048±0.0006136
6	Curcumin	0.02	0.06296±0.02424
7	Curcumol	0.024	0.115±0.02331
8	Syringin	0.625	1.128±0.5339
9	Oxymatrine	2	8.31±0.3896
10	Caffeic acid	0.0125	0.2581±0.07863
11	Apigenin	0.01	0.04817 ±0.01821
12	Glycyrrhizic acid	0.25	0.7162±0.04979
13	Psoralen	0.012	0.05549±0.01850

Figures




Figure 1

The chemical structure of 13 natural compounds.




Figure 2

Dose-response curves of all tested compounds in cytotoxicity assay. 8 of gradient dilutions of each compound was prepared by 2-fold serial dilution and incubated with cells for 60 h. Dose-response curve of the tested compounds used in PK-15 shown the more cytopathic effect were observed with the increased concentration of compounds.




Figure 3

Anti-PCV2 activities of Paeonol, Cepharanthine or Curcumin detected by qPCR and Western blot. The anti-PCV2 effect of compounds were evaluated by qPCR and Western blot. Cells were incubated with 104.4 TCID₅₀ of PCV2 for 2 h, then tested compounds were added respectively and cultured for 48 h. Cells were collected and relative assay perform. (A) The expression of the Cap gene was detected by qPCR; (B) The expression of the Cap protein was detected by Western blot. GraphPad Prism 5.0 was used for the

statistical analysis. All data were presented in Mean \pm SEM. Different letters to indicate significant differences between groups ($p < 0.05$).




Figure 4

Inhibition of PCV2-induced cell apoptosis by Cepharanthine or Curcumin. (A and B) After cells were infected with 104.4 TCID₅₀ of PCV2 for 2 h, cells were treated with Cepharanthine, Curcumin or Ribavirin for 48 h. The apoptosis rate was evaluated by Annexin V/PI staining followed by flow cytometry. The data on the graph indicated the total number of cells at the late apoptosis of the right upper quadrant and the early apoptosis of the right lower quadrant. The results indicated apoptosis rates were significantly decreased in the treated group with Cepharanthine, Curcumin or Ribavirin Compared to the PCV2-infected group except the group of 0.005 mg/mL Curcumin ($P < 0.05$).




Figure 5

Inhibition of PCV2-induced cell apoptosis by Cepharanthine or Curcumin via the mitochondria pathway. (A and B) The changes of MMP in each group after being staining with JC-1 kit were detected by flow cytometry. R2 represents the value of MMP. The results indicated MMP values were significantly decreased in the treated group with Cepharanthine, Curcumin or Ribavirin Compared to the PCV2-infected group ($P < 0.05$).




Figure 6

The mechanism of inhibition of PCV2-induced mitochondrial apoptosis by Cepharanthine or Curcumin. After cells were infected with 104.4 TCID₅₀ of PCV2 for 2 h, cells were treated with Cepharanthine, Curcumin or Ribavirin for 48 h. (A-H) The expression of the apoptin were analyzed by Western blot and grayscale analysis. (H-I) The results of flow cytometry showed that ROS values were significantly

decreased in the treated group with Cepharanthine at 0.003 and Curcumin at 0.02, 0.01 and 0.005 mg/mL Compared to the PCV2-infected group ($P<0.05$).




Figure 7

The PCV2-induced Mitochondrial apoptosis was mainly remitted by Cepharanthine and Curcumin. The “red arrow” represents the promoting effect and the red “T-shape” indicates the inhibiting effect. The PCV2 replication is inhibited by Cepharanthine and Curcumin, selected from 13 natural compounds, through mitochondrial apoptosis pathway.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile.3.docx](#)
- [AdditionalFile.2.docx](#)
- [AdditionalFile.1.docx](#)