

# An outbreak of intestinal schistosomiasis, alongside increasing urogenital schistosomiasis prevalence, in primary school children on the shoreline of Lake Malawi, Mangochi District, Malawi

**Sekeleghe Kayuni**

Liverpool School of Tropical Medicine

**Angus M. O'Ferrall**

Liverpool School of Tropical Medicine

**Hamish Baxter**

Liverpool School of Tropical Medicine

**Josie Hesketh**

Liverpool School of Tropical Medicine

**Bright Mainga**

Malawi College of Health Sciences

**David Lally**

University of Malawi College of Medicine

**Mohammad H. Al-Harbi**

Liverpool School of Tropical Medicine

**E. James LaCourse**

Liverpool School of Tropical Medicine

**Lazarus Juziwelo**

Government of Malawi

**Janelisa Musaya**

University of Malawi College of Medicine

**Peter Makaula**

Government of Malawi Department of Fisheries

**John Russell Stothard** (✉ [russell.stothard@lstmed.ac.uk](mailto:russell.stothard@lstmed.ac.uk))

Department of Parasitology <https://orcid.org/0000-0002-9370-3420>

---

## Research Article

**Keywords:** Emergence, *Schistosoma mansoni*, Urine CCA-dipsticks, Faecal occult blood, Co-infection, Morbidity, COVID-19

**DOI:** <https://doi.org/10.21203/rs.3.rs-26662/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

**Background:** Intestinal schistosomiasis was not considered endemic in Lake Malawi until November 2017 when populations of *Biomphalaria pfeifferi* were first reported; in May 2018, emergence of intestinal schistosomiasis was confirmed. This emergence was in spite of ongoing control of urogenital schistosomiasis by preventive chemotherapy. Our current study sought to ascertain whether intestinal schistosomiasis is transitioning from emergence to outbreak, to judge if stepped-up control interventions are needed.

**Methods:** During late-May 2019, three cross-sectional surveys of primary school children for schistosomiasis were conducted using a combination of rapid diagnostic tests, parasitological examinations and applied morbidity-markers; **1)** schistosomiasis dynamics were assessed at Samama (n=80) and Mchoka (n=80) schools, where *Schistosoma mansoni* was first reported, **2)** occurrence of *S. mansoni* was investigated at two non-sampled schools, MOET (Mangochi Orphan Education and Training) (n=60) and Koche (n=60) schools, where *B. pfeifferi* was nearby, and **3)** rapid mapping of schistosomiasis, and *B. pfeifferi*, conducted across a further 8 shoreline schools (n=240).

**Results:** In total, 520 children from 12 lakeshore primary schools were examined, mean prevalence of *S. mansoni* by 'positive' urine CCA-dipsticks was 31.5% [95% CI 27.5 – 35.5]. Upon comparisons of infection prevalence in May 2018, significant increases at Samama (RR=1.7 [95% CI 1.4 – 2.2]) and Mchoka (RR=2.7 [95% CI 1.7 – 4.3]) schools were observed. Intestinal schistosomiasis was confirmed at MOET (18.3%) and Koche (35.0%) schools, and in all rapid mapping schools, ranging from 10.0% to 56.7%. Several populations of *B. pfeifferi* were confirmed, with two new eastern shoreline locations noted. Mean prevalence of urogenital schistosomiasis was 24.0% [95% CI 20.3 – 27.7].

**Conclusions:** We notify that intestinal schistosomiasis, once considered non-endemic in Lake Malawi, is now transitioning from emergence to outbreak. Once control interventions can resume after COVID-19, we recommend stepped-up preventive chemotherapy, with increased community-access to treatments, alongside renewed efforts in appropriate environmental control.

## Background

Lake Malawi is the world's fourth largest freshwater lake, an important aquatic hotspot of global biodiversity but with urogenital schistosomiasis being endemic along many parts of its shoreline [1]. In Mangochi District, Malawi, the prevalence of *Schistosoma haematobium* infection in school children warrants preventive chemotherapy. This is achieved by annual mass drug administration (MDA) of praziquantel [2] as provided by the Malawi National Schistosomiasis and Soil-Transmitted Helminthiasis Control Programme (<https://www.health.gov.mw/index.php/schistosomiasis-sth-control-programme>). MDA is typically guided upon country-wide mapping information which is usually developed from inspection of 5 schools per district [3]. By contrast, intestinal schistosomiasis, caused by *Schistosoma mansoni*, is not considered endemic within the lake, as being congruent with the absence of *Biomphalaria pfeifferi*, an obligatory intermediate snail host and keystone snail species for parasite transmission [1,4,5].

This appraisal was revised in May 2018 as, since November 2017 *B. pfeifferi* has been repeatedly encountered in the lake, alongside emergence of intestinal schistosomiasis documented in 3 local primary schools [6]. Prevalence of infection by 'trace/positive' urine CCA-dipsticks was 34.3% [95% CI 27.9 - 41.3], with ova-patent *S. mansoni* in stool noted at Samama and Mchoka schools [6]. Even with ongoing annual MDA for urogenital schistosomiasis control, the dynamics of intestinal schistosomiasis need further scrutiny here, for this disease could transition from emergence to outbreak.

Transitions from emergence to outbreak are often driven by expansions in the distributions of intermediate snail hosts which, like elsewhere in Africa, can instigate, for example, new transmission foci [7]. Even though an outbreak terminology is rather vaguely defined, common with the epidemiology of other water-borne diseases [8], it is more so for schistosomiasis as its transmission dynamics also involve unsafe water contact, with per-cutaneous (and oral) entry and infection routes. However, the use of outbreak vernacular can be appropriate, foremost, to spur commensurate public health actions, for example in stepped-up surveillance for the intermediate hosts or with intensified control interventions. This was evidenced in Senegal for intestinal schistosomiasis [9] and more recently in Corsica for urogenital schistosomiasis [10] which were each urged by the use of outbreak terminologies.

To seek an appropriate public health response here on the shoreline of Lake Malawi, our investigation had three linked objectives: **1)** to resample Samama and Mchoka schools, ascertaining the dynamics of schistosomiasis infection and morbidity after annual MDA, **2)** to confirm intestinal schistosomiasis, also noting faecal occult blood (FOB), at two previously non-sampled schools, MOET (Mangochi Orphan Education and Training) and Koche schools, where in 2018 *B. pfeifferi* was found nearby and **3)** to conduct a wider rapid mapping survey for schistosomiasis at 8 further schools (St Augustine II, Ndembo, Chikomwe, Chipeleka, Sungusya, St Martins, Makumba and Mtengeza) to judge if an outbreak of intestinal schistosomiasis was occurring.

## Methods

Study design and sample size determination for each objective

A cross-sectional study design was used to achieve the three study objectives, see STROBE checklist within supplemental materials. Based on previous epidemiological information [6], a sample size calculation with single population proportion formula (<http://www.raosoft.com/samplesize.html>) showed that a total sample size of 520 was sufficient to estimate overall prevalence of intestinal and urogenital schistosomiasis with  $< \pm 5\%$  precision and 95% confidence.

Based on prevalence data provided by the authors of the May 2018 study [6], a Fisher's exact test was used to show that sampling of 80 children from each of Samama and Mchoka schools in June 2019 was sufficient to detect a 25-percentage-point rise in prevalence of each *Schistosoma* species at each school ( $\alpha < 0.05$ ,  $\beta < 0.20$ ) (objective 1). To ensure detection of *S. mansoni* if present at MOET and Koche schools (objective 2), 60 children were sampled from each. For objective 3, according to WHO recommendations

for rapid mapping, 30 children per school were sampled per school [11]. Random sampling was used at each school following stratification by age and gender. A study flow diagram is included, Figure 1.

*<please insert Figure 1 near here>*

## Study area

At each school, global position system (GPS) coordinates were taken using an Oregon 650 receiver (Garmin, Olathe, Kansas, USA). The GPS locations for each school in decimal degrees are as follows: Samama (-14.417465°, 35.217580 °), Mchoka (-14.439481 °, 35.220644 °), MOET (-14.320776 °, 35.131558 °), Koche (-14.330917 °, 35.146186 °), St Augustine II (-14.473926 °, 35.279613 °), Ndembo (-14.456385 °, 35.273794 °), Chikomwe (-14.422136 °, 35.265088 °), Chipeleka (-14.385387 °, 35.292935 °), Sungusya (-14.386472 °, 35.311398 °), St Martins (-14.351401 °, 35.294435 °), Makumba (-14.319806 °, 35.286104 °) and Mtengeza (-14.288932 °, 35.264073 °). A location map of the 12 schools is shown, Figure 2.

*<please insert Figure 2 near here>*

## Inclusion/exclusion criteria, diagnostics and praziquantel treatment of participants

The surveys took place during late May/June 2019; after obtaining written informed parental consent for each child, a total of 520 children, aged 6-15, of balanced gender, were enrolled. Children not attending school and acutely unwell children were excluded. Participants could withdraw consent at any point. On the appointed day of survey, each school child provided a mid-morning urine sample and when requested, a stool sample, alongside undertaking a brief interview by questionnaire documenting place of birth, recent travel, water-contact habits and praziquantel treatment history. If found infected, upon ova patent infection or 'trace/positive' urine CCA-dipstick test, each child was provided with praziquantel (IDA Foundation, Amsterdam, The Netherlands) at 40 mg/kg.

For detection of intestinal schistosomiasis, two drops of urine were applied to a CCA-dipstick (Rapid Medical Diagnostics, Pretoria, South Africa). Results were scored visually against a reference colour photograph as negative (-ve), trace (tr) or positive (+ve) and cross-checked [12]. To augment urine CCA-dipsticks, on-site inspection of collected stool was performed with parasitological methods; at Mchoka, Samama, MOET and Koche schools, all children were asked to provide a stool sample with a total of 265 specimens obtained, Table 1. Following our rapid mapping protocol at 8 remaining schools, stool was only requested from urine CCA-dipstick +ve children, obtaining 70 specimens, Table 1.

To visualize helminth ova in stool, individual specimens were filtered across a 212 µm metal mesh then applied to produce duplicate thick (41.7 mg) Kato-Katz [11] smears as examined for lateral spine *S. mansoni* ova by microscopy (x100). Intensity of *S. mansoni* infection as eggs per gram (epg) was classified as: light (1 – 99 epg), medium (100 – 399 epg) and heavy ( $\geq$  400 epg) according to WHO guidelines [11]. To assess putative pathology associated with intestinal schistosomiasis [5], stools were screened for FOB using ALLTEST<sup>®</sup> cassettes (Access Diagnostic Tests UK Ltd, Aylsham, UK).

For detection of urogenital schistosomiasis, 10 ml of well-mixed urine was filtered by syringe across a circular nylon mesh of 1.5 cm diameter, with 20 µm pore size (Plastok<sup>®</sup> [Meshes and Filtration] Ltd, Birkenhead, UK). The mesh was stained with Lugol's iodine, then inspected by microscopy (x100) to count terminal spine *S. haematobium* ova. Infection intensity was classified as light (< 50 ova per 10 ml) or heavy ( $\geq$  50 ova per 10 ml) according to WHO guidelines [11]. Putative pathology associated with urogenital schistosomiasis was assessed by Siemens Multistix<sup>®</sup> 10 SG reagent strips (Medisave UK Ltd, Weymouth, UK) for microhematuria [5].

## Malacological surveillance

During May/June 2019, all known locations where *B. pfeifferi* was found were re-surveyed, alongside several new locations as visited on the eastern shoreline of the lake, based upon convenience sampling from in-field observations of human water contact. At each site, two collectors searched, for 20 minutes, for *B. pfeifferi* by hand and with metal sieves. Global position system (GPS) coordinates, altitude and location photographs were taken with an Oregon 650 receiver (Garmin, Olathe, Kansas, USA). Water temperature (°C), pH and conductivity (µS) were recorded with a HI-98129 Pocket EC/TDS and pH Tester (Hanna Instruments Ltd, Leighton Buzzard, Bedfordshire, UK). Collected snails were kept for a week and screened daily for shedding *S. mansoni* cercariae by exposure to sunlight under a dissecting microscope (x20).

## Data analyses

Demographic, questionnaire and diagnostic data were tabulated with statistical analysis carried out using IBM SPSS<sup>®</sup> Version 24 (IBM, Portsmouth, UK). Univariate analyses and Chi-square testing were first performed, then binary logistic regression undertaken, calculating adjusted odds ratios with generalised linear models, with stepwise subtraction of variables, to investigate (un)adjusted epidemiological associations.

# Results

## Prevalence and distribution of intestinal and urogenital schistosomiasis

The outline map, Figure 2, is a summary of all information obtained from urine CCA-dipsticks with the distribution of intestinal schistosomiasis displayed. When 'trace' was considered infected, mean prevalence was 82.5%. When 'trace' was considered not infected, this declined to 31.5%. Common across all school children were very high levels of reported weekly water contact (>75%), inclusive of bathing, swimming and drinking. The known distribution of *B. pfeifferi* along the western shoreline, alongside new reports on the eastern shoreline in December 2018 and May/June 2019, is shown. In locations where *B. pfeifferi* was found, water parameters ranged: pH 7.5 - 8.5, temperature 21.5 – 26.2 °C, conductivity 312 – 458 µS and total dissolved salts 155 – 244 ppm; no collected snail (n=52) was observed to shed *S. mansoni* cercariae.

<please insert Table 1 near here>

Table 1 shows ova-patent *S. mansoni* and that both medium and heavy intensity infections have been observed. Ova patent urogenital schistosomiasis was detected in all schools, ranging from 1.7% to 60.0%, inclusive of heavy intensity infections, except at Koche, St Martins and Makumba schools. Across our sample, 75 (14.4%) children were considered 'free' from schistosomiasis; if urine CCA-dipstick 'trace' was considered infected or 'trace' was considered not infected, then 109 (36.5%) or 56 (10.7%) children were judged co-infected with intestinal and urogenital schistosomiasis, respectively.

<please insert Figure 3 near here>

#### Risk factors associated with schistosomiasis-associated morbidity

Figure 3 shows significant increases of schistosomiasis at Mchoka and Samama, even though MDA treatment coverage (81.9%), as reported by interview, was good. Relative risk (RR) of infection prevalence of *S. mansoni* significantly increased at Samama (RR=1.7 [95% CI 1.4 – 2.2]) and Mchoka (RR= 2.7 [95% CI 1.7 – 4.3]) schools, indicative of substantive re-infection concurrent with increasing environmental transmission for both types of schistosomiasis.

<please insert Table 2 near here>

Table 2 shows an analysis of risk factors associated with schistosomiasis-associated morbidity. 'Positive' urine CCA-dipstick results and ova-patent *S. mansoni* were significantly associated with FOB, alongside ova-patent *S. haematobium* with microhaematuria. Neither age nor gender were associated with these morbidity indicators although a marginal protective effect of MDA, on both FOB and microhaematuria, was observed.

## Discussion

Our integrated surveillance approach was unified by three linked cross-sectional surveys, see Figure 1, and a conjoined malacological inspection. Collectively this builds a more thorough assessment of the changing epidemiology of intestinal and urogenital schistosomiasis on the Lake Malawi shoreline, Figure 2 and Table 1. Of note, is that the prevalence of both forms of schistosomiasis is increasing, Figure 3, indicative perhaps that the force of infection [13] for each parasite is rising, with intestinal schistosomiasis being of newest public health concern here.

Our study detected a mean prevalence of intestinal schistosomiasis by 'positive' urine CCA-dipstick results of 31.5% [95% CI 27.5 – 35.5]. Notably, significant increases in infection prevalence since May 2018 were observed at Samama (RR=1.7 [95% CI 1.4 – 2.2]) and Mchoka (RR= 2.7 [95% CI 1.7 – 4.3]) schools. The disease was also confirmed at MOET (18.3%) and Koche (35.0%) schools with a broader geographical footprint apparent across the 8 rapid mapping schools, with prevalence ranging from 10.0% to 56.7%, and several extant populations of *B. pfeifferi* were confirmed on the eastern and western lake shoreline. Concurrently mean prevalence of urogenital schistosomiasis was 24.0% [95% CI 20.3 – 27.7] with 109 (36.5%) or 56 (10.7%) children co-infected with intestinal schistosomiasis, as contingent upon interpretation of urine-CCA dipstick 'trace' as infection-positive or negative, respectively.

The unexpected occurrence of intestinal schistosomiasis elsewhere in Malawi, alongside the more well-known urogenital schistosomiasis, has been encountered before; the surveys conducted by Poole *et al.* in Chikhwawa during June 2012 noted that 24.9% and 9.1% of mothers and their pre-school-aged children were positive by urine CCA-dipsticks with ova-patent *S. mansoni* infections confirmed [14]. While *Biomphalaria* was not detected in their search for local snails [14], the occurrence of *B. pfeifferi*, as shown here in Figure 2, adds weight to their postulate of intermittent transmission of *S. mansoni* in Chikhwawa. They suggested that the occasional influx of upstream populations of *B. pfeifferi* in the Shire River, as being swept downstream during seasonal flooding, might then colonize temporary pools in the Lower Shire River flood plain, to spark sporadic transmission in Chikhwawa [14]. By contrast, an enduring presence of *B. pfeifferi* along Lake Malawi and Upper Shire River, gives rise to more sustained opportunities in local transmission of *S. mansoni* in Mangochi District.

In regard of this lake shoreline setting, we have shown 1) increases in the prevalence of intestinal schistosomiasis at Mchoka and Samama Schools, 2) occurrence of intestinal schistosomiasis at MOET and Koche schools and 3) endemic intestinal schistosomiasis occurring along a 80 km section of Lake Malawi and Shire River shoreline, noting additional populations of *B. pfeifferi* on the lake's eastern shoreline, Figure 2. Of particular note is the strong association of *S. mansoni* infection, as detected by

urine CCA-dipsticks, with FOB in 16.2% of examined children, see Table 2, indicative of overt intestinal pathology [15]. Combined with the observations of ova-patent infections of moderate- and heavy-intensities at Koche and Ndembo, as well as, ova-patent infections at a further 5 schools, this is pervasive evidence of more sustained local transmission of intestinal schistosomiasis.

Whilst the debate on how to interpret 'trace' reactions of urine-CCA dipsticks continues, a 'positive' reaction is considered solid evidence of active intestinal schistosomiasis [12]. With no association detected between urogenital schistosomiasis and urine-CCA in our study, we conclude that urine-CCA tests are highly specific for *S. mansoni* detection, with 'trace' results indicating light sub-clinical infections, with sub-patent egg outputs. Therefore, 31.5% [95% CI 27.5 – 35.5] of our sampled children were suffering from intestinal schistosomiasis but if a 'trace' reaction was considered diseased then a total of 82.5% [95% CI = 79.2 – 85.8] were infected or, at the very least, at-risk. Of particular note in this light is intestinal schistosomiasis at Ndembo and St Augustine II schools, see Figure 2, where the prevalence of 'positive' urine-CCA dipsticks was > 50% and ova-patent *S. mansoni* infections were encountered, being of light and moderate infection intensities, Table 1; moreover, moderate and heavy ova-patent *S. mansoni* infections were detected at Koche school where the prevalence of 'positive' urine-CCA dipsticks was 35.0%, with *B. pfeifferi* found nearby.

Our rapid disease mapping surveillance across 8 schools, currently augments district-level information of the NCP, critically revising scientific appraisals concerning the previous absence of intestinal schistosomiasis [1], and better demonstrates the newly defined endemicity of intestinal schistosomiasis along the Mangochi District shoreline. When taken as a whole, we judge that there is now sufficient evidence to notify that an outbreak of intestinal schistosomiasis is occurring. This has immediate bearing on the health of the local populace and tourists who may visit here, as well as, in health advice or diagnostic testing undertaken in local or international medical clinics presently unaware of this new risk of intestinal schistosomiasis.

<please insert Figure 4 near here>

In terms of environmental surveillance, it is worthy to note that the lake is undergoing ecological change, most easily seen with lake level changes through time, see Figure 4. Its dynamic shoreline and lake level are manifest, perhaps creating new habitats for *B. pfeifferi* to colonize and or were facilitating collection of this snail in locations previously too deep to be retrieved by hand. The dispersion of this snail, a keystone species for *S. mansoni*, like in Senegal [9] or in Ethiopia [7], is a critical epidemiological driver of intestinal schistosomiasis transmission.

Control of schistosomiasis needs a multisectoral approach and it is often debated how control tactics should be changed [16] or better tailored to aquatic environments [17]. To respond to this outbreak of intestinal schistosomiasis, we propose that current MDA efforts should be intensified, adopting biannual

treatment cycles in schools, which has been successfully implemented elsewhere [18], alongside expanded access to praziquantel for all community members with intestinal schistosomiasis, in need of regular treatment throughout the year [19]. From recent surveys of adult fishermen who have urogenital schistosomiasis, making specific reference to male genital schistosomiasis, co-infection with *S. mansoni* has been noted alongside re-infections within a calendar year [20, 21]. To augment MDA and community-access to praziquantel, it is important to strengthen health education and outreach with suitable WASH interventions [20, 22], better appropriate to this lakeshore setting, noting that even focal application of molluscicides is inappropriate [17], given this lake's global importance in biodiversity.

A significant limitation of our study was the exclusion of certain demographic groups in our survey. This included pre-school-aged and out-of-school children, as well as, more vulnerable adults [23]. However, with increased future resourcing inspection of these groups is important to better assess how they are each afflicted by this outbreak. To do so, we recommend a combination of both rapid urine and faecal sampling methods with inspection of a more extensive range of point-of-contact morbidity markers to provide a better insight into individual disease progression(s) [24]. Future use of 20m shuttle-run tests to assess children's aerobic capacity in relation to *S. mansoni* infection could be insightful, as recently shown elsewhere [25]. However, with COVID-19 suspending annual MDA, we should expect and better prepare for increasing severity of intestinal schistosomiasis in following years.

## Conclusion

Our three main study objectives were achieved: demonstration of increasing prevalence of intestinal and urogenital schistosomiasis at Samama and Mchoka schools, newly confirmed intestinal schistosomiasis at previously non-sampled schools near reported *B. pfeifferi* sites (MOET and Koche schools), and detection of intestinal schistosomiasis at a further 8 sampled schools along the shoreline. Despite ongoing annual MDA of praziquantel for urogenital schistosomiasis, we conclude that an outbreak of intestinal schistosomiasis is occurring in Mangochi District, Malawi. Increased vigilance for *B. pfeifferi*, especially along the lake's eastern shores and in downstream locations on the Shire River, is needed with additional epidemiological inspections of adjacent schools and communities to better gauge the full footprint of intestinal schistosomiasis. Due to the COVID-19 pandemic, this outbreak will continue to expand unchecked, but once control activities can resume, we strongly recommend stepping-up MDA treatment cycles, i.e. from annual to biannual, increasing community access to praziquantel treatment throughout the year, with renewed efforts to mitigate environmental transmission with health education and appropriate WASH interventions.

## Abbreviations

CCA: circulating cathodic antigen; FOB: faecal occult blood; MDA: Mass drug administration; EPG: eggs per gram; RR: relative risk.

## Declarations

## **Acknowledgments**

We are grateful to the local health and education authorities of Malawi with specific thanks to the headteachers, teachers, children and parents who participated in our study; to our friend and colleague Father Henry Chagoma and his staff of Montfort Mission Lake House for their hospitality and convivial company. We thank Dr Michelle Stanton, LSTM for assistance in interpretation of remote sensing imagery presented in Figure 3. Urine CCA dipsticks were supplied by Rapid Medical Diagnostics, South Africa with manufacturer batch number 180907091, expiry date 09/2020. We thank the two anonymous referees who improved this manuscript with their insightful comments.

## **Funding**

SAK and MHAI-H are funded by PhD scholarships from the Commonwealth Scholarship Commission and Ministry of Health, Kingdom of Saudi Arabia, respectively, and JRS, EJLaC by the Higher Education Funding Council for England (HEFCE).

## **Availability of data and materials**

Data used for the analysis are available from the corresponding author upon reasonable request.

## **Authors contributions**

SAK, EJLaC, LJ, JM, PM and JRS designed the study; SAK, AMO'F, HB undertook the parasitological fieldwork with laboratory support from BM and DL. Malacological studies were undertaken by PM, JH, MHAI-H and JRS. Data entry and analyses were undertaken by SAK, AMO'F, HB and JH as overseen by EJLaC, LJ, JM, PM and JRS. All authors read and approved the manuscript for publication.

## **Ethics approval and consent to participate**

Research approvals were granted in Malawi by the National Health Sciences Research Committee (1805), Mangochi District Health Office Research Committee (26.04.2019) and in the UK by LSTM Research Ethics Committee (30.04.2019). Written informed guardian consent was obtained for each school child before participation in surveys.

## **Consent for publication**

All authors have provided consent for publication of the manuscript.

## **Competing interests**

The authors declare that they have no competing interests.

## **Author details**

<sup>1</sup> Department of Tropical Disease Biology, Liverpool School of Tropical Medicine, Liverpool, L3 5QA, UK. <sup>2</sup> Medi Clinic Limited, Medical Aid Society of Malawi (MASM), 22 Lower Sclatter Road, P.O. Box 1254, Blantyre, Malawi. <sup>3</sup> Laboratory Department, Mangochi District Hospital, P.O. Box 42, Mangochi, Malawi. <sup>4</sup> Malawi Liverpool Wellcome Trust Programme of Clinical Tropical Research, Queen Elizabeth Central Hospital, College of Medicine, P.O. Box 30096, Blantyre, Malawi. <sup>5</sup> Ministry of Health, Qassim, Kingdom of Saudi Arabia. <sup>6</sup> National Schistosomiasis and STH Control Programme, Ministry of Health, Lilongwe, Malawi. <sup>7</sup> Department of Basic Medical Sciences, College of Medicine, University of Malawi, Blantyre, Malawi. <sup>8</sup> Research for Health Environment and Development, P.O. Box 345, Mangochi, Malawi.

## References

1. Makaula P, Sadalaki JR, Muula AS, Kayuni S, Jemu S, Bloch P. Schistosomiasis in Malawi: A systematic review. *Parasite Vector* 2014; 7.
2. Kayuni S, Peeling R, Makaula P. Prevalence and distribution of *Schistosoma haematobium* infection among school children living in southwestern shores of Lake Malawi. *Malawi Med J* **2017**; 29:16-23.
3. WHO. Schistosomiasis: Progress report 2001–2011 and strategic plan 2012–2020. World Health Organization, Geneva, 2013.
4. Bowie C, Purcell B, Shaba B, Makaula P, Perez M. A national survey of the prevalence of schistosomiasis and soil-transmitted helminths in Malawi. *BMC Infect Dis* **2004**; 4.
5. Mtethiwa AHN, Nkwengulila G, Bakuza J, Sikawa D, Kazembe A. Extent of morbidity associated with schistosomiasis infection in Malawi: A review paper. *Infect Dis Pov* 2015; 4.
6. Alharbi MH, Condemine C, Christiansen R, LaCourse EJ, Makaula P, Stanton MC et al. *Biomphalaria pfeifferi* snails and intestinal schistosomiasis, Lake Malawi, Africa, 2017-2018. *Emerg Infect Dis* 2019; 25:613-5.
7. Bekana T, Hu W, Liang S, Erko B. Transmission of *Schistosoma mansoni* in Yachi areas, southwestern Ehtiopia: new foci. *Infect Dis Pov* 2019; 8:1.
8. Mari L, Casagrandi R, Rinaldo A, Gatto M. Epidemicity thresholds for water-borne and water-related diseases. *J Theoret Biol.* 2018 447:126-38.
9. Talla I, Kongs A, Verle P, Belot J, Sarr S, Coll AM. Outbreak of intestinal schistosomiasis in the Senegal River Basin. *Ann Soc Belge Med Trop* 1990; 70:173-80.
10. Boissier J, Grech-Angelini S, Webster BL, Allienne JF, Huyse T, Mas-Coma S, et al. Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case study. *Lancet Infect Dis.* 2016; 16:971-9.
11. WHO. Helminth control in school age children: A guide for managers of control programmes. 2nd ed. Geneva: World Health Organization. 2011.
12. Bärenbold O, Garba A, Colley DG, Fleming FM, Haggag AA, Ramzy RM, et al. Translating preventive chemotherapy prevalence thresholds for *Schistosoma mansoni* from the Kato-Katz technique into the

- point-of-care circulating cathodic antigen diagnostic test. PLOS Neglect Trop D. 2018; 12(12).
13. French MD, Churcher TS, Webster JP, Fleming FM, Fenwick A, Kabatereine NB, et al. Estimation of changes in the force of infection for intestinal and urogenital schistosomiasis in countries with schistosomiasis control initiative-assisted programmes. Parasite Vector 2015; 8.
  14. Poole H, Terlouw DJ, Naunje A, Mzembe K, Stanton M, Betson M, et al. Schistosomiasis in pre-school-age children and their mothers in Chikhwawa district, Malawi with notes on the characterization of schistosomes and snails. Parasites Vector 2014; 7:153.
  15. Stothard JR, Stanton MC, Bustinduy AL, Sousa-Figueiredo JC, Van Dam GJ, Betson M, et al. Diagnostics for schistosomiasis in Africa and Arabia: a review of present options in control and future needs for elimination. Parasitol 2014; 141:1947-1961.
  16. Tchuente LAT, Rollinson D, Stothard JR, Molyneux D. Moving from control to elimination of schistosomiasis in sub-Saharan Africa: time to change and adapt strategies. Infect Dis Pov 2017;6:42.
  17. Stothard JR, Campbell SJ, Osei-Atweneboana MY, Durant T, Stanton MC, Biritwum NK et al. Towards interruption of schistosomiasis in sub-Saharan Africa: developing an appropriate environmental surveillance framework to guide and to support 'end game' interventions. Infect Dis Pov 2017; 6:10.
  18. Knopp S, Stothard JR, Rollinson D, Mohammed KA, Khamis IS, Marti H, et al. From morbidity control to transmission control: Time to change tactics against helminths on Unguja Island, Zanzibar. Acta Trop 2013; 128:412-22.
  19. Toor J, Rollinson D, Turner HC, Gouvras A, King CH, Medley GF et al. Achieving elimination as a public health problem for *Schistosoma mansoni* and *S. haematobium*: When is community-wide treatment required? J Infect Dis 2020; corrected proof.
  20. Kayuni SA, LaCourse EJ, Makaula P, Lampiao F, Juziwelo L, Fawcett J, et al. Case Report: Highlighting male genital schistosomiasis (MGS) in fishermen from the southwestern shoreline of Lake Malawi, Mangochi District. Am J Trop Med Hyg 2019: 101; 1331-1335.
  21. Kayuni SA, Corstjens PLAM, LaCourse EJ, Bartlett KE, Fawcett J, Shaw A, et al. How can schistosome circulating antigen assays be best applied for diagnosing male genital schistosomiasis (MGS): an appraisal using exemplar MGS cases from a longitudinal cohort study among fishermen on the south shoreline of Lake Malawi. Parasitol 2019:146;1785-1795.
  22. Campbell SJ, Biritwum NK, Woods G, Velleman Y, Fleming F, Stothard JR. Tailoring Water, Sanitation, and Hygiene (WASH) targets for soil-transmitted helminthiasis and schistosomiasis control. Trends Parasitol 2018; 34:53-63.
  23. Faust CL, Osakunor DNM, Downs JA, Lamberton PHL, Reinhard-Rupp J, Rollinson D. Schistosomiasis control: Leave no age group behind. Trends Parasitol 2020; 36:582-591.
  24. Webster JP, Koukounari A, Lamberton PHL, Stothard JR, Fenwick A. Evaluation and application of potential schistosome-associated morbidity markers within large-scale mass chemotherapy programmes. Parasitol 2009; 136: 1789-1799.
  25. Smith C, McLachlan G, Al-Shehri H, Adriko M, Arinaitwe M, Atuhaire A, Tukahebwa EM, LaCourse EJ, Stanton MC, Stothard JR, Bustinduy AL. *Schistosoma mansoni* infection as a predictor of low aerobic

## Tables

**Table 1.** Occurrence of ova-patent *S. mansoni* in stool and prevalence and intensity of *S. haematobium* infections by school.

School [sample size]	Stool: Kato-Katz ( <i>S. mansoni</i> )					Urine: filtration ( <i>S. haematobium</i> )		
	Number of stool samples collected	Prevalence (%) [95% CI]	Infection intensity <sup>a</sup> (eggs per gram) [% of positives]			Prevalence (%) [95% CI]	Infection intensity <sup>b</sup> (eggs per 10 ml) [% of positives]	
			Light	Medium	Heavy		Light	Heavy
<b>TOTAL</b> [n = 520]	335	*	20 [74.1]	4 [14.8]	3 [11.1]	24.0 [20.3 - 27.7]	80 [64.0]	45 [36.0]
All collected stools, irrespective of CCA status, were examined								
<b>Mchoka</b> [n = 80]	73	1.4 [0.0 - 4.1]	1 [100.0]	0 [0.0]	0 [0.0]	18.8 [10.2 - 27.4]	11 [73.3]	4 [26.7]
<b>Samama</b> [n = 80]	77	5.2 [0.2 - 10.2]	4 [100.0]	0 [0.0]	0 [0.0]	56.3 [45.4 - 67.2]	30 [66.7]	15 [33.3]
<b>MOET</b> [n = 60]	56	3.6 [0.0 - 8.5]	2 [100.0]	0 [0.0]	0 [0.0]	8.3 [1.3 - 15.3]	3 [60.0]	2 [40.0]
<b>Koche</b> [n = 60]	59	15.3 [6.1 - 24.5]	3 [33.3]	3 [33.3]	3 [33.3]	1.7 [0.0 - 5.0]	1 [100.0]	0 [0.0]
Only selective stools from CCA +ve children were examined								
<b>St Augustine II</b> [n = 30]	15	*	3 [100.0]	0 [0.0]	0 [0.0]	43.3 [25.6 - 61.0]	8 [61.5]	5 [38.5]
<b>Ndembo</b> [n = 30]	15	*	6 [85.7]	1 [14.3]	0 [0.0]	60.0 [42.5 - 77.5]	7 [38.9]	11 [61.1]
<b>Chikomwe</b> [n = 30]	10	*	0 [-]	0 [-]	0 [-]	10.0 [0.0 - 20.7]	1 [33.3]	2 [66.7]
<b>Chipeleka</b> [n = 30]	3	*	0 [-]	0 [-]	0 [-]	26.7 [10.9 - 42.5]	4 [50.0]	4 [50.0]
<b>Sungusya</b> [n = 30]	7	*	0 [-]	0 [-]	0 [-]	16.7 [3.4 - 30.0]	4 [80.0]	1 [20.0]
<b>St Martins</b> [n = 30]	4	*	0 [-]	0 [-]	0 [-]	3.3 [0.0 - 9.7]	1 [100.0]	0 [0.0]
<b>Makumba</b> [n = 30]	6	*	1 [100.0]	0 [0.0]	0 [0.0]	6.7 [0.0 - 15.6]	2 [100.0]	0 [0.0]
<b>Mtengeza</b> [n = 30]	10	*	0 [-]	0 [-]	0 [-]	30.0 [13.6 - 46.4]	8 [88.9]	1 [11.1]

\* unable to report prevalence due to selective stool sampling (8.1% of total stool collected was ova-patent; 15.7% of stool collected in selective sampling was ova-patent)

<sup>a</sup> intensity by Kato-Katz: light = 1 - 99 epg; medium = 100 - 399 epg; heavy =  $\geq$  400 epg

<sup>b</sup> intensity by urine filtration: light = < 50 ova per 10 mL; heavy =  $\geq$  50 ova per 10 mL


**Table 2.** Risk factors analyses for morbidity associated with urogenital and intestinal schistosomiasis upon detection of microhematuria and FOB, respectively.

<sup>α</sup> all total of 200 FOB tests were available being used at Samama, Mchoka and MOET schools

<sup>b</sup> a trace result was considered here as not infected, only +ve urine CCA-dipstick scorings were considered infected; our conservative approach was based upon correlates of urine CCA-dipsticks and duplicate Kato-Katz comparisons, with ova-patent prevalence of *S. mansoni* being  $\geq 20\%$ , see Bärenbold et al. [12].


		Microhematuria		FOB	
Prevalence (%) [95% CI]		31.5 [27.5 - 35.5]		16.2 [11.0 - 21.4]	
Sample size		n = 520		n = 191 <sup>a</sup>	
		Unadjusted odds ratio (95% CI) [P-value]	Adjusted odds ratio (95% CI) [P-value]	Unadjusted odds ratio (95% CI) [P-value]	Adjusted odds ratio (95% CI) [P-value]
Urine-CCA test <sup>b</sup>	-ve	1	1	1	1
	+ve	2.0 (1.4 - 3.0) [0.001]	1.2 (0.6 - 2.6) [0.607]	12.9 (4.3 - 38.7) [0.000]	9.2 (3.0 - 28.6) [0.000]
Ova-patent intestinal schistosomiasis (Kato-Katz)	-ve	1	1	1	1
	+ve	2.2 (1.0 - 4.7) [0.059]	3.0 (1.0 - 8.6) [0.043]	11.4 (3.9 - 33.3) [0.000]	6.7 (2.0 - 22.6) [0.002]
Ova-patent urogenital schistosomiasis (urine filtration)	-ve	1	1	1	1
	+ve	42.1 (23.2 - 76.5) [0.000]	47.9 (22.6 - 101.5) [0.000]	1.6 (0.7 - 3.8) [0.248]	1.5 (0.5 - 4.9) [0.492]
Praziquantel treatment in last 12 months	No	1	1	1	1
	Yes	0.7 (0.5 - 1.1) [0.159]	0.7 (0.3 - 1.8) [0.499]	0.5 (0.2 - 1.3) [0.185]	0.8 (0.3 - 2.3) [0.651]
Gender	Male	1	1	1	1
	Female	1.0 (0.7 - 1.4) [0.850]	0.9 (0.5 - 1.8) [0.816]	1.1 (0.5 - 2.3) [1.000]	1.0 (0.4 - 2.4) [0.974]
Age	6 - 10	1	1	1	1
	11 - 15	0.9 (0.6 - 1.4) [0.706]	1.2 (0.6 - 2.3) [0.629]	1.1 (0.507 - 2.4) [0.811]	0.9 (0.3 - 2.3) [0.784]

Figures




**Figure 1**

A study flow-chart of the objectives, sample size and methods used during this investigation.




**Figure 2**

Schematic map showing the prevalence of intestinal schistosomiasis in June 2019, by sampled school, by urine CCA-dipsticks. Freshwater sites inspected for *B. pfeifferi* over the November 2017 – December 2019 period are also shown [Note that schools denoted with a flag represent locations where ova-patent *S. mansoni* infection was observed, and the schools associated with each objectives(1-3). The black arrow labelled 'A' denotes the bay area as shown in the Figure 4 where the shoreline has changed during the 2005-2016 most likely due to lowering lake levels and local sedimentation, where numerous *B. pfeifferi* ( $n \geq 10$ ) have been consistently found].



**Figure 3**

The year-on-year increase of prevalence of urogenital (by urine filtration) and intestinal (by 'trace/positive' urine CCA-dipsticks) schistosomiasis despite annual MDA across the two schools Mchoka and Samama as sampled in 2018 and 2019. Error bars indicate 95% confidence intervals.



**Figure 4**

Composite satellite map, modified from GoogleEarth imagery, that illustrates the changing shoreline of the lake in 2005, 2012, 2013 and 2016. The featured area is the bay indicated by the black arrow labelled 'A' in Figure 1. The green circle 'M12' denotes the sampling location where numerous *Biomphalaria* have been found during all malacological inspections from November 2017 to December 2019. The changing shoreline is most likely resultant from lowering lake levels, see 4B, as well as, upon influx of sediments from the seasonal river in the bottom part of this image. Figure 4B. Annual changes in the lake level during 1992 – 2019 period (see [https://ipad.fas.usda.gov/cropexplorer/global\\_reservoir/gr\\_regional\\_chart.aspx?regionid=eafrika&reservoir\\_name=Malawi](https://ipad.fas.usda.gov/cropexplorer/global_reservoir/gr_regional_chart.aspx?regionid=eafrika&reservoir_name=Malawi)), as detected by remote altimetry, denoting two particularly low-

level periods, in 1996-1998 and 2017-2019, which helps explain the changing shoreline shown in 3A as the lake recedes.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [STROBEchecklist.docx](#)