

Age and APOE genotype affect the relationship between objectively measured physical activity and power in the alpha band, a marker of brain disease.

Jaisalmer de Frutos-Lucas (✉ jaisalmer.dfl@gmail.com)

Edith Cowan University <https://orcid.org/0000-0003-2970-4338>

Pablo Cuesta

Universidad Complutense de Madrid

Federico Ramirez-Toraño

Universidad Complutense de Madrid

Alberto Nebreda

Universidad Complutense de Madrid

Esther Cuadrado-Soto

Universidad Complutense de Madrid

África Peral-Suarez

Universidad Complutense de Madrid

David Lopez-Sanz

Universidad Complutense de Madrid

Ricardo Bruña

Universidad Complutense de Madrid

Silvia Marcos-de Pedro

Hospital Rey Juan Carlos

María L Delgado-Losada

Universidad Complutense de Madrid

Ana M López-Sobaler

Universidad Complutense de Madrid

Inmaculada C Rodríguez-Rojo

Centro Universitario Villanueva

Ana Barabash

Hospital Clinico Universitario San Carlos

Juan M Serrano Rodríguez

Universidad Autonoma de Madrid

Simon M Laws

Edith Cowan University

Alberto Marcos

Hospital Clinico Universitario San Carlos

Ramon Lopez-Higes

Universidad Complutense de Madrid

Belinda M Brown

Murdoch University

Fernando Maestú

Universidad Complutense de Madrid

Research

Keywords: APOE ϵ 4, magnetoencephalography, Alzheimer's disease, physical activity, alpha power

DOI: <https://doi.org/10.21203/rs.3.rs-27040/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

BACKGROUND: Electrophysiological studies show that reductions in power within the alpha band are associated with the Alzheimer's disease (AD) continuum. Physical activity (PA) is a protective factor that has proved to reduce AD risk and pathological brain burden. Previous research has confirmed that exercise increases power in the alpha range. However, little is known regarding whether other non-modifiable risk factors for AD, such as increased age or *APOE* ϵ 4 carriage, alter the association between PA and power in the alpha band.

METHODS: The relationship between PA and alpha band power was examined in a sample of 113 healthy adults using magnetoencephalography. Additionally, we explored whether ϵ 4 carriage and age modulate this association. The correlations between alpha power and gray matter volumes and cognition were also investigated.

RESULTS: We detected a parieto-occipital cluster in which PA positively correlated with alpha power. The association between PA and alpha power remained following stratification of the cohort by genotype. Younger and older adults were investigated separately, and only younger adults exhibited a positive relationship between PA and alpha power. Interestingly, when four groups were created based on age (younger-older adult) and *APOE* (E3/E3-E3/E4), only younger E3/E3 (least predicted risk) and older E3/E4 (greatest predicted risk) had associations between greater alpha power and higher PA. Among older E3/E4, greater alpha power in these regions was associated with improved memory and preserved brain structure.

CONCLUSION: PA could protect against the slowing of brain activity that characterizes the AD continuum, where it is of benefit for all individuals, especially E3/E4 older adults.

1. Introduction

As life expectancy increases worldwide, we are witnessing an increase in the prevalence of age-related diseases, such as Alzheimer's disease (AD) (Prince et al., 2015). AD is a neurodegenerative disease of unknown aetiology characterized by progressive cognitive impairment that prevents the individual from engaging in an active, independent life and ultimately causes death (Förstl and Kurz, 1999; Rathmann and Conner, 2007). Although the clinical onset of AD usually occurs at 70 years on average (Corder et al., 1993), strong evidence suggests that the underpinning neuropathological process begins up to 20 years before the appearance of the first symptoms (Braak and Braak, 1991; Chen and Mobley, 2019; Jack et al., 2019, 2014). AD is a multifactorial disease that is influenced by a combination of environmental and genetic factors, where age and the carriage of the apolipoprotein E (*APOE*) ϵ 4 allele are considered the greatest non-modifiable risk factors (Farrer, Lindsay et al., 1997; Lindsay et al., 2002).

Several neuroimaging techniques have been utilised to investigate AD disease progression. This work has identified a slowing of background brain activity throughout the AD continuum, mainly over occipital and parietal regions, as measured through magnetoencephalography (MEG) or electroencephalography (EEG)

(Babiloni et al., 2014, 2011; Moretti et al., 2004; Susi et al., 2019). More specifically, decreases in the peak and median frequencies of the power spectrum have been consistently described (Escudero et al., 2009; Poza et al., 2007), which is thought to be driven by both an enhancement of low frequencies (2-8Hz) and a decrease in high frequencies (8-45 Hz) (Berendse et al., 2000; Osipova et al., 2005; Poza et al., 2008a, 2008b). Particularly relevant is the activity within the alpha band (8-12Hz), which usually comprises the peak of the power spectrum. Power in the alpha band has been the focus of attention in many studies targeting different stages of preclinical and prodromal AD, as well as clinically symptomatic AD. Alterations of this kind of activity have been detected in asymptomatic $\epsilon 4$ carriers (Frutos-lucas et al., 2018), in individuals with subjective cognitive decline (SCD) (López-Sanz et al., 2016) and mild cognitive impairment (MCI) (Garcés et al., 2013), and correlate with cognitive worsening. Power reduction in the alpha band is thought to be the result of cumulative synaptic damage, mostly of cholinergic connections, which impedes the synchronization of activity at higher frequencies (Ebert and Kirch, 1998; Osipova et al., 2005).

While there is currently no cure for AD, several lifestyle factors have been identified to prevent or slowdown disease progression. Physical activity (PA) is one of the most relevant protective factors within this field of study (Kivipelto et al., 2018; Yuri Di Marco et al., 2014). Several studies suggest that PA triggers a series of neuroprotective mechanisms in the brain that effectively reduce AD risk and neuropathological burden (Brown et al., 2018; Erickson et al., 2012; Kennedy et al., 2017; Lindsay et al., 2002; Okonkwo et al., 2014). Additionally, previous literature has shown that both acute and chronic PA are associated with higher peak frequencies and power in the alpha band (Frutos-lucas et al., 2018; Gutmann et al., 2015; Kubitz and Pothakos, 1997). Less well-known is how risk factors for AD, such as increased age and *APOE* $\epsilon 4$ carriage, influence this relationship. In this line, $\epsilon 4$ carriage has been associated with lower age at onset and has a greater impact on brain pathology at earlier disease stages (Corder et al., 1993; Yamazaki et al., 2019). It is crucial to understand whether individuals who are more likely to develop AD can also benefit from a more active lifestyle. We have previously shown that highly active older adults (age 60 – 80) present with higher alpha peak frequencies, although this relationship only existed among $\epsilon 4$ non-carriers (Frutos-lucas et al., 2018).

In this study, we aim to provide a detailed characterization of the relationship between PA and activity in the alpha band, and the role of age and *APOE* $\epsilon 4$ carriage as potential moderators of this relationship. With that purpose, we will examine MEG recordings at rest from a sample of younger (48-59 years) and older adults (60+) with available objective measures of PA, classified as *APOE* E3/E3 or E3/E4. Based on current literature, we hypothesize that PA will be associated with higher power in the alpha band in posterior brain regions. However, we believe that E3/E4 carriage could determine the time window during which PA could more effectively impact brain function, so that this relationship will be stronger in younger E3/E4, compared with older E3/E4.

2. Materials And Methods

2.1. Participants

A sample of 262 individuals participated in a research project aimed to study the neurophysiological features of healthy and pathological aging, with a particular interest in the recruitment of individuals at increased risk for AD. Participants were invited to join the study from local hospitals and associations and through several dissemination talks. A team of expert neuropsychologists ensured that individuals willing to participate met inclusion criteria. The detailed list of exclusion criteria have been detailed previously (de Frutos-Lucas et al., 2018). Participants were asked to provide signed informed consent. The Institutional Review Board Ethics Committee at Hospital Universitario San Carlos Ethics Committee approved the study protocol and the procedure was performed following the Helsinki declaration and national and European Union regulations.

From the original cohort, we only included participants who had available and valid data regarding our main variables of interest ($n = 191$; Mini Mental State Examination, MMSE, score, genetic information and validated magnetic resonance imaging, MRI, MEG and actigraphy data). Additionally, we excluded anyone aged less than 45 years ($n = 6$), with an MMSE score less than 26 ($n = 7$), or carrying less frequent *APOE* genotypes ($\epsilon 2\epsilon 2$, $n = 2$; $\epsilon 2\epsilon 3$, $n = 14$; $\epsilon 2\epsilon 4$, $n = 2$; $\epsilon 4\epsilon 4$, $n = 5$). Also, participants aging less than 60 were considered young adults (45–59) and those aging 60 and above were considered old adults (60–82). Accordingly, the remaining 155 participants were then categorized into one of four groups: young E3/E4 ($n = 30$), old E3/E4 ($n = 16$), young E3/E3 ($n = 61$) and old E3/E3 ($n = 48$). To ensure all groups were matched with regard to PA levels, sex, educational level, MMSE and body mass index, a subset of participants from each group were selected for inclusion: young E3/E4 ($n = 20$), old E3/E4 ($n = 16$), young E3/E3 ($n = 44$) and old E3/E3 ($n = 33$). Additionally, we made sure that genotype groups would match in age (E3/E3 and E3/E4) and age groups would match in the percentage of E3/E4 (younger adults and older adults). Reasons to match the sample according to all these relevant variables instead of using them as covariates in subsequent analyses have been reported previously (de Frutos-Lucas et al., 2020).

The final sample was composed of 113 healthy adults, aged 48–82 years. A detailed list of the sample characteristics can be found in Table 1, including scores extracted from the neuropsychological tests: Geriatric Depression Scale (Yesavage et al., 1982), the anxiety subscale from the Goldberg Anxiety and Depression Inventory (Herrero et al., 2003) and the Digit Span (an index was created using forward and backward spans), Logical Memory II (an index was created using gist immediate and delayed recall) subscales from the Weschler Adult Intelligence Scale IV (WAIS-IV, (Weschler, 2008)).

Table 1
Descriptive measures of the final sample

<i>Variable</i>	<i>WHOLE SAMPLE</i>	<i>YOUNG34</i>	<i>OLD34</i>	<i>YOUNG33</i>	<i>OLD33</i>	<i>Group Comparison</i>
Sex (M; F)	31; 82	4; 16	3; 13	12; 32	12; 21	No sig. diffs. (p > 0.05 all groups).
Age	59.92 ± 7.52	55.05 ± 2.76	66.56 ± 5.67	54.48 ± 2.9	66.91 ± 6.27	No sig. diffs. within age groups (p > 0.05 Y33-Y34 & O33-O34).
Education	4.60 ± 0.62	4.55 ± 0.69	4.8 ± 0.41	4.64 ± 0.57	4.48 ± 0.71	No sig. diffs. (p > 0.05 all groups).
MMSE	29.16 ± 0.94	29.25 ± 0.97	29.25 ± 0.86	29.07 ± 0.87	29.18 ± 1.07	No sig. diffs. (p > 0.05 all groups).
Depression	1.29 ± 1.39	1.42 ± 1.64	1.27 ± 1.10	1.38 ± 1.41	1.10 ± 1.37	No sig. diffs. (p > 0.05 all groups).
Anxiety	1.82 ± 2.12	1.80 ± 2.19	2.11 ± 2.67	1.89 ± 2.14	1.63 ± 1.96	No sig. diffs. (p > 0.05 all groups).
BMI	24.98 ± 3.56	24.53 ± 2.74	24.42 ± 3.85	25.53 ± 4.16	24.76 ± 2.98	No sig. diffs. (p > 0.05 all groups).
Physical Activity	0.012 ± 0.012	0.011 ± 0.010	0.010 ± 0.012	0.015 ± 0.013	0.011 ± 0.012	No sig. diffs. (p > 0.05 all groups).
Total Gray Matter (x10³)	5.73 ± 0.51	5.80 ± 0.40	5.48 ± 0.47	5.87 ± 0.55	5.62 ± 0.48	Diffs. across age groups (p < 0.05 for Y34-O34, O34-Y33 & Y33-O33).
Precuneus (x10³)	8.46 ± 1.04	8.62 ± 1.06	8.10 ± 0.93	8.82 ± 1.13	8.05 ± 0.75	Diffs. across age groups (p < 0.05 for Y34-O33, O34-Y33 & Y33-O33).
Hippocampus (x10³)	3.71 ± 0.41	3.79 ± 0.39	3.44 ± 0.38	3.83 ± 0.38	3.63 ± 0.40	Diffs. across age groups (p < 0.05 for Y34-O34, O34-Y33 & Y33-O33).

<i>Variable</i>	<i>WHOLE SAMPLE</i>	<i>YOUNG34</i>	<i>OLD34</i>	<i>YOUNG33</i>	<i>OLD33</i>	<i>Group Comparison</i>
Episodic Memory	24.28 ± 5.13	24.35 ± 4.97	23.09 ± 6.32	24.05 ± 4.78	25.16 ± 5.49	No sig. diffs. (p > 0.05 all groups).
Working Memory	10.26 ± 2.10	10.37 ± 2.19	10.00 ± 2.31	10.20 ± 1.95	10.39 ± 2.21	No sig. diffs. (p > 0.05 all groups).

INSERT Table 1 ABOUT HERE

2.2. Physical Activity Measurement

An ActiGraph GT3X + accelerometer (LLC, Pensacola, FL) was provided to every participant and they were requested to wear the device on their right hip for 7 complete days. They were advised to only take the ActiGraph off during water-based activities (Chomistek et al., 2017; Chudyk et al., 2017). ActiLife software (6.13.3) (LLC, Pensacola, FL) was used to clean and process the acquired data. To meet validation criteria, each individual should have worn the accelerometer for a minimum of ten hours per day during at least 3 weekdays and 1 weekend day (Chomistek et al., 2017). Non-wear time was defined as ≥ 60 min of continuous zeroes, allowing for up to 2 min of ≤ 100 counts (Troiano et al., 2008). For this study, we considered a standardized measure of total PA volumes (TPA) calculated by ActiLife: Total Time In Freedson Bouts. TPA was normalized by total wear time.

2.3. APOE Genotyping

Genomic DNA was extracted from 10 ml blood samples in ethylenediaminetetraacetic acid (EDTA). TaqMan assays were used on an Applied Biosystems 7500 Fast Real Time PCR machine (Applied Biosystems, Foster City, CA) to determine single nucleotide polymorphisms (SNPs) rs7412 and rs429358 genotypes and establish *APOE* haplotypes accordingly. As mentioned above, only $\epsilon 3\epsilon 3$ and $\epsilon 3\epsilon 4$ individuals were included in this study.

2.4. MRI acquisition and Volumetric Analyses

T1-weighted MRI images from each participant were generated using a General Electric 1.5 T system and applying a high-resolution antenna and a homogenization PURE filter (Fast Spoiled Gradient Echo sequence, TR/TE/TI = 11.2/4.2/450 ms; flip angle 12°; 1 mm slice thickness, 256 × 256 matrix and FOV 25 cm). Freesurfer software (version 6.1.0) was used for automated cortical parcellation and subcortical segmentation (Fischl et al., 2002). The measures that were included in further analyses were total grey matter, precuneus and hippocampus (in mm³). The volumes of bilateral structures were collapsed in order to obtain a single measure for each region.

2.5. Magnetoencephalography

Neurophysiological data were acquired using a whole-head Elekta-Neuromag MEG system with 306 channels (Elekta AB, Stockholm, Sweden) at the Center for Biomedical Technology (Madrid, Spain). MEG

data were collected at a sampling frequency of 1000 Hz and online band-pass filtered between 0.1 and 330 Hz.

All subjects underwent a 5 minutes eyes- closed resting-state MEG recording while sitting comfortably inside of a magnetically shielded room. They were requested to stay awake and to minimise their body movements. Each subject's head shape was defined relative to three anatomical locations (nasion and bilateral preauricular points) using a 3D digitizer (Fastrak, Polhemus, VT, USA) and head motion was tracked through four head-position indicator (HPI) coils attached to the scalp. These HPI coils continuously monitored the subjects' head movements, while eye movements were monitored by a vertical electrooculogram assembly (EOG) composed of a pair of bipolar electrodes. Raw recording data was first introduced to Maxfilter software (v 2.2, correlation threshold = 0.9, time window = 10 seconds) to remove external noise using the temporal extension of the signal space separation method with movement compensation (Taulu and Simola, 2006). Then, magnetometers data (Garcés et al., 2017) was automatically examined to detect ocular, muscle, and jump artifacts using Fieldtrip software (Oostenveld et al., 2011), which were visually confirmed by an MEG expert. The remaining artifact-free data was sectioned into four-second segments. Afterwards, independent component analysis-based procedure (ICA) was applied to remove heart magnetic field artifacts and EOG components. Only those recordings with at least 20 clean segments (80 seconds of brain activity) were utilized in subsequent analyses.

MEG clean time series were band-pass filtered (2 seconds padding) between 2 and 45 Hz. Source reconstruction was carried out using a regular grid of 1 cm spacing in the Montreal Neurological Institute (MNI) template. The resulting model comprised 2459 sources homogeneously distributed across the brain. This model was linearly transformed to each subject's space. The leadfield was calculated using a single shell model (Nolte, 2003). Sources time series were reconstructed using a Linearly Constrained Minimum Variance beamformer (Van Veen et al., 1997). Power spectrum of each grid's node was computed by means of Fast Fourier Transform using Hanning tapers with 0.25 Hz smoothing. For each node, relative power was calculated by normalizing by total power over the 1.5- to 45-Hz range. Since power in the alpha band was the focus of the study, only frequency steps (25 in total) within the interval [8-14Hz] were considered for the analyses. The source template with 2459 nodes in a 10 mm spacing grid was segmented into 72 regions of the Automated Anatomical Labeling atlas (AAL, Tzourio-Mazoyer et al., 2002), excluding the cerebellum, basal ganglia, thalamus, amygdala, insula, and olfactory cortices. Those 72 regions of interest included 1202 of the original 2459 nodes. Trials were averaged across subjects ending up with a source-reconstructed power matrix of 1202 nodes x 25 frequency steps x 113 participants. This final power matrix was employed in the correlation analysis (see below).

2.6. Statistical analyses

The aim of this study was the detection of any robust correlation between power values derived from clusters of nodes localized in certain brain regions and TPA. Such analysis relied on the network-based statistics (NBS) (Fornito et al., 2016; Zalesky et al., 2010). Clusters consisted of several adjacent nodes, which systematically showed a significant partial correlation (with age as covariate) in at least 3 consecutive frequency steps between their corresponding power values and TPA (Spearman correlation p-

value < 0.01). Importantly, all nodes within a cluster must have shown the same sign of the correlation coefficient, this indicating that the cluster might be deemed as a functional unit. Only clusters involving at least 1% of the nodes (i.e. a minimum of 12 nodes) in each frequency step were considered. Cluster-mass statistics were assessed through the sum of the Spearman rho values across all nodes and significant frequency steps.

Then, to control for multiple comparisons, the entire analysis pipeline was repeated 5000 times shuffling the correspondence between power estimates and TPA across subjects. At each repetition, the maximum statistic of the surrogate clusters (in absolute value) was kept creating a maximal null distribution that would ensure control of the family-wise error rate (FWER) at the cluster level. Cluster-mass statistics on each cluster in the original data set was compared with the same measure in the randomized data. The NBS p-value represented the proportion of the permutation distribution with cluster-mass statistic values greater or equal than the cluster-mass statistic value of the original data.

Power values were averaged across all nodes and frequencies that belonged to the cluster. Such average was considered as the representative power marker value for that cluster, and further subjected to new correlation analyses. Therefore, the statistics presented in the results section derived from the correlation between the averaged power value of each significant cluster and the corresponding TPA for each participant. As it has been mentioned above, correlations were first performed within the whole sample. In a second step, correlations between power and TPA scores were performed independently for all subgroups within the sample (E3/E3, E3/E4, younger adults, older adults, young E3/E3, young E3/E4, old E3/E3 and old E3/E4). Additionally, moderation analyses were carried out to study the possible influence exerted by either age (younger vs older adults) and *APOE* (E3/E4 vs E3/E3) in the reported relationship between average power and TPA. To this aim, we employed multiple regression analysis and calculated the increase in variance explained by our model after including the interaction term in two separate models (i.e., *APOE**TPA and age*TPA, model_1 and model_2 respectively). These models used TPA and *APOE* (model_1) / age (model_2) as predictors to linearly estimate average power in the significant cluster. In a second step, the interaction term for each specific model was added (TPA**APOE* or TPA*age). The p value for this interaction term is interpreted as the moderating effect significance.

Finally, average power values were correlated with measures of memory performance (working and episodic memory) and grey matter volumes (total grey matter, precuneus and hippocampus), traits that are known to be affected early in AD. These analyses were only carried out for those subgroups showing a significant correlation between TPA and average power (the complete list is shown in Table 2). Statistical analyses were carried out using Matlab R2017a (Mathworks Inc).

Table 2
AAL ROIs that were partially captured by the significant cluster

<i>ROI</i>	<i>% of ROI occupied</i>
Right Precentral gyrus	88.89
Right Postcentral gyrus	63.64
Left Precuneus	67.86
Left Middle Occipital lobe	58.62
Right Superior Parietal gyrus	83.33
Left Superior Parietal gyrus	81.25
Right Angular gyrus	72.22
Left Postcentral gyrus	35.29
Right Inferior Parietal gyrus	100.00
Right Superior Frontal gyrus	25.81
Right Precuneus	38.10
Right Inferior Frontal gyrus, Opercular	46.15
Left Superior Occipital lobe	54.55
Right Cingulate gyrus, Middle part	33.33
Left Cuneus	45.45
Right Supramarginal gyrus	50.00

3. Results

A significant cluster was found in the frequency interval [10.75–13Hz] mainly comprising posterior regions of the brain (see Fig. 1A and Table 2). The power in all frequencies of this interval positively correlated with TPA across the whole sample ($\rho = 0.360$, $p < 0.0001$). The maximum cluster size was found at 11.75 Hz (220 nodes). The cluster size oscillates between a minimum of 67 nodes at the beginning of the frequency range and 14 at the end of that frequency range, see Fig. 1B. 11.75 Hz showed the highest average correlation coefficient value across all nodes of the cluster; $\rho = 0.3029$.

INSERT FIGURE 1 ABOUT HERE

INSERT Table 2 ABOUT HERE

The correlation between TPA and power in the [10.75–13Hz] interval within the cluster generated in the previous step remained significant for both genotype groups (E3/E3: $\rho = 0.326$, $p = 0.004$, E3/E4: $\rho =$

0.442, $p = 0.007$; see Fig. 1C) and for younger adults ($\rho = 0.456$, $p < 0.001$), although not for older adults ($\rho = 0.234$, $p = 0.105$; Fig. 1D). In this line, only the moderator effect of age, but not that of E3/E4, was significant (respectively, $p = 0.030$ and $p = 0.923$). When both factors were simultaneously considered, only young E3/E3 ($\rho = 0.487$, $p < 0.001$) and old E3/E4 ($\rho = 0.603$, $p = 0.013$) presented a significant correlation (Fig. 1E). Importantly, there were no significant differences in power in the [10.75–13Hz] range among the four subgroups.

The average power value of the significant cluster was used for new correlation analyses in younger E3/E3 and older E3/E4. Alpha power in posterior brain regions correlated with preserved brain structure and improved working and episodic memory, but only among older E3/E4. The complete set of correlations is displayed in Table 3.

Table 3
Average Power Correlations

VARIABLE	OLD34			YOUNG33		
	n	Rho	p value	n	Rho	p value
Total Gray Matter	16	0.624	*0.012	43	-0.017	0.915
Precuneus	16	0.515	*0.044	42	0.117	0.460
Hippocampus	16	0.271	0.310	43	-0.081	0.605
Episodic Memory	11	0.881	*<0.001	43	0.181	0.246
Working Memory	16	0.587	*0.017	43	-0.220	0.156

INSERT Table 3 ABOUT HERE

4. Discussion

The slowing of background brain activity, usually defined as a shift to the left in the power spectrum, is a consolidated marker of aging and brain disease which has been used to monitor pathological progression along the AD continuum. In this study, we aimed to characterize the relationship between PA and activity in the alpha band and the role of age and *APOE* $\epsilon 4$ carriage as potential moderators of this relationship. In a previous publication, we reported that greater self-reported PA was significantly associated with greater alpha peak frequencies in older adults. Moreover, we were able to observe that such relationship remained significant only among E3/E3, although the small sample size in the E3/E4 group did not allow us to reach a definite conclusion (de Frutos-Lucas et al., 2018). In this study, taking a different approach, we have been able to detect a positive relationship between objectively measured PA (in terms of total volume) and power in the alpha band in a cluster comprising mainly occipital and parietal regions, which are classic sources of this brain rhythm. More importantly, this effect persisted in both E3/E4 and E3/E3. However, when we split the sample into younger and older adults, only younger adults exhibited greater alpha power at increased PA volumes. Interestingly, when we further divided the

sample considering both risk factors combined (age and genotype), only young E3/E3 and old E3/E4 showed the beneficial association with PA.

Late onset Alzheimer's disease is a multifactorial disease, which means that different risk and protective, environmental and genetic factors interact throughout the lifespan to determine the likelihood that an individual will develop the disease. PA has been well established as an important neuroprotective element that has the potential to decrease AD risk and improve AD-related neuropathological burden (Brown et al., 2018; Erickson et al., 2012; Okonkwo et al., 2014). In fact, both greater power in the alpha band and a more active lifestyle have been reported to be associated to an individual's capacity to generate long-term potentiation-like synaptic plasticity (Schättin et al., 2018). PA presents the advantage of being a modifiable lifestyle variable. Individuals can purposefully increase their PA levels in order to improve their health outcomes. On the other side, age and *APOE* ϵ 4 carriage are considered the two non-modifiable risk factors for AD that have the greatest impact on the probability that an individual will develop AD (Farrer, Lindsay et al., 1997; Lindsay et al., 2002). For this reason, it is important to understand how these elements interplay to shape brain structure and function.

As it has already been mentioned, increasing age is associated with greater risk for AD and lower power in the alpha band at rest. Here we report that the potentially protective association between PA and greater alpha power is stronger in younger adults (48–59 years). This finding does not necessarily contradict the previous finding showing that the alpha peak frequency at rest was higher in highly active older adults. Both features (alpha peak and alpha power) are closely related and they are affected by the same processes that cause the loss of (mainly cholinergic) synapses (Ebert and Kirch, 1998; Mufson et al., 2008; Osipova et al., 2005). Still, they reflect distinct aspects of the same phenomenon, which could be differentially modulated by various external factors along the disease continuum. Power is highly influenced by the number of simultaneously active synapses. Even in the absence of brain disease, aging is characterized by a reduction in power in high frequencies (Gómez et al., 2013). Therefore, PA might not be able to increase power as effectively in older adults, due to neuronal damage inherent to aging. However, it seems to still be able to exert a positive effect, as reflected by a positive association with the peak frequency. In fact, the power frequency range we found to be associated with PA levels (10.75-13Hz) does not overlap with (it is higher than) the average alpha peak previously reported for healthy adults (Gaál et al., 2010).

Other studies, however, have reported that acute PA (e.g., a single bout of aerobic exercise) has a greater effect on spectral measures (Dimitrova et al., 2017) and cognitive outcomes (Chang et al., 2012) in older adults, although Moraes et al. (2011) only found significant effects in young adults. The acute neurophysiological response to PA, especially the enhancement of theta activity, is believed to serve a cognitive function to overcome exercise-related cognitive challenges (Horne, 2013). In these studies, older individuals presented lower baseline power across different frequency ranges (alpha, theta and delta) than younger individuals. The increase in power produced by acute PA, which immediately reverts after exercise ceases (Kuo et al., 2010; Li et al., 2008), was greater in older individuals, while still not reaching the power levels featured by younger participants. Therefore, there seems to be more room for

improvement among older adults, which is promoted by acute PA. However, habitual engagement in PA could possibly have greater potential to constrain the slowing of background (resting) activity among younger individuals. Thus, it is reasonable to suggest that the immediate and long-term beneficial effects of PA in the brain could be mediated by different mechanisms which might not be equally prominent at different life stages. For example, different neurotransmitters could be responsible for such diverse effects, which in turn could possibly be differentially altered in aging and disease (Mattson and Magnus, 2006; Moraes et al., 2011).

Alternatively, it is possible that with aging there is a shift from posterior to anterior background activity sources. In fact, a few studies showing that alpha power could increase in older adults with acute, habitual physical exercise or after an exercise-based intervention, focused on frontal activity sources (Choi et al., 2018; Sanchez-Lopez et al., 2018; Vogt et al., 2010). Here, when we studied the effect of age we focused on a cluster comprising parieto-occipital brain regions where the relationship between PA and alpha power was maximized across the whole sample. Hence it remains possible that older adults present a weaker association limited to the brain sources explored here. It is also important to bear in mind that in previous studies both the intensity and the duration of acute exercise have been found to influence the specific neural response evoked (Mechau et al., 1998; Woo et al., 2009), which could also explain certain differences across studies.

APOE ϵ 4 carriage poses the greatest genetic risk factor for AD. There is controversy in the field regarding whether ϵ 4 carriers and non-carriers benefit to the same extent from PA engagement. Many studies conclude that physical inactivity is particularly harmful for ϵ 4 carriers (Brown et al., 2013; Head et al., 2012; Luck et al., 2014; Rovio et al., 2005; Smith et al., 2013) Yet, a considerable number of publications fail to find benefits among ϵ 4 carriers, suggesting that the presence of this allele could impair the mechanisms through which PA exerts its action (Chang et al., 2010; Fenesi et al., 2017; Podewils et al., 2005). Nevertheless, these apparently conflicting results are a consequence of the employment of different study designs to evaluate a wide variety of outcomes. In the present study, we were able to replicate the association between PA and alpha power described for the sample as a whole when looking separately at E3/E4 and E3/E3. Although ϵ 4 carriage can worsen cholinergic deficits (Poirier et al., 1995), which could explain previously reported alterations in alpha activity, PA is equally associated with increased alpha power regardless of genetic burden.

Here, we describe for the first time that only young E3/E3 and old E3/E4 present a significant positive relationship between PA and alpha power. Such findings contradict our original hypothesis that, in E3/E4, PA would be associated with better outcomes at earlier stages in life. We would have thought that the presence of this risk allele would accelerate brain aging and pathology, hence bringing forward the time window during which PA could perform its protective role. It seems though that PA in midlife is associated with greater alpha power only among ϵ 4 E3/E3. One possible explanation is that the benefits of PA are maximal in individuals at opposite ends of the risk spectrum. On the one end, young individuals who do not present increased genetic risk for AD are able to generate a positive response to exercise, since the machinery that mediates these effects is in optimal conditions. On the other end, old individuals

at increased genetic risk are at a more vulnerable position where even the slightest changes could result in meaningful improvements. This is coherent with the idea that only in this group power correlated with improved memory and better-preserved brain structure. In other words, local increases in alpha power in older $\epsilon 4$ E3/E4 could serve a compensatory purpose. However, mean age in our sample, even in the older groups, was still below previously reported mean age at disease onset (Corder et al., 1993). Therefore, it is possible that the beneficial effects that E3/E4s show in their 60 s could be observable in E3/E3 at later stages (they might not need to activate compensation mechanisms yet), when pathological burden among E3/E4 might be too advanced to allow circuit reorganization. More research is needed to fully elucidate the processes that give rise to these seeming interaction between age and *APOE* genotype.

The present study combines a series of strengths that confer it a novel perspective. Accelerometers were used to obtain objective estimates of PA, which were later used as continuous measures, thus avoiding the loss of statistical power that comes with the stratification in different PA levels. In this line, over-recruitment of E3/E4 enabled the careful examination of the interplay among a series of modifiable and non-modifiable factors that impact dementia risk. Age and genetic risk for AD were considered separately and, for the first time, jointly as potential limiting factors that could constrain the relationship between PA and synaptic function. In addition, we used a completely data-driven methodology that enables the automatic delimitation of the spatial and frequential dimensions of the clusters. Finally, brain activity data was collected through MEG, which offers ample temporal and spatial resolutions (López-Sanz et al., 2019). Neurophysiological approaches (i.e. MEG and EEG) in the study of AD characterization and prevention present the advantage of being completely non-invasive techniques that can easily be repeated over time. Therefore, they offer great potential to be used in the process of monitoring disease progression or assessing the impact of different kinds of interventions.

5. Limitations

Overall, this study endeavours to deepen our understanding of how PA could serve a protective role in the prevention of AD, even in the presence of non-modifiable risk factors. Still, numerous questions remain unanswered which future studies should address. For example, the cross-sectional nature of this study does not enable the establishment of a causal relationship between PA and brain activity. Also, we did not include $\epsilon 4\epsilon 4$ homozygotes, due to the small sample size of this group in our cohort, and therefore we could not explore dosage effects. In this line, a greater sample size would have enabled us to perform a three-way interaction analysis to properly assess the combined moderating effect of *APOE* and age on PA. Finally, although objective measures of PA suppose an improvement with respect to self-reported PA in terms of precision, they do not reflect individual levels of exertion (Gil-Rey et al., 2019), which are more closely related to the physiological response to exercise (Kennedy et al., 2017)

6. Conclusions

The current study provides new insights and builds on previous literature on the *APOE* $\epsilon 4$ modulation of the neural effects of PA. In particular, this study provides evidence that age should be considered in the

attempt to disentangle seemingly mixed results from earlier investigations. Our results show, for the first time, that only young E3/E3 and old E3/E4 carriers present a significant positive relationship between PA and alpha power and that, in the latter group, this increase in alpha power is also associated with better memory performance and gray matter preservation. Above all, this study highlights the importance of implementing public policies that promote physical activity in mid-life and late-life.

7. Declarations

7.1. Ethics approval and consent to participate

The procedure was performed following current guidelines and regulations, and the study was approved by the Hospital Universitario San Carlos Ethics Committee under the code 15/382-E_BS. Every participant signed an informed consent.

7.2. Consent for publication

Not applicable.

7.3. Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

7.4. Competing interests

None of the contributors to this study have any conflict of interest to declare.

7.5. Funding

This study was funded by the Spanish Ministry of Economy and Competitiveness under the Grant PSI2015-68793-C3-1-R [D601] and by the project B2017/BMD-3760 from NEUROCENTRO. Complimentary, it was supported by a predoctoral fellowship from La Caixa Foundation to J.F.L., a post-doctoral fellowship from the Spanish Ministry of Economy and Competitiveness to P.C. (IJC2018-038404-I), predoctoral grant by the Spanish Ministry of Economy (BES-2016-076869) to F.R.T. and a predoctoral grant by the Spanish Ministry of Economy (BES-2016-076869) to A.N.

7.6. Authors' contributions

8. References

- Babiloni, C., Del Percio, C., Lizio, R., Marzano, N., Infarinato, F., Soricelli, A., Salvatore, E., Ferri, R., Bonforte, C., Tedeschi, G., Montella, P., Baglieri, A., Rodriguez, G., Famà, F., Nobili, F., Vernieri, F., Ursini, F., Mundi, C., Frisoni, G.B., Rossini, P.M., 2014. Cortical sources of resting state electroencephalographic alpha rhythms deteriorate across time in subjects with amnesic mild cognitive impairment. *Neurobiol. Aging* 35, 130–142. <https://doi.org/10.1016/j.neurobiolaging.2013.06.019>
- Babiloni, C., Frisoni, G.B., Vecchio, F., Lizio, R., Pievani, M., Cristina, G., Fracassi, C., Vernieri, F., Rodriguez, G., Nobili, F., Ferri, R., Rossini, P.M., 2011. Stability of clinical condition in mild cognitive impairment is related to cortical sources of alpha rhythms: An electroencephalographic study. *Hum. Brain Mapp.* 32, 1916–1931. <https://doi.org/10.1002/hbm.21157>
- Berendse, H., Verbunt, J.P., Scheltens, P., van Dijk, B., Jonkman, E., 2000. Magnetoencephalographic analysis of cortical activity in Alzheimer's disease: a pilot study. *Clin. Neurophysiol.* 111, 604–612. [https://doi.org/10.1016/S1388-2457\(99\)00309-0](https://doi.org/10.1016/S1388-2457(99)00309-0)
- Braak, H., Braak, E., 1991. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol.* 82, 239–259. <https://doi.org/10.1007/BF00308809>
- Brown, B.M., Peiffer, J.J., Taddei, K., Lui, J.K., Laws, S.M., Gupta, V.B., Taddei, T., Ward, V.K., Rodrigues, M.A., Burnham, S., Rainey-Smith, S.R., Villemagne, V.L., Bush, A., Ellis, K.A., Masters, C.L., Ames, D., Macaulay, S.L., Szoek, C., Rowe, C.C., Martins, R.N., 2013. Physical activity and amyloid- β plasma and brain levels: results from the Australian Imaging, Biomarkers and Lifestyle Study of Ageing. *Mol. Psychiatry* 18, 875–881. <https://doi.org/10.1038/mp.2012.107>
- Brown, B.M., Rainey-Smith, S.R., Dore, V., Pfeiffer, J.J., Burnham, S.C., M, L.S., Taddei, K., Ames, D., Masters, C.L., Rowe, C.C., Martins, R.N., Villemagne, V.L., 2018. Self-Reported Physical Activity is Associated with Tau Burden Measured by Positron Emission Tomography. *J. Alzheimer's Dis.* 63, 1299:1305. <https://doi.org/10.3233/JAD-170998>
- Chang, M., Jonsson, P. V, Snaedal, J., Bjornsson, S., Saczynski, J.S., Aspelund, T., Eiriksdottir, G., Jonsdottir, M.K., Lopez, O.L., Harris, T.B., Gudnason, V., Launer, L.J., 2010. The Effect of Midlife Physical Activity on Cognitive Function Among Older Adults: AGES—Reykjavik Study. *JOURNALS Gerontol. Ser. A-BIOLOGICAL Sci. Med. Sci.* 65, 1369–1374. <https://doi.org/10.1093/gerona/glq152>
- Chang, Y.K., Labban, J.D., Gapin, J.I., Etnier, J.L., 2012. The effects of acute exercise on cognitive performance: A meta-analysis. *Brain Res.* 1453, 87–101. <https://doi.org/10.1016/j.brainres.2012.02.068>

- Chen, X., Mobley, W.C., 2019. Alzheimer Disease Pathogenesis: Insights From Molecular and Cellular Biology Studies of Oligomeric A β and Tau Species. *Front. Neurosci.* 13. <https://doi.org/10.3389/fnins.2019.00659>
- Choi, M.R., Kim, J.Y., Yi, E.S., 2018. Development and validation of exercise rehabilitation program for cognitive function and activity of daily living improvement in mild dementia elderly. *J. Exerc. Rehabil.* 14, 207–212. <https://doi.org/10.12965/jer.1836176.088>
- Chomistek, A.K., Yuan, C., Matthews, C.E., Troiano, R.P., Bowles, H.R., Rood, J., Barnett, J.B., Willett, W.C., Rimm, E.B., Bassett, D.R., 2017. Physical Activity Assessment with the ActiGraph GT3X and Doubly Labeled Water. *Med. Sci. Sports Exerc.* 49, 1935–1944. <https://doi.org/10.1249/MSS.0000000000001299>
- Chudyk, A.M., McAllister, M.M., Cheung, H.K., McKay, H.A., Ashe, M.C., 2017. Are we missing the sitting? Agreement between accelerometer non-wear time validation methods used with older adults' data. *Cogent Med.* 4. <https://doi.org/10.1080/2331205X.2017.1313505>
- Corder, E.H., Saunders, A.M., Strittmatter, W.J., Schmechel, D.E., Gaskell, P.C., Small, G.W., Roses, A.D., Haines, J.L., Pericak-Vance, M.A., 1993. Gene Dose of Apolipoprotein E Type 4 Allele and the Risk of Alzheimer's Disease in Late Onset Families. *Science (80-)*. 261, 921–923.
- de Frutos-Lucas, J. De, López-Sanz, D., Zuluaga, P., Rodríguez-Rojo, I.C., Luna, R., María Eugenia, L., Delgado-losada, M.L., Marcos, A., Barabash, A., López-higes, R., Maestú, F., Fernández, A., 2018. Clinical Neurophysiology Physical activity effects on the individual alpha peak frequency of older adults with and without genetic risk factors for Alzheimer ' s Disease: A MEG study. *Clin. Neurophysiol.* 129, 1981–1989. <https://doi.org/10.1016/j.clinph.2018.06.026>
- de Frutos-Lucas, J., Cuesta, P., López-Sanz, D., Peral-Suárez, Á., Cuadrado-Soto, E., Ramírez-Toraño, F., Brown, B.M., Serrano, J.M., Laws, S.M., Rodríguez-Rojo, I.C., Verdejo-Román, J., Bruña, R., Delgado-Losada, M.L., Barabash, A., López-Sobaler, A.M., López-Higes, R., Marcos, A., Maestú, F., 2020. The relationship between physical activity, apolipoprotein E ϵ 4 carriage and brain health. 19 March 2020 PREPRINT (Version 2). [https://doi.org/Available at Research Square DOI:10.21203/rs.2.22674/v1](https://doi.org/Available%20at%20Research%20Square%20DOI:10.21203/rs.2.22674/v1)
- Dimitrova, J., Hogan, M., Khader, P., O'Hora, D., Kilmartin, L., Walsh, J.C., Roche, R., Anderson-Hanley, C., 2017. Comparing the effects of an acute bout of physical exercise with an acute bout of interactive mental and physical exercise on electrophysiology and executive functioning in younger and older adults. *Aging Clin. Exp. Res.* 29, 959–967. <https://doi.org/10.1007/s40520-016-0683-6>
- Ebert, U., Kirch, W., 1998. Scopolamine model of dementia: Electroencephalogram findings and cognitive performance. *Eur. J. Clin. Invest.* 28, 944–949. <https://doi.org/10.1046/j.1365-2362.1998.00393.x>
- Erickson, K.I., Weinstein, A.M., Lopez, O.L., 2012. Physical Activity , Brain Plasticity , and Alzheimer ' s Disease. *Arch. Med. Res.* 43, 615–621. <https://doi.org/10.1016/j.arcmed.2012.09.008>

- Escudero, J., Hornero, R., Abásolo, D., Fernández, A., 2009. Blind source separation to enhance spectral and non-linear features of magnetoencephalogram recordings. Application to Alzheimer's disease. *Med. Eng. Phys.* 31, 872–9. <https://doi.org/10.1016/j.medengphy.2009.04.003>
- Farrer, Lindsay, A., Cupples, L.A., Haines, J.L., Hyman, B., Kukull, W.A., Mayeux, R., Myers, R.H., Pericak-Vance, M.A., Risch, N., van Duijn, C.M., 1997. Effects of Age, Sex, and Ethnicity on the Association Between Apolipoprotein E Genotype and Alzheimer Disease. *JAMA - J. Am. Med. Assoc.* 278, 22–29.
- Fenesi, B., Fang, H., Kovacevic, A., Oremus, M., Parminder, R., Heisz, J.J., 2017. Physical Exercise Moderates the Relationship of Apolipoprotein E (APOE) Genotype and Dementia Risk: A Population-Based Study. *J. Alzheimer's Dis.* 56, 297–303. <https://doi.org/10.3233/JAD-160424>
- Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., Kouwe, A. Van Der, Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A.M., 2002. Whole Brain Segmentation: Neurotechnique Automated Labeling of Neuroanatomical Structures in the Human Brain. *Neuron* 33, 341–355.
- Fornito, A., Zalesky, A., Bullmore, E.T., 2016. *Fundamentals of Brain Network Analysis*. Elsevier, London.
- Förstl, H., Kurz, A., 1999. Clinical features of Alzheimer's disease. *Eur. Arch. Psychiatry Clin. Neurosci.* 249, 288–290. <https://doi.org/10.1007/s004060050101>
- Gaál, Z.A., Boha, R., Stam, C.J., Molnár, M., 2010. Age-dependent features of EEG-reactivity-Spectral, complexity, and network characteristics. *Neurosci. Lett.* 479, 79–84. <https://doi.org/10.1016/j.neulet.2010.05.037>
- Garcés, P., López-Sanz, D., Maestú, F., Pereda, E., 2017. Choice of magnetometers and gradiometers after signal space separation. *Sensors (Switzerland)* 17, 1–13. <https://doi.org/10.3390/s17122926>
- Garcés, P., Vicente, R., Wibral, M., Pineda-Pardo, J. ángel, López, M.E., Aurtenetxe, S., Marcos, A., de Andrés, M.E., Yus, M., Sancho, M., Maestú, F., Fernández, A., 2013. Brain-wide slowing of spontaneous alpha rhythms in mild cognitive impairment. *Front. Aging Neurosci.* 5, 1–7. <https://doi.org/10.3389/fnagi.2013.00100>
- Gil-Rey, E., Maldonado-Martín, S., Gorostiaga, E.M., 2019. Individualized Accelerometer Activity Cut-Points for the Measurement of Relative Physical Activity Intensity Levels. *Res. Q. Exerc. Sport* 90, 327–335. <https://doi.org/10.1080/02701367.2019.1599801>
- Gómez, C., M Pérez-Macías, J., Poza, J., Fernández, A., Hornero, R., 2013. Spectral changes in spontaneous MEG activity across the lifespan. *J. Neural Eng.* 10. <https://doi.org/10.1088/1741-2560/10/6/066006>
- Gutmann, B., Mierau, A., Hülsdünker, T., Hildebrand, C., Przyklenk, A., Hollmann, W., Strüder, H.K., 2015. Effects of physical exercise on individual resting state EEG alpha peak frequency. *Neural Plast.* 2015.

<https://doi.org/10.1155/2015/717312>

Head, D., Bugg, J.M., Goate, A.M., Fagan, A.M., Mintun, M.A., Benzinger, T., Holtzman, D.M., Morris, J.C., 2012. Exercise engagement as a moderator of APOE effects on amyloid deposition. *Arch Neurol* 69, 636–643. <https://doi.org/10.1001/archneurol.2011.845.Exercise>

Herrero, M.J., Blanch, J., Peri, J.M., de Pablo, J., Pintor, L., Bulbena, A., 2003. A validation study of the hospital anxiety and depression scale (HADS) in a Spanish population. *Gen. Hosp. Psychiatry* 25, 277–283.

Horne, J., 2013. Exercise benefits for the aging brain depend on the accompanying cognitive load: Insights from sleep electroencephalogram. *Sleep Med.* 14, 1208–1213. <https://doi.org/10.1016/j.sleep.2013.05.019>

Jack, C.R., Wiste, B.A., H.J., Weigand, S.D., Rocca, W.A., Knopman, D.S., Mielke, M.M., Lowe, V.J., Senjem, M.L., Gunter, J.L., Preboske, G.M., Pankratz, V.S., Vemuri, P., Petersen, R.C., 2014. Age-specific population frequencies of amyloidosis and neurodegeneration among cognitively normal people age 50-89 years: a cross-sectional study. *Lancet Neurol* 13, 997–1005. <https://doi.org/10.3174/ajnr.A1256.Functional>

Jack Jr, C.R., Therneau, T.M., Weigand, S.D., Wiste, H.J., Knopman, D.S., Vemuri, P., Lowe, V.J., Mielke, M.M., Roberts, R.O., Machulda, M.M., Graff-radford, J., Jones, D.T., Schwarz, C.G., Gunter, J.L., Senjem, M.L., Rocca, W.A., Petersen, R.C., 2019. Prevalence of Biologically vs Clinically Defined Alzheimer Spectrum Entities Using the National Institute on Aging–Alzheimer’s Association Research Framework. *JAMA Neurol.* 76, 1174–1183. <https://doi.org/10.1001/jamaneurol.2019.1971>

Kennedy, G., Hardman, R.J., Macpherson, H., Scholey, A.B., Pipingas, A., 2017. How Does Exercise Reduce the Rate of Age-Associated Cognitive Decline? A Review of Potential Mechanisms. *J. Alzheimer’s Dis.* 55, 1–18. <https://doi.org/10.3233/JAD-160665>

Kivipelto, M., Mangialasche, F., Ngandu, T., 2018. Lifestyle interventions to prevent cognitive impairment, dementia and Alzheimer disease. *Nat. Rev. Neurol.* <https://doi.org/10.1038/s41582-018-0070-3>

Kubitz, K.A., Pothakos, K., 1997. Does Aerobic Exercise Decrease Brain Activation? *J. Sport. Exerc. Psychol.* 19, 291–301. <https://doi.org/10.1017/CBO9781107415324.004>

Kuo, T.B.J., Li, J.Y., Shen-Yu Hsieh, S., Chen, J.J., Tsai, C.Y., Yang, C.C.H., 2010. Effect of aging on treadmill exercise induced theta power in the rat. *Age (Omaha).* 32, 297–308. <https://doi.org/10.1007/s11357-010-9143-y>

Li, J.Y., Kuo, T.B.J., Hsieh, S.S.Y., Yang, C.C.H., 2008. Changes in electroencephalogram and heart rate during treadmill exercise in the rat. *Neurosci. Lett.* 434, 175–178. <https://doi.org/10.1016/j.neulet.2008.01.052>

- Lindsay, J., Laurin, D., Verreault, R., Hébert, R., Helliwell, B., Hill, G.B., McDowell, I., 2002. Risk Factors for Alzheimer's Disease: A Prospective Analysis from the Canadian Study of Health and Aging. *Am. J. Epidemiol.* 156, 445–453. <https://doi.org/10.1093/aje/kwf074>
- López-Sanz, D., Bruña, R., de Frutos-Lucas, J., Maestú, F., 2019. Magnetoencephalography applied to the study of Alzheimer's disease. *Prog. Mol. Biol. Transl. Sci.* 165, 25–61. <https://doi.org/10.1016/bs.pmbts.2019.04.007>
- López-Sanz, D., Bruña, R., Garcés, P., Camara, C., Serrano, N., Rodríguez-Rojo, I.C., Delgado, M.L., Montenegro, M., López-Higes, R., Yus, M., Maestú, F., 2016. Alpha band disruption in the AD-continuum starts in the Subjective Cognitive Decline stage: a MEG study. *Sci. Rep.* 6, 37685. <https://doi.org/10.1038/srep37685>
- Luck, T., Luppá, M., Wiese, B., Köhler, M., Jessen, F., Bickel, H., Weyerer, S., Pentzek, M., König, H.-H., Prokein, J., Ernst, A., Wagner, M., Mosch, E., Werle, J., Fuchs, A., Brettschneider, C., Scherer, M., Maier, W., 2014. Apolipoprotein E epsilon 4 genotype and a physically active lifestyle in late life: analysis of gene – environment interaction for the risk of dementia and Alzheimer's disease dementia. *Psychol. Med.* 44, 1319–1329. <https://doi.org/10.1017/S0033291713001918>
- Mattson, M.P., Magnus, T., 2006. Ageing and neuronal vulnerability. *Nat. Rev. Neurosci.* 7, 278–294. <https://doi.org/10.1038/nrn1886>
- Mechau, D., Mücke, S., Weiß, M., Liesen, H., 1998. Effect of increasing running velocity on electroencephalogram in a field test. *Eur. J. Appl. Physiol. Occup. Physiol.* 78, 340–345. <https://doi.org/10.1007/s004210050429>
- Moraes, H., Deslandes, A., Silveira, H., Ribeiro, P., Cagy, M., Piedade, R., Pompeu, F., Laks, J., 2011. The effect of acute effort on EEG in healthy young and elderly subjects. *Eur. J. Appl. Physiol.* 111, 67–75. <https://doi.org/10.1007/s00421-010-1627-z>
- Moretti, D. V., Babiloni, C., Binetti, G., Cassetta, E., Dal Forno, G., Ferreric, F., Ferri, R., Lanuzza, B., Miniussi, C., Nobili, F., Rodriguez, G., Salinari, S., Rossini, P.M., 2004. Individual analysis of EEG frequency and band power in mild Alzheimer's disease. *Clin. Neurophysiol.* 115, 299–308. [https://doi.org/10.1016/S1388-2457\(03\)00345-6](https://doi.org/10.1016/S1388-2457(03)00345-6)
- Mufson, E.J., Counts, S.E., Perez, S.E., Ginsberg, S.D., 2008. Cholinergic system during the progression of Alzheimer's disease: therapeutic implications. *Expert Rev. Neurother* 8, 1703–1718.
- Nolte, G., 2003. The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. *Phys. Med. Biol.* 48, 3637–3652. <https://doi.org/10.1088/0031-9155/48/22/002>

Okonkwo, O.C., Schultz, S.A., Oh, J.M., Larson, J., Edwards, D., Cook, D., Kosciak, R., Gallagher, C.L., Dowling, N.M., Carlsson, C.M., Rowley, H.A., Christian, B.T., Hermann, B.P., Johnson, S.C., Sager, M.A., 2014. Physical activity attenuates age-related biomarker alterations in preclinical AD. *Neurology* 83, 1760.

Oostenveld, R., Fries, P., Maris, E., Schoffelen, J.-M., 2011. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Comput. Intell. Neurosci.* 2011, 156869. <https://doi.org/10.1155/2011/156869>

Osipova, D., Ahveninen, J., Jensen, O., Ylikoski, A., Pekkonen, E., 2005. Altered generation of spontaneous oscillations in Alzheimer's disease. *Neuroimage* 27, 835–841. <https://doi.org/10.1016/J.NEUROIMAGE.2005.05.011>

Podewils, L.J., Guallar, E., Kuller, L.H., Fried, L.P., Lopez, O.L., Carlson, M., Lyketsos, C.G., 2005. Physical Activity, APOE Genotype, and Dementia Risk: Findings from the Cardiovascular Health Cognition Study. *Am. J. Epidemiol.* 161, 639–651. <https://doi.org/10.1093/aje/kwi092>

Poirier, J., Delisle, M.C., Quirion, R., Aubert, I., Farlow, M., Lahiri, D., Hui, S., Bertrand, P., Nalbantoglu, J., Gilfix, B.M., Gauthier, S., 1995. Apolipoprotein E4 allele as a predictor of cholinergic deficits and treatment outcome in Alzheimer disease. *Proc. Natl. Acad. Sci. U. S. A.* 92, 12260–12264. <https://doi.org/10.1073/pnas.92.26.12260>

Poza, J., Hornero, R., Abásolo, D., Fernández, A., García, M., 2007. Extraction of spectral based measures from MEG background oscillations in Alzheimer's disease. *Med. Eng. Phys.* 29, 1073–1083. <https://doi.org/10.1016/j.medengphy.2006.11.006>

Poza, J., Hornero, R., Abásolo, D., Fernández, A., Mayo, A., 2008a. Evaluation of spectral ratio measures from spontaneous MEG recordings in patients with Alzheimer's disease. *Comput. Methods Programs Biomed.* 90, 137–47. <https://doi.org/10.1016/j.cmpb.2007.12.004>

Poza, J., Hornero, R., Escudero, J., Fernández, A., Sánchez, C.I., 2008b. Regional analysis of spontaneous MEG rhythms in patients with Alzheimer's disease using spectral entropies. *Ann. Biomed. Eng.* 36, 141–152. <https://doi.org/10.1007/s10439-007-9402-y>

Prince, M., Wimo, A., Guerchet, M., Ali, G.-C., Wu, Y.-T., Prina, M., 2015. World Alzheimer Report 2015 The Global Impact of Dementia, Alzheimer's Disease International. London.

Rathmann, K.L., Conner, C.S., 2007. Alzheimer's disease: clinical features, pathogenesis, and treatment. *Ann. Pharmacother.* 41, 1499–1504. <https://doi.org/10.1345/aph.140065>

Rovio, S., Kareholt, I., Helkala, E.-L., Viitanen, M., Winblad, B., Tuomilehto, J., Soininen, H., Nissinen, A., Kivipelto, M., 2005. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol* 4, 705–711. [https://doi.org/10.1016/S1474-4422\(05\)70198-8](https://doi.org/10.1016/S1474-4422(05)70198-8)

- Sanchez-Lopez, J., Silva-Pereyra, J., Fernández, T., Alatorre-Cruz, G.C., Castro-Chavira, S.A., González-López, M., Sánchez-Moguel, S.M., 2018. High levels of incidental physical activity are positively associated with cognition and EEG activity in aging. *PLoS One* 13, 1–18. <https://doi.org/10.1371/journal.pone.0191561>
- Schättin, A., Gennaro, F., Egloff, M., Vogt, S., de Bruin, E.D., 2018. Physical activity, nutrition, cognition, neurophysiology, and short-time synaptic plasticity in healthy older adults: A cross-sectional study. *Front. Aging Neurosci.* 10, 1–15. <https://doi.org/10.3389/fnagi.2018.00242>
- Smith, J.C., Nielson, K. a, Woodard, J.L., Seidenberg, M., Rao, S.M., 2013. Physical activity and brain function in older adults at increased risk for Alzheimer’s disease. *Brain Sci.* 3, 54–83. <https://doi.org/10.3390/brainsci3010054>
- Susi, G., Frutos-Lucas, J. De, Niso, G., Ye-chen, S.M., Toro, L.A., Vilca Chino, N.B., Maestú, F., 2019. Healthy and Pathological Neurocognitive Aging: Spectral and Functional Connectivity Analyses Using Magnetoencephalography. *Oxford Res. Encycl. Psychol.* <https://doi.org/10.1093/acrefore/9780190236557.013.387>
- Taulu, S., Simola, J., 2006. Spatiotemporal signal space separation method for rejecting nearby interference in MEG measurements. *Phys. Med. Biol.* 51, 1759–1768. <https://doi.org/10.1088/0031-9155/51/7/008>
- Troiano, R.P, Berrigan, D., Dodd, K.W., Mâsse, L.C., Tilert, T., Mcdowell, M., 2008. Physical activity in the United States measured by accelerometer. *Med. Sci. Sports Exerc.* 40, 181–188. <https://doi.org/10.1249/mss.0b013e31815a51b3>
- Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., Joliot, M., 2002. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15, 273–89. <https://doi.org/10.1006/nimg.2001.0978>
- Van Veen, B.D., van Drongelen, W., Yuchtman, M., Suzuki, A., Veen, B.D. Van, Drongelen, W. Van, Yuchtman, M., Suzuki, A., 1997. Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans. Biomed. Eng.* 44, 867–880. <https://doi.org/10.1109/10.623056>
- Vogt, T., Schneider, S., Brümmer, V., Strüder, H.K., 2010. Frontal EEG asymmetry: The effects of sustained walking in the elderly. *Neurosci. Lett.* 485, 134–137. <https://doi.org/10.1016/j.neulet.2010.09.001>
- Weschler, D., 2008. Wechsler adult intelligence scale–Fourth Edition (WAIS–IV), Fourth Edi. ed. Pearson, San Antonio, TX.
- Woo, M., Kim, S., Kim, J., Petruzzello, S.J., Hatfield, B.D., 2009. Examining the exercise-affect dose-response relationship: Does duration influence frontal EEG asymmetry? *Int. J. Psychophysiol.* 72, 166–

172. <https://doi.org/10.1016/j.ijpsycho.2008.12.003>


Yamazaki, Y., Zhao, N., Caulfield, T.R., Liu, C.C., Bu, G., 2019. Apolipoprotein E and Alzheimer disease: pathobiology and targeting strategies. *Nat. Rev. Neurol.* 15, 501–518. <https://doi.org/10.1038/s41582-019-0228-7>

Yesavage, J.A., Brink, T.L., Rose, T.L., Lum, O., Huang, V., Adey, M., Leirer, V.O., 1982. Development and validation of a geriatric depression screening scale: a preliminary report. . *J. Psychiatr. Res.* 17, 37–49.

Yuri Di Marco, L., Marzo, A., Munoz-Ruiz, M., Ikram, M.A., Kivipelto, M., Ruefenacht, D., Venneri, A., Soininen, H., Wanke, I., Ventikos, Y.A., Frangi, A.F., 2014. Modifiable Lifestyle Factors in Dementia : A Systematic Review of Longitudinal Observational Cohort Studies. *J. Alzheimer’s Dis.* 42, 119–135. <https://doi.org/10.3233/JAD-132225>

Zalesky, A., Fornito, A., Bullmore, E.T., 2010. Network-based statistic: Identifying differences in brain networks. *Neuroimage* 53, 1197–1207. <https://doi.org/10.1016/j.neuroimage.2010.06.041>

Figures



3 Figure 1 presents: The brain regions comprised within the significant cluster (depicted in red, 1A); the evolution
 4 of the cluster size through the different frequency steps (1B, maximum size at 11.75 Hz) and the scatter plots
 5 showing the correlation between the cluster's average power (AvgPow) and total physical activity (TPA) and
 6 each genotype subgroup of the sample (APOE $\epsilon 4$ carriers - APOE $\epsilon 4$ - and non-carriers - APOE $\epsilon 3$ -; 1C), each
 7 age group (younger and older adults, 1D) and all four subgroups (1E).

Figure 1

Significant cluster at 10.75 – 13Hz