

Clinical Utilization of Point-of-care Ultrasound After Training: A Prospective Study

Wan-Ching Lien

National Taiwan University Hospital

Chih-Heng Chang

National Taiwan University Hospital

Kah-Meng Chong

National Taiwan University Hospital

Meng-Che Wu

National Taiwan University Hospital

Cheng-Yi Wu

National Taiwan University Hospital

Hsiu-Po Wang (✉ wanghp@ntu.edu.tw)

National Taiwan University Hospital

Research Article

Keywords: point-of-care ultrasound, post-graduate year resident, global rating

DOI: <https://doi.org/10.21203/rs.3.rs-270802/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: This study aimed to investigate the clinical utilization of PoCUS of the first post-graduate year residents after a PoCUS curriculum.

Methods: It was conducted at the emergency department of the National Taiwan University Hospital between July 2015 and October 2017. Every resident had one-month ED training and a curriculum was implemented during the first week. The post-training objective structured clinical examination (OSCE) global ratings were obtained, as well as the sonographic examinations by the residents. The participants were categorized into 4 groups: group 1 performed PoCUS before and after the curriculum; group 2 performed only after the curriculum; group 3 performed only before the curriculum; the last did not perform any examinations.

Results: Two hundred and thirty-nine residents participated. The median global rating was 4 (interquartile ranges, 4-5). A significantly increasing number of residents integrated PoCUS into patient care (64 vs. 170, $p=0.037$) with acceptable image quality after the curriculum. Nearly 30% of residents did not use PoCUS although no differences existed in the OSCE global ratings. Group 1 performed more cases with better image quality and using ≥ 2 ultrasound applications (8/100 vs. 82/359, $p=0.0009$) after the curriculum. Group 1 performed more cases with better image quality than group 2 after the curriculum. In groups 3 and 4, "no suitable cases" (25/69), and "choosing other imaging priorities" (43/69) were the main feedbacks for not performing PoCUS.

Conclusions: A focused training had a substantial positive impact on the utilization of PoCUS of the residents. The global ratings could not predict whether to integrate PoCUS into clinical practice. Further studies would be needed for the sustained effects of the curriculum and how to encourage these residents in groups 3 and 4 to use PoCUS.

Clinical trial registration: NCT03738033.

Background

Point-of-care ultrasound (PoCUS), referring to "the stethoscope of the 21st century" [1], is defined as ultrasonography performed by the physician at the bedside [2]. It provides real-time, dynamic images that physicians can correlate with the patient's presenting symptoms, serving as a valuable adjunct for patient care [3]. The use of PoCUS has been increasing robustly in emergency medicine (EM) in recent decades. Proliferating PoCUS use and an evolving evidence base for different applications [4] have promoted the incorporation of ultrasound (US) training into the medical education of EM.

PoCUS education is a new scope of EM education. The Accreditation Council for Graduate Medical Education (ACGME) has recommended that all EM residents attain competency in the use of US [5]. Objective structured clinical examination (OSCE) is recommended by the American College of Emergency Physicians (ACEP) policy statement to assess for US competency [6]. However, the sub-competency in the

EM Milestones document suggests that a resident should consistently utilize and integrate appropriate US applications into clinical management [7]. Little has been known about the actual use of PoCUS by the residents [8]. The results of the OSCE would not readily translate to the use of PoCUS in the clinical decision. Whether the seamless penetration into clinical practice could occur after a US curriculum in a high-volume emergency department (ED), is uncertain.

In Taiwan, after completing medical school, graduates must participate in the one-year postgraduate general medicine training as the first postgraduate year (PGY-1) residents before they start a minimum of 3-year residency in a specialty. One-month ED training was included. Since 2013, a novel PoCUS curriculum was started for the PGY-1 residents in the ED of the National Taiwan University Hospital.

This study aimed to investigate the clinical utilization of PoCUS of the PGY-1 residents.

Methods

Study setting and design

The prospective observational study was conducted at the ED of the National Taiwan University Hospital, a tertiary medical center in Taiwan, between July 2015 and October 2017. The ED has an annual uptake of approximately 85,000 patients. The study protocol was approved by the institutional review board of the hospital (201412004RIND) and registered at ClinicalTrials.gov (NCT03738033). Written informed consent was obtained from the participants.

Eight to ten PGY-1 residents attended the ED training every month. Every PGY-1 resident had 18 eight-hour working shifts during their training month and evaluated 10-15 non-critical patients under the supervision of the certified emergency physicians on each shift. A novel PoCUS curriculum including 30-min didactics and 2-hour hands-on training on a live healthy model volunteer was implemented for the PGY-1 residents during the first week. Because abdominal discomfort was the leading ED symptoms, the content of the curriculum included the extended focused assessment of sonography for trauma (eFAST) to detect intraperitoneal fluid, pericardial effusion, pleural effusion, and pneumothorax, and sonography for urinary tract to detect hydronephrosis, gall bladder to detect acute cholecystitis, and abdominal aorta to detect abdominal aortic aneurysm. The ratio of the instructor to participant is less than 1:5. The instructors are expert sonographers, board-certified by the Taiwan Society of Ultrasound in Medicine, and had over 10 years of experience in sonographic examinations. A US machine (SSA-660A, Canon, Japan) equipped with a 2-5 MHz curvilinear transducer was used for training.

The PGY-1 residents completed the curriculum and received an OSCE for post-curriculum assessment immediately. It consisted of standardized questions for image acquisition and interpretation with points for technique, image quality, and correct interpretation of anatomy of FAST, kidney, gall bladder, and aorta. A global rating score using a Likert 5-point scale (unsatisfactory=1, needs improvement=2, satisfactory=3, high satisfactory=4, outstanding=5)[9] was given by the instructor at the scene and by the

other instructor that did not involve in training through video review independently. The faces of the PGY-1 residents in the video were covered. The score given by the two instructors was averaged.

There were two US machines (SSA-550A, SSA-660A, Canon, Japan) equipped with 2-5 MHz curvilinear transducers kept ready in use in the ED clinics. Also, the reporting documentation was put in a plastic bag, along with the machine. The sonographic examinations performed by the PGY residents were obtained at the end of the one-month ED training, including indication, scanning targets, sonographic findings, sonographic diagnosis, and management. The accuracy of sonographic diagnosis was defined as the agreement between the sonographic diagnosis made by the PGY and by the attending physician. The images were reviewed by another two instructors not involving in training blindly and independently. The quality of the images was categorized using a 5-point Likert rating scale. Point 1 indicated no recognizable structures, no objective data can be gathered; point 2 indicated minimally recognizable structures but insufficient for diagnosis; point 3 indicated minimal criteria met for diagnosis, recognizable structures but with some technical or other flaws; point 4 indicated minimal criteria met for diagnosis, all structures imaged well and diagnosis easily supported; point 5 indicated minimal criteria met for diagnosis, all structures imaged with excellent image quality and diagnosis completely supported [10]. Another instructor not responsible for the curriculum would interview the PGY-1 residents who did not perform clinical sonographic examinations after the curriculum and their feedbacks was obtained.

Selection of Participants

The PGY-1 residents attending the curriculum were included. Those not attending the curriculum were excluded.

Data collection

The demographic data of the PGY-1 residents were obtained, including age, gender, and prior US experience. The global ratings of the OSCE were collected, as well as the sonographic examinations including indication, scanning targets, sonographic findings, sonographic diagnosis on shifts. Based on the clinical sonographic examinations, the residents could be categorized into 4 groups. Group 1 indicated the residents performed PoCUS before and after the curriculum; Group 2 indicated that those performed PoCUS only after the curriculum; Group 3 indicated that those performed PoCUS only before the curriculum; the last indicated those did not perform any examinations. The scanning targets were categorized into a single application (ex, FAST) and more than 2 applications (ex, FAST+gall bladder) according to the ACEP statement [6].

Outcomes

The primary outcome was the utilization of PoCUS of the PGY-1 residents during clinical practice.

Statistical analysis

All data were analyzed by SAS software (SAS 9.4, Cary, North Carolina, USA). Categorical data were expressed in counts and proportions, while continuous data were expressed in medians and interquartile ranges (IQRs). Categorical variables were compared using a Chi-square test and ANOVA. Continuous variables were examined using Wilcoxon's rank-sum test. Intraclass correlation (ICC) with one-way random effects was used to assess inter-rater reliability for those global rating scores and imaging quality scores by two evaluators.

The linear regression models were applied to identify the factors associated with numbers and the accuracy of the sonographic examinations after the curriculum. The polytomous regression models were applied to investigate the factors associated with the image quality of the sonographic examinations after the curriculum. The covariates included age, gender, prior US experience, and global ratings. Also, the factors associated with the global ratings were investigated using the polytomous regression models. The covariates included age, sex, and prior experience. A p-value of less than 0.05 was considered statistically significant.

Results

During the study period, 239 PGY-1 residents attended one-month ED training, and all of them (100%) participated in this study and completed the OSCE. A total of 901 sonographic examinations were collected and reviewed. There was a good ICC coefficient of 0.86 (95% CI, 0.68-0.96) for the OSCE global rating score and 0.84 (95% CI, 0.63-0.95) for the image quality of the sonographic examinations.

Table 1 lists the demographic data of residents. The majority were novice sonographers. The median OSCE global rating score was 4. An increasing number of PGY-1 residents integrate PoCUS into patient care on their shifts after the curriculum (64 vs. 170, $p=0.037$). Also, more increasing case numbers with better image quality and more accurate diagnoses were noted after the curriculum (Table 2). After adjusting age, gender, and prior US experience, a higher global rating (coefficient, 1.38 ± 1.16 , $p=0.03$) was significantly associated with the better image quality of the sonographic examinations after the curriculum. However, there was no impact on the numbers of the cases and the accuracy.

Based on the clinical sonographic examinations, the participants were categorized into 4 groups (Table 3). Among the groups, there were no differences in the global ratings. Group 1 residents performed more cases with better image quality after the curriculum, compared with those before the curriculum. Also, group 1 residents performed the examinations using ≥ 2 US applications after the curriculum (before vs. after, 8/100 vs. 82/359, $p=0.0009$) (Table 4). Comparing with the sonographic examinations after the curriculum in groups 1 and 2, group 1 performed more cases ($p<0.0001$) with better image quality ($p=0.03$) although the accuracy was unremarkable. Tables 4 and 5 list the US indication and scanning targets of the sonographic examinations in groups 1 and 2.

Sixty-nine residents in groups 3 and 4 did not perform sonographic examinations after the curriculum. "No suitable cases for examinations" (25/69), and "having other imaging priorities" (43/69) were the main feedbacks of these residents.

Discussion

While clinical integration is essentially addressed during PoCUS education, little has been known about the actual use of PoCUS by the residents after training [8]. This study provides evidence regarding the real US performance of the inexperienced PGY-1 residents after a short US curriculum. The results showed the US curriculum had a substantial effect on increasing the clinical integration of PoCUS. A significantly increased number of PGY-1 residents used PoCUS during clinical practice after the curriculum. Also, the global ratings had no obvious correlation to the clinical integration of PoCUS. Nearly 30% of them did not use PoCUS although their OSCE performance was good.

Time constraints would be a barrier for ED medical education. Previous studies suggested that short courses of focused US training were of acceptable quality, even in paramedics with no prior ultrasound experience [11-13]. Also, US skills would decay at different times for different organ images such as a significant decline of the ability to identify A-lines of lung over time, but no decline in cardiac image recognition [14]. The content of the US curriculum was designated as the combination of unique abdominal US techniques because abdominal disorders were frequently encountered in our ED.

The introduction of the OSCE showed comparable improvements in trainee performance and understanding feeling confident in their ability to utilize US [7]. There is evidence that global rating scales capture diverse levels of proficiencies better than checklists, and are easy for examiners to use [15]. Also, global ratings were shown to yield reliable data when compared with checklist scores for the assessment of residents in the field of medicine [13, 16]. The skills needed for US are complex and include hand-to-eye coordination, image optimization, and image interpretation [17], and clinical correlation. Therefore, this study used global ratings for OSCE performance instead of checklist scores due to the psychometric properties [16] and higher levels of expertise on US performance.

There were 4 different patterns of the behavior of doctors after US training. This short US curriculum helped inexperienced doctors to integrate PoCUS into clinical practice, especially in groups 1 and 2. The residents used self-formatted combinations of US techniques that the FAST and the renal US were the most common contents. The results showed more than 70% of residents had a desire to change the behavior to be willing to integrate PoCUS into daily practice. Although residents in groups 1 and 2 had progressed, group 1 residents seemed to be more active and motivated in performing sonographic examinations, compared with those in group 2.

The ED is a unique learning environment different from any other setting in the hospital. The high volume and acuity of patients in the ED provide opportunities to repeatedly practice and to facilitate the accumulation of clinical experience for young staffs, crossing the gap from the level 2 (learning level) to the level 3 and 4 (behavior and result levels) easily [18]. However, the characteristics of ED patients that each resident encountered are diverse and varied. The time spent with an individual patient is limited. In our ED, the PGY-1 residents could evaluate 200-250 patients during the training month. Although the number of scanning cases performed by the PGY-1 residents increased after the curriculum, the median

was 3 cases. It implied that the PGY-1 residents used PoCUS in highly selected, maximally beneficial patients.

The potential barriers exist for utilizing PoCUS in clinical settings. The frequent challenges are US equipment and lack of confidence or experience in clinical integration [8, 19]. Also, the lack of evidence for clinical utility for certain patients, a trade-off between performing US and other clinical processes, and the inability to use US for decision-making were reported [8, 20]. In our study, the facility was settled that the US machines were easily accessible. All of the sonographic examinations were under the supervision of the certified ED physicians. The working climate was designated to be friendly for using US, but some barriers still existed. As the feedbacks from groups 3 and 4, the residents could choose other imaging modalities such as computed tomography or magnetic resonance imaging for patient care, whatever autonomously or by the attending physician's suggestion. It implied the gap between PoCUS and comprehensive imaging modalities would be hardly crossed in a time-limited and stressful environment. Moreover, how to encourage groups 3 and 4 residents to use PoCUS in clinical practice would be an important issue in future studies.

This study had some limitations. First, it was conducted in an academic medical center with active US training. External generalization would be further explored. Although the distribution of ED patients varies from hospital to hospital, however, this study provided novel evidence regarding how inexperienced residents integrate PoCUS in their primary care after a tailored educational curriculum. The design and contents of the curriculum were based on the frequently encountered complaints in our ED, and more than 70% of residents used the techniques in their clinical practice. Second, this study investigated the short-term effects of the US curriculum. Insights into long-term retention of US skills and subsequent time to skill degradation were not evaluated. Future longitudinal studies would be needed for these residents for their specialties after one-year general medicine training, especially in groups 3 and 4. Whether the specialty would have an impact on the utilization of PoCUS was wondered. Third, the sonographic examinations were based on written documentation. The percentage of the residents utilizing US would be under-estimated if they performed the sonographic examinations but the results couldn't be seen in the paper records or the electronic medical records. Fourth, the change in the management plan after PoCUS was not evaluated in this study because the plan was influenced by the sonographic findings and made by the attending physician. Last, a more structured survey for groups 3 and 4 would be needed for a more detailed understanding of the reasons for not using PoCUS.

Conclusions

A focused PoCUS training had a substantial positive impact on the utilization of PoCUS of PGY-1 residents. The global ratings could not predict whether to integrate PoCUS into clinical practice. Although the working climate was designated to be friendly for utilizing US, nearly 30% of residents did not use PoCUS. Further studies would be needed for the sustained effects of the curriculum and how to encourage these residents in groups 3 and 4 to use PoCUS.

Abbreviations

Point-of-care ultrasound=PoCUS;

First post-graduate year=PGY-1;

Emergency department=ED;

Emergency medicine=EM

Ultrasound=US;

Accreditation Council for Graduate Medical Education=ACGME;

Objective structured clinical examination=OSCE;

American College of Emergency Physicians=ACEP;

National Taiwan University Hospital=NTUH;

Extended focused assessment of sonography for trauma=eFAST;

Intraclass correlation=ICC.

Declarations

Ethics approval and consent to participate: The study was approved by the institutional review board of the Research Ethics Committee of the National Taiwan University Hospital (201412004RIND). All methods were performed in accordance with the relevant guidelines and regulations by the Research Ethics Committee of the National Taiwan University Hospital. Written informed consent was obtained from the participants.

Consent for publication: Not Applicable.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Competing interests: The authors declare that they have no competing interests.

Funding: no funding.

Authors' contributions: WC and HP conceived the study and designed the trial. WC, CH, KM, MC, CY, and HP acquisition of the data. WC and HP analysis and interpretation of the data. WC drafted the manuscript, and all authors contributed substantially to its revision. HP critically revised the manuscript

for important intellectual content and takes responsibility for the paper as a whole. All authors read and approved the final manuscript.

Acknowledgments: Not applicable.

Meetings or presentations :none

Grant or financial support: none

Conflicts of interest : All authors declared no conflicts of interest.

Acknowledgments: None.

References

1. Sekiguchi H: **Tools of the Trade: Point-of-Care Ultrasonography as a Stethoscope.***Semin Respir Crit Care Med* 2016, **37**(1):68-87.
2. Moore CL, Copel JA: **Point-of-care ultrasonography.** *N Engl J Med* 2011, **364**(8):749-757.
3. Weile J, Frederiksen CA, Laursen CB, Graumann O, Sloth E, Kirkegaard H: **Point-of-care ultrasound induced changes in management of unselected patients in the emergency department - a prospective single-blinded observational trial.***Scand J Trauma Resusc Emerg Med* 2020, **28**(1):47.
4. ACEP: **Ultrasound Guidelines: Emergency, Point-of-Care and Clinical Ultrasound Guidelines in Medicine.***Ann Emerg Med* 2017, **69**(5):e27-e54.
5. Beeson MS, Carter WA, Christopher TA, Heidt JW, Jones JH, Meyer LE, Promes SB, Rodgers KG, Shayne PH, Wagner MJ *et al*: **Emergency medicine milestones.***J Grad Med Educ* 2013, **5**(1 Suppl 1):5-13.
6. ACEP: **Emergency ultrasound guidelines.***Ann Emerg Med* 2009, **53**(4):550-570.
7. Nelson M, Abdi A, Adhikari S, Boniface M, Bramante RM, Egan DJ, Fields JM, Leo MM, Liteplo AS, Liu R *et al*: **Goal-directed Focused Ultrasound Milestones Revised: A Multiorganizational Consensus.***Acad Emerg Med* 2016, **23**(11):1274-1279.
8. Haney RM, Halperin M, Diamond E, Ratanski D, Shokoohi H, Huang C, Liteplo AS: **Clinical Integration of Point-of-care Ultrasound by Emergency Medicine Residents: A Single-center Mixed-methods Study.***AEM Educ Train* 2020, **4**(3):212-222.
9. Dinh VA, Frederick J, Bartos R, Shankel TM, Werner L: **Effects of ultrasound implementation on physical examination learning and teaching during the first year of medical education.***J Ultrasound Med* 2015, **34**(1):43-50.
10. ACEP: **Emergency ultrasound standard reporting guidelines.** 2018.
11. Rooney KP, Lahham S, Anderson CL, Bledsoe B, Sloane B, Joseph L, Osborn MB, Fox JC: **Pre-hospital assessment with ultrasound in emergencies: implementation in the field.** *World J Emerg Med* 2016, **7**(2):117-123.

12. Medico MD, Altieri A, Carnevale-Maffè G, Formagnana P, Casella F, Barchiesi M, Bergonzi M, Vattiato C, Casazza G, Cogliati C: **Pocket-size ultrasound device in cholelithiasis: diagnostic accuracy and efficacy of short-term training.***Intern Emerg Med* 2018, **13**(7):1121-1126.
13. Breslin R, Collins K, Cupitt J: **The use of ultrasound as an adjunct to peripheral venous cannulation by junior doctors in clinical practice.***Med Teach* 2018, **40**(12):1275-1280.
14. Rappaport CA, McConomy BC, Arnold NR, Vose AT, Schmidt GA, Nassar B: **A Prospective Analysis of Motor and Cognitive Skill Retention in Novice Learners of Point of Care Ultrasound.** *Crit Care Med* 2019, **47**(12):e948-e952.
15. Rajiah K, Veettil SK, Kumar S: **Standard setting in OSCEs: a borderline approach.***Clin Teach* 2014, **11**(7):551-556.
16. Regehr G, MacRae H, Reznick RH, Szalay D: **Comparing the psychometric properties of checklists and global rating scales for assessing performance on an OSCE-format examination.** *Acad Med* 1998, **73**(9):993-997.
17. Nicholls D, Sweet L, Hyett J: **Psychomotor skills in medical ultrasound imaging: an analysis of the core skill set.** *J Ultrasound Med* 2014, **33**(8):1349-1352.
18. Kirkpatrick DL, Kirkpatrick JD: **Evaluating training programs: The four levels**, 3rd edn. San Francisco: Berrett-Koehler Publication; 2006.
19. Bhagra A, Tierney DM, Sekiguchi H, Soni NJ: **Point-of-Care Ultrasonography for Primary Care Physicians and General Internists.***Mayo Clin Proc* 2016, **91**(12):1811-1827.
20. Schnittke N, Damewood S: **Identifying and Overcoming Barriers to Resident Use of Point-of-Care Ultrasound.** *West J Emerg Med* 2019, **20**(6):918-925.

Tables

Table 1. The characteristics of the post-graduate year (PGY) residents.

Characteristics	PGY (n=239)
Age, years*	26 (26)
Male, n	166 (69%)
Prior US [†] experience	
≤10 scans	224 (94%)
> 10 scans	15 (6%)
OSCE global rating score*	4 (4-5)
Sonographic examinations during clinical practice	
Before the US curriculum	64 (27%)
After the US curriculum	170 (71%)

*Presented as median (interquartile ranges, IQRs).

[†]US, ultrasonography; OSCE, Objective Structured Clinical Examination.

Table 2. The sonographic examinations performed clinically by the first post graduate year (PGY-1) residents.

Variables	Before the Curriculum (129 cases)	After the curriculum (772 cases)	p-Value
Cases per resident, n*	0 (0-1)	3 (0-5)	<0.0001
Image quality*	3 (2-3)	3 (3)	<0.0001
Accuracy, %	100 (100)	100 (100)	<0.0001

*Presented as median (interquartile ranges, IQRs).

Table 3. The sonographic examinations during the clinical practice of the first post-graduate year (PGY-1) residents.

Group	OSCE global ratings**†	Before the curriculum	After the curriculum	p-Value‡
Group 1 (n=52)	4 (4-5)			
Cases, n*		1 (1-3)	5 (2-8.5)	<0.0001
Imaging quality*		3 (2-3)	3 (3)	<0.0001
Accuracy*		100 (100)	100 (100)	0.379
Group 2 (n=118)	4 (4)			
Cases, n*		-	3 (1-5)	
Imaging quality*		-	3 (3)	
Accuracy*		-	100	
Group 3 (n=12)	4.5 (4-5)			
Cases, n*		1.5 (1, 2.5)	-	
Imaging quality*		3 (2-3)	-	
Accuracy		100	-	
Group 4 (n=57)	4 (4-5)			
Cases, n		-	-	
Imaging quality		-	-	
Accuracy		-	-	

*Presented as median (interquartile ranges, IQRs).

†OSCE, Objective Structured Clinical Examination.

‡Comparing between the performance before and after the curriculum in the group 1.

Table 4. The sonographic examinations performed by group 1 residents clinically.

Indications	Before the curriculum (100 cases)	After the curriculum (359 cases)
Abdominal pain	14	95
Scanning targets		
FAST*	3 (1) [†]	12
GB*	4	27 (4)
Kidney	2	15 (1)
FAST+GB	1	3 (1)
FAST+kidney+aorta	2 (1)	6
FAST+bowel	2 (2)	2 (1)
FAST+aorta	-	2
FAST+kidney	-	5
FAST+liver	-	2
Kidney+aorta	-	5
Kidney+GB	-	7
GB+aorta	-	2
GB+liver	-	2 (1)
FAST+GB+aorta	-	4 (1)
Kidney+GB+aorta	-	1
Abdominal distention	26	61
Scanning targets		
FAST	6	17
Kidney	20 (2)	22 (1)
FAST+kidney	-	20
FAST+bowel	-	1
FAST+GB	-	1 (1)
Fever	5	26
Scanning targets		
FAST	-	5

GB	-	2
Kidney	5	10
Liver	-	1
FAST+kidney	-	4
FAST+GB	-	1
Kidney+liver	-	2
FAST+kidney+GB	-	1
Flank pain	26	78
Scanning targets		
Kidney	23 (3)	71 (5)
FAST+kidney	1	4
Kidney+aorta	1	2
FAST+kidney+aorta	1	1
FAST+kidney+GB	-	1
Hematuria	7	26
Scanning targets		
Kidney	7 (1)	23
FAST+kidney	-	3
Dysuria/oliguria	4	15
Scanning targets		
Kidney	4	15 (1)
Dyspnea	6	15
Scanning targets		
FAST	6	15
Trauma	4	26
Scanning targets		
FAST	4	24
FAST+kidney	-	2
Miscellaneous	8	17 (2)

*FAST, focused assessment of sonography for trauma; GB, gall bladder.

†presented as number (inaccurate case number).

Table 5. The sonographic examinations performed by group 2 residents clinically.

Indications	After the curriculum (413 cases)
Abdominal pain	82
Scanning targets	
FAST*	15
GB*	14 (1) [†]
Kidney	22 (1)
FAST+GB	2
FAST+bowel	1 (1)
FAST+aorta	1
FAST+kidney	8
FAST+liver	1
Kidney+aorta	1
Kidney+GB	2
GB+aorta	1
FAST+GB+aorta	8 (1)
FAST+kidney+GB	5 (3)
FAST+kidney+GB+aorta	1
Abdominal distention	73
Scanning targets	
FAST	28
Kidney	37 (1)
FAST+kidney	2
FAST+GB	2
Kidney+GB	1 (1)
FAST+kidney+GB	2
Kidney+GB+aorta	1
Fever	30

Scanning targets	
FAST	9
Kidney	17 (1)
Liver	1
GB+liver	1
FAST+aorta+bowel	1
Flank pain	97
Scanning targets	
Kidney	85 (8)
FAST+kidney	4
Kidney+aorta	6
FAST+kidney+aorta	1 (1)
FAST+kidney+GB+aorta	1 (1)
Hematuria	11
Scanning targets	
Kidney	9
FAST+kidney	2
Dysuria/oliguria	15
Scanning targets	
Kidney	15
Dyspnea	11
Scanning targets	
FAST	11
Trauma	52
Scanning targets	
FAST	45
FAST+kidney	7 (1)

*FAST, focused assessment of sonography for trauma; GB, gall bladder.

†presented as number (inaccurate case number).