

Effect of improved cookstove intervention on childhood acute lower respiratory infection in Northwest Ethiopia: Cluster randomized controlled trial

CURRENT STATUS: Under Review

 Implementation Science  BMC

Mesafint Molla Adane, Getu Degu Alene, Seid Tiku Mereta, Kristina L Wanyonyi

Mesafint Molla Adane
Bahir Dar University

 Mesafint.Molla@bdu.edu.et **Corresponding Author**
ORCID: <https://orcid.org/0000-0002-9453-5027>

Getu Degu Alene
Bahir Dar University

Seid Tiku Mereta
Jimma University

Kristina L Wanyonyi
Queen Mary University of London



Prescreen

[10.21203/rs.3.rs-28518/v1](https://doi.org/10.21203/rs.3.rs-28518/v1)

Subject Areas

Health Economics & Outcomes Research* ***Epidemiology**

Keywords

Acute lower respiratory infection, biomass-fueled, childhood, household air pollution, improved baking stove, improved cookstove

Abstract

Background

Household air pollution exposure is linked with over 3.5 million premature deaths every year, ranking highest among environmental risk factors globally. Children are uniquely vulnerable and sensitive to the damaging health effects of household air pollution which includes childhood acute lower respiratory infection. The use of improved cookstoves has been widely encouraged to reduce these health burdens. It is, however not clear whether it is possible to prevent household air pollution-related disease burdens with locally made improved cookstoves, and the evidence regarding the health effect of improved cookstove interventions among children still attracts wide debate. Therefore, we investigated the child health effect of improved baking stove intervention compared with the continuation of the customary open burning traditional baking stove.

Methods

A cluster randomized controlled trial was conducted to assess the health effect of improved baking stove intervention. A total of 100 clusters were randomly allocated to both arms at a 1:1 ratio, and intervention was delivered to all households allocated into the intervention arm. A total of four follow-up visits were carried out within one year at approximately 3-month intervals immediately after the delivery of the intervention. The intervention effect was estimated using odds ratio as a measure of effect following a Generalized Estimating Equations modeling approach among the intention-to-treat population.

Results

A total of 5508 children were enrolled in the study across 100 randomly selected clusters in both arms, among which data were obtained from a total of 5333 participants for at least one follow-up visit which establishes the intention-to-treat population dataset for analysis. The intervention was not found to have a statistically significant effect on the outcome with an estimated odds ratio of 0.95 (95% CI: 0.89–1.02) compared with the continuation of an open burning traditional baking stove method. Harms: There was no difference in the overall incidence rates of cooking-related burns between arms with an incidence rate ratio of 0.80 (95% CI: 0.53–1.21), and there were no grounds to stop the trial early for adverse events.

Conclusions

We found no evidence that an intervention comprising biomass-fuelled improved baking stove reduced the risk of childhood acute lower respiratory infection compared with the continuation of an open burning traditional baking stove method. Therefore, effective cooking solutions are needed to avert the adverse health effect of household air pollution, particularly, childhood acute lower respiratory infection.

1. Introduction

Household air pollution (HAP) exposure from cooking with solid fuels is linked with over 3.5 million premature deaths every year; including half a million deaths among children under the age of 5 years from pneumonia worldwide, and it is ranked highest among the environmental risk factors globally (1); predominantly, in Sub-Saharan Africa countries (2). Its burden is very high in the Sub-Saharan Africa region due to the widespread use of polluting household cooking technologies (3). Although HAP affects all stages of life (4), children are uniquely vulnerable and sensitive to the damaging health effects of HAP (5) due to both their physiology as well as to the extent of their exposure (6). There is substantial evidence that HAP exposure from solid-fuel use increases the risk of childhood respiratory infections (7) such as childhood acute lower respiratory infection (ALRI) (8).

The improved cookstove (ICS) intervention is a common practice to reduce the burden of HAP-related diseases (4). It is, however, unclear as to whether it is possible to prevent HAP-related disease burdens with locally made biomass-fuelled ICSs (9), and the evidence on the health effect of ICS interventions is still open to questions. In

particular, the evidence regarding the effect of biomass-fuelled IBS intervention on childhood ALRI is still debatable. For example, a study based on a field experiment in Guatemala reported that a 63.3% reduction in respiratory symptoms among children following wood-fueled ICS intervention (10), and another randomized controlled trial reported a significant risk reduction in childhood pneumonia following ICS intervention with a rate ratio of 0.82 (95% CI: 0.70, 0.98) among intervention group in Guatemala (11). In contrast,

based on experimental evidence, a recent policy research working paper reported no evidence that “Mirt” IBS intervention reduced the risk of respiratory symptoms among older children in rural Ethiopia (12). Correspondingly, a recent ICS trial found no evidence that biomass-fuelled ICS intervention reduced the risk of pneumonia among children in rural Malawi with an incidence rate ratio of 1.01 (95% CI: 0.91, 1.13) (13). In general, several recent systematic reviews concluded that evidence concerning the health effect of ICS interventions in low-and-middle-income countries (LMICs) is still sparse, heterogeneous, and inconclusive (13–15).

In Ethiopia, despite some pilot trial efforts on HAP and respiratory symptoms reduction (12, 16) and energy efficiency tests (17), trial evidence on the health impact of ICS intervention is negligible, and recent systematic reviews on this subject matter reported that such type of evidence is lacking and should be a research priority area in Ethiopia (17–19). In addition, the ICS trial evidence from other LMICs (11, 13) cannot be directly generalized to the Ethiopian perspective from a relevance point of view, such as the particular variation of cooking device requirements and trial stove designs (20). Beyond the debatable biomass-fuelled ICS trial reports worldwide (11, 13), little attention has been paid so far to the health effect of biomass-fuelled ICS in Ethiopia.

The health effect of the biomass-fuelled “Mirt” improved baking stove (IBS), is of keen interest, as it is a well-known commercially distributed type of IBS in Ethiopia (21, 22). Research into the health effect of the “Mirt” stove had not been satisfactorily dealt with in earlier studies (12) emphasizing the need for further investigation on most vulnerable populations such as children who are often exposed to this intervention. More precisely, a crucial question whether the Ethiopian IBS intervention leads to a decrease in the occurrence of childhood acute lower respiratory infection (ALRI) must be answered through further trial investigation as recommended by a previous study (23). Thus, given the high magnitude of HAP problems in Ethiopia (19, 24, 25) and lack of IBS trials (17–19), there is a need to test the health effect of IBS intervention that could potentially reduce the negative health impacts of HAP.

Because of that, we conducted a community-level cluster randomized controlled trial to assess the effect of biomass-fuelled IBS intervention on childhood ALRI at the individual child level in Northwest Ethiopia compared with the continuation of an open burning traditional baking stove (TBS). The major rationale for adopting a cluster-randomized trial (CRT) design was to prevent contamination (26) or unintentional spill-over of intervention effects from one treatment group to another of the trial if individual household randomization was used, as children would inevitably be exposed to the risk or intervention in neighbor households. The other rationales were to increase administrative effectiveness, minimize costs, and to eliminate potential ethical problems (27).

The benefits of the study go beyond addressing an important research gap. The benefits of this study can be grouped into three categories; population health, policy, and economical benefits. When considering population health, since the primary outcome sought is a reduction in childhood ALRI from an effective biomass smoke exposure reduction intervention, it would achieve a high health impact on most vulnerable children living in developing countries like Ethiopia. The study would also achieve a high policy and operational benefits by generating new evidence needed by local, national and international policy-makers and donors regarding health effects seen when households adopt IBS technology as well as it can also inform governments in LMICs how to allocate the limited resources to focus their efforts on the most effective strategies to improve health.

2. Materials And Methods Of The Trial

2.1. Study locations and context

Ethiopia is located in the Northeastern part of Africa, known as the Horn of Africa, and occupies an area of 1.1 million square kilometers ranging from 4,620 meters above sea level at Ras Dashen Mountain to 148 meters below sea level at the Danakil depression (28). It possesses three major topographic-induced climatic zones, the hot lowlands (“*Kolla*”) located below 1500, the temperate (“*Wayna Dega*”) which range 1500–2400, and the cool temperate highlands (“*Dega*”) located above 2400 meter above sea level (28, 29). The mean annual temperature is around 15–20 °C and 25–30 °C for highlands and lowlands respectively (29).

The location and context of the specific study area, this trial was conducted in a low-income rural community of the Mecha Health and Demographic Surveillance System (MHDSS) site. MHDSS site is a field research center established in 2013 by Bahir Dar University to conduct and support postgraduate level studies in the region. It is located 525 km away from the capital city of Ethiopia, Addis Ababa, towards Northwest and 40 km far away from the capital city of Amhara Regional State, Bahir Dar. According to the official population profile report of MHDSS, the study area comprises 132 clusters/ “*Gots*” with a total of 65, 086 populations within 20631 households at the end of 2016. Out of which children less than 4 years old were accounted for 13.3% of the total population, and biomass-fuelled open burning traditional baking stove (TBS) method was the major (94.5%) household energy source for baking purposes in the study locality (23).

2.2. Trial design

A community-level cluster randomized controlled trial with two arms of equal size was used to compare the effect of biomass-fuelled IBS intervention with TBS on childhood ALRI at an individual child level. Clusters were the small villages, termed as “*Gots*” in Amharic (both national & local language), are the lowest administrative units in Ethiopia, and used as the smallest unit of enumeration areas by the Ethiopian national census authority. Each cluster (“*Got*”) was comprised about 55 eligible children on average, and all eligible children were enrolled as control and intervention households for baseline and repeated follow-up visits approximately every 3 months for one year after receiving the intervention. With this design, childhood ALRI outcome was measured before installation of IBS, and again in the same households, 4 times after the intervention households received the IBS. The households with TBS method were served as a control arm.

2.3. Eligibility criteria

All clusters/ “*Gots*” under MHDSS site were eligible for participation in the trial; and to ensure a minimum of one year repeated visit for longitudinal data collection before the child’s 5th birthday, all households with at least one less than 4 years old child and who were exclusive users of TBS were eligible for participation in the trial. Merely households who did not have any enclosed baking quarter (kitchen) structure and children who were born during the course of the study were excluded.

2.4. Sample size determination

To estimate the effect of IBS intervention on child ALRI risk reduction over one year follow-up period, the sample size was calculated by applying the two-sample comparison of proportions formula using STATA. We calculated the required sample size by assuming childhood ALRI prevalence of 21% from a previous study in rural Ethiopia (30) and 0.05 two-tailed alpha to detect a 25% reduction in the prevalence of childhood ALRI with a power of 80%. Accordingly, the estimated sample size (n_1) was 891 children per arm with equal numbers to both arms under individual randomization.

However, since this trial was randomized the intervention over clusters instead of individual households, the sample size was calculated using the standard formulae for unequal size cluster randomized controlled trials as $n_c = n_1 [1 + (m' - 1) ICC]$ (31); where, n_c = sample size with cluster randomization, n_1 = sample size under individual randomization, m' = an average number of eligible children within each cluster, and ICC = intra-cluster correlation coefficient for cluster level ICC of childhood ALRI. Thus, the required sample size was 2068 per arm under equal allocation by assuming an ICC value of 0.03; which is an estimated value for child pneumonia outcome in the cookstove intervention study (13), and an average number of eligible children of 55 within each cluster from the updated data of MHDSS. The number of clusters (K) required in each arm for unequal cluster sizes was also determined using the formula; $K = n_1 [1 + ((CoV^2 + 1)m' - 1) ICC] / m'$ (31); where, CoV = coefficient

of variation. The number of clusters per arm became 40, and this was caused to increase the sample size to 4400 (2200 per arm) by considering an effective ICC value of 0.03 for cluster-level ICC value of childhood ALRI, a CoV value of 0.3 for cluster size and an average number of eligible children of 55 within each cluster.

Finally, considering the nature of cluster sampling, we added 25% of the sample to account for lost to follow-up (LTF) and any other unpredictable events in the field (32, 33), the required sample size was increased to about 2750 within 50 clusters per arm, which would provide a total of 100 clusters containing about 5500 eligible children randomized in equal numbers to both arms.

2.5. Randomization and Masking

Clusters were randomly allocated to intervention and control arms at a 1:1 ratio by an independent epidemiologist using a computer-generated randomization schedule, which was revealed after all baseline measurements had been completed as well as all study households recruited and assigned to their respective arm to ensure the allocation sequence was concealed from those assigning the arms. Both study participating households and data collectors were blinded to intervention status during study enrollment and baseline data collection. In addition, all eligible children/households within the clusters were included in the study to minimize the risk of selection bias; however, because of the distinctive feature of cluster design and nature of the intervention under study, blinding of the households receiving the IBS intervention was not possible.

2.6. Sampling method and recruitment of participants

The cluster sampling method was used to randomly select 100 clusters (50 clusters per arm) among the total 132 clusters in the MHDSS site, and all eligible households were included within the selected cluster (complete enumeration). The potential, 5500 eligible households with less than 4 years old child were chosen from the update report record of MHDSS. The list of clusters/"Gots" and households with children less than 4 years old, was established from the MHDSS record, and the selected households were identified using the permanent MHDSS site house number. Then, a child less than 4 years old was recruited from each household. In situations where there were two or more children less than 4 years old living in the same household, only the youngest child was included in the study.

The actual participants were recruited at the household level by field workers during the baseline survey after ensuring whether the households met the eligibility criteria. A screening questionnaire was used by field data collectors upon their first visit to each household to ensure that the household was appropriate and willing to participate. When the household meets the eligibility criteria, the study was explained to the index child parents, and they were asked whether the household is willing to participate in the study and use improved baking stove technology for at least 12 months. Then, when the parents of the household agreed to be involved in the study, the field staff administered a written consent form at that time and the consent procedure was conducted in Amharic (both national & local language). It was also explained that the allocation to intervention or the control group was based on the concept of a "lottery" method, and several special efforts were applied during recruitment to facilitate the enrolment process as described within our earlier research report (23), and there was no any household unwilling to participate in the study.

2.7. Intervention

2.7.1. Trial descriptions

Replacing the open burning TBS with "*Mirt*" IBS, the well-known commercially distributed type of IBS in Ethiopia, was the intervention for this study. The "*Mirt*" IBS intervention can save a considerable quantity of fuel-wood compared to the TBS method (34), and it is a biomass-fuelled without chimney stove designed by the Ethiopian Energy Studies Research Center for cooking the staple food of Ethiopia called "*Injera*" (i.e., a unique type of yeast-risen flatbread, consumed widely in Ethiopia) (21, 22). All households who were randomized to the intervention arm were identified using the permanent MHDSS house number for a convenient appointment date, and the intervention was delivered to all eligible households in 50 randomly allocated clusters at the beginning of the study period, and the control households were continued to use the customary open burning TBS method equally in 50 randomly allocated clusters. All the "*Mirt*" stoves were manufactured by a local licensed firm and

installed on-site by the installation teams.

Demonstration in the use of the IBS was also provided to each household during the time of installation, and the IBS intervention was promoted regularly throughout the follow-up period by the local energy experts' team of IBS monitors. Concerning study duration, since the life span of "Mirt" IBS is about 5 years (21, 22), the length of both the intervention and the follow-up period was one year safely to account for seasonal factors that have a major effect on the magnitude of both ALRI and HAP in Ethiopia (35), and to maintain a sufficiently short follow-up period to decrease attrition.

2.7.2. Trial adherence and compliance monitoring

Participants' adherence to the intervention assigned to them were assessed through self-report and direct observation by trained field enumerators along with local energy experts' team in both arms. At each follow-up visit, the enumerators observed and recorded the type and condition of baking stove currently being used (i.e., no observed breakage resulting in no use). Additionally, the primary cook was asked whether the baking stove intervention was in good working order (i.e., no reported breakage resulting in no use). Accordingly, timely response to trial-related difficulties such as rapid maintenance or replacement of defective baking stove were maintained and replaced by the installation teams as needed to improve adherence to intervention protocol and avoid the potentially detrimental effects of non-adherence. In addition, trial protocol compliance was checked by the local energy experts' team of stove monitors' through unannounced visual inspection visits in homes of both arms to enhance data validity.

2.7.3. Participant retention strategies

Once the households were enrolled, reasonable efforts were made to promote participant retention and complete follow-up for the entire study period by working on active community engagement through the Ethiopian health extension program as well as through the local health development army team structure to prevent missing data and avoid the associated complexities in analysis and interpretation. In this regard, interest in the study was maintained through periodic communications about the intervention protocol adherence during the regular local health development army team meetings as well as during home visits by field workers, health extension workers, and local energy experts. Furthermore, home visits were scheduled to limit the participants' burden related to follow-up visits, and at the start of the trial, control households were informed that they would receive the IBS intervention at the end of the study period to maintain justice and achieve a high level of post-recruitment participant retention.

2.7.4. Trial safety monitoring

Even though the "Mirt" IBS intervention (21, 22), which was tested by this trial, was not involved any drug or medical procedure; and not known to increase the risk of any adverse event, masked interim analyses were included in the protocol for safety and efficacy monitoring. Since the standard "Mirt" IBS intervention (21, 22) is expected to reduce HAP related health effects (22), it was likely to be safer than the open burning TBS method. Any adverse events data deemed related to the trial intervention (i.e., such as cooking-related burn events data) were collected, and reported immediately for appropriate medical action and for further assessment of seriousness and expectedness to inform the conduct of the ongoing trial. The collected data were reviewed for safety in December 2018 by an independent data Safety, and Monitoring Board to determine whether there were grounds to stop the trial for adverse events. Nevertheless, the Board found no grounds to stop the trial early due to adverse events.

2.8. Outcome assessments

To identify the occurrence of childhood ALRI, we used the definition of Integrated Management of Childhood Illnesses (IMCI) pneumonia algorithm developed by the World Health Organization (WHO) (36, 37). Since nurses can effectively diagnose childhood pneumonia, and the IMCI-pneumonia assessment protocol is a common diagnostic criterion at health facilities in Ethiopia (38), the outcome variable was assessed in the same manner in both arms by field nurses, who were trained in IMCI-pneumonia algorithm, and not involved in either randomization or intervention delivery. In this trial, the term childhood ALRI was used as a synonym for the IMCI-

pneumonia (36, 37), because it is the preferred term for childhood pneumonia in peculiar to the developing countries (39).

2.9. Data collection methods

Baseline data were collected in each household before intervention delivery, and a total of four follow-up visits were carried out, immediately after the delivery of the intervention, in the same week at approximately 3-month intervals for longitudinal data collection by trained local field nurses. The duration of the follow-up period was one year to cover the major Ethiopian periods of annual weather changes and to account for seasonal factors that might have a major effect on the magnitude of both ALRI and HAP in Ethiopia (35).

2.10. Data quality assurance

Various appropriate measures were taken to address the validity and reliability of the study by ensuring the quality of data. To start with the outcome variable, childhood ALRI was assessed in the same manner in both arms by field nurses who were trained in the standard IMCI-pneumonia algorithm. The study team was in regular contact with the study site team with scheduled meetings and additional communication as needed for quality control. About 5% of randomly selected home visits were done in duplicate by supervisors as cross-checking mechanism to ensure the validity of the collected data, and feedback sessions were done on about a three-month basis.

To minimize the risk of bias, clusters were randomly allocated to intervention and control arms, all eligible households within the clusters were included in the study, location sequence was concealed from those assigning participants to groups, primary outcome assessors were blind to the intervention, and outcome measurement was performed in the same manner in both arms. In addition, a similar licensed firm was manufactured all the IBS and similarly trained installation teams were administered the intervention. Furthermore, all initially randomized participants were analyzed in the groups they were assigned to (i.e., intention-to-treat/ ITT analysis principle). Furthermore, the methodological soundness such as large sample size, longitudinal study design, and baseline data collection on the primary outcome and risk factors to be adjusted through GEE modeling can help us to achieve an effective balance of confounders in both arms. Finally, this manuscript was reported following both the guidelines of Consolidated Standards of Reporting Trials (CONSORT) 2010 statement extension to cluster randomized trials (40) to address the essential study design components of this trial report (See additional file 1) and Template for Intervention Description and Replication (TIDieR) checklist (41) for better reporting of the intervention aspects.

2.11.

Statistical analysis methods

Data were analyzed using the Statistical Package for Social Sciences (SPSS) for Windows version 22, and all statistical tests were two-sided with p -value < 0.05 considered statistically significant. To quantify the magnitude of clustering for childhood ALRI outcome, cluster-level ICC value was calculated using the standard formulae as $ICC = \text{Var}(U_o) / \text{Var}(U_o) + \Pi^2/3$ (42, 43); where, $\text{Var}(U_o)$ is the random intercept variance, that is, the estimated cluster-level (i.e., level-two) variance component, and $\Pi^2/3$ refers to the individual-level (i.e., level-one) variance component which is fixed to be 3.29 (43). We took this fixed value of 3.29 for the individual level variance component, as there is no individual-level (level-one) residual for binary (discrete) outcome variable in the multilevel logistic analysis model (42, 43). Using this formula, the estimated magnitude of cluster-level ICC value for childhood ALRI outcome was found to be not different from zero, which indicates that only well below 1% of the total variability in the chance of acquiring childhood ALRI is explained by the between cluster-level variation. Therefore, we safely treated individual children as the sole unit of analysis and interpretation in determining the effect of IBS intervention on childhood ALRI (42) compared with the continuation of the open burning TBS.

The effect of IBS intervention on the repeated response of childhood ALRI between the two arms using adjusted Odds Ratios (AORs), with the respective 95% CIs, as measures of effect following a Generalized Estimating Equations modeling approach among the ITT population by considering the underlying design of the study and nature of the outcome variable under investigation. The GEE analysis method is the preferred method for

longitudinal data analysis due to its computational simplicity and robustness to misspecification of the repeated measures correlation structure (44). Besides, this analysis technique can take both the village-level clustering of childhood ALRI outcome and the longitudinal sampling method into account, and it can handle missing data values without the need for explicit imputation by considering all participants with at least one follow-up visit through automatically excluding the missing values by the GEE analysis model (44).

Concerning model fitness, the best working correlation structure (WCS) of the outcome variable was chosen as autoregressive (AR1) based on the quasi-likelihood under the independence model criterion, and by means of a critical examination of the working correlation matrix to uphold the goodness of model fitness, and hence to get a more precise estimation of the IBS intervention effect. As a final point, our GEE analysis model has included a binary outcome variable of childhood ALRI and a binary indicator of treatment allocation (i.e., control versus intervention) as well as other indicator variables measured at baseline such as gender, age, childhood ALRI, location of cooking quarter, stove type for other cooking/boiling water purposes and frequency of “Injera” baking event; and the results are presented next in texts and tables.

3. Results

Among the total 5830 children assessed for eligibility at baseline survey, 5508 (94.48%) children (i.e. one child in each surveyed household) were enrolled in the study from June to August in 2018 across 100 randomly selected clusters in both arms. The trial was comprised of 2750 (49.9%) and 2758 (50.1%) children in the intervention and control arm respectively, and the prevalence of childhood ALRI among the primarily enrolled children was 19.1% at baseline (19.1% and 19.2% in intervention and control arms respectively) as displayed in Table 1. A total of 322 (5.52%) children were excluded at baseline due to not meeting the inclusion criteria; and 175 (91 in intervention and 84 in the control arm) children were permanently LTF at the first follow-up visit after initial enrollment (Fig. 1).

Table 1

Baseline characteristics of enrolled households and children

Child characteristic		Treatment arm		Total (%)
		Control (%)	Intervention (%)	
Gender of child	Female	1293 (49.0)	1346 (51.0)	2639 (47.9)
	Male	1465 (51.1)	1404 (48.9)	2869 (52.1)
Age of child at entry in years (months)	< 1 Year old (0-11)	764 (48.5)	811 (51.5)	1575 (28.6)
	1 Year old (12-23)	774 (50.9)	748 (49.1)	1522 (27.6)
	2 Years old (24-35)	745 (51.9)	690 (48.1)	1435 (26.1)
	3 Years old (36-47)	475 (48.7)	501 (51.3)	976 (17.7)
Child ALRI status at baseline	No	2229 (50.0)	2225 (50.0)	4454 (80.9)
	Yes	529 (50.2)	525 (49.8)	1054 (19.1)
Location of cooking quarter	Inside the living house	1008 (50.9)	972 (49.1)	1980 (35.9)
	Separate kitchen	1750 (49.6)	1778 (50.4)	3528 (64.1)
Cookstove type	Traditional stove	2539 (50.1)	2524 (49.9)	5063 (91.9)
	Improved stove	219 (49.2)	226 (50.8)	445 (8.1)
Frequency of "Injera" baking event	One or more per day	383 (50.7)	372 (49.3)	755 (13.7)
	Every other day	303 (50.9)	292 (49.1)	595 (10.8)
	Every 3 days	2014 (49.7)	2035 (50.3)	4049 (73.5)
	Every 4 or more days	58 (53.2)	51 (46.8)	109 (2.0)
Total (%)		2758 (50.1)	2750 (49.9)	5508 (100)

The first follow-up visit was carried out from September to November 2018; the second took place from December to February 2019; the third occurred from March to May 2019, and the fourth visit was occurred from June to August in 2019 at the entire 100 clusters, and this trial study was terminated at the planned target of one year after the last participant had been randomized. Among the overall follow-up visits of 22032 in the enrolled population, data were obtained from a total of 19792 (89.8%) child observations (9860 and 9932 in the

intervention and control arms respectively) (Table 2).

Table 2

Lost to follow-up events during the entire follow-up period among enrolled participants

Characteristics		Treatment arm		Total (%)	χ^2 (P-value)
		Control (%)	Intervention (%)		
LTF	Yes	1092 (9.9)	1148 (10.4)	2240 (10.2)	1.65 (0.199)
	No	9932 (90.1)	9860 (89.6)	19792 (89.8)	
	Total	11024 (100)	11008 (100)	22032 (100)	
Reasons for LTF	Moved out of study area	558 (51.1)	549 (47.8)	1107 (49.4)	3.94 (0.268)
	Consent withdrawal	318 (29.1)	377 (32.8)	695 (31.0)	
	Death of child	148 (13.6)	148 (12.9)	296 (13.2)	
	Not at home	68 (6.2)	74 (6.5)	142 (6.4)	
	Total*	1092 (100)	1148 (100)	2240 (100)	
LTF: lost to follow-up.					
Notes: Among the *Total LTF of 2240, only 1014 (503 in the control, 511 in the intervention arm) were recorded as new incidence during the entire follow-up period; whereas 1226 (589 in the control and 637 in the intervention arm) were registered as repeated LTF observations in the preceding rounds.					

Among the total data of 19792; 5313, 5055, 4828, and 4596 data were obtained from the first, second, third, and fourth visits respectively. In addition, there were no entire cluster level losses and exclusions in both arms among the initially enrolled participants. Nevertheless, a total of 2240 (10.2%); (10.4% from the intervention and 9.9% from the control arm) participant visits were LTF after randomization during the entire follow-up period due to moving out of study area, consent withdrawals, child deaths or not at home after a repeated home visit, and reasons for attrition were similar between the intervention and control arms (Table 2). Hence, since the reasons for LTF were not related to the given study in both arms, the entire 2240 LTF child observations were omitted from the analysis and the observed 19792 data can be viewed as a random sample from the population.

Concerning harms, there was 92 episode of childhood cooking-related burn events among the ITT population during the entire follow-up period (41 in the intervention and 51 in the control group); and there was no difference in incidence rates of cooking-related burn events between the intervention and control groups with an incidence rate ratio (IRR) of 0.80 (95% CI: 0.53–1.21) as demonstrated below in Table 3.

Table 3

Incidence of cooking-related burn events among the intention-to-treat population

Intervention (n = 9860)		Control (n = 9932)		IRR (95% CI)
Number of burn events	Incidence rate	Number of burn events	Incidence rate	
41	0.42	51	0.51	0.80 (0.54-1.22)

Among the enrolled participants (5508) in the beginning, follow-up data were obtained from a total of 5333 (96.82%) children (2659 from the intervention and 2674 from the control group) for at least one follow-up visit within 50 clusters in each arm. This establishes the ITT population dataset within 100 clusters in both arms and included in each analysis. However, 175 children were completely omitted from the analysis due to LTF between the baseline and first follow-up visit (91 from the intervention, and 84 from the control arm) (Fig. 1). For the ITT population, the average cluster size was 53.33 (53.18 for intervention and 53.48 for control arm); and the childhood ALRI prevalence was 19.31% (95% CI: 18.30-20.40) at baseline with 19.18% and 19.45% in the intervention and control arms respectively. Besides, the estimated magnitude of cluster-level ICC value for childhood ALRI outcome was found to be very small as depicted in Table 4.

Table 4

Baseline characteristics of clusters and participants in the intention-to-treat population

Characteristics	Intervention arm	Control arm	Both arm
Number of clusters	50	50	100
Number of children	2659	2674	5333
Mean cluster size (SD)	53.18 (12.13)	53.48 (12.39)	53.33 (12.2)
CoV for cluster size	0.228	0.232	0.229
Childhood ALRI prevalence rate	19.2	19.5	19.3
Cluster-level variance			2.336E-11
Individual-level variance			3.29 (42)
Cluster level ICC			7.10E-12

ALRI-acute lower respiratory infection; ICC: Intra-cluster correlation coefficient; CoV-coefficient of variation.

As to childhood ALRI outcome estimations, there were 3540 episodes of childhood ALRI in the overall ITT population (1732 in the intervention and 1808 in the control group) giving a longitudinal childhood ALRI prevalence rate of 17.9% (95% CI: 17.4-18.4) during the entire follow-up period (17.5% and 18.3% in the intervention and control arm respectively). The childhood ALRI prevalence rate distribution by round was found to be 18.3% for the first round, 17.7% for the second, 17.5% for the third, and 17.9% for the fourth round as displayed in Fig. 2 by arm.

Concerning the predicted effect of the intervention, the IBS intervention was not found to have a statistically significant effect on the longitudinal ALRI among under 5 years old children with an estimated AOR of 0.95 (95% CI: 0.89–1.02) compared with the continuation of an open fire TBS as demonstrated by the GEE model in Table 5.

Table 5

Effect of the stove intervention on childhood acute lower respiratory infection

Parameters		B	S.E	95% Wald CI		AOR (95% CI)	P-value
				lower	upper		
Intercept		-0.890	0.0664	-1.020	-0.760	0.41 (0.36,0.47)*	0.001
Treatment arm	Intervention	-0.047	0.0354	-0.117	0.022	0.95 (0.89,1.02)	0.180
	Control					1	
Gender	Female	0.067	0.0353	-0.003	0.136	1.07 (1.0,1.15)	0.059
	Male					1	
Age of child at entry	< 1 Year old					1	
	1 Year old	-0.074	0.0442	-0.161	0.013	0.93 (0.85,1.01)	0.095
	2 Years old	-0.342	0.0477	-0.435	-0.248	0.71 (0.65,0.78)*	0.001
	3 Years old	-0.615	0.0591	-0.731	-0.499	0.54 (0.48,0.61)*	0.001
Baseline ALRI	Yes	0.219	0.0433	0.134	0.304	1.24 (1.14,1.36)*	0.001
	No					1	
Location of cooking quarter	Inside living house					1	
	Separate kitchen	-0.252	0.0355	-0.322	-0.183	0.78 (0.73,0.83)*	0.001
Cookstove type	Traditional					1	
	Improved	-0.484	0.0899	-0.660	-0.308	0.62 (0.52,0.74)*	0.001

Frequency of baking event	1 or more per day					1	
	Every other day	-0.278	0.0639	-0.403	-0.153	0.76 (0.67,0.86)*	0.001
	Every 3 days	-0.321	0.0464	-0.412	-0.230	0.73 (0.66,0.79)*	0.001
	Every 4 or more days	-0.426	0.1374	-0.696	-0.157	0.65 (0.50,0.86)*	0.002
Visit	First					1	
	Second	-0.041	0.0521	-0.143	0.061	0.96 (0.87,1.06)	0.430
	Third	-0.053	0.0536	-0.158	0.052	0.95 (0.85,1.05)	0.324
	Fourth	-0.026	0.0529	-0.130	0.077	0.97 (0.88,1.08)	0.619
AOR: adjusted odds ratio; B: beta coefficients; C.I.: confidence interval; GEE: generalized estimation equation; ITT: intention-to-treat; S.E: standard error.							
*Significantly associated at p-value < 0.05.							
Notes: The table reports the results of the health effect of the intervention between arms using AORs as a measure of effect using the GEE analysis modeling approach. The GEE analysis model has simultaneously included a binary outcome variable of childhood ALRI and an indicator of treatment allocation as well as other indicator variables measured at baseline (gender, age, childhood ALRI, location of cooking quarter, stove type for other cooking purposes and frequency of "Injera" baking event).							

4. Discussions

This large sample, community-level cluster randomized controlled closed cookstove trial was aimed at determining the effect of IBS intervention on childhood ALRI at the individual child level in Northwest Ethiopia compared with the continuation of open fire TBS. The biomass-fuelled TBS was a major (94.5%) household energy source for baking purpose in the study locality; and the majority (95.8%) of the households practice extra indoor burning events such coffee ceremony, burning incense inside the house as well as outdoor burning events such as burning rubbish and charcoal production was common (38.1%) observable practices in the study area (23).

In this trial, data were obtained from 5333 children (2659 from the intervention and 2674 from the control group) for at least one follow-up visit, and this was used as an ITT population dataset within 100 clusters in both arms. The longitudinal childhood ALRI prevalence rate distribution by round was showed a steep reduction followed by a slight increase in similar fashion in both arms (Fig. 2) which might be explained by time variation of the data collection rounds undertaken which is linked with seasonal factors that have a major effect on the magnitude of both ALRI and HAP in Ethiopia (35).

The predicted effect of the biomass-fuelled IBS intervention was not found to have a statistically significant effect on the risk of repeated childhood ALRI response among under 5 years old children in Northwest Ethiopia with an estimated AOR of 0.95 (95% CI: 0.89-1.02) compared with the continuation of an open burning TBS method. This finding mirrors the previous cross-sectional study, conducted as baseline part of this stove trial

project, which reported that the odds of childhood ALRI did not show a significant association among children living in households with IBS when compared with children living in households with the TBS type with an estimated AOR of 0.78 (95% CI: 0.52–1.15) (23). Likewise, a recent policy research working paper reported no evidence that “Mirt” IBS intervention reduced the risk of respiratory symptoms among older children in rural Ethiopia (12). This finding is also similar to a recent large sample size stove trial in Malawi, which found no evidence that an improved biomass-fuelled cookstove intervention reduced the risk of pneumonia among young children in rural Malawi with an incidence rate ratio of 1.01 (95% CI: 0.91, 1.13) with incidence rates of 15.76 (95% CI 14.89–16.63) and 15.58 (95% CI 14.72–16.45) per 100 child-years in the intervention and control arms respectively (13). The finding of this stove trial is also in agreement with the Guatemala randomized controlled stove trial which reported no statistically significant reduction in childhood pneumonia among intervention groups following ICS intervention with a rate ratio of 0.84 (95% CI: 0.63–1.13) (11).

Our findings do, however; contradict the findings of two previous biomass-burning stove intervention studies in Rwanda and Guatemala which found evidence of health effects among children following biomass-burning ICS interventions (10, 45). The Rwanda large-scale stove trial reported that a natural draft rocket-style cookstove intervention effectively reduced acute respiratory infection (ARI) prevalence by 25% among children < 5 years with a prevalence ratio of 0.75 (95% CI: 0.60–0.93) (45). Correspondingly, the field experiment carried out in rural Guatemala reported a 63.3% reduction in respiratory symptoms among children following wood-burning ICS intervention (10). The possible explanation for the difference between our findings and the findings of previous biomass-burning stove intervention trials might be due to the particular variation of traditional cooking device requirements and trial stove design among others. For instance, unlike our trial stove (21, 22), the Guatemala trial stove reduces smoke exposure mainly by removing emissions to the outdoor environment with a chimney rather than by improving combustion efficiency (10). In addition, since the effectiveness of cookstoves differs from an ethnic group, geographical location, and local fuel resources (46), the possible explanation for the difference might be the reflection of the difference in geographical location and cultural variations.

Generally, in our trial, the lack of intervention effect on childhood ALRI might be due to the failure of the biomass-fuelled IBS intervention that could not sufficiently reduce HAP to have a preventive health effect. Thus, the most important implication of this finding is that there is inadequate evidence to recommend the biomass-fuelled IBS intervention for the prevention of childhood ALRI compared to the use of open burning TBS. Consequently, in countries with limited available resources like Ethiopia, an improved stove intervention has to be rigorously proven to be highly effective in reducing the burden of HAP related diseases; and, an effective cooking solution that could potentially reduce smoke exposure sufficiently by removing emissions to the outdoor environment with a chimney rather than by improving combustion efficiency (10) is needed to avert the adverse health effect of HAP, particularly, acute lower respiratory infection among children, who are the most vulnerable and sensitive segment of the population to the damaging health effects of HAP (6, 7).

Finally, even though the IRR was not statistically significant, there was a 20% reduction in the risk of childhood cooking-related burn event in the intervention group, suggesting the IBS intervention might offer a considerable safety advantage over the open TBS method (IRR = 0.80: 95% CI: 0.53–1.21). Besides, the IBS intervention would achieve economic and environmental benefits by reducing household fuel consumption and by the direct reduction of the amount of wood burned which can, in turn, decrease environmental destruction and pollution (47).

Limitations: We acknowledge several limitations in our trial such as the un-blinded nature of the stove intervention; and the lack of intervention effect on childhood ALRI might be explained by exposure to other sources of HAP pollution in the study area, including other cooking activities and presence of extra indoor burning events such as the Ethiopian coffee ceremony with burning incense in the household as well as the presence of outdoor burning events such as charcoal production and rubbish burning near study households that could have diluted any potential effects of the IBS intervention.

Generalisability: This fairly large-sample cookstove trial was completed effectively from a methodological and practical point of view; and our sample was analogous to the wider population of households that uses biomass-fuelled TBS as a major household energy source for cooking purposes throughout Ethiopia, which is typically representative of the great majority (95%) of biomass fuel user households (48) and (80%) traditional stove user

population in Ethiopia (49) and elsewhere in low and middle-income countries. Therefore, the findings of this trial are generalizable to similar settings in Ethiopia and other low and middle-income countries at an individual child level.

Conclusion

In conclusion, we found no evidence that an intervention comprising biomass-fuelled IBS reduced the risk of repeated childhood ALRI at the individual child level in Northwest Ethiopia compared with the continuation of an open burning TBS. Therefore, effective cooking solutions that could potentially reduce the negative health impact of HAP are needed to avert the adverse health effect of HAP, particularly, childhood ALRI in the study area. Also, the results of this large-sample trial should have important policy implications in Ethiopia and beyond. Besides, further researches are needed to establish a longer-term health effect of locally made IBS interventions by addressing the limitations of our trial.

Abbreviations

- ALRI
- Acute Lower Respiratory Infection
- AOR
- Adjusted Odds Ratio
- CI
- Confidence Interval
- COR
- Crude Odds Ratio
- CoV
- coefficient of variation
- GEE
- Generalized Estimation Equation
- HAP
- Household Air Pollution
- IBS
- Improved baking Stove
- ICC
- Intra-cluster Correlation Coefficient
- ICS
- Improved Cookstove

- IMCI
- Integrated Management of Childhood Illnesses
- IRR
- Incidence Rate Ratio
- ITT
- Intention-to-Treat
- LMICs
- Low-and Middle-Income Countries
- LTF
- Lost to Follow-up
- MHDSS
- Mecha Health and Demographic Surveillance System
- TBS
- Traditional Baking Stove
- WCS
- Working Correlation Structure
- WHO
- World Health Organization

Declarations

Ethics approval and consent to participate

The study protocol was reviewed and obtained approval on April 04/2018 from the Scientific and Ethical Review Committee of Bahir Dar University at the College of Medicine and Health Sciences with ethics committee reference number: 088/18-04, and the trial was registered on clinical trials.gov registry database. Moreover, a Data Safety, and Monitoring Board was appointed to review and monitor the study procedures and potential adverse events. The study was being performed following good clinical practice standards, applicable local regulatory requirements, and the recommendations of the Declaration of Helsinki as reported in-depth within the baseline study of this stove trial project (23). Written informed consent (or thumbprint witnessed by someone independent of the study for those unable to read and write) was obtained from every parent or guardian for children to participate in the study following the distribution of written information sheets in “Amharic” (both official & study area language) to all participants. Moreover, the standard IBS device was manufactured and installed by a licensed company, and the design; management, analysis, and reporting of the study results were entirely independent of the manufacturer of the IBS.

Consent for publication

Not applicable

Availability of data and material

The data that support the findings of this study are available from the corresponding author on reasonable request and with permission of the “Mecha” Health and Demographic Surveillance research center at Bahir Dar University in Ethiopia.

Competing interests

None declared under financial, general, and institutional competing interests.

Funding

This work was supported by Bahir Dar University through the research funding system of the “Mecha” Health and Demographic Surveillance System [grant number RCS/253/09]. The funder played no role in study design; collection, analysis, and interpretation of data; preparation of the manuscript or decision to publish. The corresponding author had full access to all the study data and final responsibility for the decision to submit for publication.

Authors' contributions

MMA was the principal investigator of the study and took the leading responsibility starting from origin, design and supervising data collection process to the final analysis and preparation of the manuscript. GDA and STM contribute in supervising the study, reviewing the document and providing critical comments from the design and data collection process to the final analysis and preparation of the manuscript. KLW contributes to reviewing the document and providing critical comments from the design and final analysis to the preparation of the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to extend our gratitude to field data collectors, supervisors as well as all family members of the households who participated in this study.

Authors' information

1. Mesafint Molla Adane (MPH, BSc.), Bahir Dar University, College of Medicine and Health Sciences, School of Public Health, Department of Environmental Health, Bahir Dar, Ethiopia: Email: Mesafint.Molla@bdu.edu.et or mesafintmolla@yahoo.com
2. Professor Getu Degu Alene (Ph.D., MSc., BSc.), Bahir Dar University, College of Medicine and Health Sciences, School of Public Health, Department of Epidemiology and Biostatistics, Bahir Dar, Ethiopia: Email: adgetu123@yahoo.com.
3. Seid Tiku Mereta (Ph.D., MSc.BSc.), Jimma University, Department of Environmental Health Sciences and Technology, Jimma, Ethiopia: Email: seidtiku@yahoo.com
4. Kristina L Wanyonyi (BDS, MPH, Ph.D, FHEA, DDPH/RCS Eng), Queen Mary University of London, Institute of Dentistry, Barts and The London School of Medicine and Dentistry, London, United Kingdom: Email: k.wanyonyi@qmul.ac.uk.

OTHER INFORMATION

Registration:

The trial was retrospectively registered on August 2, 2018 at clinical trials.gov registry database (<https://clinicaltrials.gov/>) with NCT03612362 registration identifier number.

Protocol:

The full trial protocol is available from the corresponding author, IRB office and School of Public at Bahir Dar University in Ethiopia.


References

1. Smith K, Bruce N, Balakrishnan K, Adair-Rohani H, Balmes J, Chafe Z, et al. Millions Dead: How Do We Know and What Does It Mean? Methods Used in the Comparative Risk Assessment of Household Air Pollution. *Annual Review of Public Health*. 2014 Mar 18;35 (1):185–20610. doi:10.1146/annurev-publhealth-032013-182356.
2. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012 Dec 15;380(9859):2224–60. PubMed PMID: 23245609. Pubmed Central PMCID: 4156511.
3. World Health Organization. *Burning opportunity: clean household energy for health, sustainable development, and wellbeing of women and children*. Switzerland: Geneva; 2016.
4. Apte K, Salvi S. *Household air pollution and its effects on health*. F1000Research. 2016;5.
5. World Health Organization. *Effects of air pollution on children's health and development: a review of the evidence*. Denmark: WHO Regional Office for Europe, Copenhagen; 2005.
6. Po JY, FitzGerald JM, Carlsten C. Respiratory disease associated with solid biomass fuel exposure in rural women and children: systematic review and meta-analysis. *Thorax*. 2011 Mar;66(3):232–9. PubMed PMID: 21248322.
7. Rees N. *Clear the air for children: the impact of air pollution on children*. New York: UNICEF Division of Data, Research and Policy; October 2016.
8. Nigel B, Annette P-U, Dan P, Heather AR, Eva R. WHO IAQ guidelines: household fuel combustion - Technical paper on evidence review methods, 2014. available at: <http://www.who.int/indoorair/guidelines/hhfc>.
9. Baumgartner J, Smith KR, Chockalingam A. Reducing CVD through improvements in household energy: implications for policy-relevant research. *Glob Heart*. 2012;7:243–47.
10. Ludwinski D, Moriarty K, Wydick B. Environmental and health impacts from the introduction of improved wood stoves: evidence from a field experiment in Guatemala. *Environ Dev Sustain*. 2011;13(4):657–76.
11. Smith KR, McCracken JP, Weber MW, Hubbard A, Jenny A, Thompson LM, et al. Effect of reduction in household air pollution on childhood pneumonia in Guatemala (RESPIRE): a randomised controlled trial. *Lancet*. 2011 Nov 12;378(9804):1717–26. PubMed PMID: 22078686.
12. LaFave D, Beyene AD, Bluffstone R, Dissanayake ST, Gebreegziabher Z, Mekonnen A, et al. *Impacts of Improved Biomass Cookstoves on Child and Adult Health: Experimental Evidence from Rural Ethiopia*. The World Bank; 2019.
13. Mortimer K, Ndamala CB, Naunje AW, Malava J, Katundu C, Weston W, et al. A cleaner burning biomass-fuelled cookstove intervention to prevent pneumonia in children under 5 years old in rural Malawi (the Cooking and Pneumonia Study): a cluster randomised controlled trial. *The Lancet* January. 2017;389(10065):167–75. doi:10.1016/S0140-6736(16)32507-7.
14. Daniel P, Nigel B, Mukesh D, Kirstie J, Eva R. Real-life effectiveness of 'improved' stoves and clean fuels in reducing PM2.5 and CO: systematic review and meta-analysis. *Environ Int*. 2017;101:7–18.
15. Quansah R, Semple S, Ochieng CA, Juvekar S, Armah FA, Luginaah I, et al. Effectiveness of interventions to reduce household air pollution and/or improve health in homes using solid fuel in low-and-middle income countries: A systematic review and meta-analysis. *Environment International*. 2017.
16. Pennise D, Brant S, Agbeve SM, Quaye W, Mengesha F, Tadele W, et al. Indoor air quality impacts of an improved wood stove in Ghana and an ethanol stove in Ethiopia. *Energy Sustain Dev*. 2009;13(2):71–6.
17. Kumie A, Samet J, Berhane K. *Air, Pollution, Occupational Health and Safety, and Climate Change: Findings, research needs and policy implications establishing a GEOHealth Hub for East Africa. Situational analysis and needs assessment for Ethiopia*. August 2014.
18. Berhane K, Kumie A, Samet J. *Health Effects of Environmental Exposures, Occupational Hazards and Climate Change in Ethiopia: Synthesis of Situational Analysis, Needs Assessment and the Way Forward*. *Ethiopian Journal of Health Development*. 2016;30(1):50–6.
19. Tefera W, Asfaw A, Gilliland F, Worku A, Wondimagegn M, Kumie A, et al. *Indoor and Outdoor Air Pollution-related Health Problem in Ethiopia: Review of Related Literature*. *Ethiopian Journal of Health*

- Development. 2016;30(1):5-16.
20. Beyene AD, Bluffstone R, Gebreegziabher Z, Martinsson P, Mekonnen A, Vieider F. The improved biomass stove saves wood, but how often do people use it? Evidence from a randomized treatment trial in Ethiopia: The World Bank; 2015.
 21. Damte A, Koch SF. Clean fuel saving technology adoption in urban Ethiopia. South Africa: Department of Economics Working Paper, University of Pretoria. 2011.
 22. World Bank. Household cookstoves, environment, health, and climate change: A new look at an old problem. 2011.
 23. Adane MM, Alene GD, Mereta ST, Wanyonyi KL. Prevalence and risk factors of acute lower respiratory infection among children living in biomass fuel using households: a community-based cross-sectional study in Northwest Ethiopia. *BMC Public Health*. 2020 March 19;20(1):363. Doi:10.1186/s12889-020-08515-w.
 24. Keil C, Kassa H, Brown A, Kumie A, Tefera W. Inhalation exposures to particulate matter and carbon monoxide during Ethiopian coffee ceremonies in Addis Ababa: a pilot study. *Journal of environmental and public health*. 2010;2010.
 25. Sanbata H, Asfaw A, Kumie A. Indoor air pollution in slum neighbourhoods of Addis Ababa, Ethiopia. *Atmos Environ*. 2014;89:230-4.
 26. Chakraborty H. The design and analysis aspects of cluster randomized trials. *Statistical Advances in the Biomedical Sciences: Clinical Trials, Epidemiology, Survival*.
 27. Biswas A, Datta S, Fine JP, Segal MR. *Statistical advances in the biomedical sciences: clinical trials, epidemiology, survival analysis, and bioinformatics*. 2007.
 28. Kloos H, Woldu Z. The physical and biotic environment: its features and health implications. In: Berhane Y, Haile-Mariam D, Kloos H, editors. *Epidemiology and Ecology of Health and Disease in Ethiopia*. Addis Ababa/Ethiopia: Shama Books; 2006.
 29. World Bank. *Economics of Adaptation to Climate Change, Synthesis Report*. Washington, DC: 2010a.
 30. Desalegn B, Suleiman H, Asfaw A. Household fuel use and acute respiratory infections among younger children: an exposure assessment in Shebedino Wereda, Southern Ethiopia. *African Journal of Health Sciences*. 2011;18(1-2):31-6.
 31. Hemming K, Girling AJ, Sitch AJ, Marsh J, Lilford MJ. Sample size calculations for cluster randomized controlled trials with a fixed number of clusters. *BMC Med Res Methodol*. 2011;11:102.
 32. Edwards R, Hubbard A, Khalakdina A, Pennise D, Smith KR. Design considerations for field studies of changes in indoor air pollution due to improved stoves. *Energy for Sustainable Development* June 2007;11(2).
 33. Smith KR, Dutta K, Chengappa C, Gusain P, Masera O, Berrueta V, et al. Monitoring and evaluation of improved biomass cookstove programs for indoor air quality and stove performance: conclusions from the Household Energy and Health Project. *Energy Sustain Dev*. 2007;11(2):5-18.
 34. Shanko M, Abebe T, Lakew H. A report on Mirt biomass injera stove market penetration and sustainability study in Amhara. Ethiopia: Oromiya and Tigray National Regional States. GTZ Sun Energy, Addis Ababa; 2009.
 35. Kumie A, Emmelin A, Wahlberg S, Berhane Y, Ali A, Mekonnen E, et al. Sources of variation for indoor nitrogen dioxide in rural residences of Ethiopia *Environ Health*. 2009;8(51).
 36. World Health Organization. *Handbook IMCI. Integrated management of childhood illness*: World Health Organization. Geneva, Switzerland, 2005.
 37. World Health Organization. *Revised WHO classification and treatment of pneumonia in children at health facilities: Evidence summaries, quick reference guide*. Geneva: World Health Organization; 2014.
 38. Federal Democratic Republic of Ethiopia Ministry of Health. *Integrated management of newborn and childhood illness*. Ethiopia: Addis Ababa; 2011.
 39. Ranganathan SC, Sonnappa S. Pneumonia and other respiratory infections. *Pediatr Clin North Am*. 2009;56(1):135-56.
 40. Campbell MK, Piaggio G, Elbourne DR, Altman DG. Consort 2010 statement: extension to cluster randomised trials. *Bmj*. 2012;345:e5661.
 41. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *Bmj*. 2014;348:g1687.
 42. Sommet N, Morselli D. Keep Calm and Learn Multilevel Logistic Modeling: A Simplified Three-Step

- Procedure Using Stata, R, Mplus, and SPSS. *International Review of Social Psychology*. 2017;30:203–18.
43. Snijders T, Bosker AB. RJ. *Multilevel analysis: an introduction to basic and advanced multilevel modeling*. London: Sage Publications; 1999.
 44. Liang K-Y, Zeger SL. *Longitudinal data analysis using generalized linear models*. *Biometrika*. 1986;73(1):13–22.
 45. Kirby MA, Nagel CL, Rosa G, Zambrano LD, Musafiri S, de Dieu Ngirabega J, et al. Effects of a large-scale distribution of water filters and natural draft rocket-style cookstoves on diarrhea and acute respiratory infection: A cluster-randomized controlled trial in Western Province, Rwanda. *PLoS Med*. 2019;16(6):e1002812.
 46. Mohammadreza S, Hesamoddin S. A comprehensive review of technical aspects of biomass cookstoves. *Renew Sustain Energy Rev*. 2017;70:656–65.
 47. Gebreegziabher Z, Van Kooten GC, Van Soest DP. Technological innovation and dispersion: Environmental benefits and the adoption of improved biomass cookstoves in Tigray, northern Ethiopia. *Energy Economics*. 2017 Sep 7;67:337–45. doi:10.1016/j.eneco.2017.08.030.
 48. Central Statistical Agency. *Welfare Monitoring Survey*. Addis Ababa, Ethiopia: 2011.
 49. Ethiopian Ministry of Water and Energy. *Energy Sector Mapping and Database Development*. Addis Ababa, Ethiopia: 2013a.

Figures






Figure 1

Flow diagram of the study progress from eligibility assessment to enrollment, follow-up, and analysis.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [_Additionalfile1CompletedtableofCONSORT2010checklist.doc](#)