

Maternal and perinatal outcomes and pharmacological management of Covid-19 infection in pregnancy: a systematic review protocol

CURRENT STATUS: Under Review

 Systematic Reviews  BMC

Binny Thomas, Moza Al Hail, Pallivalappila Abdulrouf, Wessam Elkassem, Asma Tarannum, Fatema Al Hail, Mohammed Rijims, Hussain Parappil, Arabo Ibrahim Bayo, Shamsa Ahmad, Zachariah Nazar, Derek Stewart

Binny Thomas

Robert Gordon University

 binnyinhmc@gmail.com **Corresponding Author**
ORCID: <https://orcid.org/0000-0003-1065-8343>

Moza Al Hail

Hamad Medical Corporation

Pallivalappila Abdulrouf

Hamad Medical Corporation

Wessam Elkassem

Hamad Medical Corporation

Asma Tarannum

Hamad Medical Corporation

Fatema Al Hail

Hamad Medical Corporation

Mohammed Rijims

Hamad Medical Corporation

Hussain Parappil

Hamad Medical Corporation

Arabo Ibrahim Bayo

Hamad Medical Corporation

Shamsa Ahmad

Hamad Medical Corporation

Zachariah Nazar

Qatar University College of Pharmacy

Derek Stewart

Qatar University College of Pharmacy



Prescreen

[10.21203/rs.3.rs-28635/v1](#)

Subject Areas

Infectious Diseases* *Maternal & Fetal Medicine

Keywords

COVID-19, maternal, perinatal, pharmacological management, pregnant

Abstract

Background

Over 4.2 million confirmed cases and more than 285,000 deaths, COVID-19 pandemic continues to harm significant number of people worldwide. Several studies have reported the impact of COVID-19 in general population, however, there is scarcity of information related to pharmacological management, maternal and perinatal outcomes during the pandemic. Altered physiological, anatomical and immunological response during pregnancy makes it more susceptible to infections. Furthermore, during pregnancy, a woman undergoes multiple interactions with the health care system that increases her chance of getting infected, therefore managing pregnant population presents a unique challenge.

Research Question(s)

This systematic review seeks to answer the following questions in relation to COVID-19,

What are the different clinical characteristics presented in maternal and perinatal population?

What are the different maternal and perinatal outcome measures reported?

What are the distinct therapeutic interventions reported to treat Covid-19?

Is it safe to use 'medications' used in the treatment of COVID-19 during antenatal, perinatal, postnatal and breastfeeding?

Method

The search will follow a comprehensive, sequential three step search strategy. Several databases relevant to COVID-19 and its impact on pregnancy including Medline, CINAHL, LitCovid etc. will be searched from the inception of the disease until the completion of data collection. The quality of this search strategy will be assessed using Peer Review of Electronic Search Strategies Evidence-Based Checklist (PRESS EBC). An eligibility form will be developed for a transparent screening and inclusion/exclusion of studies. All studies will be sent to Refworks and abstraction will be independently performed by two researchers. Risk of bias will be assessed using Cochrane Risk of Bias tool for randomized controlled trials, Newcastle-Ottawa Quality Assessment Scale for non-randomized studies and for case reports Murad et al tool will be used. Decision to conduct metanalysis will be based on several factors including homogeneity and outcome measures reported, otherwise a narrative synthesis will be deemed appropriate.

Discussion

This systematic review will summarise the existing data on effect of COVID-19 on maternal and perinatal population. Furthermore, to the best of our knowledge, this is the first systematic review addressing therapeutic management and safety of medicines to treat COVID-19 during pregnancy and breastfeeding.

Systematic review registration

Prospero ([CRD42020181163](https://doi.org/10.1111/1747-7528.12163))

Introduction

The novel coronavirus now known as SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) has spread globally (1). On 12th March 2020, the World Health Organization (WHO) declared it as a 'pandemic' outbreak of utmost international concern (2,3). The first coronavirus patient was identified in the Wuhan City, China in December 2019 (4,5). Previous infectious outbreaks such as, H1N1 influenza virus, Zika virus, severe acute

respiratory syndrome corona virus (SARS-CoV) and Middle East respiratory syndrome corona virus (MERS-CoV) have had significant impact on maternal as well as perinatal outcomes (6,7).

Epidemiology, transmission and symptoms

Ever since the first case reported in China, the infection has been spreading at an accelerating pace, across 213 countries, infecting over 3.7 million people, 261,380 cases of death (as of 05th May, 2020), at an estimated case fatality ranging 0.5% - 6% (3). Adults and older people are more susceptible to the infection in comparison to younger age group, while the infection is predominantly reported among men, the rate of infection among children is scarce (8,9). Furthermore, co-morbidities such as diabetes, hypertension, cardiovascular or respiratory disorders are significantly associated with poor outcomes and substantially increased mortality rate (10,11).

While vast majority of cases are categorised as 'mild' or 'uncomplicated', almost 14% of patients require hospitalization and/or oxygen support and approximately 5% required intensive care admission to survive (12). Common clinical manifestations reported among hospitalized patients were fever (83-100%), cough (59-82%), myalgia (11-35%), headache (7-8%), and diarrhea (2-10%). Almost all patients had abnormal radiographic chest imaging (5,13,14).

The National Institutes of Health (NIH) in the United States has since developed a tool to categorize the severity of the disease, based on the presenting symptoms (Table 1).

The transmission routes of COVID-19 include direct transmission, droplet inhalation transmission, and contact transmission. (15).

COVID-19 and its impact on pregnancy

The 'obstetric' population is considered vulnerable as different stages of pregnancy involves multiple interactions with the health care system, therefore managing pregnant population presents a unique challenge during this pandemic. Additionally, the physiological changes and partial immune suppression during pregnancy makes pregnant women and newborn babies susceptible to several infections. Post-partum hemorrhage, maternal sepsis, preeclampsia, premature rupture of membrane are the most common COVID-19 induced adverse events reported among pregnant women (16).

Anecdotal evidence suggests that pregnant women do not appear to be different than the general population in terms of disease transmission and to date there is no evidence of vertical transmission from mother to fetus.

There are currently limited data on first trimester or early pregnancy COVID-19 infections. One case from Iran reported poor maternal outcomes at 30-week gestation causing maternal and fetal death (17). Data from Intensive Care National Audit and Research Centre in the UK reported a marginal difference in the rate of current/recent pregnancy amongst all individuals admitted to critical care (2.3%) compared to the reported rate for non-COVID viral pneumonia during 2017-19 (3.3%). In one study, 47% (15/32) of pregnant women with COVID-19 infections delivered preterm, another study reported 7/15 women delivered preterm by Cesarean section (17). Cesarean section is the most commonly reported mode of delivery for patients confirmed with COVID-19 (17). Several studies have shown no detrimental effect on the fetal growth where COVID-19 positive pregnant women deliver within 13 days after the onset of symptoms (18-21).

Another study followed 13 pregnant women diagnosed with COVID19, mostly in their third trimester (11 patients < 28 weeks gestation). Three were discharged and continued to their pregnancy, however, eight underwent cesarean section due to reasons including fetal distress (3), premature rupture of the membrane or PROM (1), and stillbirth (1), multiple organ dysfunction (1), and 6 cases delivered preterm (18). Whilst state of pregnancy, reduced mobility and hospital admissions are considered as risk factors for hypercoagulable disorders, COVID-19 infection during pregnancy is more likely to cause maternal venous-thromboembolism (16).

To date, there are no valid evidence suggesting COVID-19 infection in the amniotic fluid, umbilical cord blood, or breastmilk samples.

COVID-19 and its impact on perinatal and breastfeeding

While there is scarcity of evidence demonstrating effect of COVID-19 on the fetus, factors such as mother-to-child vertical transmission and its long term and short-term effect on the offspring remain unclear. Wang [et.al](#) reported a case with positive qRT-PCR in both the mother and the neonate. The mother was admitted at 40 weeks gestation and computerized tomography (CT) images were remarkable, suggesting for COVID-19, following which, she underwent emergency cesarean section delivering the baby with normal Apgar scores. Hours after birth, the neonate had lymphocytopenia, deranged liver function, elevated creatine kinase and a pharyngeal swab collected 36 hours after birth turned positive (20).

Pharmacological management of pregnant women with COVID-19

Ever since the outbreak, the healthcare team and researchers across the globe have proposed several pharmacological interventions that could prove effective in treating COVID-19 (22). Various treatment options include, but are not limited to, antivirals and combinations, angiotensin receptor blockers, antimalarial, antihistamines, steroids, antipyretics (23). Whilst COVID-19 treatment strategies vary across different countries, the WHO and the Centers for Disease Control and Prevention (CDC) guidelines are very general, and both advise to manage pregnant and pediatric patients with caution.

Rationale for this review

Infecting over 4 million people, the global pandemic is anticipated to affect considerable number of pregnant women worldwide. The alarmingly increasing mortality rates warrants an early identification and protection of this vulnerable population. Despite favorable maternal and perinatal outcomes reported in most cases, the scientific evidence in managing this vulnerable population remains unclear (24). There is a scarcity of evidence demonstrating pharmacological management, maternal and perinatal outcomes during COVID-19.

The existing studies have predominantly originated from one geographical location, or as single case or case series. Furthermore, fewer sample size, varied findings and continuously evolving treatment protocols and ambiguity in reliability of these results make the findings difficult to interpret. Hence collating scientific evidence concerning the pandemic is imperative. Therefore, a systematic review evaluating the maternal and perinatal outcomes and therapeutic management of pregnant women diagnosed with COVID-19 is highly recommended.

Research Question(s)

This systematic review seeks to answer the following questions in relation to COVID-19,

- What are the different clinical characteristics presented in maternal and perinatal population?
- What are the different maternal and perinatal outcome measures reported?
- What are the distinct therapeutic interventions reported to treat Covid-19?
- Is it safe to use 'medications' used in the treatment of COVID-19 during antenatal, perinatal, postnatal and breastfeeding?

Methods / Design

Study registration

This systematic review was registered on the International Prospective Register of Systematic Reviews (PROSPERO) ([CRD42020181163](#)) on 23rd of April 2020 (25). (Additional file 1) This protocol is prepared in accordance with the Preferred Reporting Item for Systematic Review and Meta-analysis (PRISMA-P) statement (26) (Additional file 2), the PROSPERO record will be updated simultaneously in case of any important amendments.

Eligibility criteria

Inclusion

Type of studies

Study designs, particularly, case reports, case series, observational studies, randomized and quasi-randomized controlled trials (RCTs/CCTs), controlled before and after studies (CBAs) and interrupted time series (ITSs), will be included. Previous systematic reviews will be analyzed for cross-referencing. Only studies published in English will be included in this review.

Type of participants

The current review will include all studies reporting clinical characteristics, outcomes, treatment options, and any reported adverse events, in pregnant women diagnosed with COVID-19 and/or the fetuses and infants of mothers who are COVID-19 positive. Infants and children exposed *in-utero*, during delivery or breastfeeding will be included. There will be no restriction with respect to ethnicity, setting or location.

Type of interventions (or exposure)

Pregnant women infected and diagnosed (laboratory confirmed infection) with COVID-19 infection (exposure) will be considered as an intervention.

Type of comparators

We do not anticipate any RCTs for pregnant women diagnosed with COVID-19, however, if any, uninfected pregnant women will be the comparator/control.

Outcome measures

Outcome 1 (clinical characteristics): Clinical characteristics as illustrated by the NIH definition, will be classified as mild, moderate and severe. See table 1

Outcome 2 (maternal and perinatal outcomes): The maternal mortality and morbidity, other complications such as post-partum hemorrhage, maternal sepsis, and preeclampsia etc., mode of delivery, PPRM, ventilator support or ICU. Perinatal outcomes, APGAR scores, neonatal birth outcomes, birth weight, infections, IUGR and mortality etc.).

Outcome 3 (safety of therapeutic interventions): the primary outcome in terms of safety of the medications used in the treatment of covid-19 in pregnancy, post pregnancy and breastfeeding are presence of malformation, still birth or child death etc. secondary outcome includes drug interactions, drug disease interactions, any adverse drug reactions, other minor adverse drug events or medication errors reported due to the pharmacological agents used in the treatment of covid19 infections among pregnant women.

Exclusion

Studies that do not report/present a laboratory confirmed infection will be excluded. Studies published in any other language other than English will be excluded. Studies reporting covid-19 infections in non-pregnant or women with gynecological complications. Data obtained from reviews, personal opinions, commentaries, conference presentations will be excluded. Also studies published before 12 December 2019 will be excluded

Databases and search strategy

Literature search

The search will follow a comprehensive, sequential three step search strategy.

The first step will be an initial limited search that will include two most relevant online databases (MEDLINE/PubMed and CINAHL), we will identify the key words in the title and abstract of retrieved papers, and the index terms used to describe the articles.

In the second search, we will use all identified keywords and index terms to search across different databases. All search terms used for the search will be added in the appendix. Various boolean operators (AND/OR/NOT), truncation, wildcards etc. either individually or in combination will be used to ensure the comprehensiveness of the search process.

The electronic database that will be searched are,

1. MEDLINE®
2. PubMed®
3. The Cumulative Index of Nursing and Allied Health Literature (CINAHL®)
4. LitCovid
5. ScienceDirect
6. The Cochrane Database of Systematic Reviews (CDSR)
7. The Centre for Reviews and Dissemination (CRD)
8. Joanna Briggs Institute Library (JBI)

Thirdly, the reference list of identified, relevant articles will be searched for additional studies. In case of missing information/incomplete data, the authors of primary studies will be contacted.

Full text articles reporting information about maternal or perinatal outcomes during Covid-19, therapeutic interventions used in treatment and evidence demonstrating its safety during antenatal, perinatal, postnatal or breastfeeding while using COVID-19 therapeutic options.

All studies published in English from the inception of this disease (December 2019) until the completion of this review will be included.

The quality of this search strategy will be assessed by at least two independent researchers using the Peer Review of Electronic Search Strategies Evidence-Based Checklist (PRESS EBC) (27)

Study selection

A study eligibility form will be developed for screening the titles, abstracts and potentially relevant full-text articles. The form will be developed by BT & DS and shared across the team, the form will be tested and revised if necessary. Inter-rater reliability will be calculated from the pilot-test and screening will only commence after high agreement (e.g., kappa statistic $\geq 60\%$) is observed. All the articles will be individually screened by 2 authors, BT/DS, any disagreements will be clarified by contacting a third reviewer (PAR). The studies will be classified into three categories: (1) included, (2) excluded, and (3) pending. Pending articles will be considered as those articles that initially appear to incompletely fit the exclusion and/or the inclusion criteria. A decision to include or exclude the pending articles will be made through discussion and mutual consensus. All screening procedures will be presented using a PRISMA flow diagram.

Data extraction

All studies identified will be sent to RefWorks and the duplicated will be removed. Data abstraction will be performed by two independent reviewers (BT and PAR) using a standardized template to independently apply the inclusion and exclusion criteria, any disagreements will be resolved by discussion between the two, or else a third reviewer (FAH) will be contacted if no consensus are reached. The extraction will include four main categories,

(i) General Characteristics of the reviewed studies such as, country, month/year of publication, study objective(s), study design, period of study, and type of hospital, number of beds, and number of patients.

(ii) Maternal outcomes: number of patients, age, parity, pregnancy outcome, ethnicity, presence of comorbidities, and any other pregnancy related complications, maternal mortality, and other outcome related data.

Perinatal outcomes: gestational age, neonatal compromise (APGAR scores), neonatal birth outcomes, birth

weight, infections, mal-presentation, intrauterine growth restriction and mortality etc.), the study will also include data on (vertical transmission if available).

(iii) Pharmacological interventions: drug name, pharmacological class, dose, route, frequency, concurrent medications used etc.

(iv) Safety of medications used: if any reported drug interactions, adverse drug reactions, adverse drug events or any other complications due to pharmacological agents used.

Assessment of risk of bias

In order to achieve a consistency in assessing the risk of bias, the review team will pre-assess a sample of selected studies, and the result of the assessment will be shared among the assessors as well as the review team and discussed. Pairs of reviewers will independently assess quality of each of the studies (BT & PAR). The Cochrane Risk of Bias tool (28) will be used for randomized controlled trials (if any) and for non-randomized studies we plan to use the Newcastle–Ottawa Quality Assessment Scale (NOS) (29). NOS is a three-dimensional appraisal tool to assess the methodological quality that includes, (i) selected population, (ii) comparability of groups, and (iii) outcome of interest. Considering scarcity of information around this topic, no studies will be excluded based on the assessment of risk of bias. The study team may also include Murad et al tool for assessing quality of case reports (30).

Data synthesis and analysis

If the studies are found heterogeneous, a narrative synthesis method will be considered as appropriate, all included studies will be summarized in the form of a detailed commentary in accordance to the data extraction form summarized in the previous section. Tables and graphs will be created to illustrate the key study characteristics such as (population characteristics, maternal or perinatal outcomes, sample sizes, settings, medications used, results, and any other important aspect related to each research question of interest). Descriptive analysis will be performed for all categorical variables using frequency, mean and standard deviation wherever possible. If the studies are found similar enough to be pooled, a random effect meta-analysis will be applied. Statistical heterogeneity between studies will be assessed using τ^2 and I^2 statistics and $P < 0.10$ and $I^2 > \text{than } 50\%$, will be considered as a high level of statistical heterogeneity between the studies.

Quality of evidence

Two authors will independently assess the quality of evidence for each outcome. The quality of evidence will be determined by Grading of Recommendations Assessment, Development, and Evaluation (GRADE) system (31). The studies will be rated based on their evidence ranging high, moderate, low, or very low level. Individual outcomes will be assessed for following aspects, limitations, inconsistency, indirectness, imprecision, and publication bias.

Discussion

As pandemic continues to affect a large number of pregnant women worldwide, the findings of this systematic review will contribute to improve to the current state of knowledge about effect of COVID-19 on maternal and perinatal population. Furthermore, to the best of our knowledge, this is the first systematic review addressing therapeutic management and safety of medicines to treat COVID-19 during pregnancy and breastfeeding.

Pooling of data for meta-analysis seems difficult as we anticipate a considerable degree of heterogeneity between the studies in terms of study design, exposure measurement, characteristics of participants, and outcomes.

To ensure a wider dissemination, we will present the interim findings at local and international conferences, and publish the findings in high impact open access journals. Furthermore, we will brief the findings to key stakeholders, policymakers, antimicrobial stewardship committee, hospital infection control committee etc.

Strengths and limitations of this study

- To the best of our knowledge, this will be the first review to use a validated systematic and transparent methodological process to appraise and summarize the existing literature on COVID-19 associated pregnancy and perinatal outcomes.
- To the best of our knowledge, this systematic review is the first to address the pharmacological management in COVID-19 and its impact on maternal and perinatal outcomes.
- The proposed systematic review is in accordance to the Preferred Reporting Items for Systematic Reviews and Meta Analyses guidelines (PRISMA), ensuring consistency and uniformity in reporting the full systematic review.
- Two independent reviewers will screen the eligible studies and perform extraction and quality assessment to minimize potential reviewer bias.
- One key limitation of this review is it will only include studies published in English, while majority of cases were reported from China, and there are chances that we may miss information published in Chinese or other regional languages. This limitation may cause bias.
- There is a possibility of reporting bias, as we plan to exclude no studies based on the quality and there is a possibility that findings originate from low and inconsistent studies.

This systematic review is anticipated to summarise the available evidence that examined the impact of COVID 19 infections in maternal and perinatal outcomes. Finally the results will be most likely to be used to provide recommendations to the current topic of concern and provide endorsements to future research.

Declarations

Ethics

No ethics approval is required for this protocol as it is based on information from the published data.

Acknowledgements

The study team would like to acknowledge PROSPERO International prospective register of systematic reviews and the editorial for an expedited review.

Author's contribution

All authors have made substantial contributions to the conception or design, acquisition, analysis, or interpretation of data for the review. All authors reviewed and approved the final version of the manuscript prior to submission.

Funding

The project will seek funding from the Hamad Medical Corporation's Medical Research Center.

Conflict of interest

All authors declare no conflict of interest.

References

- (1) Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, et al. World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). *International Journal of Surgery*. 2020;.
- (2) The World Health Organisation. *Coronavirus disease (COVID-19) outbreak*. [homepage on the Internet]. 2020 [updated May/05; cited 2020 May/05]. Available from: <https://www.who.int/westernpacific/emergencies/covid-19>.

- (3) The World Health Organisation. *Coronavirus disease (COVID-19) Pandemic*. [homepage on the Internet]. World Health Organisation; 2020 [updated April/26; cited 2020 April/26]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>.
- (4) World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19-11 March 2020. *Geneva, Switzerland*. 2020;.
- (5) Guo Y, Cao Q, Hong Z, Tan Y, Chen S, Jin H, et al. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Military Medical Research*. 2020; 7(1):1-10.
- (6) Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Da Fonseca EB, et al. Characterizing the pattern of anomalies in congenital Zika syndrome for pediatric clinicians. *JAMA pediatrics*. 2017; 171(3):288-295.
- (7) Rasmussen SA, Jamieson DJ, Honein MA, Petersen LR. Zika virus and birth defects—reviewing the evidence for causality. *New England Journal of Medicine*. 2016; 374(20):1981-1987.
- (8) Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*. 2020; 395(10223):497-506.
- (9) Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical characteristics of coronavirus disease 2019 in China. *New England journal of medicine*. 2020; 382(18):1708-1720.
- (10) Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*. 2020; 323(13):1239-1242.
- (11) Yang S, Cao P, Du P, Wu Z, Zhuang Z, Yang L, et al. Early estimation of the case fatality rate of COVID-19 in mainland China: a data-driven analysis. *Annals of Translational Medicine*. 2020; 8(4).
- (12) Surveillances V. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19)—China, 2020. *China CDC Weekly*. 2020; 2(8):113-122.
- (13) Adhikari SP, Meng S, Wu Y, Mao Y, Ye R, Wang Q, et al. Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: a scoping review. *Infectious diseases of poverty*. 2020; 9(1):1-12.
- (14) Lai C, Shih T, Ko W, Tang H, Hsueh P. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and corona virus disease-2019 (COVID-19): the epidemic and the challenges. *International journal of antimicrobial agents*. 2020;:105924.
- (15) Mullins E, Evans D, Viner R, O'Brien P, Morris E. Coronavirus in pregnancy and delivery: rapid review. *Ultrasound in Obstetrics & Gynecology*. 2020;.
- (16) The Royal College of Obstetricians and Gynaecologists (RCOG). *Coronavirus (COVID-19) Infection in Pregnancy*. London, United Kingdom: Royal College of Obstetricians and Gynaecologists; 2020.
- (17) Karami P, Naghavi M, Feyzi A, Aghamohammadi M, Novin MS, Mobaien A, et al. Mortality of a pregnant patient diagnosed with COVID-19: A case report with clinical, radiological, and histopathological findings. *Travel Medicine and Infectious Disease*. 2020;:101665.
- (18) Liu Y, Chen H, Tang K, Guo Y. Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy. *The Journal of infection*. 2020;.
- (19) Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *The Lancet*. 2020; 395(10226):809-815.

- (20) Wang1a X, Zhou2a Z, Zhang J, Zhu F, Tang Y, Shen X, et al. A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clin Infect Dis*. 2020;.
- (21) Zhu H, Wang L, Fang C, Peng S, Zhang L, Chang G, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Translational pediatrics*. 2020; 9(1):51.
- (22) Tobaiqy M, Qashqary M, Al-Dahery S, Mujallad A, Hershian AA, Kamal MA, et al. Therapeutic Management of COVID-19 Patients: A systematic review. *Infection Prevention in Practice*. 2020;:100061.
- (23) Gautret P, Lagier J, Parola P, Meddeb L, Mailhe M, Doudier B, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International journal of antimicrobial agents*. 2020;:105949.
- (24) Della Gatta AN, Rizzo R, Pilu G, Simonazzi G. COVID19 during pregnancy: a systematic review of reported cases. *American Journal of Obstetrics and Gynecology*. 2020;.
- (25) Thomas B, Al Hail M, Abdulrouf P, Elkassem W, Tarannum A, Al Hail F, Rijims M, Parappil H, Stewart D. Clinical characteristics, maternal and perinatal outcomes and safety of therapeutic interventions during Covid-19: a systematic review protocol. 23 April. [serial on the Internet] 2020 PROSPERO International prospective register of systematic reviews(CRD42020181163):5th March 2020. Available from: https://www.crd.york.ac.uk/prospERO/display_record.php?RecordID = 181163.
- (26) Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. 2015; 4(1):1.
- (27) McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS peer review of electronic search strategies: 2015 guideline statement. *Journal of clinical epidemiology*. 2016; 75:40-46.
- (28) Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ (Clinical research ed.)*. 2011; 343:d5928.
- (29) Wells G, Shea B, O'connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Quality Assessment Scale (NOS) for assessing the quality of nonrandomized studies in meta-analyses. *Clin Epidemiol [Internet]*. 2017;:1-2.
- (30) Murad MH, Sultan S, Haffar S, Bazerbachi F. Methodological quality and synthesis of case series and case reports. *BMJ evidence-based medicine*. 2018; 23(2):60-63.
- (31) GRADE Working Group. *Grading of recommendations assessment, development and evaluation*. 2017.

Table

Severity	Symptoms
Asymptomatic	Positive test for SARS-CoV-2 but no symptoms
Mild illness	Any signs and symptoms (eg, fever, cough, sore throat, malaise, headache, muscle pain) without shortness of breath, dyspnea, or abnormal chest imaging.
Moderate illness	Evidence of lower respiratory disease by clinical assessment or imaging and a saturation of oxygen (SaO ₂) >93 percent on room air at sea level.
Severe illness	Respiratory frequency >30 breaths per minute, SaO ₂ ≤93 percent on room air at sea level, ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO ₂ /FiO ₂) <300, or lung infiltrates >50 percent.
Critical illness	Respiratory failure, septic shock, and/or multiple organ dysfunction.

Table 1: NIH COVID-19 severity assessment scale.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CRD42020181163.pdf](#)
- [PRISMA2009checklist.doc](#)