

# The Effect of Maternal Body Mass Index on In Vitro Fertilization-Intracytoplasmic Sperm Injection Treatment: A Prospective Comparative Study

Zeynep Ozturk Inal

Konya Education and Research Hospital

Hasan Ali Inal (✉ [dr.hasanaliinal@yahoo.com](mailto:dr.hasanaliinal@yahoo.com))

Konya Education and Research Hospital <https://orcid.org/0000-0002-8361-7908>

---

## Research Article

**Keywords:** assisted reproductive techniques, body mass index, ovulation induction, clinical pregnancy rates

**Posted Date:** April 12th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-298772/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

# Abstract

**Purpose:** To evaluate whether or not maternal body mass index (BMI) has an effect on pregnancy rates following in vitro fertilization-intracytoplasmic sperm injection (IVF-ICSI) treatment.

**Methods:** A total of 869 patients who had undergone IVF-ICSI treatment between 2012 and 2017 were included in this study. The participants were stratified according to maternal BMI as Group 1 ( $\text{BMI} < 25 \text{ kg/m}^2$ ;  $n=394$ ), Group 2 ( $25 \text{ kg/m}^2 \leq \text{BMI} < 30 \text{ kg/m}^2$ ;  $n=303$ ), and Group 3 ( $\text{BMI} \geq 30 \text{ kg/m}^2$ ;  $n=172$ ). Basal parameters and IVF-ICSI outcomes were compared between the groups.

**Results:** While there were no differences between the groups in terms of age, smoking status, etiology of infertility, thyroid-stimulating hormone, prolactin levels, antral follicle count, and stimulation protocol ( $p > 0.05$ ), there was significant statistical difference ( $p < 0.05$ ) in terms of BMI, duration of infertility, baseline follicle-stimulating hormone, luteinizing hormone, estradiol ( $E_2$ ), duration of stimulation, total gonadotropin dose required, peak  $E_2$  levels, progesterone levels, endometrial thickness on hCG administration, and cycle cancellation rate. In addition, the numbers of MII and 2PN oocytes retrieved and the rates of clinical pregnancy, live births, and miscarriages were also different between the groups ( $p < 0.05$ ).

**Conclusion:** Our data suggest that there is an inverse impact of increased BMI on laboratory and reproductive outcome parameters of IVF-ICSI treatment. Taking cost-effectiveness into consideration, weight loss should be suggested before ovulation is induced.

## Introduction

Obesity is a serious public health problem in developed and developing countries. Historically more prevalent in more advanced age groups, it is now seen with increasing frequency in reproductive ages where endocrinological effects such as hypothalamic-pituitary ovarian axis disorder can cause changes in the secretion of pulsatile gonadotropin, sex hormone-binding globulin, and ovarian androgen. This can prompt menstrual irregularity, anovulation, insulin resistance, have negative psychological and social effects, and can increase the risk of infertility by three times (1-3). It has also been demonstrated that weight loss can return women to spontaneous ovulation and pregnancy without any additional treatment (4-6). Elsewhere, it has been reported that obesity reduces fecundity and increases the rate of miscarriage through a negative effect on endometrial receptivity (7). The adverse effects of pre-pregnancy obesity on perinatal (e.g., preeclampsia, gestational diabetes mellitus, preterm labor, and surgical delivery) and neonatal (e.g., macrosomic fetus) outcomes have also been reported (2). Additionally, obesity has been found to have a negative effect on serum testosterone and estrogen in men and to decrease sperm motility and quality (4,5).

There are conflicting results in the literature regarding the effects of maternal obesity on the success of assisted reproductive techniques (ART). Although some studies have found that maternal body mass index (BMI) has no negative effect on ART outcomes (8-11), others show that a higher BMI increases the amount of gonadotropin required, produces fewer oocytes, increases IVF-ICSI cancellation rate, decreases clinical pregnancy and live birth rates, and increases miscarriage rate (12-14). Since the effects of obesity on ART outcomes have not been fully elucidated, the current study sought to evaluate whether or not maternal BMI has an effect on pregnancy rates following in vitro fertilization-intracytoplasmic sperm injection (IVF-ICSI) treatment.

## Materials And Method

### Study participants and data collection

This prospective study was carried out at Ali Kemal Belviranlı Women's Health and Children's Hospital, IVF Unit. Outcomes of 757 fresh ICSI cycles were reviewed between January 2012 and December 2017. Inclusion criteria were participants aged 20–44 years, body mass index (BMI) between 18 and 35  $\text{kg/m}^2$ , regular menstrual cycles, no uterine abnormalities in the

ultrasound, and normal baseline hormonal levels. Participants were excluded from the study if they were  $\geq 45$  years, any diseases that affect the outcome of IVF/ICSI, such as hydrosalpinx and endometriosis. The ethical board approval was given from the institutional review board (2012/57). Written and oral informed agreement was given from the participants.

Data were obtained for age, BMI ( $\text{kg}/\text{m}^2$ ), smoking status, infertility period, cause of infertility, the baseline at day 3 for follicle-stimulating hormone (FSH), luteinizing hormone (LH), and estradiol ( $\text{E}_2$ ) levels, thyroid-stimulating hormone (TSH), prolactin, antral follicle count, stimulation parameters, cycle cancellation rate, IVF-ICSI outcomes, CPR, live birth, and miscarriage rates.

### **Ovarian stimulation and oocyte retrieval**

Controlled ovulation stimulation was negotiated using the gonadotropin-releasing hormone agonist (GnRHa) or the flexible gonadotropin-releasing hormone antagonist (GnRHant) protocol.

### **ET Procedure**

Two senior physicians performed the ETs accompanied ultrasonographic appearance (Logiq 200 Pro, General Electric, Seoul, South Korea) using an embryo transfer catheter system. A sterile speculum was introduced to the vagina in the lithotomy position and the vagina and the cervix were cleared using sterile cotton swabs.

An embryologist loaded the embryos into a soft transfer catheter which was advanced to the ET physician who deposited the embryos approximately 10 mm from the uterine fundus under USG. The catheter was gently removed after 5 seconds. In cases of ET with external guidance, an initial catheter with inner sheath was inserted into the external cervical os, and then advanced through the cervical canal and internal os to 10 mm of the uterine fundus using USG. The internal sheath was withdrawn, and a second catheter loaded with embryos was introduced in its place and advanced to approximately 10 mm from the uterine fundus where the embryos were deposited. Difficult transfers required the use of a stylet in addition to this form of external guidance.

All catheters were immediately checked for retained embryos, blood, and the patient remained in the Trendelenburg position for about 10 minutes. Patients in whom tenaculum were excluded from the study. Luteal phase support was provided with progesterone in the form of Crinone 8% gel (Serono, Istanbul, Turkey) at a daily dose of 90 mg. Baseline parameters and IVF-ICSI outcomes were compared between the groups. Biochemical pregnancy was detected with a by hCG levels in venous blood tests performed 12-14 days after embryo transfer, and clinical pregnancy was accepted as those with a gestational sac accompanying fetal heart-beat on ultrasound examination at 4-5 weeks after embryo transfer. Live birth was defined as the birth of a live fetus after 22 weeks of gestational age. The subjects were stratified according to the maternal BMI as Group 1 ( $\text{BMI} < 25 \text{ kg}/\text{m}^2$ ;  $n = 394$ ), Group 2 ( $25 \text{ kg}/\text{m}^2 \leq \text{BMI} < 30 \text{ kg}/\text{m}^2$ ;  $n = 303$ ), and Group 3 ( $\text{BMI} \geq 30 \text{ kg}/\text{m}^2$ ;  $n = 172$ ). Basal parameters, clinical and laboratory IVF-ICSI outcomes, and reproductive outcome parameters were compared between the groups.

### **Statistical analysis**

The statistical analyses were performed using SPSS 15.0 for Windows (SPSS, Chicago, IL, USA). The Kolmogorov-Smirnov test was used for examining the continuous variables with normal and non-normal distributions. The one-way analysis of variance (ANOVA) for normally distributed variables and the Kruskal-Wallis test for not-normally distributed variables were used to compare groups. Categorical data were examined by Pearson's chi-square test, and Fisher's exact test was applied if the expected frequency was less than 5 in  $>20\%$  of all cells. The continuous variables were presented as the mean  $\pm$  standard deviation (SD) and the categorical variables were demonstrated as the number of cases and percentages. The Bonferroni-adjustment was used to control the type I errors for all possible multiple comparisons. A  $p < 0.05$  value was established as statistically significant.

## Results

A total of 51 patients were excluded from the study, specifically those with age  $\geq 45$  (n=19), BMI  $\geq 35$  kg/m<sup>2</sup> (n=14), systemic disease (n=9), endocrine or metabolic disorders (n=6), and concomitant medication (n=3). The remaining 869 participants were classified into the three groups and their outcomes analyzed (Figure 1).

A comparison of the sociodemographic and stimulation characteristics of the participants is provided in Table 1. While there were no differences between the groups in terms of age, smoking status, etiology of infertility, thyroid-stimulating hormone, prolactin levels, antral follicle count, and stimulation protocol ( $p > 0.05$ ), there was significant statistical difference ( $p < 0.05$ ) in terms of BMI, duration of infertility, baseline follicle-stimulating hormone, luteinizing hormone, estradiol (E<sub>2</sub>), duration of stimulation, total gonadotropin dose required, peak E<sub>2</sub> levels, progesterone levels, endometrial thickness on hCG administration, and cycle cancellation rate.

The laboratory and reproductive outcomes of the participants are summarized in Table 2. The numbers of MII oocytes and 2PN retrieved and the rates of clinical pregnancy, live births, and miscarriages were also different between the groups ( $p < 0.05$ ).

## Discussion

We found that overweight and obese patients with higher BMIs had worse responder rates, lower peak E<sub>2</sub> levels, and less endometrial thickness and required higher gonadotropin doses than the normal weight group. In addition, lower numbers of MII and 2PN oocytes were retrieved, the clinical pregnancy and live birth rates were lower, and the miscarriage rate was higher.

Obesity has historically been observed more frequently in adult and advanced age groups. It is now a global epidemic and has become an important public health problem in younger age groups, too (2). Infertile women, including obese patients, in the reproductive age group benefit from ART to fulfill their fertility requirements, and so the possible effects of increased BMI on ART are of great importance for the clinician, the patient, and public health (3).

Endocrinological and paracrinological factors play a role in the interaction between embryo and endometrium for successful implantation and live birth (15). Hyperandrogenemia, insulin resistance, and abnormal hormone levels that occur with increased BMI can negatively affect this process (1,16). A higher BMI also affects the levels of inflammatory markers such as insulin-like growth factors, tumor necrosis factor-alpha, and interleukine-6 which play roles in cell differentiation and differentiation, folliculogenesis, oocyte maturation, and embryo development. As a result, embryo implantation can be negatively affected and the risk of miscarriage can increase (17-19). Previous studies have shown that increased BMI is associated with poor IVF-ICSI outcomes through the effect of these endocrinological factors (2,20-21) with one study reporting that a reduction of one BMI unit can increase the chance of pregnancy by 19% (22). It has also been found that advanced maternal age and smoking negatively affect live birth rates (23), although the mean age and smoking rates were similar between the groups in our study.

In the literature, conflicting results exist regarding the effects of increased BMI on ART outcomes. For example, Fedorcsek et al. (13) evaluated 5019 IVF-ICSI cycles and found no significant difference in live birth rates between obese and normal-weight women (41.4% vs 50.3%). Similarly, Wittemer et al. (24) and Dokras et al. (25) show that BMI has no negative effect on rates of clinical pregnancy or live birth. On the other hand, however, Luke et al. (26) evaluated approximately 45,000 embryo transfers and show that an increased BMI decreases clinical pregnancy and live birth rates and that this effect is especially pronounced in women under 35 years of age. A separate meta-analysis of 33 studies and 47,967 IVF-ICSI cycles found that obese and overweight patient groups had poorer outcomes compared to normal weight women and that the obese group was worse than the overweight patient group (1). These findings were in agreement with our results. Sartorius et al. (25) demonstrate that an increased BMI can have other negative perinatal outcomes such as preeclampsia, preterm birth, and surgical delivery, as well as reducing live birth rates. Relatedly, Pinborg et al. (6) evaluate 1,417 IVF-ICSI cycles and show that

the cancellation rate increases with increased BMI, the two key reasons being that obesity makes the oocyte pick-up procedure more difficult and insufficient follicles are developed despite the use of high gonadotropin doses.

The possible negative effects of an increased BMI on ART should be explained to overweight and obese women who are scheduled for IVF-ICSI treatment. Before the process begins, it should also be explained that weight loss can increase the chance of success in terms of pregnancy and live birth. Overweight and obese women should consequently be encouraged to lose weight, and clinicians planning ART should implement weight loss programs involving diet and exercise. In addition, it should be understood that the gonadotropin dose required will increase with higher BMIs meaning that treatment costs will also grow despite the increasingly negative perinatal outcomes.

The strong point of the current study consist of its prospective arrangement, the adequate number of subjects in each group, and the prototypical sample from central Turkey. The results can be generalized to most of the country's population. However, the potential limitations of the study are that it was conducted in a tertiary single care institution and that the cumulative CPR was not evaluated because no frozen ETs were included.

In conclusion, this study found that an increased BMI has a negative effect on ART outcomes as shown in decreasing clinical pregnancy and live birth rates and increasing miscarriage rates. Further studies with more participants are needed to elucidate this effect.

## Declarations

### Compliance with ethical standards

**Ethics Committee Approval:** Ethics committee approval was received for this study from the local ethics committee of Necmeddin Erbakan University Medical Faculty (reference number: 2012/57). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Informed Consent:** Written informed consent was obtained from patients who participated in this study.

**Author Contributions:** Concept – Z.O.I., H.A.I.; Design – Z.O.I., H.A.I.; Supervision – Z.O.I., H.A.I.; Materials – Z.O.I., H.A.I.; Data Collection and/or Processing – Z.O.I., H.A.I. and/or Interpretation - Z.O.I., H.A.I.; Literature Review – Z.O.I., H.A.I.; Writer – Z.O.I., H.A.I.; Critical Review - Z.O.I., H.A.I.

**Conflict of interest:** Zeynep Ozturk Inal and Hasan Ali Inal declare that they have no conflict of interest.

**Financial Support:**None

**Presented at a meeting:** Presented as an oral presentation at 16th National Congress of Gynecology and Obstetrics, 9-13 May 2018, in Antalya, Turkey.

## References

1. Rittenberg V, Seshadri S, Sunkara SK, Sobaleva S, Oteng-Ntim E, El-Toukhy T. Effect of body mass index on IVF treatment outcome: an updated systematic review and meta-analysis. *Reprod Biomed Online*. 2011;23:421-39.
2. Wang X, Hao J, Zhang F, Li J, Kong H, Guo Y. Effects of female and male body mass indices on the treatment outcomes and neonatal birth weights associated with in vitro fertilization/intracytoplasmic sperm injection treatment in China. *Fertil Steril*. 2016;106:460-6.
3. Gorkem U, Inal ZO, Inal HA, Bostanci MO. Serum progranulin levels are elevated in infertile women with obesity. *Endokrynol Pol*. 2018;69:661-6.

4. Davidson LM, Millar K, Jones C, Fatum M, Coward K. Deleterious effects of obesity upon the hormonal and molecular mechanisms controlling spermatogenesis and male fertility. *Hum Fertil (Camb)* 2015;18:184–93.
5. Inal HA, Kahyaoglu I, Turkmani A, Tuzluoglu D, Yilmaz N. Retrospective comparison of intracytoplasmic sperm injection outcomes of sperm retrieved from a testicular biopsy and freshly ejaculated semen in oligozoospermia. *Rev Int Androl*. 2018;16:131-6.
6. Pinborg A, Gaarslev C, Hougaard CO, Nyboe Andersen A, Andersen PK, Boivin J, Schmidt L. Influence of female bodyweight on IVF outcome: a longitudinal multicentre cohort study of 487 infertile couples. *Reprod Biomed Online*. 2011;23:490-9.
7. Sharifzadeh F, Kashanian M, Jouhari S, Sheikhsari N. Relationship between pre-pregnancy maternal BMI with spontaneous preterm delivery and birth weight. *J Obstet Gynaecol* 2015;35:354–7.
8. Maheshwari A, Stofberg L, Bhattacharya S. Effect of overweight and obesity on assisted reproductive technology a systematic review. *Hum. Reprod. Update* 2007;13: 433–4.
9. Lashen H, Ledger W, Lopez Bernal A, Barlow D. Extremes of body mass do not adversely affect the outcome of superovulation and in vitro fertilization. *Hum. Reprod* 1999;14:712–5.
10. Dechaud H, Anahory T, Reyftmann L, Loup V, Hamamah S, Hedon B. Obesity does not adversely affect results in patients who are undergoing in vitro fertilization and embryo transfer. *Eur J Obstet Gynecol Reprod Biol* 2006;127:88–93.
11. Matalliotakis I, Cakmak H, Sakkas D, Mahutte NG, Koumantakis G, Arici A. Impact of body mass index on IVF and ICSI outcome: a retrospective study. *Reprod Biomed Online* 2008;16: 778–83.
12. Wang JX, Davies M, Norman RJ. Body mass and probability of pregnancy during assisted reproduction treatment: retrospective study. *BMJ* 2000;321:1320–1.
13. Fedorcsak P, Dale PO, Storeng R, Ertzeid G, Bjercke S, Oldereid N, Omland AK, Abyholm T, Tanbo T. Impact of overweight and underweight on assisted reproduction treatment. *Hum Reprod* 2004;19:2523–8.
14. Veleza Z, Tiitinen A, Vilska S, Hyde'n-Granskog C, Toma's C, Martikainen H, Tapanainen JS. High and low BMI increase the risk of miscarriage after IVF/ICSI and FET. *Hum Reprod* 2008;23:878–84.
15. Inal ZO, Inal HA, Erdem S. The effect of serum and follicular fluid secreted frizzled-related protein-5 on in vitro fertilization outcomes in patients with polycystic ovary syndrome. *Mol Biol Rep*. 2018;45:2037-2044.
16. Brewer C, Balen A. The adverse effects of obesity on conception and implantation. *Reproduction* 2010;140:347–64.
17. Ozturk Inal Z, Yilmaz N, Inal HA, Hancerliogullari N, Coskun B. Are there any differences between antagonist administration on days <6 and ≥6 of Controlled Ovarian Hyperstimulation on assisted reproductive technique outcomes? *J Chin Med Assoc*. 2018;81:53-7.
18. Ma Y, Zhu MJ, Zhang L, Hein SM, Nathanielsz PW, Ford SP. Maternal obesity and overnutrition alter fetal growth rate and cotyledonary vascularity and angiogenic factor expression in the ewe. *Am J Physiol Regul Integr Comp Physiol* 2010;299:249–58.
19. Dimitriadis E, Nie G, Hannan NJ, Paiva P, Salamonsen LA. Local regulation of implantation at the human fetal–maternal interface. *Int J Dev Biol* 2010;54:313–22.
20. Bellver J, Melo MA, Bosch E, Serra V, Remohí J, Pellicer A. Obesity and poor reproductive outcome: the potential role of the endometrium. *Fertil Steril* 2007;88:446–51.
21. Zhang JJ, Feret M, Chang L, Yang M, Merhi Z. Obesity adversely impacts the number and maturity of oocytes in conventional IVF not in minimal stimulation IVF. *Gynecol Endocrinol* 2015;31:409–13.
22. Ferlitsch K, Sator MO, Gruber DM, Rucklinger E, Gruber CJ, Huber JC. Body mass index, follicle-stimulating hormone and their predictive value in in vitro fertilization. *J Assist Reprod Genet* 2004;21:431–6.
23. Lintsen AM, Pasker-de Jong PC, de Boer EJ, Burger CW, Jansen CA, Braat DD, et al. Effects of subfertility cause, smoking and bodyweight on the success rate of IVF. *Hum Reprod* 2005;20:1867–75.

24. Wittemer C, Ohl J, Bailly M, Bettahar-Lebugle K, Nisand I. Does body mass index of infertile women have an impact on IVF procedure and outcome? J Assist Reprod Genet 2010;17:547–52.

25. Dokras A, Baredziak L, Blaine J, Syrop C, VanVoorhis BJ, Sparks A. Obstetric outcomes after in vitro fertilization in obese and morbidly obese women. Obstet Gynecol 2006;108:61–9.

26. Luke B, Brown MB, Stern JE, Missmer SA, Fujimoto VY, Leach R. Female obesity adversely affects assisted reproductive technology (ART) pregnancy and live birth rates A SART Writing Group. Hum Reprod 2011;26:245–52.

27. Sartorius GA, Nieschlag E. Paternal age and reproduction. Hum Reprod Update 2010;16:65–79.

Tables

**Table 1.** Demographic and stimulation characteristics of the patients.

		BMI<25 (Group 1) (n=394)	25≤BMI<30 (Group 2) (n=303)	BMI≥30 (Group 3) (n=172)	p		
					1 vs 2	1 vs 3	2 vs 3
Age (years)		30.03±4.55	30.31±4.96	30.94±4.56	0.109		
BMI (kg/m²)		21.9±81.99	27.23±1.43	32.88±2.38	<0.001	<0.001	<0.001
Smoking rate (%)		7.4%	7.3%	5.8%	0.775		
Duration of infertility (years)		5.51±3.33	6.20±3.78	7.39±3.89	0.038	<0.001	0.004
Etiology of infertility (%)	Male factor	39.4%	38.1%	29.7%	0.116		
	Tubal factor	1.3%	2.6%	1.7%			
	Unexplained	34.4%	36.8%	36.1%			
	Poor responder	24.9%	22.5%	32.6%			
Baseline-FSH (IU/mL)		7.45±2.28	7.14±2.52	6.74±2.45	0.205	0.003	0.187
Baseline-LH (IU/mL)		5.93±2.77	5.14±2.80	5.08±3.11	0.001	0.003	0.975
Baseline-Estradiol (pg/mL)		46.50±16.65	42.69±16.79	42.63±15.28	0.007	0.027	0.996
Antral follicle count		6.06±2.60	6.41±2.72	5.67±2.55	0.064		
TSH (μIU/mL)		2.22±1.12	2.09±1.03	2.29±1.23	0.144		
Prolactin (ng/mL)		16.70±9.40	16.0±28.01	15.34±9.37	0.231		
Stimulation protocol (%)	Long	17.8%	19.5%	20.5%	0.875		
	Antagonist	81.4%	80.1%	78.9%			
	Microdose	0.8%	0.3%	0.6%			
Duration of stimulation (days)		9.76±1.52	9.62±1.51	10.17±1.86	0.528	0.147	0.001
Gonadotropin dose (IU)		1929.63±912.79	2008.26±883.51	2234.60±1019.92	0.511	0.001	0.030
Estradiol levels on day hCG (pg/mL)		2015.65±1140.77	1802.17±1063.92	1705.64±1323.19	0.043	0.010	0.660
Progesterone levels on day hCG (pg/mL)		0.89±0.39	0.79±0.37	0.69±0.37	0.002	<0.001	0.031
Endometrial thickness on day hCG (mm)		10.3±91.62	10.28±1.74	9.86±1.65	0.651	0.002	0.028
Endometrial thickness on transfer day (mm)		10.52±1.66	10.44±1.83	10.02±1.74	0.042	0.006	0.019
Cycle cancellation rate (%)		1.7%	2.6%	6.9%	0.444	0.004	0.031

BMI: body mass index; FSH: follicle stimulating hormone; LH: luteinizing hormone; TSH: thyroid stimulating hormone; hCG: human chorionic gonadotropine

p<0.05 is statistically significant




**Table 2.** Laboratory and reproductive outcome parameters of the patients.

		BMI<25 (Group 1) (n=387)	25≤BMI<30 (Group 2) (n=295)	BMI≥30 (Group 3) (n=160)	p		
					1 vs 2	1 vs 3	2 vs 3
Number of oocytes retrieved		9.41±6.01	8.97±5.11	7.90±5.50	0.576	0.011	0.126
Number of MII oocytes		7.46±4.45	7.19±4.22	6.33±4.58	0.712	0.017	0.115
2 Pronucleus		5.11±3.42	4.98±3.24	4.06±2.88	0.868	0.010	0.030
Fertilization rate (%)		68.48±24.5	68.80±23.67	67.09±25.17	0.768		
Grade I embryo (%)		67.2%	66.5%	65.1%	0.231		
Number of embryo transfers (%)	Single	82.7%	78.2%	75.2%	0.105		
	Multiple	17.3%	21.8%	24.8%			
The days of embryo transfer (%)	2	3.9%	5.1%	10.1%	0.117		
	3	84.8%	85.6%	79.9%			
	5	11.3%	9.3%	10.0%			
The embryo transfer technique (%)	Easy transfer with a soft catheter	22.1%	21.5%	16.8%	0.173		
	After external guidance transfer	71.9%	69.1%	78.5%			
	Difficult transfer with a stylet	6.0%	9.4%	4.7%			
Clinical pregnancy rate (%)		40.1%	33.2%	23.8%	0.042	<0.001	0.041
Live birth rate (%)		33.6%	23.7%	13.9%	0.005	<0.001	0.017
Miscarriage rate (%)		17.7%	28.6%	44.7%	0.045	0.001	0.042

BMI: Body mass index

p&lt;0.05 is statistically significant

## Figures



**Figure 1**

Flowchart of the Study. Enrollment and follow-up of the study subjects.