

An Ozonolysis Based Method and Applications for the Non-lethal Modification of Insect Cuticular Hydrocarbons

Benjamin Alexander Savage (✉ benalexandersavage@gmail.com)

Michigan State University College of Agriculture and Natural Resources <https://orcid.org/0000-0001-6916-2743>

Henry Chung

Michigan Agricultural College: Michigan State University

Susan Masten

Michigan State University College of Engineering

Zinan Wang

Michigan Agricultural College: Michigan State University

Matthew Grieshop

Michigan Agricultural College: Michigan State University

Research Article

Keywords: Cuticular hydrocarbons, modification, ozone, *Drosophila suzukii*, desiccation resistance

DOI: <https://doi.org/10.21203/rs.3.rs-299480/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Cuticular hydrocarbons (CHCs) are important, multi-function components of the insect epicuticle. In *Drosophila* spp., CHCs provide protection from desiccation and serve as semiochemicals for both intra- and interspecific communication. We developed a non-lethal method for the modification of *Drosophila* CHCs profiles through the exposure of live insects to a high dose of ozone gas (~45,000 ppm). *Drosophila suzukii* that were treated with ozone showed a 1.63-3.10 fold reduction in unsaturated hydrocarbons with these CHCs shown to regenerate over 108 h. Changes in CHCs were correlated with significantly reduced desiccation resistance in both male and female *D. suzukii* at one h after ozone treatment. Interestingly, individuals treated with ozone showed increased desiccation resistance in comparison to controls at 108 h after ozone treatment. The methodology reported in this paper provides a novel approach to investigate the biosynthesis and functions of CHCs during the lifespan of an insect.

Introduction

The *Drosophila* spp. epicuticle contains a wide range of saturated, unsaturated, and branched hydrocarbons broadly defined as cuticular hydrocarbons (CHCs) (Bartelt et al. 1986; Jallon and David 1987; Howard et al. 2003; Howard and Blomquist 2005). *Drosophila* CHCs form a waxy layer on the cuticle that reduces desiccation as well as provides inter- and intra-specific chemical signaling (Chung and Carroll 2015). It has been hypothesized that insect cuticular desiccation resistance decreases as epicuticular lipids change from a solid to liquid state (Ramsay 1935; Hadley 1994; Gibbs 1998). Unsaturated and methyl branched alkanes melt at lower temperatures compared to saturated hydrocarbons (Gibbs and Pomonis 1995). Therefore, saturated hydrocarbons are hypothesized to confer greater cuticular desiccation resistance than unsaturated hydrocarbons at higher temperatures (Gibbs 1998). For example, *D. pseudoobscura* dwelling in arid regions had a greater abundance of long-chained saturated hydrocarbons than laboratory maintained colonies, which correlates to a lower water loss rate (WLR) (Toolson and Kuper-Simbron 1989).

Previously, ozonolysis of CHCs have been used to identify double bond positions of unsaturated hydrocarbons on dipteran and hymenopteran cuticles following extraction using chemical solvents (Bartelt et al. 1982, 1986; Antony et al. 1985). Beroza and Bierl (1967) developed a method which uses ozonolysis to identify insect unsaturated hydrocarbons. This methodology involves extracting CHCs with a non-polar solvent and then treating the extract with ozone prior to gas chromatography–mass spectrometry (GC-MS) analysis (Beroza and Bierl 1967). Antony et al. (1985) demonstrated that monoenes and dienes extracted from *Drosophila melanogaster* undergo cleavage via ozonolysis to varied extents. For example, the researchers noted a “major” reduction in 7(Z)-tricosene after ozonolysis, while 9(Z)-tricosene experienced a “minor” reduction (Antony et al. 1985). However, ozonolysis of CHCs on live insects has not been previously reported. If ozonolysis can disrupt CHCs on live insects, this will allow us to investigate how changes to the CHC layer can affect desiccation resistance as well as chemical signaling in many insect species that do not have established molecular methods to disrupt these CHCs, opening the doors to further discoveries in basic and applied research. We use *Drosophila suzukii* as a model for our experiment because *D. suzukii* or spotted-wing drosophila is a fruit fly species that has invaded many temperate fruit growing areas, causing hundreds of millions of dollars of damage annually to fruit crops in the United States alone (Bolda et al. 2010). Understanding how ozone can affect *D. suzukii* may lead to novel methods to control this pest. The *D. suzukii* CHC profile is largely sexually monomorphic, but small differences in compound abundance have been observed between males and females (Dekker et al. 2015; Snellings et al. 2018). *D. suzukii* has a large variety of CHCs present, including monoenes (unsaturated hydrocarbons), dienes (unsaturated hydrocarbons) and n-alkanes (saturated hydrocarbons) with 7(Z)-tricosene being the most prevalent CHC in *D. suzukii* (Dekker et al. 2015; Snellings et al. 2018). Our study has three objectives. First, we will determine if ozonolysis affects CHCs on living *D. suzukii*. Secondly, we will evaluate the duration of these effects on *D. suzukii* and whether CHCs will be regenerated after the initial ozonolysis. Thirdly, we will determine whether modifications to CHCs via ozonolysis affects *D. suzukii* desiccation resistance. We hypothesize that ozonolysis of unsaturated hydrocarbons will occur after an ozone treatment, that CHC's will recover over time and that CHC ozonolysis will reduce desiccation resistance.

Materials And Methods

We performed two experiments to address our objectives. In experiment 1 we evaluated the effects of ozone on the CHCs of *D. suzukii* and the duration of these effects. Experiment 2 explored the effect of ozone exposure on the desiccation resistance of *D. suzukii*.

Colony Details & Maintenance

Drosophila suzukii were sourced from a colony reared out of tart cherries (*Prunus cerasus*) collected from the Trevor-Nichols Research Center located in Fennville, Michigan in 2015. Flies were maintained on 5 mL of solid diet (Dalton et al. 2011) in 50 mL polystyrene vials (Lab Express, Cat. # 8002-cs). The colony chamber was set on an 8-h dark period to a 16-h photoperiod, while maintaining an average relative humidity of 77% and temperature of 23°C.

Drosophila Handling

Newly emerged flies were collected, separated by sex, and placed into vials with 5 mL of solid diet (Dalton et al. 2011) where they were allowed to age (3-5 d or 9-11 d) prior to experiments. Thirty to sixty aged flies of a single sex were transferred using a small paintbrush and forceps into 304 stainless steel cages for ozone exposures. Steel cages were fabricated from half of a 5.33 cm diameter spherical tea infuser (Fu Store, 8541896633) that was folded back onto itself to create a "clamshell". Stainless steel was used due to its low reactivity with ozone. All tools and surfaces were cleaned with 70% ethanol between each experimental run/treatment to minimize the chance of contamination.

Flies in experiment 1 were anesthetized with carbon dioxide before being placed into stainless steel cages for treatment and, after treatment, into new vials with diet. Flies in experiment 2 were aspirated from vials containing diet into stainless steel cages for treatment and into new vials with diet after treatment application. The difference in handling procedures between experiments 1 & experiment 2 was due to the extremely low humidity of the carbon dioxide anesthetizing gas, which could directly affect the outcome of the desiccation trial (experiment 2). After ozone exposure, flies from both experiments were maintained in the colony chamber described above.

Ozone Treatment Application

An ozone generator (Absolute Ozone, NANO) applied 45,000 ppm of humidified ozone by using 99.5% oxygen as a feed gas at 14 psi and a flow rate of $5.1 \times 10^{-5} \text{ m}^3/\text{s}$. Ozone was humidified by passing it through 100 mL distilled water in a 250 mL glass gas washing bottle before the gas entered the treatment flask (Fig. 1.). Flies were exposed to ozone for 5 seconds in a specially modified, glass 250 mL Erlenmeyer flask with a dorsal 29/42 ASTM ground glass joint and a lateral 34/45 ASTM ground glass joint which acted as a port to allow insertion of stainless-steel cages (described above). Control flies were treated with humidified oxygen (99.5% purity) for 5 seconds and handled in the same fashion as the ozone treated flies, while untreated flies were not exposed to ozone nor oxygen. Gaseous ozone concentrations were measured during treatment applications using an ozone monitor (2B Technologies, Model 106-H). Relative humidity and temperature of oxygen treatments inside the treatment chamber were monitored using an electronic hygrometer sensor (Sensirion, EK-H4).

CHC Collection and Analysis

Five flies of the same treatment and sex were placed into a ½ dram glass vial (Kimble Glass Incorporated, Art. No. 60910L 12) along with 200 µL of a hexane wash. The hexane wash contained an internal standard of 25 ng/µL hexacosane (Sigma-Aldrich, #241687-5G). The hexane wash was added to the glass vials with 100 µL calibrated glass pipets (VWR International, Cat. No. 53432-921) and an aspirator (VWR International, Cat. No. 53432-921). Flies were left in hexane for 10-15 minutes. Samples were then placed on a mini-vortex (Fisher Scientific, Cat. No. 12-810-1) for 30 seconds at a vortex rate of 5 (~1,800 rpm). The hexane solution was transferred into a 0.25 mL glass insert (Supelco, Cat. No. 24717) inside a 2 mL glass vial (Supelco, Cat. No. 27330) for GCMS analysis. The 2 mL glass vials were capped with 9 mm Blue S/T Caps (Supelco, 29044-U). Samples were stored in a -20°C freezer (Fisher Scientific, Cat. No. 13986149) prior to and after GC-MS analysis.

Samples were eluted through a DB-17HT column (Agilent Technologies, Part No. 122-1831) that had a length of 30 m, a diameter of 0.25 mm, and a 0.15 µm thickness. Helium was used as a carrier gas at a rate of 1 mL/min through a GC-MS (Agilent Technologies, 5975C Series GC/MSD). Samples injected into the GC-MS were eluted after a four-minute solvent delay and a starting oven temperature of 50 C that increased by 4 C/min until a final temperature of 300 C was attained. The temperature remained constant for 10 minutes once 300 C was reached.

Integration and quantitation of peak areas were determined by using the QuanLynx program of the MassLynx MS Software version 4.2 to evaluate total ion chromatograms (Waters 2020). After determining peak areas, the total area of each peak per sample was divided by the total number of flies (5) from each sample to give a mean estimate of cuticular hydrocarbon mass per fly. The mass of each compound was calculated by referencing the peak area of hexacosane (5,000 ng), the internal standard (IS).

Experiment 1: Ozonolysis of Hydrocarbons

Comparison of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons 1 h after ozonolysis

The CHC profile of 3-5 and 9-11 d old male and female flies were compared (4 groups total). Treatments consisted of an untreated control, oxygen treatment (99.5% purity) and an ozone treatment (45,000 ppm) (Table 1). Treatments followed procedures outlined in section 2.3 Ozone Treatment Application. Total amount, in nanograms (ng), of unsaturated hydrocarbons (5(Z)-tricosene, 7(Z)-tricosene, 9(Z)-tricosene/tricosane, 5(Z)-pentacosene, 7(Z)-pentacosene, 9(Z)-pentacosene), aldehydes (heptanal, nonanal, tetradecanal, pentadecanal, hexadecanal, octadecanal) and saturated hydrocarbons (heneicosane, heptacosane, 2-methyl octacosane, nonacosane) peaks were quantified from CHC extracts from living flies 1 h after treatment application.

Collection and analysis of hydrocarbon extraction samples followed the procedures explained in section 2.4. The masses of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons were collected from five cuticular hydrocarbon extraction samples (5 flies/sample) for each treatment within a group. As the data were not normally distributed, a Kruskal-Wallis rank sum test was used for all three responses, with a separate analysis run for each sex and age for each of the three responses (12 models total). Kruskal-Wallis rank sum tests were performed by using the 'kruskal.test' function in R version 3.5.1 (R Core Team 2015). Wilcoxon rank sum tests were performed for post-hoc analyses by using the 'pairwise.wilcox.test' function in R version 3.5.1, which adjusted *p*-values by using the 'Holm' method (R Core Team 2015).

Experiment 1: Hydrocarbon Regeneration

Comparison of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons 1, 12, 36, 108 h after ozonolysis

We compared the CHC profiles of ozone treated and untreated 3-5 d and 9-11 d old male and female *D. sukuzii* at 1, 12, 36, 108 h after ozone exposure in 12 separate models (Table 1.). Ozone treatment, CHC extraction and quantification were performed as described above. A 2 x 4 factorial ANOVA model was used to analyze unsaturated and saturated CHC concentrations based on experimental treatment (ozone and untreated) and CHC extraction time after treatment (1, 12, 36, 108 h) as fixed factors. Aldehyde masses were fit to a linearized model, 'lm' function in R version 3.5.1 prior to ANOVA analysis and, subsequently, a post-hoc Tukey test (R Core Team 2015). ANOVAs and Tukey tests were performed by using the 'aov' and 'TukeyHSD' functions, respectively, in R version 3.5.1 (R Core Team 2015).

Experiment 2: Desiccation Resistance Assessment

Two desiccation resistance trials were performed on male and female *D. sukuzii* (3-5 d). In the first trial, flies were evaluated 1 h after ozonolysis and in the second, 108 h after ozonolysis. Methods were modified from Folk et al. (2001). Experimental arenas consisted of 50 mL polystyrene vials (Lab Express, Cat. # 8002-cs) containing 4.5 grams of desiccant (W. A. Hammond Drierite Co. Ltd., Stock No: 11001) at their base with a permeable plastic barrier placed above the desiccant. Nine to 11 single sex adult flies were placed into each arena and vials were capped with plastic wrap (Gordon Food Services, Item: 115193). Relative humidity and temperature were measured inside an experimental vial without flies during each trial using an electronic hygrometer sensor (Sensirion, EK-H4). Fly survival was assessed at 30-minute intervals for 10 h or until all flies had died. Fly mortality was determined by lightly shaking a vial and recording the number of individuals that reoriented to a standing position; those who did not re-orient were recorded as dead. As trials were completed when all flies had died, no censored data were included in Kaplan-Meier analyses.

Data for both desiccation trials were also analyzed by Mantel-Haenszel log-rank tests and Cox proportional hazard models with separate analyses performed for male and female flies using the R 'survival' package (Therneau et al. 2020). Trial one compared the survival of male or female *D. sukuzii* 1 hour after ozonolysis with flies treated with oxygen or an untreated control. Optimized Cox Proportional Hazard comparisons were made by first evaluating differences between the oxygen and untreated controls; if they were found to be similar, the combined oxygen and untreated controls were compared to the ozone treated flies (Crawley 2007; RICH et al. 2010). Trial two compared the survival of male or female *D. sukuzii* 108 h after ozonolysis with an untreated control of the same age, so a pairwise comparison of Cox Proportional Hazards was made. Hazard ratios (HR) were determined from Cox Proportional Hazard models. The hazard ratio, or instantaneous rate of death, represents the rate of death in comparison to control survival at any point of time.

Results

Ozonolysis affects the composition of CHCs on live D. sukuzii

Chromatograms of *D. sukuzii* CHCs for untreated controls and ozonated flies show differences 1 h following ozonolysis but not at 108 h after ozonolysis (Fig. 2 & 3). In the former, the total amount of 7-tricosene and other unsaturated hydrocarbons were reduced by approximately half and new aldehyde peaks (heptanal, nonanal, tetradecanal, pentadecanal, hexadecanal, octadecanal) were detected in the ozone treatment (Fig. 2). In the latter, peak heights for 7-tricosene and unsaturated CHC were similar to those observed for the untreated flies, and aldehydes were not observed for either group (Fig. 2). The elution peak positions of the aforementioned compounds and mass chromatograms of the aldehydes are presented in the supplementary data file (Fig. S1).

Experiment 1: Ozonolysis decreases total amount of unsaturated hydrocarbons, producing aldehydes, but do not affect the total amounts of saturated hydrocarbons.

The masses of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons at 1 h after treatment for 3-5 d and 9-11 d flies are presented in Table 2. The amounts of unsaturated hydrocarbons of 3-5 d old females were significantly reduced by ozonolysis with 2.03- and 1.89-fold reductions compared to that for the oxygen treated and untreated flies, respectively ($\chi^2 = 9.62$, $df=2$, $p=0.0081$), whereas the masses

were similar between the oxygen treated and untreated flies ($df=1, p=0.548$). A similar pattern was observed for 9-11 d old females with 2.79- and 2.63-fold reductions in ozone treated flies compared to oxygen treated and untreated flies, respectively ($\chi^2=10.22, df=2, p=0.006$), and no difference was detected between oxygen and untreated flies ($df=1, p=0.222$). Likewise, 3-5 d old ozone treated males demonstrated 2.85- and 3.10-fold reductions in unsaturated hydrocarbons compared to oxygen treated and untreated 3-5 d old males, respectively ($\chi^2=11.18, df=2, p=0.0037$), and similar unsaturated hydrocarbon amount between oxygen and untreated flies ($df=1, p=0.056$). Nine-11 d old males presented a slightly different pattern, with 1.63- and 2.28-fold reductions in unsaturated hydrocarbons in ozone treated flies compared to oxygen treated and untreated flies ($\chi^2=11.52, df=2, p=0.0031$), and a 1.4 fold reduction in levels observed from the oxygen to the untreated control flies ($df=1, p=0.032$).

The masses of aldehydes extracted from 3-5 d old females, 9-11 d old females, 3-5 d old males and 9-11 d old males were significantly increased by ozonolysis compared to oxygen treated and untreated flies ($\chi^2=13.29, df=2, p=0.0013, \chi^2=13.29, df=2, p=0.0013, \chi^2=13.29, df=2, p=0.0013, \chi^2=13.29, df=2, p=0.0013$, respectively). No aldehydes were extracted from oxygen or untreated flies in 3-5 d old females, 9-11 d old females, 3-5 d old males and 9-11 d old males.

The masses of saturated hydrocarbons extracted from 3-5 d old females were significantly increased after ozone treatment compared to oxygen treated and untreated flies ($\chi^2=9.42, df=2, p=0.0090$), while the masses were similar between oxygen treated and untreated flies ($df=1, p=0.841$). There was no difference in the masses of saturated hydrocarbons extracted from 9-11 d old females between ozone treated, oxygen treated and untreated flies ($\chi^2=1.46, df=2, p=0.4819$). The masses of saturated hydrocarbons extracted from 3-5 d old males treated with ozone and oxygen were significantly reduced in comparison to that found in untreated flies ($\chi^2=8.96, df=2, p=0.0113$), while saturated hydrocarbon amounts were similar between ozone treated and oxygen treated flies ($df=1, p=0.548$). There was no significant difference in the mass of the saturated hydrocarbons extracted from 9-11 d old male untreated flies with those from flies treated with either ozone or oxygen ($\chi^2=4.58, df=2, p=0.1013$).

Experiment 1: Hydrocarbons regenerate within 108 hours after ozonolysis

The masses of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons at 1, 12, 36, 108 h after treatment for 3-5 d and 9-11 d flies are presented in Table 3 and Figs. 5-7. Statistical output for ANOVA models is presented in Table 4.

Ozone exposure significantly reduced the mass of unsaturated hydrocarbons for male and female flies of both ages at time zero with masses returning to levels comparable to untreated flies within 108 h (Fig 4). For female flies of both ages, regeneration was comparable to untreated flies within 36 h with 3-5 day old ozone treated flies producing significantly more unsaturated hydrocarbons than the untreated flies, with mean \pm SEM values of 2145 ± 79 and 1489 ± 112 ng, respectively (Figure 5a). In contrast, ozone treated male flies of either age did not produce comparable masses of unsaturated hydrocarbons to their respective controls until 108 h after treatment (Table 4a).

Ozone exposure resulted in the formation of aldehydes from both treated sexes at both ages, but virtually no aldehydes were recovered from the untreated flies of either sex or age at any time point (Fig 5). Extractions of ozone treated flies contained aldehydes immediately after treatment with similar patterns of aldehyde reduction over subsequent time periods observed for both sexes and age groups (Table 4b). The aldehyde content approached 0 ng for the 3-5 day and 9-11 day old males and for the 3-5 day old females at 108 h after exposure. The data for 9-11 day old females at 108 h after treatment are not presented due to contamination issues. However, the mass of aldehydes for this group was reduced from 715 ± 28 ng at 1 h to 90 ± 6 ng at 36 h after ozone treatment.

Saturated hydrocarbon measurements provided the least consistent trends across sex and age groups compared to unsaturated hydrocarbons and aldehydes (Table 4c). Significantly higher masses of unsaturated hydrocarbons were detected for ozone treated 3-5 day old females at 1, 12 and 36 h after treatment (Fig 6a). Likewise, higher masses of unsaturated hydrocarbons were detected for ozone treated 9-11 day old females at 1 and 12 h with a significant difference detected at 36 h (Fig. 6b). In contrast, no significant differences between ozone treated and untreated males of either age group were detected at any time point (Fig 6 c,d).

Experiment 2: Ozonolysis decreases desiccation resistance in D. suzukii

Kaplan-Meier survival curves for desiccation trials conducted 1h and 108 h after ozonolysis are presented in Table 5 and Fig. 7. The mean (\pm SEM) relative humidity and temperature of desiccation trials at 1 h after treatment application were 18.13% ($\pm 1.54\%$) and 28.89°C (± 0.20), respectively. The mean (\pm SEM) relative humidity and temperature of desiccation trials at 108 h after treatment application were 15.93% (± 2.12) and 28.32°C ($\pm 0.34^\circ\text{C}$), respectively. Optimized Cox Proportional Hazard models for male and female flies 1h after ozonolysis compared oxygen and untreated controls to ozone treated flies (Crawley 2007; Rich et al. 2010). Output from Cox Proportional Hazard models for the two trials is presented in Table 6.

Mantel-Haenszel log-rank tests of female and male survival 1 h after ozonolysis (trial 1) showed significant differences between ozone treated and control flies ($c^2=158$, $df=2$, $p<0.001$, $c^2=75$, $df=2$, $p<0.001$, respectively). Cox Proportional Hazard models of female and male survival 1 h after ozonolysis showed significantly reduced survival times of ozone treated flies to the combined control flies 1 h after ozonolysis ($z=12.15$, $df=1$, $p<0.001$; $z=8.411$, $df=1$, $p<0.001$, respectively). A Hazard Ratio of 4.267 was observed when comparing controls to ozonated female flies with median times of death of 4.5 h and 2.5 h, respectively (Table 6). Similarly, ozonated males had a Hazard Ratio of 2.467 when compared to control flies with median times of death of 2 h and 3 h, respectively (Table 6).

Mantel-Haenszel log-rank tests of female and male survival 108 h after ozonolysis showed significant differences between ozone treated and control flies ($c^2=4.2$, $df=1$, $p=0.04$, $c^2=22.4$, $df=1$, $p<0.001$, respectively). Cox Proportional Hazard models of female and male survival 108 h after ozonolysis showed that ozone treated flies had significantly reduced survival times compared to control flies 1 h after ozonolysis ($z=-1.972$, $df=1$, $p=0.0486$, $z=-4.942$, $df=1$, $p<0.001$, respectively). Ozonated females had a Hazard Ratio of 0.753 when compared to control flies with median times of death of 4.5 h and 4 h, respectively (Table 6). Similarly, ozonated males had a Hazard Ratio of 0.470 when compared to control flies with median times of death of 2.5 h and 2 h, respectively (Table 6).

Discussion

Our results demonstrate that ozonolysis of live *D. sukuii* significantly reduces the total amount of unsaturated hydrocarbons, increases of the mass of aldehydes but does not affect the levels of saturated hydrocarbons in hexane extracted CHCs (Fig. 4, 5 & 6) (Table 3). Furthermore, the levels of CHCs from flies treated with ozone return to untreated CHC levels within 108 h after exposure (Fig. 4, 5 & 6 and Table 3). However, flies regenerated unsaturated hydrocarbons at different rates depending on their sex and age (Fig. 4 and Table 3). Desiccation resistance was correlated to changes in CHC abundance, with an immediate decrease followed by recovery over the same time. (Fig. 7 and Table 5). Surprisingly, desiccation resistance significantly increased in comparison to untreated controls at 108 h after treatment application, although a concomitant significant increase in unsaturated hydrocarbons was only observed for females at 3-5 d at 36 h after treatment application (Fig. 4 & 7 and Table 3 & 4).

Our study is the first to quantify ozonolysis of CHCs on living insect specimens and may provide an important new method for exploring the biosynthesis, structure, and function of these important constituents of the insect epicuticle. Current methodology for the modification of living insect CHCs include genetic modification and direct CHC application via perfuming (Ferveur 1997) (Billeter et al. 2009). While these methodologies are useful for determining the function of CHCs, they are expensive and/or time intensive, requiring the genetic modification of individual species/lineages. Our method could be used to modify the unsaturated hydrocarbons of any insect and allow the measurement of CHC generation time, potential for regeneration as well as how they modify behavior and survival.

While CHC generation in *Drosophila* species has been described in previous work (Bartelt et al. 1986; Jallon and David 1987; Toolson and Kuper-Simbron 1989; Dekker et al. 2015; Snellings et al. 2018), ours is a novel study providing data on the regeneration of unsaturated hydrocarbons following their removal from living subjects. The regeneration of CHCs to untreated levels suggests that maintaining CHCs is of great importance to *D. sukuii*. Insect CHCs have been found to function as (1) pheromones, (2) to increase desiccation resistance and (3) to protect from entomo-pathogens (Quinlan and Hadley 1993; Gibbs 1998; Howard and Blomquist 2005; Blomquist and Bagnères 2010; Ortiz-Urquiza and Keyhani 2013; Chung and Carroll 2015).

Our results also provide evidence that *D. sukuii* of different sexes and ages regenerate CHCs to untreated levels, albeit point estimates of CHC's vary across these groups (Fig. 4 and Table 3). For example, 3-5 d old and 9-11 d old females regenerated unsaturated hydrocarbons to levels observed in untreated fly by 12 and 36 h after treatment application, respectively (Fig. 4 and Table 3). However, 3-5 d and 9-11 d males did not regenerate unsaturated hydrocarbons to untreated fly levels until 108 h after treatment application (Fig. 4 and Table 3). This suggests that females either have a greater capacity to generate CHCs or prioritize the regeneration of CHCs after ozonolysis.

Unexpectedly, the mass of saturated hydrocarbons increased over time after ozone exposure in female flies. We propose that the CHC compensation mechanism for the regeneration of unsaturated hydrocarbons also affects saturated hydrocarbon biosynthesis. One possible explanation may involve the over-expression of a single P450 decarboxylase, *Cyp4g1*, which is directly involved in the biosynthesis pathway of insect CHCs (Qiu et al. 2012). The over-expression of the P450 decarboxylase would allow the regeneration of unsaturated CHCs to normal levels, as well as the biosynthesis of saturated CHCs to elevated levels (Chung and Carroll 2015).

Insect CHCs have been found to decrease water permeability through the cuticle and, thus increasing desiccation resistance (Quinlan and Hadley 1993; Gibbs 1998; Blomquist and Bagnères 2010; Chung and Carroll 2015). We observed greatly reduced desiccation resistance immediately following ozone treatments with a return to pre-treatment desiccation resistance following the regeneration of unsaturated hydrocarbons (Fig. 4 & 7. and Table 5). This finding supports the hypothesis that CHCs function to reduce cuticle water permeability (Ramsay 1935; Gibbs 1998) and that *D. sukuii* unsaturated hydrocarbons play a significant role in their desiccation resistance. This is of interest

because it has been previously suggested that saturated hydrocarbons are generally more correlated with desiccation resistance due to their higher melting points (Gibbs and Pomonis 1995). Additional supporting evidence suggests that a greater abundance of long-chained saturated hydrocarbons impart greater desiccation resistance by reducing the water loss rate (WLR) in *D. pseudoobscura* and *Tibicen dealbatus* (Homoptera: Cicadidae) (Toolson 1984; Toolson and Kuper-Simbron 1989).

Our data provide strong correlative evidence for the importance of unsaturated CHC's for desiccation resistance in *D. sukukii*. Gibbs (2002) hypothesizes that alkenes and alkanes form layers on the epicuticle dependently on lipid melting points. Alkenes may form liquid layers on the cuticle and allow greater permeability of water due to their lower melting temperatures (Gibbs 2002). This layered packing of alkanes and alkenes could help to explain the decreased survival rate of ozone treated flies as well as the uniform reduction of all unsaturated hydrocarbons. Furthermore, SEM images of a tick cuticle, *Rhipicephalus sanguineus* (Latreille) (Ixodida: Ixodidae), after ozone exposure qualitatively shows the damaging impact of ozone to the epicuticle layer (Moreira et al. 2018). This provides additional evidence to support the correlation of decreased desiccation resistance resulting from ozone mediated damage to the epicuticle.

While our data strongly suggest that desiccation resistance is directly linked to unsaturated hydrocarbon quantity, it is possible that observed differences were due to other ozone induced effects. The ozonolysis performed in this experiment, while largely non-lethal, did result in some mortality. Dose response curves developed in Savage (2020) predict that a CT product of 3,750 ppm-min of gaseous ozone would result in 11% and 25% mortality immediately following ozonolysis for males and females, respectively (Savage 2020). One potential, non-desiccation, source of mortality could be tracheal damage resulting from ozone exposure. However, Sousa et al. (2008) examined the respiration rates of *T. castaneum*, *R. dominica* and *O. surinamensis* and concluded that ozone toxicity and respiration rates did not correlate (Sousa et al. 2008). Our study did not directly measure the respiration rate of specimens after ozone exposure.

Potential future applications of ozonolysis of CHCs include characterizing arthropod physiology and behavior in regards to desiccation, chemical communication and entomopathogen resistance (Quinlan and Hadley 1993; Gibbs 1998; Howard and Blomquist 2005; Blomquist and Bagnères 2010; Ortiz-Urquiza and Keyhani 2013; Chung and Carroll 2015). Unsaturated hydrocarbons have been shown to be important in *Drosophila spp.* for identification of conspecifics and courtship/mating behaviors (Antony et al. 1985; Jallon and David 1987; Ferveur 1997, 2005; Howard and Blomquist 2005). For example, 7,11-heptacosadiene has been shown to be an aphrodisiac for male *D. melanogaster* (Antony et al. 1985). Cleavage of the double bonds of 7,11-heptacosadiene at the 7th and 11th carbon positions would occur after an ozone treatment using our methodology. This method could be combined with mating assays to determine how courtship and copulation are affected after ozonolysis of unsaturated hydrocarbons.

Ozonolysis of CHCs could also be combined with genetic modification of CHCs or CHC perfuming (Ferveur 1997, 2005). For example, ozonolysis of the genetically modified oenocyte-less (oe-) fly lineage of *D. melanogaster*, that produces no CHCs (Billeter et al. 2009) could be used to further elucidate whether ozonolysis effects desiccation resistance in the absence of CHCs. Additionally, courtship and copulation is shown to be mediated by unsaturated hydrocarbons, such as the anti-aphrodisiac 7-tricosene, in both *D. sukukii* flies and *D. melanogaster* males (Ferveur 1997; Snellings et al. 2018). Post ozonolysis "perfuming" of insects could be used to evaluate the relative importance of specific semiochemicals in courtship and mate selection. Ozonolysis of unsaturated hydrocarbons on female *D. sukukii* were correlated to reduced courtship and/or copulation by untreated males (unpublished data). This is counter-intuitive to the reduction of the 7-tricosene (an anti-aphrodisiac) after ozonolysis but may be explained by the interaction of ozone with other insect tissues. For example, dominant lethal chemicals have been shown to be produced after ozone exposure, which cause mutagenicity and a reduced reproductive potential in *D. virilis* (Erdman and Hernandez 1982).

In conclusion, the masses of *D. sukukii* unsaturated hydrocarbons are significantly reduced by a factor of 2-3 after ozone treatment due to the process of ozonolysis. This creates aldehydes which remain on the cuticle for between 36 and 108 h. The saturated CHC amount on flies are largely unaffected by ozone treatment, except for female flies at 3-5 d where a significant increase in saturated hydrocarbons were found. Additionally, flies demonstrated differential CHC regeneration based on sex and age. Females regenerated CHCs more quickly than males, as well as having an increased CHC regeneration rate at 3-5 d than 9-11 d. Finally, the reduction and recovery of desiccation resistance in ozone treated flies was correlated to the reduction and regeneration of unsaturated hydrocarbons. However, the desiccation resistance of ozone treated flies was elevated above untreated flies after unsaturated CHC regeneration. These findings provide novel methodology for insect CHC reduction/modification, evidence for CHC regeneration after reduction/modification and evidence supporting the contribution of unsaturated hydrocarbons in desiccation resistance.

Declarations

Funding: Organic Pest Management lab of the Entomology Department at Michigan State University.

Conflicts of interest/Competing interests: NA

Availability of data and material: Available upon request.

Code availability: Available upon request.

Authors' contributions:

- Benjamin Savage designed experiments, executed experiments, collected data, performed statistical analysis, wrote code for statistical analysis, developed figures and wrote the article.
- Zinan Wang provided methodological support for cuticular extractions, GC/MS set-up, and developed figures.
- Susan Masten provided ozone concentration measurement equipment and scientific advice
- Henry Chung provided experimental design support.
- Matthew Grieshop provided scientific leadership, equipment, and funding.

References

1. Antony C, Davis TL, Carlson DA, et al (1985) Compared behavioral responses of male *Drosophila melanogaster* (Canton S) to natural and synthetic aphrodisiacs. *J Chem Ecol* 11:1617–1629. <https://doi.org/10.1007/BF01012116>
2. Bartelt RJ, Arnold MT, Schaner AM, Jackson LL (1986) Comparative analysis of cuticular hydrocarbons in the *Drosophila virilis* species group. *Comp Biochem Physiol Part B Comp Biochem* 83:731–742. [https://doi.org/10.1016/0305-0491\(86\)90138-0](https://doi.org/10.1016/0305-0491(86)90138-0)
3. Bartelt RJ, Jones RL, Kulman HM (1982) Hydrocarbon components of the yellowheaded spruce sawfly sex pheromone. *J Chem Ecol* 8:95–114. <https://doi.org/10.1007/BF00984008>
4. Beroza Morton, Bierl BA (1967) Rapid determination of olefin position in organic compounds in microgram range by ozonolysis and gas chromatography. Alkylidene analysis. *Anal Chem* 39:1131–1135. <https://doi.org/10.1021/ac60254a038>
5. Billeter J-C, Atallah J, Krupp JJ, et al (2009) Specialized cells tag sexual and species identity in *Drosophila melanogaster*. *Nature* 461:987–991. <https://doi.org/10.1038/nature08495>
6. Blomquist GJ, Bagnères A-G (2010) *Insect Hydrocarbons: Biology, Biochemistry, and Chemical Ecology*. Cambridge University Press
7. Bolda MP, Goodhue RE, Zalom FG (2010) Spotted Wing *Drosophila*: Potential Economic Impact of a Newly Established Pest. 4
8. Chung H, Carroll SB (2015) Wax, sex and the origin of species: Dual roles of insect cuticular hydrocarbons in adaptation and mating. *BioEssays* 37:822–830. <https://doi.org/10.1002/bies.201500014>
9. Crawley MJ (2007) *The R Book*. John Wiley & Sons
10. Dalton DT, Walton VM, Shearer PW, et al (2011) Laboratory survival of *Drosophila suzukii* under simulated winter conditions of the Pacific Northwest and seasonal field trapping in five primary regions of small and stone fruit production in the United States. *Pest Manag Sci* 67:1368–1374. <https://doi.org/10.1002/ps.2280>
11. Dekker T, Revadi S, Mansourian S, et al (2015) Loss of *Drosophila* pheromone reverses its role in sexual communication in *Drosophila suzukii*. *Proc R Soc B Biol Sci* 282:. <https://doi.org/10.1098/rspb.2014.3018>
12. Erdman HE, Hernandez T (1982) Adult toxicity and dominant lethals induced by ozone at specific stages in spermatogenesis in *drosophila virilis*. *Environ Mutagen* 4:657–666. <https://doi.org/10.1002/em.2860040605>
13. Ferveur J-F (1997) The pheromonal role of cuticular hydrocarbons in *Drosophila melanogaster*. *BioEssays* 19:353–358. <https://doi.org/10.1002/bies.950190413>
14. Ferveur J-F (2005) Cuticular Hydrocarbons: Their Evolution and Roles in *Drosophila* Pheromonal Communication. *Behav Genet* 35:279. <https://doi.org/10.1007/s10519-005-3220-5>
15. Gibbs A, Pomonis JG (1995) Physical properties of insect cuticular hydrocarbons: The effects of chain length, methyl-branching and unsaturation. *Comp Biochem Physiol B Biochem Mol Biol* 112:243–249. [https://doi.org/10.1016/0305-0491\(95\)00081-X](https://doi.org/10.1016/0305-0491(95)00081-X)
16. Gibbs AG (1998) Water-Proofing Properties of Cuticular Lipids. *Integr Comp Biol* 38:471–482. <https://doi.org/10.1093/icb/38.3.471>
17. Gibbs AG (2002) Lipid melting and cuticular permeability: new insights into an old problem. *J Insect Physiol* 48:391–400. [https://doi.org/10.1016/S0022-1910\(02\)00059-8](https://doi.org/10.1016/S0022-1910(02)00059-8)
18. Hadley NF (1994) Ventilatory Patterns and Respiratory Transpiration in Adult Terrestrial Insects. *Physiol Zool* 67:175–189. <https://doi.org/10.1086/physzool.67.1.30163841>

19. Howard RW, Blomquist GJ (2005) Ecological, Behavioral, and Biochemical Aspects of Insect Hydrocarbons. *Annu Rev Entomol* 50:371–393. <https://doi.org/10.1146/annurev.ento.50.071803.130359>
20. Howard RW, Jackson LL, Banse H, Blows MW (2003) Cuticular Hydrocarbons of *Drosophila birchii* and *D. serrata*: Identification and Role in Mate Choice in *D. serrata*. *J Chem Ecol* 29:961–976. <https://doi.org/10.1023/A:1022992002239>
21. Jallon J-M, David JR (1987) Variation in Cuticular Hydrocarbons Among the Eight Species of the *Drosophila melanogaster* Subgroup. *Evolution* 41:294–302. <https://doi.org/10.2307/2409139>
22. Moreira LH, Figueiredo TFB, Alves LP, et al (2018) Effect of Ozone as Acaricide: Action of the Ozone on the Cuticle and Respiratory Spiracle of Tick *Rhipicephalus sanguineus sensu lato*. *Ozone Sci Eng* 40:183–190. <https://doi.org/10.1080/01919512.2017.1403306>
23. Ortiz-Urquiza A, Keyhani NO (2013) Action on the Surface: Entomopathogenic Fungi versus the Insect Cuticle. *Insects* 4:357–374. <https://doi.org/10.3390/insects4030357>
24. Qiu Y, Tittiger C, Wicker-Thomas C, et al (2012) An insect-specific P450 oxidative decarboxylase for cuticular hydrocarbon biosynthesis. *Proc Natl Acad Sci* 109:14858–14863. <https://doi.org/10.1073/pnas.1208650109>
25. Quinlan MC, Hadley NF (1993) Gas Exchange, Ventilatory Patterns, and Water Loss in Two Lubber Grasshoppers: Quantifying Cuticular and Respiratory Transpiration. *Physiol Zool* 66:628–642. <https://doi.org/10.1086/physzool.66.4.30163812>
26. R Core Team (2015) R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria
27. Ramsay JA (1935) The Evaporation of Water from the Cockroach. *J Exp Biol* 12:373–383
28. Rich JT, Neely JG, Paniello RC, et al (2010) A Practical Guide To Understanding Kaplan-Meier Curves. *Otolaryngol–Head Neck Surg Off J Am Acad Otolaryngol-Head Neck Surg* 143:331–336. <https://doi.org/10.1016/j.otohns.2010.05.007>
29. Savage BA (2020) Exploring The Insecticidal Potential Of Aqueous And Gaseous Ozone Using Spotted Wing *Drosophila*, *Drosophila suzukii* (Matsumura) (Diptera-drosophilidae) As A Model Organism. Michigan State University
30. Snellings Y, Herrera B, Wildemann B, et al (2018) The role of cuticular hydrocarbons in mate recognition in *Drosophila suzukii*. *Sci Rep* 8:4996. <https://doi.org/10.1038/s41598-018-23189-6>
31. Sousa AH, Faroni LRD, Guedes RNC, et al (2008) Ozone as a management alternative against phosphine-resistant insect pests of stored products. *J Stored Prod Res* 44:379–385. <https://doi.org/10.1016/j.jspr.2008.06.003>
32. Toolson EC (1984) Interindividual Variation in Epicuticular Hydrocarbon Composition and Water Loss Rates of the Cicada *Tibicen dealbatus* (Homoptera: Cicadidae). *Physiol Zool* 57:550–556. <https://doi.org/10.1086/physzool.57.5.30163947>
33. Toolson EC, Kuper-Simbron R (1989) Laboratory Evolution of Epicuticular Hydrocarbon Composition and Cuticular Permeability in *Drosophila pseudoobscura*: Effects on Sexual Dimorphism and Thermal-Acclimation Ability. *Evolution* 43:468–473. <https://doi.org/10.2307/2409222>
34. Waters (2020) MassLynx MS Software. In: Waters. https://www.waters.com/waters/en_US/MassLynx-MS-Software/nav.htm?cid=513662&locale=en_US. Accessed 1 Apr 2020

Tables

Table 1 Experiment 1, 2 & 3 data collection times, treatments, fly age and replication. An “*” in the ‘Replication (Male, Female)’ column indicates the same number of replications at every data collection time in the ‘Data Collection Hour after Treatment Application’ column and a ‘;’ indicates a separation of replication numbers consistent with the data collection times in the ‘Data Collection Time after Treatment Application (h(s))’ column.

EXPERIMENT SET-UP					
Experiment	CHC Extraction Hour after Treatment Application	Treatment	Fly Age (days) at Treatment application	Replication	(Male, Female)
Experiment 1	1	Oxygen	3-5	25, 25	
	1, 12, 36, 108	Untreated	3-5	*25, 25	
	1, 12, 36, 108	Ozone	3-5	*25, 25	
	1	Oxygen	9-11	25, 25	
	1, 12, 36	Untreated	9-11	*25, 25	
	1, 12, 36	Ozone	9-11	*25, 25	
Experiment 2	1	Oxygen	3-5	140, 141	
	1, 108	Untreated	3-5	138, 140 ; 100, 99	
	1, 108	Ozone	3-5	141, 140 ; 101, 99	

Table 2 Mean (ng) and standard error of mean (SEM) of unsaturated hydrocarbons, aldehydes and saturated hydrocarbons extracted from female and male flies at 1 h after treatment application (3-5 & 9-11 days old). Disparate letters signify differences within a population-based Wilcoxon rank sum test with Holm p-adjustment.

CUTICULAR COMPOUND MASS (NG) OF <i>D. SUZUKII</i> AT 1 H AFTER TREATMENT						
Unsaturated hydrocarbons						
Females						
	3-5 d			9-11 d		
Treatment	Mean	SEM	CLD	Mean	SEM	CLD
Ozone	721	52	b	1017	85	b
Oxygen	1462	101	a	2832	78	a
Untreated	1366	45	a	2664	108	a
Males						
	3-5 d			9-11 d		
	Mean	SEM	CLD	Mean	SEM	CLD
Ozone	586	5	b	894	110	c
Oxygen	1674	32	a	1453	132	b
Untreated	1817	76	a	2036	64	a
Aldehydes						
Females						
	3-5 d			9-11 d		
Treatment	Mean	SEM	CLD	Mean	SEM	CLD
Ozone	574	18	a	715	28	a
Oxygen	0	0	b	0	0	b
Untreated	0	0	b	0	0	b
Males						
	3-5 d			9-11 d		
	Mean	SEM	CLD	Mean	SEM	CLD
Ozone	382	11	a	443	32	a
Oxygen	0	0	b	0	0	b
Untreated	0	0	b	0	0	b
Saturated hydrocarbons						
Females						
	3-5 d			9-11 d		
Treatment	Mean	SEM	CLD	Mean	SEM	CLD
Ozone	495	12	a	716	45	a
Oxygen	333	24	b	668	35	a
Untreated	319	9	b	648	32	a
Males						
	3-5 d			9-11 d		
	Mean	SEM	CLD	Mean	SEM	CLD

Ozone	344	7	a	503	37	a
Oxygen	350	10	a	378	38	a
Untreated	397	18	b	485	20	a

Table 3 Unsaturated hydrocarbons, aldehydes and saturated hydrocarbons mean amount extracted from flies (3-5 & 9-11 days old). Cuticular hydrocarbon extractions occurred at 1 h, 12 h, 36 h and 108 h after treatment application. CLD marks differences between "Treatment - Hours" means based on an ANOVA p-values. Disparate letters only signify differences within a sex and age.

CUTICULAR COMPOUND AMOUNT (NG) OF <i>D. SUZUKII</i> AT 1, 12, 36 AND 108 H AFTER TREATMENT													
		3-5 d						9-11 d					
		Female			Male			Female			Male		
	Treatment - Hour	Mean	SEM	CLD	Mean	SEM	CLD	Mean	SEM	CLD	Mean	SEM	CLD
Unsaturated	Ozone - 1	721	52	e	586	5	d	1017	85	c	894	110	d
	Untreated - 1	1366	45	d	1817	76	bc	2664	108	a	2036	64	be
	Ozone - 12	1460	100	cd	1635	55	c	1837	82	b	1314	104	cd
	Untreated - 12	1929	215	bc	2659	45	a	2635	140	a	2072	124	be
	Ozone - 36	2145	79	ab	2130	97	b	2660	170	a	1631	88	ce
	Untreated - 36	1489	112	cd	2871	72	a	2659	88	a	2092	28	b
	Ozone - 108	2417	74	a	2786	128	a				2928	90	a
	Untreated - 108	2052	59	ab	2700	85	a				2832	126	a
Aldehyde	Ozone - 1	574	18	a	382	11	a	715	28	a	443	32	a
	Untreated - 1	0	0	c	0	0	d	0	0	d	0	0	c
	Ozone - 12	273	31	b	214	9	b	345	12	b	190	20	b
	Untreated - 12	0	0	c	0	0	d	0	0	d	0	0	c
	Ozone - 36	57	6	c	59	9	c	90	6	c	52	3	c
	Untreated - 36	0	0	c	0	0	d	0	0	d	0	0	c
	Ozone - 108	3	1	c	4	0	d				7	2	c
	Untreated - 108	0	0	c	0	0	d				0	0	c
Saturated	Ozone - 1	495	12	bc	344	7	c	716	45	ab	503	37	b
	Untreated - 1	319	9	e	397	18	c	648	32	b	485	20	b
	Ozone - 12	600	30	ab	607	9	b	794	21	ab	519	21	b
	Untreated - 12	424	43	cde	613	15	b	677	36	b	481	42	b
	Ozone - 36	654	27	a	632	25	ab	860	53	a	489	21	b
	Untreated - 36	370	34	de	708	31	a	659	37	b	460	17	b
	Ozone - 108	582	23	ab	714	22	a				786	25	a

Untreated - 108	483	20	bcd	694	23	ab	711	30	a
--------------------	-----	----	-----	-----	----	----	-----	----	---

Table 4 Results of 2 x 4 factorial ANOVA models comparing treatment to CHC extraction time after treatment (1, 12, 36 and 108 h) on unsaturated CHCs, adlehydes and saturated CHCs. A '*' indicates a significant main effect (* 0.05 > ** 0.01 > *** 0.001).

ANOVA MODELS COMPARING CHC MASS (NG) OF D. SUZUKII OVERTIME						
Unsaturated hydrocarbons						
Females						
a)	3-5 d			9-11 d		
Factors	df	F	p	df	F	p
Treatment	1	0.094	0.761	1	73.1	*** <0.001
Hour	3	43.8	*** <0.001	2	24.6	*** <0.001
Treatment:Hour	3	18	*** <0.001	2	24.9	*** <0.001
Males						
	3-5 d			9-11 d		
	df	F	p	df	F	p
Treatment	1	173	*** <0.001	1	68.5	*** <0.001
Hour	3	150	*** <0.001	3	83.4	*** <0.001
Treatment:Hour	3	27.4	*** <0.001	3	14.5	*** <0.001
Aldehydes						
Females						
b)	3-5 d			9-11 d		
Factors	df	F	p	df	F	p
Treatment	1	618	*** <0.001	1	1320	*** <0.001
Hour	3	201	*** <0.001	2	297	*** <0.001
Treatment:Hour	3	201	*** <0.001	2	297	*** <0.001
Males						
	3-5 d			9-11 d		
c)	df	F	p	df	F	p
Treatment	1	1492.	*** <0.001	1	340.8	*** <0.001
Hour	3	396.9	*** <0.001	3	109.5	*** <0.001
Treatment:Hour	3	396.9	*** <0.001	3	109.5	*** <0.001
Saturated hydrocarbons						
Females						
	3-5 d			9-11 d		
Factors	df	F	p	df	F	p
Treatment	1	93.4	*** <0.001	1	16.6	*** <0.001
Hour	3	8.87	*** <0.001	2	2.12	0.143
Treatment:Hour	3	4	* 0.016	2	1.50	0.243
Males						
	3-5 d			9-11 d		
	df	F	p	df	F	p

Treatment	1	3.98	0.055 .	1	4.04	0.053 .
Hour	3	111	*** <0.001	3	43.3	*** <0.001
Treatment:Hour	3	2.30	0.096 .	3	0.383	0.766

Table 5 Kaplan-Meier survival analyses of desiccation resistance from flies (3-5 days old). Flies (10) were placed into a plastic vial with a desiccant, drierite, and sealed. Flies were considered living until they failed to re-orient to a standing position after a shake of the vial. Flies that did not re-orient were marked as deceased. No censored data were included in the analyses.

KAPLAN-MEIER SURVIVAL ANALYSIS					
Sex	Time (h)	Treatment	Sample Size	\bar{x} (Median h)	95% CL (Lower-Upper)
Female	1	Untreated	141	5	4.5-5.5
		Oxygen	140	4	4-5
		Ozone	140	2.5	2.5-5
		Untreated/Oxygen	281	4.5	4.5-5
	108	Untreated	99	4	3.5-4
		Ozone	99	4.5	4-5
Male	1	Untreated	140	3	3-3.5
		Oxygen	138	3	2.5-3.5
		Ozone	141	2	2-2
		Untreated/Oxygen	278	3	3-3.5
	108	Untreated	100	2	2-2.5
		Ozone	101	2.5	2.5-3

Table 6 Cox proportional hazard analysis models performed on fly survival within a sex and h after treatment application. All model parameters compared to untreated fly survival within the same sex and h. The ' β ' is the parameter estimate. The 'SE' is the standard error of the ' β '. The 'HR' is the hazard ratio. The 'CL' is the confidence level.

SURVIVAL MODELS: COX PROPORTIONAL HAZARD REGRESSION ANALYSIS									
		Parameters	β	SE	df	z	P	HR	95% CL (Lower-Upper)
Sex	H(s)	Baseline:Comparison							
Female	1	Untreated+Oxygen:Ozone	1.451	0.119	1	12.150	<0.001	4.267	3.377-5.392
	108	Untreated:Ozone	-0.284	0.144	1	-1.972	0.049	0.753	0.567-0.998
Male	1	Untreated+Oxygen:Ozone	0.903	0.107	1	8.411	<0.001	2.467	1.999-3.045
	108	Untreated:Ozone	-0.756	0.153	1	-4.942	<0.001	0.470	0.348-0.634