

# Type of discrepancy between glycosylated hemoglobin and fasting plasma glucose is associated with in-hospital outcomes in patients with acute coronary syndrome and diabetes: findings from the Improving Care for Cardiovascular Disease in China - Acute Coronary Syndrome (CCC-ACS) Project

**Nan Ye**

Beijing Anzhen Hospital, Capital Medical University <https://orcid.org/0000-0001-9339-9361>

**Lijiao Yang**

Beijing Anzhen Hospital, Capital Medical University

**Guoqin Wang**

Beijing Anzhen Hospital, Capital Medical University

**Weijing Bian**

Beijing Anzhen Hospital, Capital Medical University

**Fengbo Xu**

Beijing Anzhen Hospital, Capital Medical University

**Changsheng Ma**

Beijing Anzhen Hospital, Capital Medical University

**Dong Zhao**

Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases

**Jing Liu**

Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases

**Yongchen Hao**

Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases

**Jun Liu**

Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases

**Na Yang**

Beijing Anzhen Hospital, Capital Medical University, Beijing Institute of Heart, Lung and Blood Vessel Diseases

Hong Cheng (✉ [drchengh@163.com](mailto:drchengh@163.com))

<https://orcid.org/0000-0002-7908-3333>

---

## Research article

**Keywords:** acute coronary syndrome, diabetes, fasting plasma glucose, glycosylated hemoglobin

**DOI:** <https://doi.org/10.21203/rs.3.rs-29995/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

The discrepancy between glycosylated hemoglobin (HbA<sub>1c</sub>) and fasting plasma glucose (FPG) in clinical practice may be related to factors such as acute stress, renal dysfunction, and anemia, and its relationship with in-hospital outcomes is uncertain. The aim of this study is to investigate the association between the type of discrepancy between HbA<sub>1c</sub> and FPG and in-hospital outcomes in patients with acute coronary syndrome (ACS) and diabetes.

## Methods

The Improving Care for Cardiovascular Disease in China - Acute Coronary Syndrome (CCC-ACS) project is a national, hospital-based quality improvement project with an ongoing database. Patients with ACS, diabetes and complete HbA<sub>1c</sub> and FBG values at admission were included. The consistent group included patients with HbA<sub>1c</sub> < 6.5% and FBG < 7.0 mmol/L or HbA<sub>1c</sub> ≥ 6.5% and FBG ≥ 7.0 mmol/L. The discrepancy group included patients with HbA<sub>1c</sub> ≥ 6.5% and FBG < 7.0 mmol/L (increased HbA<sub>1c</sub> group) or HbA<sub>1c</sub> < 6.5% and FBG ≥ 7.0 mmol/L (increased FBG group).

## Results

A total of 7,762 patients were included in this study. The numbers of patients in the consistent group and discrepancy group were 5,490 and 2,272 respectively. In the discrepancy group, increased HbA<sub>1c</sub> accounted for 77.5% of discrepancies, and increased FPG accounted for 22.5% of discrepancies. After adjusting for confounders, patients in the increased FPG group had a 1.6-fold increased risk of heart failure (OR, 1.62; 95% CI, 1.08–2.44), a 1.6-fold increased risk of composite cardiovascular death and heart failure (OR, 1.63; 95% CI, 1.09–2.43), and a 1.6-fold increased risk of composite major adverse cardiovascular and cerebrovascular events (MACCEs) and heart failure (OR, 1.56; 95% CI, 1.08–2.24).

## Conclusions

Patients with an increased FPG but normal HbA<sub>1c</sub> had a higher risk of in-hospital adverse outcomes than those with increased HbA<sub>1c</sub> but normal FPG. This result may indicate that when HbA<sub>1c</sub> and FPG are inconsistent in patients with ACS and diabetes, the increased FPG that may be caused by stress hyperglycemia may have a more substantial adverse effect than increased HbA<sub>1c</sub>, which may be caused by chronic hyperglycemia. These high-risk patients should be given more attention and closer monitoring in clinical practice.

# Background

Cardiovascular disease is the leading cause of both death and premature death in China and is the cause of 40% of deaths in the Chinese population [1]. Acute coronary syndrome (ACS) is an acute manifestation of cardiovascular disease with a high risk of mortality that can lead to critical conditions such as cardiogenic shock and cardiac arrest. Patients with ACS and diabetes usually have worse clinical outcomes than those with normal blood glucose [2–8], regardless of in-hospital or long-term outcomes. Indicators commonly used for evaluating blood glucose include intravenous blood glucose, glycosylated hemoglobin (HbA<sub>1c</sub>), and glycosylated serum albumin. Glucose was first used in the diagnosis of diabetes, including fasting plasma glucose (FPG), oral glucose tolerance tests and random blood glucose. In 2013, the American Diabetes Association approved the use of HbA<sub>1c</sub> to diagnose diabetes [9]. In addition to their diagnostic value, FPG and HbA<sub>1c</sub> are also associated with clinical outcomes. Several studies have shown that abnormal blood glucose is an important factor associated with clinical outcomes in patients with ACS and diabetes [10–17].

However, we noticed that some patients have a discrepancy between HbA<sub>1c</sub> and FPG in clinical practice. This condition may be related to factors such as acute stress, renal dysfunction, and anemia that may affect FPG and HbA<sub>1c</sub>. The discrepancy can be an increased FPG with a normal HbA<sub>1c</sub> or an increased HbA<sub>1c</sub> with a normal FPG. We decided to explore which discrepancy indicates worse in-hospital outcomes. There are few studies focusing on this issue.

The Improving Care for Cardiovascular Disease in China - Acute Coronary Syndrome (CCC-ACS) project is a national, hospital-based quality improvement project with an ongoing database, aiming to increase adherence to ACS guidelines in China and to improve patient outcomes. We conducted this study based on the CCC-ACS project to investigate the types of discrepancies between HbA<sub>1c</sub> and FPG and their relationships to in-hospital outcomes.

## Methods

### Research Design

Details of the design and methodology of the CCC-ACS project have been published [18], and the study was registered at ClinicalTrials.gov (NCT02306616). In brief, the CCC-ACS is a national, hospital-based quality improvement project with an ongoing database, aiming to increase adherence to ACS guidelines in China and to improve patient outcomes. It was launched in 2014 as a collaborative initiative of the American Heart Association and Chinese Society of Cardiology. A total of 240 hospitals were recruited representing the diversity of ACS care in hospitals in China, including 150 tertiary hospitals in phase I and phase II and 82 secondary hospitals and 8 tertiary hospitals in phase III (from July 2017) and phase IV (from November 2018). Clinical data were collected via a web-based data collection platform (Oracle Clinical Remote Data Capture, Oracle Corporation). Trained data abstractors entered the data elements abstracted from medical charts. Eligible patients were consecutively reported to the CCC-ACS database

for each month before the middle of the following month. Approximately 5% of reported cases were randomly selected and compared with the original medical records. An audit by a third party was performed to ensure that cases were reported consecutively rather than selectively.

## Research Population

A total of 104,516 inpatients with ACS, identified based on their principal diagnosis at discharge, were enrolled from 240 hospitals across China from November 2014 to December 2019. Patients with diabetes and complete HbA<sub>1c</sub> and FPG values at admission were included in this study. Only patients from July 2017 to December 2019 were included in this study because the FPG value at admission was not included in the database before July 2017. Patients were divided into a consistent group and a discrepancy group based on the HbA<sub>1c</sub> and FPG values at admission. The consistent group included patients with HbA<sub>1c</sub> < 6.5% and FPG < 7.0 mmol/L or patients with HbA<sub>1c</sub> ≥ 6.5% and FPG ≥ 7.0 mmol/L. The discrepancy group included patients with HbA<sub>1c</sub> ≥ 6.5% and FPG < 7.0 mmol/L or patients with HbA<sub>1c</sub> < 6.5% and FPG ≥ 7.0 mmol/L. The discrepancy group was further divided into an increased HbA<sub>1c</sub> but normal FPG group (HbA<sub>1c</sub> ≥ 6.5% and FPG < 7.0 mmol/L) and an increased FPG but normal HbA<sub>1c</sub> group (HbA<sub>1c</sub> < 6.5% and FPG ≥ 7.0 mmol/L). Institutional review board approval was granted for the use of an aggregate data set for research and quality improvement by the Ethics Committee of Beijing Anzhen Hospital, Capital Medical University. No informed consent was required.

## Definition Of Variables

Diabetes was defined as having a history of diabetes, receiving glucose-lowering agents before hospitalization, having diabetes listed in the medical records as a secondary discharge diagnosis, or having HbA<sub>1c</sub> ≥ 6.5% at admission. Hypertension was defined as having a history of hypertension, receiving antihypertensive medication, or having systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg at admission. The ACS classification was based on the primary diagnosis at discharge in the medical record. Non-ST-segment elevation ACS was defined as non-ST-segment elevation myocardial infarction (STEMI) or unstable angina. All the laboratory testing values were the values tested the first time after admission. The estimated glomerular filtration rate (eGFR) was calculated according to the equation developed by the Chronic Kidney Disease Epidemiology Collaboration [19]. Medication was prescribed after admission.

## In-hospital Outcomes

The outcomes of this study included major adverse cardiovascular and cerebrovascular events (MACCEs), heart failure, composite of cardiovascular death and heart failure, composite of MACCEs and

heart failure, and death from any cause. MACCEs were defined as cardiovascular death, cardiac arrest, cardiogenic shock, recurrent myocardial infarction, stent thrombosis, and stroke.

## Statistical analysis

Continuous variables were presented as the mean and standard deviation or median and interquartile range when the distribution and variance met the appropriate conditions. Categorical variables were presented as percentages. The comparisons between groups of continuous variables were performed by an unpaired t-test or a Mann-Whitney U test (Kruskal-Wallis), and the chi-square test was used to compare the categorical variables. A multivariate logistic regression model was used to determine the association between the type of discrepancy and in-hospital outcomes by controlling for potential confounders. Candidate adjustment factors included age, sex, blood pressure, heart rate, eGFR, hemoglobin, heart function, type of ACS, medical history, glucose-lowering drug use, in-hospital management, and referral status. The heterogeneity of effects on the in-hospital outcomes across subgroups was estimated using random effects meta-analysis. For data with missing values lower than 15% (Additional file 1: Table S1), the sequential regression multiple imputation method implemented by IVEware software version 0.2 (Survey Research Center, University of Michigan, Ann Arbor, MI, USA) was used to impute the missing values. All P values were 2-tailed, and  $P < 0.05$  was considered statistically significant. Statistical analyses were performed using SPSS 23.0 (SPSS Inc., Chicago, IL) and Stata/IC 15.1.

## Results

### Characteristics Of Patients In The Discrepancy Group

A total of 7,762 patients were included in this study (Additional file 1: Figure S1). The mean age was 64.4 ( $\pm 11.6$ ) years, and males accounted for 68.8% of the study population. The mean hemoglobin level was 135.1 ( $\pm 21.3$ ) g/L, and the mean eGFR was 81.7 ( $\pm 25.4$ ) ml min<sup>-1</sup> (1.73 m)<sup>-2</sup>. A total of 53.3% of patients were treated with at least one class of oral glucose-lowering drugs or insulin. The numbers of patients in the consistent group and the discrepancy group were 5,490 (70.7%) and 2,272 (29.3%), respectively. Patients in the discrepancy group were more likely to have lower eGFR (Additional file 1: Figure S2). In the discrepancy group, increased HbA<sub>1c</sub> but normal FPG accounted for 77.5% of the discrepancies (1,761/2,272), and increased FPG but normal HbA<sub>1c</sub> group accounted for 22.5% of the discrepancies (511/2,272). The baseline characteristics of patients in the increased HbA<sub>1c</sub> but normal FPG group and increased FPG but normal HbA<sub>1c</sub> group are shown in Table 1. Patients in the increased FPG but normal HbA<sub>1c</sub> group were more likely to have lower eGFR, higher heart rate, poorer heart function, STEMI and hypertension, and be treated with glucose-lowering agents.

Table 1  
 Characteristics of patients with discrepancy between HbA<sub>1c</sub> and FPG.

	<b>Increased HbA<sub>1c</sub> group (n = 1761)</b>	<b>Increased FPG group (n = 511)</b>	<b>P value</b>
Age (years, mean [SD])	65.4(11.2)	65.4(10.9)	0.947
Male (n [%])	1184(67.2)	348(68.1)	0.713
Systolic blood pressure (mmHg, mean [SD])	135.0(22.8)	135.0(25.2)	0.895
Diastolic blood pressure (mmHg, median [IQR])	78.0(70.0, 87.0)	78.0(70.0, 89.0)	0.442
Heart rate (bpm. Median [IQR])	78.0(68.0, 87.0)	80.0(70.0, 90.0)	< 0.001
Current smoker (n [%])	508(28.8)	153(29.9)	0.047
Family history of CHD (n [%])	71(4.0)	21(4.1)	0.937
Hypertension (n [%])	1300(73.8)	403(78.9)	0.021
Previous acute myocardial infarction (n [%])	214(12.2)	58(11.4)	0.623
Previous coronary artery bypass grafting (n [%])	16(0.9)	4(0.8)	0.789
Atrial fibrillation history (n [%])	53(3.0)	14(2.7)	0.751
Heart failure history (n [%])	58(3.3)	19(3.7)	0.640
Cerebrovascular disease history (n [%])	185(10.5)	57(11.2)	0.675
Peripheral artery disease history (n [%])	31(1.8)	10(2.0)	0.769
Killip class (n [%])			0.289

ACE, angiotensin-converting enzyme; ACS, acute coronary syndrome; CHD, coronary heart disease; eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA<sub>1c</sub>, glycosylated hemoglobin; HDL, high density lipoprotein; IQR, interquartile range; LDL, low density lipoprotein; NSTEMI, non-ST-segment elevation acute coronary syndrome; SD, standard deviation; STEMI, ST-segment elevation myocardial infarction.

	Increased HbA <sub>1c</sub> group (n = 1761)	Increased FPG group (n = 511)	<i>P</i> value
I or II	1505(85.5)	427(83.6)	
III or IV	256(14.5)	84(16.4)	
Types of ACS (n [%])			< 0.001
STEMI	716(40.7)	276(54.0)	
NSTE-ACS	1045(59.3)	235(46.0)	
HbA <sub>1c</sub> (%; mean [SD])	8.2(14.7)	5.8(0.8)	< 0.001
FPG (mmol/L; mean [SD])	5.6(1.2)	9.5(2.6)	< 0.001
eGFR (ml min <sup>-1</sup> [1.73m] <sup>-2</sup> ; mean [SD])	79.1(25.1)	76.4(27.3)	0.046
Hemoglobin (g/l; mean [SD])	132.1(20.4)	132.4(23.6)	0.192
Total cholesterol (mmol/L; median [IQR])	4.3(3.5, 5.1)	4.3(3.5, 5.1)	0.721
HDL-cholesterol (mmol/L; median [IQR])	1.0(0.8, 1.2)	1.0(0.8, 1.2)	0.931
LDL-cholesterol (mmol/L; median [IQR])	2.6(2.0, 3.2)	2.5(2.0, 3.2)	0.326
Triglyceride (mmol/L; median [IQR])	1.6(1.1, 2.4)	1.5(1.0, 2.3)	0.027
Oral glucose-lowering agents or insulin use before admission (n [%])	809(45.9)	288(56.4)	< 0.001
Therapy during hospitalization (n [%])			

ACE, angiotensin-converting enzyme; ACS, acute coronary syndrome; CHD, coronary heart disease; eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA<sub>1c</sub>, glycosylated hemoglobin; HDL, high density lipoprotein; IQR, interquartile range; LDL, low density lipoprotein; NSTE-ACS, non-ST-segment elevation acute coronary syndrome; SD, standard deviation; STEMI, ST-segment elevation myocardial infarction.

	Increased HbA <sub>1c</sub> group (n = 1761)	Increased FPG group (n = 511)	<i>P</i> value
Percutaneous coronary intervention	1170(66.4)	337(65.9)	0.836
Aspirin	1644(93.4)	482(94.3)	0.432
P2Y <sub>12</sub> inhibitors	1603(91.0)	473(92.6)	0.276
Statins	1653(93.9)	477(93.3)	0.669
β-blockers	1146(65.1)	291(56.9)	0.001
ACE inhibitor/angiotensin receptor blocker	917(52.1)	248(48.5)	0.159
Patients with referral (n [%])	587(33.3)	172(33.7)	0.891
ACE, angiotensin-converting enzyme; ACS, acute coronary syndrome; CHD, coronary heart disease; eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA <sub>1c</sub> , glycosylated hemoglobin; HDL, high density lipoprotein; IQR, interquartile range; LDL, low density lipoprotein; NSTEMI, non-ST-segment elevation acute coronary syndrome; SD, standard deviation; STEMI, ST-segment elevation myocardial infarction.			

The prevalence of the increased FPG but normal HbA<sub>1c</sub> discrepancy was higher in patients over 65 years of age, hemoglobin less than 120 g/L, eGFR less than 60 ml min<sup>-1</sup> (1.73 m)<sup>-2</sup>, Killip class III or IV, and STEMI and who were treated with oral glucose-lowering drugs or insulin (Fig. 1, Additional file 1: Table S2).

## Type Of Discrepancy And In-hospital Outcomes

The comparison of in-hospital outcomes between the increased HbA<sub>1c</sub> but normal FPG group and the increased FPG but normal HbA<sub>1c</sub> group is shown in Fig. 2. The rates of all the in-hospital outcomes were higher in the increased FPG but normal HbA<sub>1c</sub> group than in the increased HbA<sub>1c</sub> but normal FPG group. A logistic regression model was performed to explore the relationship between the type of discrepancy and in-hospital outcomes, except for death from any cause because of the small event number. In univariate logistic regression analysis, a significantly higher risk of all the in-hospital outcomes was observed in patients with increased FPG but normal HbA<sub>1c</sub> (Table 2). After adjusting for confounders in the multivariate logistic regression model, patients in the increased FPG but normal HbA<sub>1c</sub> group had a significant 1.6-fold increased risk of heart failure (OR, 1.62; 95% CI, 1.08–2.44), a 1.6-fold increased risk of composite cardiovascular death and heart failure (OR, 1.63; 95% CI, 1.09–2.43), and a 1.6-fold

increased risk of composite MACCEs and heart failure (OR, 1.56; 95% CI, 1.08–2.24) (Table 2). The effect on MACCEs was not significant (OR, 1.49; 95% CI, 0.85–2.63) (Table 2).

Table 2  
Logistic regression analysis for in-hospital outcomes in increased FPG group compared with increased HbA<sub>1c</sub> group.

	<b>Unadjusted OR (95% CI)</b>	<b>P value</b>	<b>Adjusted OR (95% CI)*</b>	<b>P value</b>
MACCE	1.94(1.13–3.34)	0.016	1.49(0.85–2.63)	0.165
Heart failure	2.09(1.42–3.07)	< 0.001	1.62(1.08–2.44)	0.020
Cardiovascular death or heart failure	2.11(1.44–3.07)	< 0.001	1.63(1.09–2.43)	0.018
MACCE or heart failure	1.98(1.41–2.79)	< 0.001	1.56(1.08–2.24)	0.018
* ORs were adjusted for age, gender, systolic blood pressure, heart rate, current smoker, hypertension, hemoglobin at admission, eGFR at admission, Killip class, type of acute coronary syndrome, and glucose-lowering drug use.				
FPG, fasting plasma glucose; HbA <sub>1c</sub> , glycosylated hemoglobin; MACCE, major adverse cardiovascular and cerebrovascular event.				

Subgroup analysis was performed based on age, sex, medical history, Killip class, hemoglobin, eGFR, type of ACS, and glucose-lowering drug use before hospitalization. A higher risk of all the in-hospital outcomes was observed in patients with increased FPG but normal HbA<sub>1c</sub>, which was consistent in all subgroups (all P for interaction > 0.05), except for the eGFR subgroup for MACCEs (Fig. 3), which showed that increased FPG but normal HbA<sub>1c</sub> increased the risk of MACCEs to a greater extent in patients with eGFR  $\geq 60 \text{ ml min}^{-1} (1.73 \text{ m})^{-2}$ .

## Discussion

In this study, we investigated the type of discrepancy between HbA<sub>1c</sub> and FPG in patients with ACS and diabetes. We found that nearly one third of patients had a discrepancy between HbA<sub>1c</sub> and FPG. Of the patients with discrepancies, the patients with increased FPG had a higher risk of in-hospital adverse cardiovascular outcomes than those with increased HbA<sub>1c</sub>.

Discrepancies between HbA<sub>1c</sub> and FPG have been reported by some studies. A study of the risk of hypertension in patients with prediabetes demonstrated the discrepancy between HbA<sub>1c</sub> and FPG [20]. A study using data from residents of Yunnan Province, China, showed that a discrepancy between HbA<sub>1c</sub> and FPG was present in approximately 30% of participants [21]. In our study, the discrepancy between

HbA<sub>1c</sub> and FPG was also found in patients with ACS and diabetes. We found that the discrepancy group, composed of 77.5% patients with increased HbA<sub>1c</sub> but normal FPG and 22.5% patients with increased FPG but normal HbA<sub>1c</sub>, accounted for 29% of the total study population. Patients often experience hyperglycemia in the acute phase of many diseases, such as ACS, which is called stress hyperglycemia. HbA<sub>1c</sub> reflects average glycemia over approximately 3 months, so an increase in HbA<sub>1c</sub> usually indicates chronic hyperglycemia. We found that patients in the increased FPG but normal HbA<sub>1c</sub> group were more likely to have lower eGFR and be treated with glucose-lowering agents. A higher proportion of glucose-lowering agent use may be related to well-controlled blood glucose and lower HbA<sub>1c</sub>. Furthermore, changes in the metabolism of glucose-lowering drugs, insulin clearance, and the uremic environment in patients with renal function insufficiency may also reduce HbA<sub>1c</sub> values [22]. From our study, not only was a discrepancy between HbA<sub>1c</sub> and FPG be found in patients with chronic kidney disease but the proportion of the increased FPG group was found to be significantly higher than that of the increased HbA<sub>1c</sub> group.

There is a strong association between cardiovascular disease, diabetes and chronic kidney disease. People with diabetes and chronic kidney disease have a substantially increased risk of all-cause mortality, cardiovascular mortality, and kidney failure [23, 24]. Furthermore, we analyzed the relationship between the type of discrepancy and in-hospital outcomes. We know that HbA<sub>1c</sub> and FPG are both closely related to in-hospital outcomes. Most previous studies have shown that increased HbA<sub>1c</sub> or FPG was significantly associated with poor in-hospital outcomes in patients with ACS and diabetes. An observational study that included 250 patients with ACS found that coronary atherosclerosis was more advanced in patients with HbA<sub>1c</sub>  $\geq$  5.7% than in those with HbA<sub>1c</sub>  $<$  5.7% [17]. Goyal et al [25] conducted a post hoc analysis including two randomized controlled trials of acute myocardial infarction with ST-segment elevation, involving 30,536 subjects with diabetes history, and showed that patients with in-hospital glucose  $\geq$  144 mg/dL had a very high risk of death. However, in clinical practice, some conditions, such as acute stress, renal dysfunction, and anemia, can cause uncertainty in the measured values of FPG, HbA<sub>1c</sub>, and the discrepancy between FPG and HbA<sub>1c</sub>. Until now, the association of in-hospital outcomes with the discrepancy between HbA<sub>1c</sub> and FPG in patients with ACS and diabetes has not been clear. There are few studies focusing on this issue. From our study, we can conclude that patients in the increased FPG group, who were more likely to have a higher heart rate, poorer heart function, a higher incidence of STEMI and hypertension, had a higher risk of in-hospital cardiovascular adverse outcomes than those with increased HbA<sub>1c</sub>. Stress hyperglycemia, which is a reflection of high free fatty acids, insulin resistance, and steroid hormones, affects the course of the disease in an adverse way [26]. From another study, we learned that the level of stress hyperglycemia often correlates with the severity of disease and can predict mortality [27]. In our study, we also found that patients with severe clinical conditions, such as a higher heart rate and poorer heart function, were more likely to have increased FPG. As a result, stress hyperglycemia may have a greater adverse effect on patients with ACS and diabetes than chronic hyperglycemia.

The findings of this study may have some important implications for clinical practice. The HbA<sub>1c</sub> test is a major tool for assessing glycemic control and has strong predictive value for diabetes complications [28]. Chronic hyperglycemia is an important risk factor for cardiovascular disease and mortality [24], although the variability in HbA<sub>1c</sub> in patients with renal insufficiency should be considered. However, in patients with ACS and diabetes, increased FPG may be associated with a higher risk of adverse in-hospital outcomes, even if HbA<sub>1c</sub> is well controlled. These patients, especially those with renal insufficiency, should be given more attention and closer monitoring in clinical practice.

The major strength of our study is that it is based on a nationally representative registry and aimed at investigating the discrepancy between HbA<sub>1c</sub> and FPG and the influence of the discrepancies on the in-hospital outcomes of patients with ACS and diabetes, which has rarely been reported until now. Our study also has certain limitations. First, all-cause mortality was not included in the logistic regression analysis because of very limited events. Second, we could not collect all information related to glucose metabolism in this real-world study of ACS patients based on medical records, thus contributing to some residual confounding from unmeasured confounders. Last, fasting status, blood sample collection and testing methods were difficult to unify, as this was a real-world multicenter study.

## Conclusions

In summary, our study showed that the discrepancy between HbA<sub>1c</sub> and FPG accounts for nearly 30% of discrepancies among patients with ACS and diabetes. Patients with an increased level of FPG had a higher risk of in-hospital cardiovascular adverse outcomes than those with an increased level of HbA<sub>1c</sub>. This result may indicate that when HbA<sub>1c</sub> and FPG are inconsistent in patients with ACS and diabetes, the increased FPG that may be caused by stress hyperglycemia may have a more substantial adverse effect than increased HbA<sub>1c</sub>, which may be caused by chronic hyperglycemia. These high-risk patients, especially those with renal insufficiency, should be given more attention and closer monitoring in clinical practice.

## Abbreviations

ACS, acute coronary syndrome; CCC-ACS, Improving Care for Cardiovascular Disease in China - acute coronary syndrome; eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA<sub>1c</sub>, glycosylated hemoglobin; MACCEs, major adverse cardiovascular and cerebrovascular events; STEMI, ST-segment elevation myocardial infarction

## Declarations

## Ethics approval and consent to participate

Institutional review board approval was granted for the aggregate data set for research and quality improvement by the Ethics Committee of Beijing Anzhen Hospital, Capital Medical University. Being a large population-based study, approval from Ethics Committee of Beijing Anzhen Hospital, Capital Medical University included a waiver of informed consent.

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

The datasets used and analyzed during the current study are available from the principal investigator of CCC-ACS on reasonable request.

### **Competing interests**

The authors declare that they have no competing interests.

## **Funding**

The CCC-ACS project is a collaborative program of the American Heart Association and Chinese Society of Cardiology. The American Heart Association was funded by Pfizer and AstraZeneca for the quality improvement initiative through an independent grant for learning and change. The funding bodies had no role in the design of this study and collection, analysis, and interpretation of data and in writing the manuscript.

## **Authors' contributions**

HC, NY, DZ, CSM and JL designed the study; NY, LJY, GQW, WJB, FBX, YCH, JL and NY cleaned the data; NY and LJY analyzed the data; NY and HC wrote the manuscript. All authors reviewed and edited the manuscript. All authors read and approved the final manuscript.

## **Acknowledgments**

We acknowledge all participating hospitals for their contributions to the CCC-ACS project (Additional file 1: Table S3). We would like to thank AJE ([www.aje.cn](http://www.aje.cn)) for English language editing.

## **References**

1. Zhao D, Liu J, Wang M, Zhang X, Zhou M. Epidemiology of cardiovascular disease in China: current features and implications. *Nat Rev Cardiol.* 2019;16:203–12. doi:10.1038/s41569-018-0119-4.


2. Lenzen M, Ryden L, Ohrvik J, Bartnik M, Malmberg K, Scholte Op Reimer W, et al. Diabetes known or newly detected, but not impaired glucose regulation, has a negative influence on 1-year outcome in patients with coronary artery disease: a report from the Euro Heart Survey on diabetes and the heart. *Eur Heart J*. 2006;27:2969–74. doi:10.1093/eurheartj/ehl363.
3. Figueiredo VN, Godoi FC, Martins NS, Quinaglia e Silva JC, Nadruz W Jr, Coelho OR, et al. Diabetes mellitus unawareness is a strong determinant of mortality in patients manifesting myocardial infarction. *Curr Med Res Opin*. 2013;29:1423–7. doi:10.1185/03007995.2013.832186.
4. Aggarwal B, Shah GK, Randhawa M, Ellis SG, Lincoff AM, Menon V. Utility of Glycated Hemoglobin for Assessment of Glucose Metabolism in Patients With ST-Segment Elevation Myocardial Infarction. *Am J Cardiol*. 2016;117:749–53. doi:10.1016/j.amjcard.2015.11.060.
5. Rawshani A, Rawshani A, Franzen S, Sattar N, Eliasson B, Svensson AM, et al. Risk Factors, Mortality, and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med*. 2018;379:633–44. doi:10.1056/NEJMoa1800256.
6. Zhou M, Liu J, Hao Y, Liu J, Huo Y, Smith SC Jr, et al. Prevalence and in-hospital outcomes of diabetes among patients with acute coronary syndrome in China: findings from the Improving Care for Cardiovascular Disease in China-Acute Coronary Syndrome Project. *Cardiovasc Diabetol*. 2018;17:147. doi:10.1186/s12933-018-0793-x.
7. Bjarnason TA, Hafthorsson SO, Kristinsdottir LB, Oskarsdottir ES, Johnsen A, Andersen K. The prognostic effect of known and newly detected type 2 diabetes in patients with acute coronary syndrome. *Eur Heart J Acute Cardiovasc Care*. 2019:2048872619849925. doi:10.1177/2048872619849925.
8. Nystrom T, Sartipy U, Contardi A, Lind M, Bellocco R, Eliasson B, et al. Glycated Hemoglobin A1c Levels in Type 1 Diabetes Mellitus and Outcomes After Myocardial Infarction. *Circulation*. 2019;139:2380–2. doi:10.1161/CIRCULATIONAHA.118.039223.
9. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2013;36:67–74. doi:10.2337/dc13-S067.
10. Dziejewicz A, Giszterowicz D, Siudak Z, Rakowski T, Dubiel JS, Dudek D. Admission glucose level and in-hospital outcomes in diabetic and non-diabetic patients with acute myocardial infarction. *Clin Res Cardiol*. 2010;99:715–21. doi:10.1007/s00392-010-0175-1.
11. Li DB, Hua Q, Guo J, Li HW, Chen H, Zhao SM. Admission glucose level and in-hospital outcomes in diabetic and non-diabetic patients with ST-elevation acute myocardial infarction. *Intern Med*. 2011;50:2471–5. doi:10.2169/internalmedicine.50.5750.
12. Sanjuán R, Núñez J, Luisa Blasco M, Miñana G, Martínez-Maicas H, Carbonell N, et al. Prognostic Implications of Stress Hyperglycemia in Acute ST Elevation Myocardial Infarction. Prospective Observational Study. *Revista Española de Cardiología. (English Edition)*. 2011;64:201–7. doi:10.1016/j.rec.2010.08.005.
13. Tsai JP, Tian J, Wang WY, Ng AC. Glycated hemoglobin vs fasting plasma glucose as a predictor of left ventricular dysfunction after ST-elevation myocardial infarction. *Can J Cardiol*. 2015;31:44–9.

doi:10.1016/j.cjca.2014.10.029.

14. Liang H, Guo YC, Chen LM, Li M, Han WZ, Zhang X, et al. Relationship between fasting glucose levels and in-hospital mortality in Chinese patients with acute myocardial infarction and diabetes mellitus: a retrospective cohort study. *BMC Cardiovasc Disord*. 2016;16:156. doi:10.1186/s12872-016-0331-2.
15. Farhan S, Clare RM, Jarai R, Giugliano RP, Likhnygina Y, Harrington RA, et al. Fasting glucose, NT-proBNP, treatment with eptifibatide, and outcomes in non-ST-segment elevation acute coronary syndromes: An analysis from EARLY ACS. *Int J Cardiol*. 2017;232:264–70. doi:10.1016/j.ijcard.2017.01.007.
16. Zhao S, Murugiah K, Li N, Li X, Xu ZH, Li J, et al. Admission Glucose and In-hospital Mortality after Acute Myocardial Infarction in Patients with or without Diabetes: A Cross-sectional Study. *Chin Med J (Engl)*. 2017;130:767–75. doi:10.4103/0366-6999.202733.
17. Acar B, Ozeke O, Karakurt M, Ozen Y, Ozbay MB, Unal S, et al. Association of Prediabetes With Higher Coronary Atherosclerotic Burden Among Patients With First Diagnosed Acute Coronary Syndrome. *Angiology*. 2019;70:174–80. doi:10.1177/0003319718772420.
18. Hao Y, Liu J, Liu J, Smith SC Jr, Huo Y, Fonarow GC, et al. Rationale and design of the Improving Care for Cardiovascular Disease in China (CCC) project: A national effort to prompt quality enhancement for acute coronary syndrome. *Am Heart J*. 2016;179:107–15. doi:10.1016/j.ahj.2016.06.005.
19. Levey ASSL, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J. CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A New Equation to Estimate Glomerular Filtration Rate. *Ann Intern Med*. 2009;150:604–12.
20. Geva M, Shlomai G, Berkovich A, Maor E, Leibowitz A, Tenenbaum A, et al. The association between fasting plasma glucose and glycated hemoglobin in the prediabetes range and future development of hypertension. *Cardiovasc Diabetol*. 2019;18:53. doi:10.1186/s12933-019-0859-4.
21. Feng L, Nian S, Zhao Y, Bai X, Luo F, Luo X, et al. Higher HbA1c and/or glucose levels alter the association patterns between glycated hemoglobin and fasting glucose levels. *Diabetes Res Clin Pract*. 2018;142:353–62. doi:10.1016/j.diabres.2018.06.011.
22. Kuo IC, Lin HY, Niu SW, Hwang DY, Lee JJ, Tsai JC, et al. Glycated Hemoglobin and Outcomes in Patients with Advanced Diabetic Chronic Kidney Disease. *Sci Rep*. 2016;6:20028. doi:10.1038/srep20028.
23. Fox CS, Matsushita K, Woodward M, Bilo HJG, Chalmers J, Heerspink HJL, et al. Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *The Lancet*. 2012;380:1662–73. doi:10.1016/s0140-6736(12)61350-6.
24. American Diabetes Association. 10. Cardiovascular Disease and Risk Management: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43:S111-S134. doi:10.2337/dc20-S010.
25. Goyal A, Mehta SR, Gerstein HC, Diaz R, Afzal R, Xavier D, et al. Glucose levels compared with diabetes history in the risk assessment of patients with acute myocardial infarction. *Am Heart J*. 2009;157:763–70. doi:10.1016/j.ahj.2008.12.007.


26. Görmeli Kurt N. Relation of Stress Hyperglycemia to Mortality in Acute Myocardial Infarction Patients. *Eurasian Journal of Medicine Oncology*. 2018;2:138–41. doi:10.14744/ejmo.2018.49469.
27. Cheung NW, Wong KYC, Kovoor P, McLean M. Stress hyperglycemia: A prospective study examining the relationship between glucose, cortisol and diabetes in myocardial infarction. *J Diabetes Complications*. 2019;33:329–34. doi:10.1016/j.jdiacomp.2018.12.015.
28. American Diabetes Association. 6. Glycemic Targets: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43:66–76. doi:10.2337/dc20-S006.

## Figures




**Figure 1**

Prevalence of discrepancy in different population. eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA<sub>1c</sub>, glycosylated hemoglobin; NSTE-ACS, non-ST-segment elevation acute coronary syndrome; STEMI, ST-segment elevation myocardial infarction.



**Figure 2**

In-hospital outcomes in patients with discrepancy between HbA<sub>1c</sub> and FPG. FPG, fasting plasma glucose; HbA<sub>1c</sub>, glycosylated hemoglobin; MACCE, major adverse cardiovascular and cerebrovascular event.



**Figure 3**

Subgroup analysis for association between the type of discrepancy and in-hospital outcomes. Panel a shows the effect of increased FPG group on MACCE compared with increased HbA1c group. Panel b shows the effect of increased FPG group on heart failure compared with increased HbA1c group. Panel c shows the effect of increased FPG group on the composite of cardiovascular death and heart failure compared with increased HbA1c group. Panel d shows the effect of increased FPG group on the composite of MACCE and heart failure compared with increased HbA1c group. eGFR, estimated glomerular filtration rate; FPG, fasting plasma glucose; HbA1c, glycosylated hemoglobin; MACCE, major adverse cardiovascular and cerebrovascular event; NSTE-ACS, non-ST-segment elevation acute coronary syndrome; STEMI, ST-segment elevation myocardial infarction.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.pdf](#)