

TABLE 5. Complications in the first year post-KT in the pairs with different 1-year proteinuria < 0,5 g/day and ≥ 0,5 g/day.

	<b>Kidney pairs (86 pts)</b>	<b>Kidney twin with 1 year PTO &lt; 0,5 g/L (43 pts)</b>	<b>Kidney twin with 1 year PTO ≥ 0,5 g/L (43 pts)</b>	<b>P value</b>
Post-Transplant Complications in 1 <sup>st</sup> post-KT year				
DGF %	24 (28)	8 (19)	16 (38)	0,03
Transplant Urologic Complications %	48 (56)	23 (53)	25 (58)	0,41
Ureteral fistula %	3 (3)	1 (2)	2 (5)	0,5
Ureteral stenosis %	7 (8)	1 (2)	6 (14)	0,05
Transplant Vascular Complications %	19 (22)	11 (26)	8 (19)	0,3
Vascular fistula %	8 (9)	5 (12)	3 (7)	0,36
Renal artery stenosis %	9 (10)	4 (9)	5 (12)	0,5
Overall infections %	66 (78)	31 (74)	35 (81)	0,28
CMV infections	29 (34)	13 (30)	16 (37)	0,32
Ischemic Cardiopathy	7 (8)	5 (12)	2 (5)	0,22
NODAT	22 (26)	9 (21)	13 (30)	0,32
Extrarenal Major Vascular Complications*	18 (21)	10(23)	8(19)	0,60
Hypertension	27 (96)	12 (44)	15 (56)	0,57
Rejection	7 (8)	0 (0)	7 (16)	<0,01
Biopsies (%)	51 (65)	18 (49)	33 (80)	<0,01
Glomerulonephritis (%)	7 (8)	1 (2)	6 (14)	0,05

DGF = delayed graft function; CMV = cytomegalovirus; NODAT=new onset diabetes after transplantation. All Biopsies are performed for cause.

\*Including stroke, arterial stenosis and thrombosis.