

Investigation of the Clinical Significance and Pathological Features of Lanthanum Deposition in the Gastric Mucosa

Shinya Nishida

Osaka Ika Daigaku

Kazuhiro Ota (✉ clash_kaz@yahoo.co.jp)

Osaka Medical College <https://orcid.org/0000-0002-5628-9662>

Kimiaki Hattori

Moriguchi Keijinkai Byouin

Taro Iwatsubo

Osaka Ika Daigaku

Yuichi Kojima

Osaka Ika Daigaku

Toshihisa Takeuchi

Osaka Ika Daigaku

Tamaki Maeda

Moriguchi Keijinkai Byouin

Masahiro Sakaguchi

Moriguchi Keijinkai Byouin

Kazuhide Higuchi

Osaka Ika Daigaku

Shimpei Kawaguchi

Second Department of Internal Medicine

Research article

Keywords: lanthanum, phosphate binder, histiocytic infiltration, whitish lesion

Posted Date: May 21st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-30322/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 23rd, 2020. See the published version at <https://doi.org/10.1186/s12876-020-01545-z>.

Abstract

Background: There are often specific endoscopic findings caused by deposition of lanthanum (La) in the gastric mucosa of patients taking lanthanum carbonate (LaC), a novel phosphate binder for patients on hemodialysis. We conducted a retrospective study to investigate the clinical significance of Ladeposition in the gastric mucosa, and the association between endoscopic features and histologic findings in the same population.

Methods:We compared background factors in patients taking LaC with and without Ladeposition in their gastroscopic biopsy specimen. We also investigated the relationship between gastric endoscopic biopsy specimens with Ladeposition and the concurrent endoscopic images.

Results:There was a significant difference in the total dose of LaC between the La-positive and La-negative groups(990 g [180–3150 g] vs. 480 g [225–1328 g]; $p=0.013$). In 27 biopsy specimens with specific whitish mucosa, 10 showed mild histiocytic infiltration and 17 showed strong infiltration. In contrast, among 24 specimens with non-whitishmucosa, five showed no histiocytic infiltration, 10 showed mild infiltration, and nine showed strong infiltration. There was a significant relationship between endoscopic features and the degree of histiocytic infiltration ($p=0.026$).

Conclusions: We demonstrated that Ladeposition in the gastric mucosa depended on the total dose of LaC and was not affected by background factors. The specific endoscopic features of Ladeposition are associated with the infiltration of histiocytes, which represents the body's normal response to foreign bodies.

The protocol was registered in the University Hospital Medical Information Network Clinical Trial Registry (UMIN000038929).

Introduction

The number of dialysis patients in Japan reached 329,609 in 2016 [1]. For dialysis patients, hyperphosphatemia is one of the critical complications because it significantly increases cardiovascular disease and mortality. To control the blood concentration of phosphoric acid, several types of oral phosphate binders have been developed, including lanthanum carbonate (LaC) (Fosrenol®; Shire Pharmaceuticals, Hampshire, UK), which has been authorized by the Ministry of Health, Labor and Welfare in Japan. LaC is the first non-calcium, non-resin phosphate binder for hyperphosphatasemia in patients with chronic kidney disease on hemodialysis. The use of LaC has increased in Japan since its launch in 2009 as it has fewer side effects than conventional medicines. Recently, it was reported that there are specific endoscopic findings caused by deposition of lanthanum (La) in the gastric mucosa of patients taking LaC [2].

We previously reported that the daily dose and total dose of LaC were significantly correlated with La deposition in the gastroduodenal mucosa [3]. However, it is still unknown why these phenomena appear in LaC users and what subsequent difficulties, if any, appear in patients with La deposition in the gastrointestinal mucosa. Although it was shown that patients on hemodialysis with anemia are at a high risk of small intestinal lesions in our previous study [4], the relationships between La deposition and other factors were not investigated.

Murakami et al. investigated seven cases with the specific endoscopic features of La deposition in the gastric mucosa and described the whitish lesions as follows: an annular whitish mucosa, diffuse whitish mucosa, and whitish spots in the gastric mucosa [5]. Although the endoscopic biopsy specimens from the above endoscopic features revealed La deposition histologically, it was uncertain whether these whitish lesions were pathologically consistent with La deposition.

We therefore conducted the two following clinical retrospective investigations: 1) comparative analysis of the clinical features of LaC users with and without La deposition in endoscopic biopsy specimens from gastric mucosa and 2) histological identification of the cause of the whitish lesions using biopsy specimens with La deposition by comparing the histological features of the endoscopic findings.

Methods

These two investigations were conducted in accordance with the 1975 Declaration of Helsinki (as revised in 1983). The protocol was approved by the ethics review committee at Moriguchi Keijinkai Hospital (No. 3104) and registered in the University Hospital Medical Information Network Clinical Trial Registry (UMIN000038929).

Patient population and study design

Study 1

This retrospective observational investigation included patients on hemodialysis taking LaC from whom biopsies from the gastric mucosa were taken via gastric endoscopy at Moriguchi Keijinkai Hospital between April 2014 and August 2019. Most dialysis patients in the Moriguchi Keijinkai Hospital are advised to undergo esophagogastroduodenoscopy regularly to screen for malignant tumors by their attending physician even if they are asymptomatic. An endoscopist judges each case based on its individual characteristics before performing endoscopic biopsy. The aims of biopsy varied: searching for malignant tumors, confirming benign lesions, and determining the cause of unknown lesions (including whitish mucosal lesion by La deposition). All biopsy specimens, which were fixed with 10% formalin and embedded in paraffin, were stained with hematoxylin and eosin. In the histological evaluation of biopsy specimens, the crystalloids dyed with eosin were diagnosed as La deposition based on a previous report [3]. Our main objective was to reveal the clinical significance of La deposition in the gastric mucosa by comparing the clinical features of patients with and without La deposition in endoscopic biopsy

specimens from the gastric mucosa. The characteristics compared were as follows: sex, age, the degree of gastric mucosal atrophy, the primary disease at time of dialysis introduction, the dialysis period, the period of taking LaC, the daily dose of LaC, the total dose of LaC, blood test findings (calcium, inorganic phosphorus, total protein, albumin, total cholesterol, HDL-cholesterol, triglyceride, white blood cell count, and hemoglobin), and concomitant drugs (non-steroidal anti-inflammatory drugs [NSAIDs] including low-dose aspirin, proton pump inhibitors [PPIs], histamine H2 receptor antagonists [H2RAs], and gastric mucoprotective drugs). All items were collected two weeks before and after endoscopic biopsy.

In addition, we compared the group of patients taking LaC with the control group of dialysis patients; the abovementioned patients taking LaC with and without gastric La deposits, and the control group of dialysis patients who were not taking LaC and underwent endoscopic biopsy in the same period.

Study 2

This retrospective investigation was performed using the gastric endoscopic biopsy specimens with La deposition as well as the concurrent endoscopic images. First, we confirmed the presence or absence of whitish lesions - the specific endoscopic feature of La deposition - in the gastric mucosa at the endoscopic biopsy sites (Fig. 1). In this study, we determined the specific whitish mucosa of the biopsy site in the following cases: when an endoscopist clearly recognized the whitish lesions and biopsied them, or when a biopsy site unexpectedly contained a white lesion. Two independent gastrointestinal endoscopists (S.N. and K.O.) performed these evaluations retrospectively. When opinions differed between the endoscopists, a third independent gastrointestinal endoscopist (Y.K.) was consulted. Second, for each biopsy specimen we evaluated the amount of deposited La and the degree of histiocytic infiltration, which are the characteristic findings for diagnosis of La deposition. These histological evaluations were performed by two pathologists (K.H. and T.M.). Finally, we contrasted the endoscopic features and histological findings and investigated the pathological significance of the whitish lesions as specific findings of La deposition.

Statistical analysis

Significant differences between groups were determined using the Mann-Whitney U test for continuous or categorical variables and the Pearson's chi-square test for binary variables. The trends in binomial proportions across the levels of a single variable were determined using the Cochran-Armitage trend test. All reported p-values were two-sided, and values < 0.05 were considered statistically significant. All calculations were performed with JMP Pro 14 software (SAS Institute Inc., Cary, NC, USA).

Results

Study 1

This analysis was performed on 42 patients who were on hemodialysis and taking LaC medication and who underwent gastric endoscopy and endoscopic biopsy from the gastric mucosa at Moriguchi Keijinkai Hospital during the study period. The number of patients with La deposition in the gastric mucosa (La-positive group) was 23, and the number without La deposition was 19 (La-negative group). There was a significant difference between the two groups regarding the total dose of LaC (La-positive group, 990 g [180–3150 g]; La-negative group, 480 g [225–1328 g]; $p = 0.013$, Mann-Whitney U test). Other variables, including the daily dose of LaC, were not significantly different in the two groups; however, there was some missing blood test data (Table 1).

Table 1
Patient characteristics in Study 1

Patient characteristics	La deposition (+), n = 23	La deposition (-), n = 19	Odds ratio (95%CI)	p value
Male sex, n (%)	14 (60.9%)	13 (68.4%)	1.39 (0.39–5.01)	0.258
Age, median (range)	72 (47–82)	68 (45–83)	-	0.909
Gastric mucosal atrophy (Mild, Medium, Severe)*, n (%)	5 (21.7%), 7 (30.4%), 9 (47.8%)	7 (36.8%), 3 (15.8%), 9 (47.8%)	-	0.413
Primary diseases of dialysis (Diabetes, Hypertension, Nephritis), n (%)	8 (34.8%), 3 (13.4%), 12 (52.2%)	10 (52.6%), 2 (10.5%), 7 (36.8%)	-	0.504
Years since start of dialysis, median (range)	7.32 (0.70–26.4)	5.83 (1.57–14.2)	-	0.092
Months of oral use of LaC, median (range)	34 (4–70)	28 (10–72)	-	0.206
Daily dose of LaC, mg, median (range)	1000 (500–1500)	750 (250–1500)	-	0.104
Total dose of LaC, g, median (range)	990 (180–3150)	480 (225–1328)	-	0.013
Ca, mg/dL, median (range), (Number of missing data: 4)	8.9 (8-10.2)	8.4 (7.2–9.9)	-	0.084
IP, mg/dL, median (range), (Number of missing data: 7)	5.5 (3.5–8.1)	5.3 (3.1–8.3)	-	0.586
TP, g/dL, median (range), (Number of missing data: 8)	6.4 (4.8–7.9)	6.35 (4.3–7.3)	-	0.479
Albumin, g/dL, median (range), (Number of missing data: 7)	3.5 (2.7-4.0)	3.6 (2.4-4.0)	-	0.752
Total cholesterol, mg/dL, median (range), (Number of missing data: 10)	166 (103–203)	151 (88–240)	-	0.454
HDL-cholesterol, mg/dL, median (range), (Number of missing data: 13)	47 (24–67)	47 (18–97)	-	0.790

* The degree of gastric mucosal atrophy is classified into mild, medium, or severe according to the Takemoto-Kimura classification [13]. Mild: C-1 and C-2, Medium: C-3 and O-1, Severe: O-2 and O-3.

Bold font represents statistical significance.

La, lanthanum; CI, confidence interval; LaC, lanthanum carbonate; Ca, calcium; IP, inorganic phosphorus; TP, total protein; WBC, white blood cell; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitor

Patient characteristics	La deposition (+), n = 23	La deposition (-), n = 19	Odds ratio (95%CI)	p value
Triglyceride, mg/dL, median (range), (Number of missing data: 9)	103 (50–336)	94.5 (35–198)	-	0.649
WBC, /μL, median (range), (Number of missing data: 6)	5460 (3370–8300)	6535 (2650–12470)	-	0.133
Hemoglobin, g/dL, median (range), (Number of missing data: 5)	10.5 (7.3–12.5)	10.55 (7.3–12.8)	-	0.976
NSAIDs (including low-dose aspirin), n (%)	10 (43.5%)	10 (52.6%)	0.69 (0.20–2.35)	0.554
PPIs, n (%)	17 (73.9%)	17 (89.5%)	0.33 (0.06–1.89)	0.201
PPIs + histamine H2 antagonists, n (%)	21 (91.3%)	17 (89.5%)	1.24 (0.16–9.71)	0.841
Gastric mucoprotective drugs, n (%)	9 (39.1%)	4 (21.1%)	2.41 (0.60–9.63)	0.207
* The degree of gastric mucosal atrophy is classified into mild, medium, or severe according to the Takemoto-Kimura classification [13]. Mild: C-1 and C-2, Medium: C-3 and O-1, Severe: O-2 and O-3.				
Bold font represents statistical significance.				
La, lanthanum; CI, confidence interval; LaC, lanthanum carbonate; Ca, calcium; IP, inorganic phosphorus; TP, total protein; WBC, white blood cell; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitor				

There was a significant difference in years since start of dialysis between the LaC taking group (n = 42) and the control dialysis group (n = 42). There were no significant differences in other characteristics between two groups (Table 2).

Table 2
Comparison of the LaC taking patients and the control dialysis patients

Patient characteristics	LaC taking patients, n = 42	Control dialysis patients, n = 42	Odds ratio (95%CI)	p value
Male sex, n (%)	15 (35.7%)	20 (47.6%)	0.61 (0.25–1.47)	0.269
Age, median (range)	68 (45–83)	70 (35–86)	-	0.400
Gastric mucosal atrophy (Mild, Medium, Severe)*, n (%)	12 (28.6%), 10 (23.8%), 20 (47.6%)	14 (33.3%), 10 (23.8%), 18 (42.9%)	-	0.879
Primary diseases of dialysis (Diabetes, Hypertension, Nephritis), n (%)	18 (42.9%), 5 (11.9%), 19 (45.2%)	13 (31.0%), 7 (16.7%), 22 (52.4%)	-	0.507
Years since start of dialysis, median (range)	6.12 (0.7–26.4)	3.58 (0.002–12.5)	-	0.0002
Ca, mg/dL, median (range)	8.75 (7.2–10.2)	8.9 (6.1–10.3)	-	0.300
IP, mg/dL, median (range)	5.5 (3.1–8.3)	4.9 (1.9–7.5)	-	0.320
TP, g/dL, median (range)	6.4 (4.3–7.9)	6.6 (3.9–7.9)	-	0.184
Albumin, g/dL, median (range)	3.5 (2.4–4.0)	3.5 (2.5–4.7)	-	0.362
Total cholesterol, mg/dL, median (range)	158.5 (88–240)	160 (93–303)	-	0.402
HDL-cholesterol, mg/dL, median (range)	47 (18–97)	49.5 (23–78)	-	0.601
Triglyceride, mg/dL, median (range)	97 (35–336)	92.5 (36–271)	-	0.721
WBC, /μL, median (range)	5720 (2650–12470)	5210 (2800–11800)	-	0.068
Hemoglobin, g/dL, median (range)	10.5 (7.3–12.8)	10.6 (7.1–12.4)	-	0.709
NSAIDs (including low-dose aspirin), n (%)	18 (42.9%)	10 (23.8%)	2.4 (0.94–6.12)	0.064

* The degree of gastric mucosal atrophy is classified into mild, medium, or severe according to the Takemoto-Kimura classification [13]. Mild: C-1 and C-2, Medium: C-3 and O-1, Severe: O-2 and O-3.

Bold font represents statistical significance.

La, lanthanum; CI, confidence interval; LaC, lanthanum carbonate; Ca, calcium; IP, inorganic phosphorus; TP, total protein; WBC, white blood cell; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitor

Patient characteristics	LaC taking patients, n = 42	Control dialysis patients, n = 42	Odds ratio (95%CI)	p value
PPIs, n (%)	34 (81.0%)	32 (76.2%)	1.32 (0.47–3.79)	0.595
PPIs + histamine H2 antagonists, n (%)	38 (90.5%)	35 (83.3%)	1.9 (0.51–7.05)	0.332
Gastric mucoprotective drugs, n (%)	13 (31.0%)	6 (14.3%)	2.79 (0.91–7.95)	0.068
* The degree of gastric mucosal atrophy is classified into mild, medium, or severe according to the Takemoto-Kimura classification [13]. Mild: C-1 and C-2, Medium: C-3 and O-1, Severe: O-2 and O-3.				
Bold font represents statistical significance.				
La, lanthanum; CI, confidence interval; LaC, lanthanum carbonate; Ca, calcium; IP, inorganic phosphorus; TP, total protein; WBC, white blood cell; NSAID, non-steroidal anti-inflammatory drug; PPI, proton pump inhibitor				

Study 2

The investigation was performed on 51 biopsy specimens from 23 patients with La deposition in the gastric mucosa and the endoscopic images corresponding to each biopsy site. The 27 biopsy specimens with the characteristic whitish gastric mucosa showed La deposition. La deposition was also detected on the 24 biopsy specimens with non-whitish mucosa.

We classified the patients into two groups according to the amount of La deposition in the biopsy specimen. One or two small fragments of La crystalloid were observed in the slice of 31 specimens (Fig. 2a). The other 20 specimens showed more crystalloids (Fig. 2b). In 20 specimens, less than 30% of the lamina propria was occupied by histiocytes (Fig. 3a), and more than 80% of the lamina propria was occupied by histiocytes in 26 specimens (Fig. 3b). The other five specimens showed no histiocyte occupation.

Of 27 biopsy specimens with the specific whitish mucosa, 14 showed a small amount of La deposition while 13 showed a large amount. In contrast, in the 24 specimens with non-whitish mucosa, 17 showed a small amount of La deposition and seven showed a large amount. There was no significant relationship between the endoscopic features and the amount of La deposition ($p = 0.166$, Pearson's chi-square test; Table 3). In the 27 biopsy specimens with specific whitish mucosa, 10 showed mild histiocytic infiltration and 17 showed strong infiltration. In contrast, among the 24 specimens with non-whitish mucosa, five showed no histiocytic infiltration, 10 showed mild infiltration, and nine showed strong infiltration. There

was a significant relationship between the endoscopic features and the degree of histiocytic infiltration (p = 0.026, Pearson's chi-square test; p = 0.013, Cochran-Armitage trend test; Table 4).

Table 3
The relationship between the endoscopic features and the amount of La deposition

La crystalloid materials	small amount	large amount
Specific whitish mucosa + 27 biopsy specimens	14 (51.9%)	13 (48.2%)
Specific whitish mucosa – 24 biopsy specimens	17 (70.8%)	7 (29.2%)
Data are presented as number (percentage).		
Pearson's chi-square test: p = 0.166		

Table 4
The relationship between endoscopic features and degree of histiocytic infiltration

histiocytic infiltration	none	mild	severe
Specific whitish mucosa + 27 biopsy specimens	0	10 (37.0%)	17 (63.0%)
Specific whitish mucosa – 24 biopsy specimens	5 (20.8%)	10 (41.7%)	9 (37.5%)
Data are presented as number (percentage).			
Pearson's chi-square test: p = 0.026, Cochran-Armitage trend test: p = 0.013			

Discussion

To the best of our knowledge, this retrospective study is the first to investigate the clinical significance of La deposition in the gastric mucosa (as in Study 1), and the association between endoscopic features and histologic findings (as in Study 2).

In Study 1, the most important finding was that there was no significant difference between patients with and without La deposition in the gastric mucosa in terms of blood test findings. In addition, there was no significant difference between the patients taking LaC and those not taking LaC. These results show that gastric La deposition may not cause clinical problems and LaC might be a safe treatment for hyperphosphatasemia in patients with chronic kidney disease on hemodialysis even if La deposits occur in the gastric mucosa. It was revealed that the total dose of LaC was significantly correlated with La

deposition in the gastric mucosa. Although our result was similar to our previous report, this study differs in that the daily dose of LaC was not related to La deposition [3]. We had predicted that the daily dose might not significantly affect La deposition when the period of LaC administration was longer. In addition, concomitant drugs (PPIs, H2RAs, NSAIDs, and gastric mucoprotective drugs) did not affect the deposition of La in the gastric mucosa. Although Shitomi et al. reported that the histological finding of *Helicobacter pylori* infection might decrease the deposition of La [5], Ban et al. reported that the lesions associated with *Helicobacter pylori* infection, such as intestinal metaplasia, regenerative changes, and foveolar hyperplasia were more frequently observed in mucosa with large quantities of La deposits [6, 7]. Given that our present study also showed that the range of atrophic changes in the gastric mucosa was not associated with La deposition, *Helicobacter pylori* infection may not be associated with the deposition of La in the gastric mucosa.

Study 2 revealed that the gastric endoscopic findings of La deposition reflected the infiltrated histiocytes, not the La deposition itself. The appearance of histiocytes must depend on the amount of La deposition, which acts as foreign matter to which histiocytes react. We suggest that the whitish lesions of La deposition reflect the existence of macrophages, similar to gastric xanthoma. La deposition was also detected from non-whitish normal mucosa. In summary, the specific features of La deposition reflect the body's normal reaction to foreign bodies. However, the reason the histiocytic infiltration was divided clearly into two groups (less than 30% and more than 80%) is uncertain.

Although gastric La deposition may not cause clinical problems and LaC might be safe for hyperphosphatasemia in patients with chronic kidney disease on hemodialysis, it has been reported that several side effects of gastrointestinal symptoms have been associated with LaC, such as nausea, vomiting, and constipation [9]. These symptoms might be caused by gastrointestinal motility dysfunction. As it has been reported that functional dyspepsia is caused, in part, by *Helicobacter pylori* infection [10], the gastrointestinal symptoms in hemodialysis patients taking LaC might be caused by La deposition even if there is no organic lesion. Although it has been reported in a pharmacological safety study of LaC that it has no significant effects on gastric emptying and intestinal transit time at high doses [11, 12], an association between La deposition in the gastric mucosa has been loosely associated with gastric peristalsis. Because the present study did not investigate the relationship between the gastrointestinal symptoms and motility function, we should plan the next study to investigate gastric motility function in hemodialysis patients undergoing LaC treatment.

The present study had some limitations. First, the sample size was small, as this was a single-center retrospective study. In addition, there was some missing blood test data. Second, there may have been patients with gastric La deposition in the La-negative group in Study 1 because Study 2 revealed that La deposition was sometimes detected on biopsy specimens from normal gastric mucosa. Third, although La is often deposited at the duodenal mucosa [8], we did not evaluate duodenal La deposition in the present study because of the low number of subjects or biopsy specimens. There might be a relationship between duodenal La deposition and some background factors.

In conclusion, we demonstrated that La deposition in the gastric mucosa depended on the total dose of LaC and was not affected by any background factors. The specific endoscopic features of La deposition are associated with the infiltration of histiocytes, which represents the body's normal response to foreign bodies.

Declarations

Acknowledgements

The authors would like to thank all participants in our study. We would like to thank Editage (www.editage.jp) for English language editing.

Disclosure statement

The authors declare no conflict of interests for this article.

References

1. Annual Dialysis Data Report 2016, JSDT Renal Data Registry
Masakane I, Taniguchi M, Nakai S, et al. Annual Dialysis Data Report 2016, JSDT Renal Data Registry. *Ren Replace Ther.* 2018;4:45.
2. Makino M, Kawaguchi K, Shimojo H, et al. Extensive lanthanum deposition in the gastric mucosa: the first histopathological report. *Pathol Int.* 2015;65:33–7.
3. Hattori K, Maeda T, Nishida S, et al. Correlation of lanthanum dosage with lanthanum deposition in the gastroduodenal mucosa of dialysis patients. *Pathol Int.* 2017;67:447–52.
4. Ozaki H, Ota K, Kojima Y, et al. Small Intestinal Lesions in Patients with Anemia on Hemodialysis: A Two-Center, Cross-Sectional Study [published online ahead of print, 2019 Aug 16]. *Digestion.* 2019;1–7.
5. Shitomi Y, Nishida H, Kusaba T, et al. Gastric lanthanosis (lanthanum deposition) in dialysis patients treated with lanthanum carbonate. *Pathol Int.* 2017;67(8):389–97.
6. Ban S, Suzuki S, Kubota K, et al. Gastric mucosal status susceptible to lanthanum deposition in patients treated with dialysis and lanthanum carbonate. *Ann Diagn Pathol.* 2017;26:6–9.
7. Iwamuro M, Urata H, Okada H. Relationship between gastric lanthanum deposition and *Helicobacter pylori*-associated chronic gastritis: is it rare or common? *Pathol Int.* 2017;67:649.
8. Iwamuro M, Tanaka T, Urata H, et al. Lanthanum phosphate deposition in the duodenum. *Gastrointest Endosc.* 2017;85:1103–4.
9. Murakami N, Yoshioka M, Iwamuro M, et al. Clinical Characteristics of Seven Patients with Lanthanum Phosphate Deposition in the Stomach. *Intern Med.* 2017;56:2089–95.

10. Moayyedi P. The health economics of *Helicobacter pylori* infection. *Best Pract Res Clin Gastroenterol.* 2007;21:347–61.
11. Damment SJ, Pennick M. Clinical pharmacokinetics of the phosphate binder lanthanum carbonate. *Clin Pharmacokinet.* 2008;47:553–63.
12. Hutchison AJ, Wilson RJ, Garafola S, et al. Lanthanum carbonate: safety data after 10 years. *Nephrology (Carlton).* 2016;21:987–94.
13. Kimura K, Takemoto T. An endoscopic recognition of the atrophic border and its significance in chronic gastritis. *Endoscopy.* 1969;1:87–97.

Figures




Figure 1

Specific endoscopic feature (red arrows) of Ladeposition in the gastric mucosa at the endoscopic biopsy sites a: Whitish spot. b:Annular whitish mucosa. c: Diffuse whitish mucosa




Figure 2

Histologic features of La crystalloid (hematoxylin-eosin stain) The bar shows 100 μ m. a: Whole specimen from the gastric mucosa with very small fragments of Lacrystalloid materials (yellow arrows). There are one or two small fragments of La crystalloid in the slice of the specimen. b: Whole specimen from the gastric mucosa with more Lacrystalloid materials (yellow arrows).




Figure 3

Histologic features of histiocytes aggregating (hematoxylin-eosin stain) The bar shows 100 μ m. a. Less than 30% of the lamina propria is occupied by histiocytes aggregating in the gastric mucosa (yellow arrows). a'. The enlarged view of the red square in Figure 3A b. More than 80% of the lamina propria is occupied by histiocytes aggregating in the gastric mucosa b'. The enlarged view of the blue square in Figure 3B.