

SUPPLEMENTARY TABLES

Supplementary Table S1. Details of treatment protocols of adjuvant SCT and SCT-CRT

Clinical trial	Number of patients	Adjuvant therapy setting	Regimens	Protocols
Oettle et al [12]	179	SCT	Gemcitabine	Six cycles of gemcitabine (1000 mg/m ²) on days 1, 8, and 15 every 4 weeks.
Neoptolemos et al [13]	551	SCT	5-FU/folinic acid	Six cycles of folinic acid (20 mg/m ²) and 5-FU (425 mg/m ²) for 5 consecutive days every 4 weeks.
Uesaka et al [19]	192	SCT	S-1	Four cycles of S-1 (40 mg, 50 mg, or 60 mg) orally administered twice a day for 4 weeks followed by a 2-week rest, every 6 weeks.
APCT trial [NCT01964430]	ongoing	SCT	Gemcitabine/nab-paclitaxel	Six cycles of nab-paclitaxel (125 mg/m ²) and Gemcitabine (1000 mg/m ²) on days 1, 8, and 15 every 4 weeks.
Laethem et al [10]	45	SCT-CRT	Gemcitabine + CRT	Two cycles of gemcitabine (1000 mg/m ²) on days 1, 8, and 15 every 4 weeks + CRT (50.4 Gy/28 Fx) with weekly gemcitabine (300 mg/m ²) for 5 to 6 weeks.
Phase II Trial [NCT01396681]	57	SCT-CRT	Gemcitabine/cisplatin + CRT + gemcitabine	Two cycles of Gemcitabine (1200 mg/m ²) on days 1 and 8 and cisplatin (60 mg/m ²) on day 1 every 3 weeks + CRT (50.4 Gy/28 Fx) with weekly gemcitabine (300 mg/m ²) for 5 to 6 weeks + (after 4-6 weeks of rest) four cycles of gemcitabine (1200 mg/m ²) on days 1 and 8 every 3 weeks
Neoptolemos et al [21]	72	SCT-CRT	CRT + 5-FU/folinic acid	Two cycles of CRT (20 Gy/10 Fx) with 5-FU (500 mg/m ²) on days 1-3 over 2 weeks and a 2-week rest + six cycles of folinic acid (20 mg/m ²) and 5-FU (425 mg/m ²) for 5 consecutive days every 4 weeks.

SCT, systemic chemotherapy; SCT-CRT, systemic chemotherapy and chemoradiation therapy