

Systemic inflammation and chronic kidney disease in a patient due to the RNASEH2B defect

Tingyan He (✉ hetingyan2017@outlook.com)

Shenzhen Children's Hospital

Yu Xia

Shenzhen Children's Hospital

Jun Yang

Shenzhen Children's Hospital

Research article

Keywords: Auto-inflammation, Autoimmunity, Aicardi-Goutieres syndrome, Chronic kidney injury, RNASEH2B

DOI: <https://doi.org/10.21203/rs.3.rs-33775/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: Aicardi-Goutières (AGS) is a rare immune dysregulated disease due to mutations in *TREX1*, *RNASEH2A*, *RNASEH2B*, *RNASEH2C*, *SAMHD1*, *ADAR1*, or *IFIH1*. Clinical features include basal ganglia calcifications, white matter abnormalities, and cerebral atrophy. Severe systemic inflammation and chronic kidney disease (CKD) are extremely rare in AGS. Herein, we report a patient presenting with systemic inflammation and CKD to broaden the clinical phenotype spectrum of the *RNASEH2B* defect.

Methods: All testing and molecular genetic analysis were performed after obtaining the informed consent of the parents. Demographic, clinical, and laboratory findings were abstracted from outpatient and inpatient encounters. Cerebral magnetic resonance imaging (MRI), computed tomography (CT) scans, and renal biopsy histopathology reports were reviewed and summarized. Whole exome sequencing (WES) was performed on peripheral blood cells. After exposure to cGAMP in vitro for 24 hours, mRNA expression of twelve IFN-stimulated cytokine genes in PBMCs was assessed. Serum cytokine levels were detected by Milliplex.

Results: A 11-year-old girl presented with recurrent aseptic fever, arthritis, chilblains, failure to thrive, mild hearing loss, and neurological manifestations. Laboratory and immunologic findings demonstrated lymphopenia, low complement levels, positive autoantibodies, elevated levels of acute-phase reactants and inflammatory cytokines. Cerebral imaging showed cerebral atrophy, white matter abnormalities, and intracranial calcification. Renal biopsy showed glomerular sclerosis in three of fourteen glomeruli, infiltration of lymphocytes and other mononuclear cells. WES revealed a homozygous and heterozygous mutations in *RNASEH2B*. Over-expression of IFN-stimulated cytokine genes was observed, including *IFI44*, *IFI27*, *IFIT1*, *IFIT2*, *IFIT3*, *ISG15*, *OAS1*, and *SIGLEC1*.

Conclusions: To date, only two cases with AGS have been reported to have renal disease. Here, we describe a patient with both homozygous and heterozygous variants in *RNASEH2B*, presenting with neurological manifestations, persistently systemic autoinflammation, and CKD. CKD has never been reported in patients with AGS due to the *RNASEH2B* defect.

Background

Aicardi-Goutières (AGS) is a rare immune dysregulated disease due to mutations in *TREX1*, *RNASEH2A*, *RNASEH2B*, *RNASEH2C*, *SAMHD1*, *ADAR1* or *IFIH1*, characterized by encephalopathy, dystonia, basal ganglia calcifications, white matter abnormalities, and cerebral atrophy [1, 2]. Although most patients experienced severe neurological dysfunction within the first year of life, some patients presented with later onset of this disease with mild neurological manifestations and normal intellectual function. Systemic inflammation is not typically persistent. Renal dysfunction has been rarely described in AGS [2]. Here, we report a patient with both homozygous and heterozygous mutations in *RNASEH2B*, presenting with later onset recurrent sterile fever, arthritis, chilblains, failure to thrive, mild hearing loss, and

neurological manifestations, which may broaden the clinical phenotype spectrum of the *RNASEH2B* defect.

Materials And Methods

Subjects

This study was approved by the Ethics Committee of Shenzhen Children's hospital. All human subjects (or their guardians) provided written informed consent. Clinical data of a patient with both homozygous and heterozygous variants in *RNASEH2B* was collected. Fifteen healthy volunteers were included as healthy controls (HCs). Venous blood (3mL) was collected from each study subject.

Whole exome sequencing (WES)

Genomic DNA was extracted from peripheral blood cells isolated from the patient and her parent. The exonic regions and flanking splicing or intronic junctions of the whole genome were captured and sequenced using an Illumina HiSeq 2000 sequencer conducted by MyGenostics (Beijing, China). The FASTQ files were mapped to the human reference genome (hg19). The functional effects of variants were predicted using three algorithms (PolyPhen-2, SIFT, and MutationTaster), and amino acid conservation among species was analyzed. Sanger sequencing was used to confirm pathogenic variants. The primers used to target human *RNASEH2B* included (forward: CAGGGATTTGAAGCTCTTTGG) and (reverse: TAGTGCTCTGTCCTGCACTGG).

Cell culture

Peripheral blood mononuclear cells (PBMCs) were isolated by Ficoll-Paque PLUS (GE Healthcare) gradient density centrifugation and ACK lysis (Quality Biological). PBMCs were resuspended in complete RPMI (cRPMI) medium (Gibco, USA) containing 10% fetal bovine serum (BI, Israel), 2mM glutamine, and penicillin-streptomycin (100U/mL each; Sigma-Aldrich, USA). Cells at 1×10^6 /mL were exposed to cyclic guanosine monophosphate-adenosine monophosphate (cGAMP, CST#35573) at the concentration of 10 ug/ml.

Real-time PCR

After exposure to cGAMP in vitro for 24 hours, mRNA expression of twelve IFN-stimulated cytokine genes in PBMCs was assessed. Total RNA was extracted from PBMCs isolated from the patient and five HCs by RNA isolation kit (DP424, TIANGEN). cDNA was derived following the GoScript Reverse Transcription System kit (A5001, Promega). Quantitative reverse transcription PCR analysis was performed with the GoTaq qPCR Master Mix (A6002, Promega). Primers for PCR included were described in the supplementary material.

Quantification of cytokine levels

Plasma samples were isolated from the patient and fifteen HCs. **Cerebrospinal fluid** (CSF) sample was collected from the patient. Blood samples were collected in vacutainers containing sodium heparin. Plasma cytokine analyses were determined on a bead-based immunoassay (Milliplex, HCYTOMAG-60K, Millipore, USA) according to the manufacturer's protocol.

Statistical analysis

Data were analyzed using an unpaired two-tailed Student t-test. All statistical analyses were conducted in GraphPad Prism 7 software (GraphPad Software, Inc., San Diego, CA).

Results

Clinical manifestations

The patient presented with recurrent fever, arthritis, movement limitation, and growth retardation at the age of 11 years. At the age of 2 years, she began to suffer from recurrent aseptic fever with an intermittent resolution by traditional Chinese medicine. At the age of 5 years, she began to present with arthritis accompanied by mild hearing loss. She was born to a non-consanguineous healthy parent. At birth, her weight was 3 Kg, crown to heel length was 49 cm, and head circumference was 34 cm. She had standard motor and language development. Manifestations of failure to thrive had been significant since she was three years old. Physical examination revealed short stature with 106 cm top(<-3SD)(Fig.1B3), **macrocephaly** with 54 cm head width, chilblains on elbows and lower limbs (Fig.1B2), swelling and deformation of inter-phalangeal and knee joints (Fig.1B1). Her Intelligence Quotient (IQ) test value was 108. Her EPQJ, CBCL, Conners, and HAMA scale tests did not demonstrate any social and psychological problems. Knee magnetic resonance imaging (MRI) revealed a thickness of the synovial capsule without invasive bone destruction(Fig.1B4). Cerebral MRI showed cerebral atrophy and white matter abnormalities(Fig.1B6). Intracranial calcification was further identified at the basal ganglia and cerebellum by CT scanning (Fig.1B5 and Fig.1B7). Laboratory findings revealed hyper-inflammation and chronic kidney disease (Fig.1C, Fig.2B, and Fig.2F). Screening tests for fungal, bacteria, and Mycobacterium tuberculosis infection were all negative. Pathology of the renal biopsy showed glomerular sclerosis in three of fourteen glomeruli, a mild proliferation of mesangial cells without deposits of any amyloid, immunoglobulin or immune complex, expansion of the tubular lumen, partial tubular atrophy, mild tubular fibrosis, infiltration of lymphocytes and other mononuclear cells (Fig.1C). Granule degeneration and calcium deposition were visible in renal tubules. Austin score index for the evaluation of activity and chronicity was two and three points, respectively.

Abnormality in clinical and immunologic phenotype

Analysis of peripheral blood leukocyte revealed persistent lymphopenia(Fig.2A). Except for rheumatoid factor (RF) and anti-cyclic peptide containing citrulline (anti-CCP), other auto-antibodies for mixed connective tissue disease were all negative, including anti-nuclear antibody (ANA), anti-neutrophil cytoplasmic antibody (ANCA), anti-SSA, anti-SSB, anti-dsDNA, anti-thyroglobulin, anti-thyroperoxidase,

and anti-TSH receptor antibodies. Other abnormal clinical and immunologic phenotypes included intermediate elevation of IgM and IgA levels (Fig.2C) and mild reduction of C3 and C4 levels (Fig.2D).

Both homozygous and heterozygous variants in *RNASEH2B*

Whole exon sequencing revealed three variants in the *RNASEH2B* gene (OMIM#610181). There was a single nucleotide homozygous variant, c.859G>T, p.A287S (Fig.1A). Predicted values of SIFT, PolyPhen_2, Mutation Taster, and GERP++ were 0.235, 0.721, 1, and 6.06, suggesting tolerated, possibly damaging, and disease-causing effects, respectively. Both parents carried a heterozygous mutation at the same locus. Another single-nucleotide heterozygous variant, c.269C>T, p.P90L, was identified (Fig.1A). The predicted values of SIFT, PolyPhen_2, Mutation Taster, and GERP++ were 0.002, 0.988, 1, and 4.69, suggesting damaging, probably damaging, and disease-causing effects, respectively. This heterozygous variation was further confirmed by Sanger in both her parents. Both variations were in the conserved domains. Pathogenic variants were not identified in other genes related to autoinflammation, autoimmunity, or inherited renal disorders (Supplemental Table).

Over-expression of IFN-stimulated cytokine genes

After exposure to cGAMP in vitro for 24 hours, mRNA expression of IFN-stimulated cytokine genes in PBMCs was detected by real-time PCR. In contrast to five healthy controls, over-expression of IFN-stimulated cytokine genes was observed in the patient, including IFI44, IFI27, IFIT1, IFIT2, IFIT3, ISG15, OAS1, and SIGLEC1. Normal mRNA expressions were found in IFN β 1 and IRF9 (Fig.3B).

Elevations in inflammatory cytokine levels

Compared to fifteen age-matched healthy controls, plasma cytokine levels were significantly elevated, including interleukin (IL)-1 β , IL-6, tumor necrosis factor- α (TNF α), interferon- γ (IFN- γ), IFN- α , IL-4, IL-10, IL-12, IL-17A and IP-10 (Fig.3A and Table 1). IFN- α level in CSF was very low.

Treatment and outcome

A one-year course of growth hormone showed no response to improve her short stature. She had received long-term treatment of ibuprofen, methotrexate, folic acid, and prednisone for more than five years. Aseptic fever relapsed intermittently. Tocilizumab was started for the high dose dependence of glucocorticoids and elevated pro-inflammatory cytokine levels. Following a 48-week course of tocilizumab, the prednisone dose was gradually reduced to 0.2–0.3mg/Kg.d with partial improvement of some abnormal laboratory findings (Fig.2B and Fig.2F). However, urine β 2 microglobulin level was persistently elevated markedly (Fig.2E). Tocilizumab was discontinued. She began to receive tofacitinib (5mg, twice every day) for the over-expression of IFN-stimulated cytokine genes. The 48-week course of tofacitinib led to partial response, including increased lymphocytes, C3 and C4 levels, reduced levels of urine β 2 microglobulin, C-reactive protein (CRP), and pro-inflammatory cytokines (Fig.2A and Fig.2D). However, chronic systemic inflammation was not completely controlled since the levels of inflammatory cytokines, erythrocyte sedimentation rate (ESR), and serum amyloid A (SAA) were still increased.

Discussion

Biallelic mutations of *RNASEH2B* are most common in AGS. While three allelic variants in *RNASEH2B* have been identified in this patient. The population frequency of the variant c.859G>T, p.A287S in East Asian is 0.02, with two homozygotes demonstrated in ExAC Browser. This variation can cause reduced enzyme activity of RNase H2 and lower stability of the RNase H2 complexes, increasing susceptibility to systemic lupus erythematosus (SLE) [3]. The heterozygous variant c.269C>T, p.P90L is highly conserved with extremely low population frequencies and no homozygotes in ExAC Browser. Its clinical significance remains uncertain in ClinVar. The tri-allelic mutations in *RNASEH2B* may cause a synergistic pathogenic effect since neither heterozygous nor homozygous variants alone can account for her skin and neurological manifestations.

This patient has demonstrated a later onset of AGS with average intelligence, presenting with chilblains, cerebral atrophy, white matter abnormalities, intracranial calcification, and over-expression of Interferon-stimulated genes. Besides, this patient has persistent systemic inflammation and chronic renal dysfunction, which are uncommon in AGS (Table 2). Mixed connective tissue diseases have been excluded by the systemic evaluation. Systemic juvenile idiopathic arthritis (SoJIA), and later onset chronic infantile neurologic, cutaneous, and arthritis (CINCA) syndrome were once suspected. Different from the clinical manifestations of this patient, chilblains and intracranial calcification are not present in SoJIA or CINCA; leukocytosis, destructive arthritis, or macrophage activation syndrome (MAS) are noted in SoJIA [4-6]; visual impairment, sensor neural deafness or progressive chronic meningitis have been commonly reported in CINCA [5]. Chronic kidney disease due to amyloidosis has been rarely reported in SoJIA, which is common in CINCA (Table 2) [8].

Renal involvement has been described in a case with a gain-of-function mutation in *IFIH1* [9]. Renal dysfunction caused by thrombotic microangiopathy has been reported in a case with C-terminal frame-shift mutation in *TREX1* [10]. Renal biopsy in our patient revealed glomerular sclerosis and tubular injury without amyloidosis. *RNASEH2B* is moderately expressed in the kidney. Pathogenic variations in *RNASEH2B* might impair the normal function of kidney directly, or secondary to chronic inflammation. Human IFN- α is filtrated by the kidney, primarily reabsorbed, most probably catabolized within the tubular epithelium, and excreted in negligible amounts with the urine [11]. A fairly high IFN- α level within the tubular epithelium due to a persistently elevated IFN- α level in plasma might amplify the activation of the interferon pathway, leading to the infiltration of lymphocytes and mononuclear cells, and local chronic inflammation. Further investigations will help to explore the distinct pathogenesis underlying chronic renal dysfunction in the *RNASEH2B* defect.

IL-6 is one of the downstream effector cytokines in the IFN signaling pathway. IL-6 blockade has good efficacy in a patient with a cerebral vasculopathy due to a homozygous *SAMHD1* mutation[12]. Tocilizumab has partial efficacy in this patient, leading to a reduction of acute-phase reactants. However, it has failed to improve the chronic renal tubular disease. Further clinical trials are required to clarify the efficacy of tocilizumab in AGS.

IFN- α and IFN- β act on type I receptors (IFNAR1/2) to activate the Janus kinase (JAK)-signal transducers and activators of the transcription (STAT) pathway. JAK inhibitors have good efficacy in patients with some type I interferonopathies, including STING-associated vasculopathy, infantile-onset (SAVI), and proteasome-associated autoinflammatory syndrome (PRAAS) [13-16]. Sustained elevated IFN- α and IFN- β levels are common in AGS. JAK inhibitors can theoretically help to reduce the autoinflammation in AGS. Ruxolitinib has reduced neuroinflammation in a patient with a heterozygous mutation in *IFIH1* [17]. Baricitinib could alleviate chilblain lesions in a patient with AGS5 [18]. Tofacitinib ameliorated aortic valve calcification in a patient with Singleton-Merten syndrome (SMS) [19]. Ruxolitinib led to an improvement of psychomotor delay with a reduction in dystonic movements in two patients with AGS2 [20]. However, ruxolitinib failed to prevent the onset of clinical signs in a patient with *RNASEH2B* mutation [21]. Tofacitinib demonstrated a partial response in this patient, failing to ameliorate autoinflammation and chronic kidney disease completely. Therefore, based on limited case reports, the efficacy of JAK inhibitors in AGS remains uncertain. The currently ongoing trial conducted at the Children's Hospital of Philadelphia (ClinicalTrials.gov number, NCT03921554) will help to explore the efficacy and safety of baricitinib in AGS and AGS-related interferonopathies.

Conclusions

we have described a patient with both homozygous and heterozygous variants in *RNASEH2B*, revealing a possible synergistic pathogenic effect among variants in the same gene. Her systemic autoinflammation and chronic kidney disease will expand the clinical phenotype spectrum of this syndrome. The pathogenesis underlying chronic renal dysfunction in this patient remains poorly understood. The efficacy of tocilizumab and JAK inhibitors in AGS remains uncertain, and further clinical researches are needed.

Abbreviations

AGS: Aicardi-Goutières; WES: Whole exome sequencing; PBMCs: Peripheral blood mononuclear cells; cGAMP: cyclic guanosine monophosphate-adenosine monophosphate; IQ: Intelligence Quotient; MRI: magnetic resonance imaging; RF: rheumatoid factor; ANA: anti-nuclear antibody; ANCA: anti-neutrophil cytoplasmic antibody; TNF α : tumor necrosis factor- α ; IFN- γ : interferon- γ ; CRP: C-reactive protein; ESR: erythrocyte sedimentation rate; SAA: serum amyloid A; SLE: systemic lupus erythematosus; SoJIA: Systemic juvenile idiopathic arthritis; CINCA: onset chronic infantile neurologic, cutaneous, and arthritis syndrome; MAS: macrophage activation syndrome; JAK: Janus kinase; SAVI: STING-associated vasculopathy, infantile-onset; PRAAS: proteasome-associated autoinflammatory syndrome; SMS: Singleton-Merten syndrome.

Declarations

Acknowledgments

The authors wish to thank all the patients, their families, and healthy controls for the participation.

Funding

This work was supported by the Sanming Project of Medicine in Shenzhen (SZSM201812002), Science and Technology Planning Project of Shenzhen Municipality (JCY20170303155201082), and Shenzhen Key Medical Discipline Construction Fund (SZGSP012).

Authors' Contribution

Tingyan He performed the main experiments, analyzed the data, and drafted the manuscript. Yu Xia collected clinical data from the patient. Jun Yang reviewed the manuscript. All authors read and approved the final manuscript.

Availability of data and materials

Clinical datasets were collected from medical records of the participated patient in Shenzhen Children's hospital.

Ethics approval and consent to participate

All participated family members were enrolled upon approval of the ethics committee of Shenzhen Children's hospital and written consent of all the families.

Consent for publication

Written consent for publication of this anonymous information was obtained from the patient's parents.

Conflict of Interests

All authors declare no conflict of interest.

References

1. Cuadrado E, Michailidou I, van Bodegraven EJ, Jansen MH, Sluijs JA, Geerts D, et al. Phenotypic variation in Aicardi-Goutieres syndrome explained by cell-specific IFN-stimulated gene response and cytokine release. *J IMMUNOL*. 2015-04-15;194(8):3623-33.
2. Crow YJ, Chase DS, Lowenstein SJ, Szykiewicz M, Forte GM, Gornall HL, et al. Characterization of human disease phenotypes associated with mutations in TREX1, RNASEH2A, RNASEH2B, RNASEH2C, SAMHD1, ADAR, and IFIH1. *AM J MED GENET A*. 2015-02-01;167A(2):296-312.
3. Claudia Günther, Barbara Kind, Martin A.M. Reijns, Nicole Berndt, Manuel Martinez-Bueno, Christine Wolf, et al. Defective removal of ribonucleotides from DNA promotes systemic autoimmunity. *J Clin Invest*. 2015;125(1):413–424.

4. Hinze CH, Holzinger D, Lainka E, Haas JP, Speth F, Kallinich T, et al. Practice and consensus-based strategies in diagnosing and managing systemic juvenile idiopathic arthritis in Germany. *Pediatr Rheumatol Online J*. 2018-01-22;16(1):7.
5. Barut K, Adrovic A, Sahin S, Tarcin G, Tahaoglu G, Koker O, et al. Prognosis, complications and treatment response in systemic juvenile idiopathic arthritis patients: A single-center experience. *INT J RHEUM DIS*. 2019-09-01;22(9):1661-9.
6. Yasin S, Schulert GS. Systemic juvenile idiopathic arthritis and macrophage activation syndrome: update on pathogenesis and treatment. *CURR OPIN RHEUMATOL*. 2018-09-01;30(5):514-20.
7. Goldbach-Mansky R. Current status of understanding the pathogenesis and management of patients with NOMID/CINCA. *CURR RHEUMATOL REP*. 2011-04-01;13(2):123-31.
8. Finetti M, Omenetti A, Federici S, Caorsi R, Gattorno M. Chronic Infantile Neurological Cutaneous and Articular (CINCA) syndrome: a review. *ORPHANET J RARE DIS*. 2016-12-07;11(1):167.
9. Zheng S, Lee PY, Wang J, Wang S, Huang Q, Huang Y, et al. Interstitial Lung Disease and Psoriasis in a Child With Aicardi-Goutières Syndrome. *Front Immunol*. 2020 May 20;11:985.
10. Komaki R, Ueda T, Tsuji Y, Miyawaki T, Kusuhara S, Hara S, **et al**. Retinal vasculopathy with cerebral leukoencephalopathy carrying TREX1 mutation diagnosed by the intracranial calcification: a case report. *Rinsho Shinkeigaku*. 2018 Feb 28;58(2):111-117.
11. Bocci V, Pacini A, Muscettola M, Paulesu L, Pessina GP, Santiano M, et al. Renal filtration, absorption and catabolism of human alpha interferon. *J Interferon Res*. 1981-01-19;1(3):347-52.
12. Henrickson M, Wang H. Tocilizumab reverses cerebral vasculopathy in a patient with homozygous SAMHD1 mutation. *CLIN RHEUMATOL*. 2017-06-01;36(6):1445-51.
13. Tonduti D, Fazzi E, Badolato R, Orcesi S. Novel and emerging treatments for Aicardi-Goutieres syndrome. *Expert Rev Clin Immunol*. 2020-02-01;16(2):189-98.
14. de Jesus AA, Brehm A, VanTries R, Pillet P, Parentelli AS, Montealegre SG, et al. Novel proteasome assembly chaperone mutations in PSMG2/PAC2 cause the autoinflammatory interferonopathy CANDLE/PRAAS4. *J Allergy Clin Immunol*. 2019-05-01;143(5):1939-43.
15. Melki I, Rose Y, Ugenti C, Van Eyck L, Fremont ML, Kitabayashi N, et al. Disease-associated mutations identify a novel region in human STING necessary for the control of type I interferon signaling. *J Allergy Clin Immunol*. 2017-08-01;140(2):543-52.
16. Sanchez G, Reinhardt A, Ramsey S, Wittkowski H, Hashkes PJ, Berkun Y, et al. JAK1/2 inhibition with baricitinib in the treatment of autoinflammatory interferonopathies. *J CLIN INVEST*. 2018-07-02;128(7):3041-52.
17. Kothur K, Bandodkar S, Chu S, Wienholt L, Johnson A, Barclay P, et al. An open-label trial of JAK 1/2 blockade in progressive IFIH1-associated neuroinflammation. *NEUROLOGY*. 2018-02-06;90(6):289-91.
18. Meesilpavikkai K, Dik WA, Schrijver B, van Helden-Meeuwsen CG, Versnel MA, van Hagen PM, et al. Efficacy of Baricitinib in the Treatment of Chilblains Associated With Aicardi-Goutieres Syndrome, a Type I Interferonopathy. *ARTHRITIS RHEUMATOL*. 2019-05-01;71(5):829-31.

19. Parra-Izquierdo I, Castanos-Mollor I, Lopez J, Gomez C, San RJ, Sanchez CM, et al. Calcification Induced by Type I Interferon in Human Aortic Valve Interstitial Cells Is Larger in Males and Blunted by a Janus Kinase Inhibitor. *Arterioscler Thromb Vasc Biol.* 2018-09-01;38(9):2148-59.
20. Tungler V, Konig N, Gunther C, Engel K, Fiehn C, Smitka M, et al. Response to: 'JAK inhibition in STING-associated interferonopathy' by Crow et al. *ANN RHEUM DIS.* 2016-12-01;75(12):e76.
21. Crow YJ, Shetty J, Livingston JH. Treatments in Aicardi-Goutieres syndrome. *DEV MED CHILD NEUROL.* 2020-01-01;62(1):42-7.

Tables

Due to technical limitations, Tables 1-2 are provided in the Supplementary Files section.

Figures




Figure 1

Clinical features and genetic analysis. A. Pedigree and a homozygous and heterozygous variants in RNASEH2B. B. Physical examinations and imaging findings showing deformed inter-phalangeal joints (B1), chilblains (B2), short stature (B3), synovial thickening (B4), cerebral atrophy and white matter

abnormalities (B6), calcifications in basal ganglia and cerebellum (B5 and B7). C. Renal biopsy findings(10x20) showing glomerular sclerosis, a mild proliferation of mesangial cells, infiltration of lymphocytes and other mononuclear cells.




Figure 2

Abnormalities in laboratory findings. A. lymphopenia. B. Elevated CRP, ESR, and SAA levels. C. Immunoglobulin levels showing intermediate elevated IgM and IgA levels. D. Reduced C3 and C4 levels. E. Persistently elevated levels of urine beta2 microglobulin. F. Intermediate mild abnormalities in Cr and BUN. The normal reference ranges are as follows: lymphocytes (800~4000 cells/ul); CRP 0~10 mg/L; ESR (0~20 mm/h); SAA (0~6 mg/L); IgG (5.28~21 g/L); IgM (0.48~2.26 g/L); IgA (0.44~3.99 g/L); C3 (0.7~2.06 g/L); C4 (0.11~0.61 g/L); urine beta2 microglobulin (0~0.3 mg/L); GFR (80~120 ml/min). Single arrow and double arrows labeled the time when tocilizumab was started and discontinued, respectively. The asterisk labeled the time when tofacitinib was started.




Figure 3

Abnormalities in plasma cytokines and IFN-stimulated cytokine genes. A. Significantly elevated levels of all plasma cytokines in the patient. B. Over-expression of IFN-stimulated cytokine genes in the patient, including IFI44, IFI27, IFIT1, IFIT2, IFIT3, ISG15, OAS1, and SIGLEC1. Normal mRNA expressions were observed in IFN β 1 and IRF9.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTableOtherVariants.xlsx](#)
- [Table1Otherlaboratoryfindings.xlsx](#)
- [Table2.xlsx](#)