

# A modified single-armed microsurgical vasoepididymostomy for epididymal obstructive azoospermia: intraoperative choice and postoperative consideration

**Nachuan Liu**

Shanghai Jiaotong University First People's Hospital

**Peng Li**

Shanghai Jiaotong University First People's Hospital

**Erlei Zhi**

Shanghai Jiaotong University First People's Hospital

**Chencheng Yao**

Shanghai Jiaotong University First People's Hospital

**Chao Yang**

Shanghai Jiaotong University First People's Hospital

**Liangyu Zhao**

Shanghai Jiaotong University First People's Hospital

**Ruhui Tian**

Shanghai Jiaotong University First People's Hospital

**Huixing Chen**

Shanghai Jiaotong University First People's Hospital Department of Orthopaedics

**Yuhua Huang**

Shanghai Jiaotong University First People's Hospital

**Yuexin Yu**

Anhui Medical University

**Zheng Li** (✉ [lizhengboshi@sjtu.edu.cn](mailto:lizhengboshi@sjtu.edu.cn))

Shanghai Jiaotong University First People's Hospital

---

## Research article

**Keywords:** obstructive azoospermia, male infertility, vasoepididymostomy, patency, pregnancy

**Posted Date:** January 20th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.12933/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on August 12th, 2020. See the published version at <https://doi.org/10.1186/s12894-020-00692-5>.

# Abstract

**Background:** To evaluate the clinical outcomes and the duration required for the sperm to return to the ejaculate after a modified single-armed 2-suture longitudinal intussusception vasoepididymostomy (SA-LIVE).

**Methods:** From March 2015 to December 2018, 134 patients with epididymal obstruction azoospermia underwent the modified single-armed vasoepididymostomy at Shanghai General Hospital. The outcomes and clinical findings were documented and evaluated. The mean follow-up period was 17 (range: 3–36) months.

**Results:** Patency was assessed by the return of sperm in the ejaculate. The overall patency rate was 55.2%, and the patency rates were 58.9%, 40.7%, 36.4%, and 58.9% for bilateral surgery, unilateral surgery, proximal anastomosis, and distal anastomosis, respectively. The average time to achieve patency was  $4.11 \pm 2.74$  months. In the first six months, 87.8% (65/74) patency patients reported sperm in the ejaculate. The overall pregnancy rate was 40.9% (29/66) at the follow-up of 3–36 months, and the natural pregnancy rate was 30.3% (20/66). The natural pregnancy rate was 32.1% post-bilateral surgery and 33.3% for the site of distal anastomosis; surprisingly, it was 0% for the site of proximal anastomosis.

**Conclusion:** Modified SA-LIVE is safe and may achieve favorable patency and pregnancy rates. When double-armed sutures are not accessible, single-armed may be preferable. The expected patency time was within one year. Moreover, because of the low natural pregnancy rate for proximal anastomosis, sperm banking is preferred to SA-LIVE.

## Background

Azoospermia affects 1% of the general population and 10–15% of infertile men [1]. Obstructive azoospermia, primarily caused by epididymal obstruction, is diagnosed in approximately 40% of azoospermic men [2] [3]. Microsurgical vasoepididymostomy (MVE) has been established as a more cost-effective alternative for men with obstructive azoospermia than direct assisted reproductive techniques (ART) [4]. MVE necessitates superior surgical skills and meticulous surgical technique. Cornell et al. first reported that the single-armed 2-suture longitudinal intussusception vasoepididymostomy (SA-LIVE) was similar to that of the double-armed procedure in an animal study [5]. Zhao et al. reported the modified single-armed VE technique with favorable patency in a human study trial [6]. Previous studies have shown that a high patency rate may be related to factors, such as epididymal fullness, unilateral or bilateral procedure, and site of anastomosis [7]. However, only a few studies have assessed the clinical outcomes and the time of sperm returning to the ejaculate after SA-LIVE. Thus, in the present study, we aimed to evaluate the fertility outcomes of MVE in 134 males with epididymal obstruction and identify the putative predictors for natural pregnancy. Also, novel intraoperative procedures and decision-making post-operation were explored.

# Methods

## Subjects and laboratory examination

From March 2015 to December 2018, 158 patients who suffered from epididymal obstruction azoospermia (EOA) underwent VE surgery in our center (four patients were misdiagnosed with NOA previously, and one patient had VE in another hospital two years ago). However, during the follow-up, we collected the data from the semen analysis of 134 patients from our center or telephonic follow-up. In this cohort, 80/134 (55.9%) of the patients had a previous history of urological or genital infection before the study, and the etiology of the other 54 cases remained unknown. None of the patients underwent a vasectomy in this study.

All patients underwent semen analysis at least three times before surgery. No sperm was detected in a centrifugal (1,500 ×g) semen assay, and ejaculate fructose tests were positive. The sex hormone levels, such as follicle-stimulating hormone (FSH) and testosterone (T), in the serum, were within normal limits. Scrotal ultrasonography showed dilation of the epididymal tube if present. These were objective measurements for obstructive azoospermia [3].

## Diagnostic clinical condition and criteria

The diagnostic criteria for inclusion in the study were as described previously [8]: obstruction was suspected when the infertile male had normal ejaculate volume with azoospermia; the physical examination showed non-atrophic testes with normal vas deferens bilaterally, slightly swollen epididymis, and bilateral or unilateral hard epididymal nodules; normal serum total T and FSH levels; the ultrasonography showed the dilation of epididymal tube without dilation of the ejaculatory duct or seminal vesicle. The patients with chromosomal or sex chromosomal abnormalities, history of vasectomy, or whose female partners were reported to be infertile, were excluded from the current study.

## Surgical procedure

1. General preparation: All patients underwent scrotal exploration under general anesthesia. Surgeries were performed by two experienced surgeons. Testis was exposed and biopsy was performed as follows: a touch-prep is made by blotting the cut surface of the testis several times on a glass slide and adding a drop of human tubal fluid and a coverslip. The samples were examined under high power using a light microscope with phase contrast that revealed the presence of sperm with tails and facilitated the assessment of motility. Subsequently, normal spermatogenesis was confirmed [9, 10]. All patients provided written informed consent before the study.
2. Microsurgical preparation: A 24-gauge angiocatheter sheath was used to cannulate the lumen of the vas deferens, and Trypan blue was injected to judge the patency of the seminal vesicle side. A sufficient length was freed to allow the most proximal convoluted portion of the vas to be later brought to the lateral aspect against the epididymis without tension.

3. Microsurgical SA-LIVE approach: A Carl Zeiss S88 operating microscope (Carl Zeiss Shanghai Co., Ltd, Shanghai, China) was used to perform the microsurgical procedure at a magnification of 8–15X. A dilative epididymal tubule was selected and dissected under microscopic guidance. Subsequently, anastomosis was performed using the modified SA-LIVE technique (Fig. 1). Two single-armed 10–0 nylon sutures (Ethicon W2790, length 200 mm, circle 3/8) were prepared for the intussusception sutures. The first suture was placed in an outside-in fashion through the mucosal layer of the vas deferens at point a1. Then, the needle was used to pierce the lateral aspect of the epididymal tubule and placed longitudinally. The second 10-0 single-armed proline suture was placed identically through point b1 on the vas deferens, parallel to the first suture on the contralateral side of the epididymal tubule. The two needles were used to parallelly pierce the epididymal tubule laterally within and longitudinally on the outside of the tubule. The epididymal tubule was opened longitudinally between the two sutures using a micro knife, and the exuded epididymal fluid was examined for sperm[10]. If the sperm or sperm fragments were present, the needles were pulled through the wall of the epididymis and placed in an inside-out manner through the full layer of the vas at positions a2 and b2. Finally, all the sutures were tied together (a1 to a2 and b1 to b2), and the epididymal tubule was gently intussuscepted into the lumen of the vas deferens. Then, the epididymal tunic was secured to the vassal muscle and adventitia with an 8-0 nylon suture. The epididymal and testicular sperms were cryopreserved during VE as a backup of the procedure.

### **Postoperative management**

Most of the patients were discharged home on the 1<sup>st</sup> day after the surgery. They were advised to refrain from any heavy lifting and sports activity for eight weeks post-surgery, and asexual abstinence was instructed for four weeks. Semen analyses were initiated at four weeks after surgery, and every month after that, until pregnancy was achieved. Patency was defined as the sperm concentration > 1 million/ml in at least one postoperative ejaculate sample. Pregnancy was defined as the establishment of a fetal heartbeat. The follow-up information was obtained by clinic visits and telephonic contacts. Patients without a postoperative semen analysis or patients lost to follow-up were excluded from the report, and the follow-up time was at least three months.

### **Statistical analysis**

The patency and pregnancy rates of each group (unilateral and bilateral MVE group and the proximal and distal anastomosis group) were calculated. The chi-square test was used for all analyses.  $P < 0.05$  indicated statistical significance.

## **Results**

From March 2015 to December 2018, a total of 158 patients diagnosed with an epididymal obstruction underwent modified SA-LIVE in our center; among these, 134 were followed up in a prospective study. Reportedly, 80 patients had a history of epididymitis or orchitis; full epididymis could be palpated easily

with inflated nodes. The diagnostic ultrasonography findings also proved the existence of inflammation in the scrotum based on the thin net-like ectasia of the epididymal tube with the inner diameter up to 0.4 mm [11]. The mean age of the included patients was  $32.1 \pm 6.7$  (range: 23–50) years, and the mean follow-up was for  $17 \pm 3.3$  (range: 3–36) months. The mean FSH and T levels were  $4.3 \pm 2.4$  mIU/mL and  $5.0 \pm 2.9$  ng/mL, respectively. The mean testicular size was  $15.4 \pm 3.3$  cm<sup>3</sup>, as measured by ultrasonography (Table 1).

A total of 132 patients were subjected to sperm banking during the operation in the case of future intracytoplasmic sperm injection (ICSI). Sperm was present in the ejaculate of 74/134 (55.2%) patients after the surgery, and the average time to achieve patency was  $4.11 \pm 2.74$  months. In the first six months post-treatment, 87.8% (65/74) patients were reported to have sperm in their ejaculate. Only 12.2% (9/74) patients claimed to achieve patency in the following six months (Figure 2). The variables related to patency for follow-up are listed in Table 2. The patency rate was 58.9%, 40.7%, 36.4%, and 58.9% for bilateral surgery, unilateral surgery, proximal anastomosis, and distal anastomosis, respectively. In this cohort, 30.3% (20/66) patients were reported to have spontaneous pregnancy after VE. The variables related to natural pregnancy are listed in Table 3. The natural pregnancy rate was 30.5% for bilateral surgery, 20% for unilateral surgery, and 33.3% for the site of distal anastomosis, but 0% for the proximal anastomosis. Furthermore, nine couples got pregnant by ICSI (seven used frozen sperm collected during the operation, and two used fresh sperm from the ejaculate). The total and natural pregnancy rate was 43.9% (29/66) and 30.3% (20/66), respectively, at the follow-up of 3–36 months. The mean time to achieve natural pregnancy was  $11.05 \pm 5.75$  (range: 3–24) Months.

## Discussion

Approximately 10–15% of infertile men suffer from azoospermia, while about 40% suffer from obstructive azoospermia (OA). This obstruction may be attributed to bilateral occlusion at any point in the reproductive ductal system, which comprises of the efferent duct, epididymis, vas deferens, and the ejaculatory ducts. Unlike the USA, EOA is rarely caused by vasectomy rather by infection in China [12, 13]. The microsurgical anastomosis, including microsurgical vasovasostomy (VV), cross vasovasostomy (CVV), and vasoepididymostomy (VE), is considered as the most successful measure for a reversal [14]. Herein, we discussed the modified single-armed 2-suture longitudinal intussusception vasoepididymostomy. The data were similar to the previously reported patency rate of 52–92% and the pregnancy rate of 11–56% [15, 16]. In the present cohort, we evaluated the pre-, intra-, and postoperative variables of individual patients that might affect the outcomes of the modified SA-LIVE.

The VE with the double-armed suture is the standard golden management for the EOA, and LIVE simplifies the anastomosis and improves the outcomes [7, 17]. However, in China, these specialized double-armed sutures for male infertility microsurgery are challenging, while access to single-armed microsurgical sutures is easy [18]. Furthermore, the cost of a single-armed sutures was cheaper than that of the double-armed microsurgical sutures. Therefore, SA-LIVE should be designated as an effective alternative when double-armed sutures are not available. When the needles are in the tubule, the two

sutures in a lower position can avoid the crossing of the sutures [6]. The two knots of the suture are left outside during the procedure, which might decrease the possibility of fibrosis and anastomotic stricture. However, the single-armed suture is time-consuming as compared to the double-armed suture, which was placed inside-out on the vas deferens to avoid back-walling the tubular lumen. Since the single-armed suture was used in this study, the needle passed through the inferior points of the vasal mucosal layer in an outside-in manner through the epididymal tubule, and finally through the superior points of the vasal mucosal layer in an inside-out fashion, which was double time-consuming than the double-armed suture, thereby increasing the risks of surgery. Also, we had to dilate the vasal lumen sufficiently wide to pass the needle through the lumen with the aid of a microneedle holder to avoid back-walling during the suture placement. Supposedly, these procedures must be completed before the presence of the sperm in the epididymal fluid [19]. However, if no sperm was detected, the procedure was repeated 2–4 times, which would significantly increase the time required for the surgery and the fatigue of the surgeon.

In this cohort, the patency and the pregnancy rates were similar to those reported previously [12]. However, no statistically significant associations were found between the patency rate and various predictors, such as bilateral or unilateral anastomosis and anastomotic site ( $P > 0.05$ ; Tables 2 and 3). Moreover, the natural pregnancy rate was 0% for the proximal anastomosis group, while it was 33.3% for the distal group, which prompted us to investigate whether patients can benefit from proximal anastomosis. Firstly, the luminal diameters of epididymal tubules in the caput were significantly smaller than those in the corpus and caudal, and hence, the modified SA-LIVE on the distal epididymis was more accessible than that on the proximal epididymis. Moreover, spermatozoa could become fully motile as well as recognize and fertilize an egg within the epididymis, which might improve the pregnancy rate. As the sample size was not sufficient for further investigation, the reason for the failure of the technique is yet to be determined.

Although there have been many clinical studies on VE, none proposed a specific follow-up time frame on SA-LIVE. The current study revealed that 87.8% of the patients achieved patency within six months post-operation, and none achieved patency since then, suggesting that SA-LIVE on patency can be followed up to 12 (average:  $4.11 \pm 2.74$ ) months. The patency rate for the first six months was 48.5% (65/134) and 11.4% (9/69) for the next six months. Previous data on the mean time to achieve patency after vasovasostomy and vasoepididymostomy ranged from 1.7–4.3 and 2.8–6.6 months, respectively [12], which is in agreement with the current results. This finding could be valuable for the clinicians and researchers to predict the outcome of patients undergoing SA-LIVE as well as the precise time to transfer to ART.

Notably, four patients were diagnosed as non-obstructive azoospermia in other hospitals previously according to the negative results of the testis histopathology; however, we found motile sperm through the routine testicular biopsy. Thus, the scheduled operation of microdissection was transferred to VE, and 3/4 patients achieved patency afterward. This suggested that the testis biopsy during the operation is essential [20].

Also, one patient, who had undergone VE in another hospital previously was subjected to VE operation due to the previous failure. Consequently, sperm appeared in the ejaculate, and finally, this couple achieved a spontaneous pregnancy. Thus, we concluded that in addition to ICSI, a second VE surgery might be a possibility for those patients who failed the first operation.

Interestingly, VE is an effective treatment for azoospermia patients with epididymal obstruction and previous failure to achieve pregnancy by sperm retrieval with ICSI [21]. Three patients diagnosed with OA chose to use the sperm from testicular puncture for ICSI in another hospital but failed possibly due to miscarriage, maturation arrest in utero, or the failure of embryo transfer [21]. After undertaking SA-LIVE in our hospital, all the patients achieved patency: one had natural pregnancy, one underwent IVF after the surgery, and one is still trying to get a natural pregnancy. Furthermore, MEV has significant advantages, such as cost-efficiency, spontaneous pregnancy possibilities, and decreasing the potential risks of congenital disability as compared to ICSI. Thus, microsurgical reconstruction is an effective treatment and should be the first choice for epididymal obstruction patients whose female partners have normal fertility features. Thus, EOA patients with prior failure of ICSI could consider taking LIVE to get pregnant.

This study revealed that if sperm did not show up in the ejaculate after one-year post-operation, ART might be the remedy for such patients. Thus, intra- or postoperative sperm cryopreservation would be helpful. Intraoperative sperm cryopreservation could avoid the additional surgeries for sperm retrieval in case of failure of microsurgery. Postoperative sperm cryopreservation allows sperm from the ejaculate to be used for ART in the event of late failure. In the current cohort, we found out that one patient who achieved recanalization of the vas experienced a recurrence of obstruction after six months at the rate of 0.07%, which was lower than that reported previously (1–50%) [12, 22]. Sperm cryopreservation was used for a total of 132 patients. Of these, 29 patients who were not successful by vas-mediated patency achieved pregnancy through ICSI by the sperm cryopreserved intraoperatively, and nine successful patients chose to use the sperm from ejaculate to have babies through ART. The pregnancy rate was improved from 30.3% to 43.9%. With the development of the ART technique in the recent decades, the influence of whether the sperm on the next generation was unclear irrespective of its origin from testis or post-epididymis. Thus, taken together, we advocate intra- or postoperative sperm cryopreservation for all OA patients.

## Conclusion

Modified SA-LIVE is preferred when double-armed sutures are not accessible. The expected patency time is within one-year, and ART may be the remedial choice for patients. Assisted with intra-operative sperm cryopreservation, the pregnancy outcomes were improved in patients. The low natural pregnancy rate for the proximal anastomosis suggested that when the obstruction occurs proximally, sperm banking is preferred for future ICSI.

## Abbreviations

SA-LIVE: modified single-armed 2-suture longitudinal intussusception vasoepididymostomy

VE: Microsurgical vasoepididymostomy

ART: assisted reproductive techniques

FSH: follicle-stimulating hormone

T: testosterone

VV: vasovasostomy

ICSI: intracytoplasmic sperm injection

OA: obstructive azoospermia

EOA: epididymal obstructive azoospermia

IVF: in-vitro fertilization

## Declarations

**Ethics approval and consent to participate:** All procedures performed

in studies involving human participants were in accordance

with the ethical standards of Institutional Ethical Review Committee of

Shanghai general hospital, Shanghai Jiaotong University School of

Medicine and with the 1964 Helsinki declaration and its later amendments

or comparable ethical standards. Informed consent was obtained

from all individual participants included in the study.

**Consent for publication:** Not applicable.

**Availability of data and materials:** The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

**Competing interests:** The authors declare that they have no competing interests.

**Funding:** This work was supported by grants from National Natural Science Foundation of China (81701524), and Strategic Priority Research Program of the Chinese Academy of Sciences

(XDA16020701), and Medical-Engineering(Science) cross-Research Fund of Shanghai Jiao Tong University(YG2017ZD04), and National Natural Science Foundation of China (81871215). The funding body had no control over the study design, or the collection and analysis of the data.

**Authors' contributions:** Peng Li, Erlei Zhi, Chencheng Yao, Chao Yang, Liangyu Zhao, Ruhui Tian, Huixing Chen, and Yuhua Huang performed the surgery, Yuexin Yu and Nachuan Liu collected the data. Nachuan Liu and Peng Li analyzed the data and wrote the manuscript. Zheng Li conceived the project, designed the analysis, and supervised the entire study. All of the authors read and approved the final manuscript.

**Acknowledgements:** Not applicable.

## References

1. De Kretser DM, Baker HW: **Infertility in men: recent advances and continuing controversies.** *J Clin Endocrinol Metab* 1999, **84**(10):3443-3450.
2. Jarow JP, Espeland MA, Lipshultz LI: **Evaluation of the azoospermic patient.** *J Urol* 1989, **142**(1):62-65.
3. Jungwirth A, Diemer T, Kopa Z, Krausz C, Tournaye H, Kelly B, Pal R: **EAU Guidelines on Male Infertility 2017;** 2017.
4. Kolettis PN, Thomas AJ, Jr.: **Vasoepididymostomy for vasectomy reversal: a critical assessment in the era of intracytoplasmic sperm injection.** *J Urol* 1997, **158**(2):467-470.
5. Monoski MA, Schiff J, Li PS, Chan PT, Goldstein M: **Innovative single-armed suture technique for microsurgical vasoepididymostomy.** *Urology* 2007, **69**(4):800-804.
6. Zhao L, Deng CH, Sun XZ, Chen Y, Wang WW, Zhao LY, Zeng LY, Tu XA: **A modified single-armed technique for microsurgical vasoepididymostomy.** *Asian J Androl* 2013, **15**(1):79-82.
7. Hong K, Zhao LM, Xu SX, Tang WH, Mao JM, Liu DF, Jiang H, Ma LL, Qiao J: **Multiple factors affecting surgical outcomes and patency rates in use of single-armed two-suture microsurgical vasoepididymostomy: a single surgeon's experience with 81 patients.** *Asian J Androl* 2016, **18**(1):129-133.
8. Chan PT, Brandell RA, Goldstein M: **Prospective analysis of outcomes after microsurgical intussusception vasoepididymostomy.** *BJU Int* 2005, **96**(4):598-601.
9. Dabaja AA, Schlegel PN: **Microdissection testicular sperm extraction: an update.** *Asian journal of andrology* 2013, **15**(1):35.
10. Wein AJ, Kavoussi LR, Novick AC, Partin AW, Peters CA, Ramchandani P: **Campbell-Walsh urology tenth edition.** In.: Elsevier, Philadelphia; 2011.
11. Cornud F, Belin X, Delafontaine D, Amar T, H el enon O, Moreau J: **Imaging of obstructive azoospermia.** *European radiology* 1997, **7**(7):1079-1085.
12. Farber NJ, Flannigan R, Li P, Li PS, Goldstein M: **The Kinetics of Sperm Return and Late Failure Following Vasovasostomy or Vasoepididymostomy: A Systematic Review.** *J Urol* 2019, **201**(2):241-

250.

13. Chen XF, Chen B, Liu W, Huang YP, Wang HX, Huang YR, Ping P: **Microsurgical vasoepididymostomy for patients with infectious obstructive azoospermia: cause, outcome, and associated factors.** *Asian J Androl* 2016, **18**(5):759-762.
14. Liang ZY, Zhang FB, Li LJ, Li JP, Wu JG, Chen C, Zhu YM: **Clinical application of cross microsurgical vasovasostomy in scrotum for atypical obstructive azoospermia.** *J Zhejiang Univ Sci B* 2019, **20**(3):282-286.
15. Zhang GX, Bai WJ, Xu KX, Wang XF, Zhu JC: **Clinical observation of loupe-assisted intussusception vasoepididymostomy in the treatment of obstructive azoospermia (analysis of 49 case reports).** *Asian J Androl* 2009, **11**(2):193-199.
16. Chan PT: **The evolution and refinement of vasoepididymostomy techniques.** *Asian J Androl* 2013, **15**(1):49-55.
17. Binsaleh S: **Two-suture single-armed longitudinal intussusception vasoepididymostomy for obstructive azoospermia: report of patients characteristics and outcome.** *Int Urol Nephrol* 2014, **46**(12):2271-2277.
18. Zhao L, Tu XA, Zhuang JT, Chen Y, Wang WW, Zeng LY, Deng CH: **Retrospective analysis of early outcomes after a single-armed suture technique for microsurgical intussusception vasoepididymostomy.** *Andrology* 2015, **3**(6):1150-1153.
19. Lyu KL, Zhuang JT, Li PS, Gao Y, Zhao L, Zhang YD, Zhou MK, Yu JW, Feng X, Sun XZ *et al*: **A novel experience of deferential vessel-sparing microsurgical vasoepididymostomy.** *Asian J Androl* 2018, **20**(6):576-580.
20. Hayden RP, Li PS, Goldstein M: **Microsurgical vasectomy reversal: contemporary techniques, intraoperative decision making, and surgical training for the next generation.** *Fertil Steril* 2019, **111**(3):444-453.
21. Peng J, Yuan Y, Zhang Z, Cui W, Song W, Gao B: **Microsurgical vasoepididymostomy is an effective treatment for azoospermic patients with epididymal obstruction and prior failure to achieve pregnancy by sperm retrieval with intracytoplasmic sperm injection.** *Hum Reprod* 2014, **29**(1):1-7.
22. Alom M, Ziegelmann M, Savage J, Miest T, Kohler TS, Trost L: **Office-based andrology and male infertility procedures-a cost-effective alternative.** *Transl Androl Urol* 2017, **6**(4):761-772.

## Tables

Table 1: Preoperative characteristics and intraoperative choices in all 134 patients

| Items                                    | Value                   |
|--|-------------------------|
| Age (year), mean                         |                         |
| Patients                                 | 32.1±6.7 (ranged 23-50) |
| Female partners                          | 27.2±3.7 (ranged 20-43) |
| Serum FSH (mIU/mL), mean                 | 4.3±2.4                 |
| Serum total testosterone (ng/mL), mean   | 5.0±2.9                 |
| Testicular size (cm <sup>3</sup> ), mean | 15.4±3.3                |
| Surgery, n (%)                           |                         |
| Bilateral                                | 107(79.9%)              |
| Unilateral                               | 27(20.1%)               |
| Anastomotic site, n (%)                  |                         |
| Proximal                                 | 22(16.4%)               |
| Distal                                   | 112(83.6%)              |


Table 2. The patency rate stratified by two variables for 134 patients.

| Variable         | Patency (n)   | P Value |
|------------------|---------------|---------|
| Surgery          |               | 0.090   |
| Bilateral        | 63/107(58.9%) |         |
| Unilateral       | 11/27(40.7%)  |         |
| Anastomosis site |               | 0.052   |
| Proximal         | 8/22(36.4%)   |         |
| Distal           | 66/112(58.9%) |         |

Table 3. The natural pregnancy rate stratified by two variables for 134 patients.


| Variable         | Natural pregnancy (n) | P Value |
|------------------|-----------------------|---------|
| Surgery          |                       | 0.442   |
| Bilateral        | 18/56(32.1%)          |         |
| Unilateral       | 2/10(20.0%)           |         |
| Anastomosis site |                       | 0.090   |
| Proximal         | 0/6(0%)               |         |
| Distal           | 20/60(33.3%)          |         |

## Figures



**Figure 1**

Placement of sutures in modified SA-LIVE. The needles were sequentially placed outside-in (a1 and b1) through the mucosal layer of the vas deferens, parallelly through the epididymal tubule, then placed inside-out (a2 and b2) through the mucosal layer of the vas deferens.



**Figure 2**

Time of sperm returning to the ejaculate. The column height of the histogram represents the patients counts within various periods to achieve patency.