

The situation of elderly with cognitive impairment living at home during lockdown in the Corona-pandemic in Germany

Jochen René Thyrian (✉ rene.thyrian@dzne.de)

German Center for Neurodegenerative Diseases (DZNE) <https://orcid.org/0000-0001-6889-693X>

Friederike Kracht

German Center for Neurodegenerative Diseases (DZNE)

Angela Nikelski

Evangelisches Klinikum Bethel, Division of Geriatric Psychiatry

Melanie Boekholt

German Center for Neurodegenerative Diseases (DZNE)

Fanny Schumacher-Schönert

German Center for Neurodegenerative Diseases (DZNE)

Anika Rädke

German Center for Neurodegenerative Diseases (DZNE)

Bernhard Michalowsky

German Center for Neurodegenerative Diseases (DZNE)

Horst Christian Vollmar

Ruhr-University Bochum (RUB)

Wolfgang Hoffmann

German Center for Neurodegenerative Diseases (DZNE)

Francisca S. Rodriguez

German Center for Neurodegenerative Diseases (DZNE)

Stefan H. Kreisel

Evangelisches Klinikum Bethel, Division of Geriatric Psychiatry

Research Article

Keywords: COVID19, pandemic, Corona, cognitive impairment, lockdown, elderly, Germany, loneliness, health care

Posted Date: July 7th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-38185/v1>

Abstract

Background: The outbreak of the Corona virus is a challenge for health care systems worldwide. Elderly people with comorbidities are at a higher risk and the different levels of government in Germany reacted jointly to the challenge with social distancing and a lockdown. Little empirical evidence exists about the psychosocial situation of elderly people during this time. Thus, the aim of this study is to analyze a) knowledge about, and feelings related to the Corona-pandemic. Describe b) loneliness, depression and anxiety, and, c) the perceived, immediate impact of the lockdown on frequency of social contacts and quality of health care provision of people with cognitive impairment during social distancing and lockdown in the primary care system and daily living at home in Germany.

Methods: This analysis is based on data from standardized telephone interviews in a convenience sample of n=141 people with known cognitive impairment in the primary care setting. Participants were recruited from currently running interventional trials. Therefore, data on e.g. cognitive and psychological status prior to the pandemic was available. Attitudes, knowledge about and perceived personal impact of the pandemic, social support, loneliness, anxiety, depression, frequency of social activities due to the pandemic and perceived impact of the pandemic on health care services were assessed during the time of lockdown.

Results: The vast majority of participants consider themselves sufficiently informed about Corona (85%) and most think that the measures taken are appropriate (64%). A total of 11% shows one main symptom of a depression according to DSM-V. The frequency of depressive symptoms has not increased between the time before pandemic and lockdown in almost all participants. The sample shows minimal (65.0%) or low levels of anxiety (25%). The prevalence of loneliness is 10%. On average seven activities have decreased in frequency due to the pandemic. Social activities related to meeting people, dancing or birthday visits have decreased significantly. Talking with friends by phone and activities like gardening have increased. Utilization of health care services like day clinics, relief services and prescribed therapies are reported to have worsened due to the pandemic. Also visits to general practitioners decreased.

Discussion: The study shows a limited impact of the pandemic on psychological variables including depression, anxiety and loneliness in the short-term. People are well informed but not especially concerned about themselves or others. There is a decrease in social activities as expected, in line with the restrictions imposed. The impact on health care provision is prominent. There is a need for longitudinal studies to assess longer-term effects of the pandemic and social distancing on elderly people with cognitive impairment living at home and their caregivers. There is also a need for qualitative, in-depth studies to further interpret the results. In general, elderly people in the setting under examination and with the limitations of this specific study seem to compensate restrictions during the time of lockdown in Germany rather well.

Background

The first cases of a novel atypical pneumonia – the infection later being named COVID-19 and shown to be caused by the corona virus strain SARS-CoV-2 – were reported in the city of Wuhan, China in December 2019. The WHO announced it officially as pandemic on March 11, 2020, after the virus had rapidly spread to numerous countries worldwide. The first symptomatic case of a coronavirus SARS-CoV-2 infection in Germany was reported at the end of January 2020. The most common symptoms are dry cough, fever, fatigue, chest tightness, diarrhea, and dizziness [1] and the most severe symptoms include respiratory distress and septic shock. Early reports had suggested that about 40% of the patients progress to severe illness within a time period of five days [2].

Scientific evidence has rapidly evolved during the last few months and is continuing to do so. However, it was clear from very early on that older adults and those with serious underlying medical conditions are at a higher risk for severe illness from COVID-19. Older people are more likely to develop severe symptoms, fever, and breathing problems than younger people [3]. Comorbidity and lymphocyte count are the strongest predictors of progression to severity [2], with hypertension, diabetes, and higher body mass index as the most frequent comorbidities among patients with severe symptoms [4]. Accordingly, people who need extra precautions are, among others, elderly people with chronic diseases. A clinical review on testing, treatment and prognosis of COVID-19 in the elderly supports the association of age and presence of comorbidities with increased disease severity and mortality [5].

With no causal or symptomatic treatment being available, the aim of all protective measures has been to stop or slow down the pandemic and prevent people from becoming infected. This has been especially challenging as many of those infected remain asymptomatic and all are infectious before symptoms appear. However, public health measures have been associated with the epidemiology of the disease [6]. General measures to encounter the pandemic were introduced in Germany starting March 22nd, 2020 when the federal government in agreement with the governments of all federal states enacted limitations in social contacts (i.e. “social distancing”) throughout the country. Targeted social distancing is the standard practice to influence infection spread, especially for vulnerable groups because spread is strongest through social contact networks [7]. While individual quarantine of infected people is an efficient type of social distancing and useful for many infectious diseases, social distancing on a broader community level aims at achieving minimal interaction between people in the population and is ethically more challenging [8]. Social distancing, also referred to as “physical distancing”, is described by the Center for Disease Control and Prevention (CDC) as “keeping space between yourself and other people outside of your home” <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/social-distancing.html>. The CDC recommends to a) stay at least a certain distance from other people, b) not gather in groups, and c) stay out of crowded places and avoid mass gatherings. Worldwide children were barred from going to schools or preschools, and university classes, religious services, major events in sports, film industry, music festivals and concerts, conferences, and fashion shows were canceled [9]. In line with this strategy, the German government recommended to reduce spatial closeness and decrease individual mobility as much as possible. In public places, a minimal distance between individuals of 1.50m was introduced. Public gatherings were restricted to two people (this rule, however, did not apply to

people residing in the same household, e.g. families with more members could move freely in the public), public and private gatherings were forbidden, but people were generally “asked” to stay at home. In contrast to other countries, there was no strict, legally binding “lock down”. Furthermore, shops selling non-essential products, restaurants, clubs, schools, universities, public buildings, recreational facilities and other institutions of public life were closed. On April 15th some of the restrictions were eased (e.g. selected shops were allowed to open under specific restrictions), but social distancing was recommended to be kept in place until May 6th. Latter measure has now been extended until further notice. Due to the federal nature of state organization in Germany, recommendations and regulations were introduced jointly as a result of an inter-governmental agreement, but nonetheless posed only a non-binding framework for the federal states (the entities legally responsible for pandemic measures). Each federal state was able to develop specific plans, set different deadlines and interpret nationwide recommendations differently. Although the initial consensus was upheld in the early stages of the lockdown, federal diversity has increased lately. Starting May 6th 2020 it was mandated that the nationwide approach in respect to preventive measures would be delegated to a more regional level and on 6th of June more restrictions were loosened. It has been shown that the provisions taken have contributed to the substantial reduction of the growth rate in Germany [10].

As the spread of COVID-19 is a specific challenge for the health care system, several measures have been introduced to deal with the pandemic in addition to the general restriction outlined above. The Robert-Koch-Institute, the German federal government’s central scientific institution for public health, has issued recommendations for hygiene for the public and the health care system in Germany. In this article, we focus on general recommendations that affect health care and might specifically affect the elderly. For example, to be prepared for an increased COVID-19 associated utilization of hospitals, the total number of beds for intensive care were increased, by e.g. postponing elective surgery and planned hospital stays. Social distancing rules were applied to health care institutions and the primary care system. For example: day care clinics were closed, ambulatory service provision decreased, treatments cancelled or restricted in case these could not meet the recommended hygienic requirements (e.g. out-patient physiotherapy).

As of yet the consequences of these measures on medical, psychological or social variables during the pandemic remain unknown. Nonetheless, an impact can be expected. For instance, while social distancing is not meant to imply a reduction in social or emotional support it can easily be misunderstood as “distancing”[11]. Further, social distancing can be associated with a loss of motivation, loss of meaning, loss of self-worth, and subsequent mental health problems [12]. Moreover, the public dissemination of preventive measures can trigger anxieties, lead to discrimination and stigma, and cause misinformation through misunderstanding [13]. People with cognitive impairments might have difficulties to comprehend the imposed restrictions or may misinterpret the protective gear worn by health care professionals as threatening [14].

There is some scientific evidence from recent pandemics (SARS, H1N1, Ebola, MERS; [15]) that highlight the impact of being “quarantined” on psychosocial factors. Among the negative outcomes were: depression, stress, irritability, sleep disorders, anger, etc. The highest prevalences were observed for

irritability and depressive symptoms. In addition, relatives of people being quarantined experienced fear, nervousness, sadness and guilt. Albeit, there also positive outcomes were reported such as happiness and relief. Factors associated with negative outcomes were length of quarantine, frustration, boredom, fear of infection, undersupply of daily goods and information deficit.

A recent rapid literature review of the psychosocial impact of quarantine during pandemics implied extensive negative consequences for mental health [16]. Even though these results may be transportable to the impact of social distancing, there is a significant difference. Quarantine requires being infected or having had contact to a person infected whereas systematic social distancing is a measure with an impact at the population level.

Lack of social support, loneliness, and restricted social contact is associated with higher mortality [17] as well as a higher risk for cardiovascular disease [18] and dementia [19]. Moreover, among cancer patients, those with more social support have better survival chances [20]. This, what is considered to be a stress-buffering effect [21], is likely to be brought about by physiologic pathways related to neuroendocrine or immunologic function [22]. Social support appears to enhance the efficiency of the immune system in dealing with inflammatory processes [23,24]. Furthermore, studies point out that neurons of socially isolated animals are less responsive to stimulation [25], emphasizing the relevance of social contacts for neurodegenerative processes. The health promoting effects of social contacts contrast with the vulnerability of older people to be lonely [26]. There is outspoken concern about the effect of isolating the elderly given the pandemic and beyond [27] and the German Association of Gerontology and Geriatrics has published a statement emphasizing the importance to maintain participation and social inclusion for the elderly during the Corona-pandemic [28].

A description and analysis of the perceived impact of the pandemic on the population at risk is needed to add scientific evidence and to ultimately gear (future) healthcare to provide appropriate measures [29]. Therefore, the aims of this study are to:

- a. Analyze knowledge about, and feelings related to the Corona-pandemic.
- b. Analyze loneliness, depression and anxiety of people with cognitive impairment during lockdown.
- c. Analyze the perceived, immediate impact of the lockdown on frequency of social contacts and quality of health care provision

in a cohort of people with cognitive impairment in the primary care system and living at home in Germany.

Methods

This analysis is based on data from a telephone-based questionnaire and semi-structured interview assessment in a convenience sample of n=141 people with cognitive impairment in the primary care setting in Mecklenburg-Western Pomerania and North-Rhine-Westphalia, respectively. Participants were recruited from currently running interventional trials intersec-CM (Supporting elderly people with cognitive

impairment during and after hospital stays with intersectoral care management) [30] and the DCM:IMPact study (Dementia Care Management: Implementation into different Care Settings), an implementation study of the dementia care management of the DelpHi-MV trial [31-33]. The assessments were conducted during the time of the lockdown during study contacts between 22nd of March and 5th of June 2020, before restrictions were lifted.

Sample

A total of n=115 participants of the intersec-CM trial received the additional assessment. Intersec-CM is a complex, longitudinal, multisite randomized controlled trial. It was designed to treat a hospital-based cohort of people above the age of 70 with an adaption of “Dementia Care Management” (DCM) – a treatment proven to be effective in primary care – to the discharge setting. As part of this, specifically trained study staff develops, implements, and monitors a treatment and care plan, based on comprehensive assessments during the hospital stay, recommendations at discharge and unmet needs identified at home. For the 3 months after discharge study staff coordinates treatment and care in close cooperation with the discharging hospital, treating physician and other care providers. The first patient from a total of n=401 was recruited in November 2018, the last in April 2020. Participants of the ongoing intersec-CM trial were asked to take part in the additional assessment only if they belonged to the original study’s intervention group or were part of the control group after completion of their individual follow-up time period.

A total of n=26 participants were interviewed in the framework of the DCM:IMPact study, a mixed-methods, multi-center, implementation study. DCM:IMPact complements the DelpHi-trial (Dementia: Life- and person-centered help). In this study, the effective [32] and cost-efficient [33] dementia care management is implemented in different care settings, including ambulatory care services, dementia care networks, and hospitals. The study aims to evaluate various care setting for need and implementation barriers for such a model of care, and in how its effects could be optimized. The DCM:IMPact study started in 2019.

Owing to this particular sampling scheme, participants come from two German federal states: Nordrhein-Westfalen and Mecklenburg-Vorpommern.

Data assessment

The benefit of interviewing a convenience sample out of established studies is that baseline sociodemographic characteristics and certain clinical variables are available. Variables that usually cannot be assessed by phone, like cognitive status, are known. The sample is described in detail in table 1. Additionally, study participants have an established relationship with the interviewers. We expect this personal contact to increase the validity. It also allows for more semi-structured qualitative information about the participants’ situation.

From end of March until 6th of June, we conducted a telephone-based questionnaire and semi-structured interview assessment. The questionnaire included variables from previously validated instruments. They cover:

- a. Attitudes, knowledge about and perceived personal impact of the pandemic
- b. social support
- c. loneliness
- d. anxiety
- e. depression
- f. change in the frequency of social activities due to the pandemic
- g. perceived impact of the pandemic on health care related services

Ad a) Participants were asked whether they felt sufficiently informed about Corona (yes/ no) and whether they knew someone infected (yes/ no) and whether they rated the measures against the pandemic as appropriate ("Yes", "No – too strict", "No – not strict enough"). They were asked to rate whether they felt "1 – strongly", "2 – moderately", "3 – not at all" worried because of Corona for themselves; "stressed out", "concerned about their health", "concerned about the health of family and friends", "worried about Corona in general". They were asked to rate their fear to infect family members, whether their everyday life has changed by the pandemic and whether the visiting restrictions for nursing homes do affect their life. The items are listed in table 2.

Ad b) Social support was measured using selected items from the German version of the Resource Generator [34]. The Resource Generator covers a wide variety of social resources that can be divided in four aspects of social capital (capital associated to: prestige and education of the network, political and financial knowledge of the network, special skills of the network, social support of the network). For our analysis and due to time restrictions in the assessment four relevant items from the aspect "instrumental social support" were chosen (i.e. is there anyone, who "supports you when repairing something", "supports you when shopping", "supports you in filling out forms", "supports you with legal and administrative challenges"). The participant was asked to rate these with: "0 – not at all", "1 – rather not", "2 – rather", "3 – exactly". Each item is analyzed separately and for the purpose of analysis, we add the scores of each item to generate a total score of social support, ranging from 0 to 12.

Ad c) Loneliness was measured using the loneliness scale of de Jong Gierveld & van Tilburg [35]. This scale consists of 6 items with statements the participant is asked to rate with "1 - exactly", "2 - rather", "3 - rather not" or "4 - not at all". The scores of each item are summed to a total score, ranging from 6 to 24; the higher the score the the more intense loneliness is present in the participant. This instrument was chosen, because it is used in the longitudinal population-based cohort of the elderly in Germany and reference scores are available for the years 2008 to 2017 [36]. We furthermore measured loneliness with a focus on social relationship using a short loneliness scale developed specifically for use in a telephone survey [37]. Three items ask how often one feels "lacking companionship", "being left out" and "being

isolated". Each item is rated "seldom", "sometimes" or "often". The sum score indicates the level of loneliness, ranging from 3 to 9.

Ad d) Anxiety is measured using the Generalized Anxiety Disorder Scale-7 (*GAD-7*) [38]. It is a screening instrument to diagnose a generalized anxiety disorder according to DSM-V and ICD 10. The psychometric properties are considered appropriate. The German version is validated for the general population [39]. Participants are asked to rate how frequent symptoms of anxiety have occurred during the time-frame of 2 weeks preceding the interview ("0 – not at all", "1 – single days", "2 – more than half of the days", "3 – nearly daily"). The GAD-7 delivers a sum score by adding each item and ranges from 0 to 21. The sum score yields whether anxiety symptoms are minimal (0-4), low (5-9), medium (10-14) or severe (15-21). Furthermore, there are cut-off scores for a generalized anxiety disorder being possible (score >10) or probable (score >15).

Ad e) The two item Patient Health Questionnaire (PHQ-2) was used for screening for depressive symptoms [40,41]. The participant is asked to categorize how often he/ she has experienced "loss of interest or pleasure in activities" and "sadness" during the 2 weeks prior to the assessment. The options are "0 – never", "1 – single days", "2 – more than half of the days", "3 – nearly daily". These two items represent the main indicators for a positive screening of depression according to DSM-V, with one of both being mandatory. Depression was considered possible if one of the two items was scored ≥ 2 . Depression was considered unlikely otherwise. For the sample of participants recruited for the intersec-CM study we included pre-pandemic data and could therefore calculate whether there was a change in frequency of symptoms during the period of lockdown ("more frequent", "equally frequent", "less frequent").

Ad f) To measure change in frequency of (social) activities due to the pandemic we use a generic list of social activities and ask whether the specific activity is carried out "more frequently", "less frequently" or "with no change in frequency" during the time of lockdown as compared to before. If an item does not apply to the participant's situation, it was marked as being "not applicable" (e.g. if a participant never "went dancing" previously, this item was marked as "not-applicable"). The list consists of 24 items (see table 3). Each item is analyzed separately. We furthermore provide three scores per person, which represent the number of activities that have "increased", "decreased", or remain "unchanged" in frequency.

Ad g) To assess the perceived impact of the pandemic on health care related services participants are asked to rate whether the provision and use of the service is "better", "worse" or "unchanged" during the lockdown in comparison to the time before. Items that were irrelevant to a participant's situation were marked "not applicable" (e.g. if a participant never utilized "podiatry" as a service, this item was marked as "not applicable"). The full list is provided in table 4. Each service is analyzed separately. We furthermore provide three scores per person, which represent the number of services where the quality is rated to be "better", "worse" and "unchanged".

Statistical analysis

In this exploratory study we provide descriptive statistics for the variables under analysis.

Results

Sociodemographic and clinical characteristics

Participants in the final sample were on average 81.5 years old and 61.7% were women. The sample includes mainly people with mild dementia (mean MMSE score of 23.1). Around one third of the participants live alone (34.1%). Women, on average, were two years older and were less likely to live alone (27.2% vs 45.1% in males).

Attitudes, knowledge and personal impact of the pandemic

The vast majority of participants report that they feel sufficiently informed about Corona (83.5%) and most think that the measures taken are appropriate (62.1%). Nonetheless, 26.2% think they are inappropriate, but this group divides about half into “too strict” and half into “not strict enough”. Most participants do generally not worry (48.5%) or worry moderately (42.7%) about Corona. In respect to their own risk, most participants feel moderately worried (51.5%) or not worried at all (36.9%). Approximately the same distribution is found for specific worries about one’s own health, family and friends, fear of infection, fear of infecting others, and change of everyday life. The detailed results are shown in table 2.

Loneliness, anxiety, and depression

Analyzing measures of psychological status (i.e. loneliness, anxiety, and depression), the cohort shows sadness “never” in more than half of the people (59.8%) or on only “single days” in another third (36.2%). There is also “never” loss of interest in 51.6% of the sample. A total of $n=15$ (11.4%) show one main symptom of depression according to DSM-V. The cohort shows minimal (65.1%) or low symptoms of anxiety (25.2%). The mean sum score on the GAD-7 is 3.2 (range: 0-18; $SD=3.2$). Possible anxiety disorder (GAD-7 score >10) is found in $n=7$ participants. According to the loneliness-score of De Jong Gierveld 10.2% of our cohort are considered lonely, the mean loneliness scores being 11.1 (range 6-20, $SD=3.2$); participants scored 4.5 on the Hughes scale (range 3-9, $SD=1.8$). Possible anxiety disorder shows a remarkable difference between the sexes: males show to be more anxious (9.8%) than females (2.4%). See table 1 for detailed results.

Social support and social activities

Instrumental social support in this sample is high. The scale ranges from 0 to 3 per dimension and the mean is higher than 2 in all social support dimensions measured. The total sum of social support shows a mean of 9.6 (range 2-12, $SD=2.3$). The participants reported on average 1.3 distinct social activities with an increase in their pursuit during the time of pandemic ($SD=1.3$), on average 6.8 activities were pursued with a decreased frequency ($SD=4.0$) and on average 5.1 ($SD=3.2$) were carried out with the same frequency as before. On average $n=10.0$ activities were indicated as not relevant for the individual participant. Among the activities that have increased in frequency are: “Talk with friends and relatives by

phone" (37.9% of the total sample), "Watch TV" (22.1%), "Spend time outside" (14.3%), "Work in the garden" (9.2%), "Being visited by family" (6.4%), "Knitting" (5.0%) and "Using computer/ tablet" (5.0%). Activities that decreased in frequency are: "Visit birthdays/ festivities" (62.9%), "Visit hairdresser" (62.1%), "Meet friends" (61.4%), "Meet relatives" (61.4%), "Go for coffee" (57.9%), and "Being visited by neighbors" (52.1%). More details are found in table 3 and illustrated in figure 1.

If one limits the analysis only to those activities that were reported to be relevant to each participant (i.e. removing individual "non-applicables") the results are remarkably different. This is visualized in figure 2. Every participant indicated that "Go dancing" had decreased due to the Corona pandemic. More than 70% indicated that "Going for coffee", "Visiting birthdays/ festivities" or "Visiting the hairdresser", "Meeting friends and relatives" and "Visits by friends and neighbors" had decreased due to Corona. On the other hand, the frequency of "Talking with friends and relatives by phone" increased in approximately 40% of the participants and did not decrease relevantly in the others.

Impact on health care related services

Services that were perceived as being worse were: "Podiatry" (21.3%), "Prescribed therapies" (14.2%) and "Relief services" (11.3%). Almost no health-care-related items were perceived as having improved during the pandemic ($m=0.2$, range 0-2, $SD=0.5$). Those that were nonetheless reported to be used/available more often were: "Relief services" (4.3%), "Ambulatory care services" (3.5%), and "Provision of medication" (2.8%). See table 4 for more details.

Again limiting the analysis only to relevant health-related services (i.e. removing individual "non-applicables"), the results change somewhat (see figure 2). Services described as having been utilized less often during the lockdown by more than 30% were "Day clinics", "Relief services", utilization of "Prescribed therapies" and "Podiatry". There is only a small proportion of participants indicating that "Ambulatory care services", "Day clinics" and "Relief services" had improved.

Discussion

Our results reflect the situation of elderly in primary care, living at home during the lockdown imposed on them by the Corona pandemic in Germany.

Knowledge about, and feelings related to the Corona-pandemic

The evidence shows that most elderly perceive their knowledge about the virus as sufficient, indicating that the public information policy was adequate. That is important as research has shown that lack of knowledge in quarantine is a risk factor for later negative health outcomes [15]. In general, the cohort is modestly worried to not worried about themselves, their health or their relatives. Of note: None of those interviewed were infected themselves, only few knew others that had been infected. This fact might partly explain the low level of reported worries. Other explanations are also possible and may reflect aspects of age and life experience. Notes from semi-structured interviews accompanying the questionnaire can be

interpreted in this light. Participants commented the situation as follows: “I am not afraid. I am 93, you can only die once! [Ich habe keine Angst, bin jetzt 93, einmal kann man nur sterben]” or “We have already lived through much worse situations [Wir haben schon Schlimmeres erlebt]”.

Nonetheless, agreement with the measures taken by the government was very high. It is unclear what drives this consensus. On the one hand, the active and transparent health communication adopted by policy makers might have contributed. On the other hand, generation-specific experiences or age-associated attitudes may have been beneficial. One participant explained it saying: “Queen Elizabeth has given courage and hope in her address [Queen Elizabeth hat in ihrer Ansprache Mut und Hoffnung gemacht]”, reflecting the, in the German media, well-publicized speech by Queen Elizabeth II on the April 5th.

Loneliness, depression and anxiety of people with cognitive impairment during lockdown

Participants of our study show levels of loneliness, anxiety and depression that are comparable to people in the general population – during “normal” times. In Germany, the prevalence of loneliness in the general population of 45-84 year olds is 9.2%, in the age group presented here it is between 7.5% and 8.1% [36]. This is comparable to the 10.2% we found in our study. Loneliness being a rather stable construct may explain this result. A short-term isolation or restriction of social contacts with a clear cause and a foreseeable end might not influence the feeling of being lonely too much. It might worsen the state of people who are already lonely, but not make people lonely that were not lonely before.

Remarkably, depression and anxiety scores were lower than would be expected in the general population – outside of the pandemic. The prevalence of any anxiety disorder according to DSM-IV has been estimated to be around 10% in the general population [42]. The prevalences in this study are much lower. The same holds for depression. The German population-based Leila75+ study showed a prevalence of depression in the 75+ year olds of 38.2% [43], the LIFE study with participants of the age from 18-80 a prevalence of 6.4% [44]. In our cohort, we assessed prevalence (11.4%) using a cutoff of 2 or more points on one item of the PHQ-2; this potentially even overestimates the true prevalence of depression in our cohort.

However, these results need interpretation. Prevalence data for Germany is population-based. Our study, on the other hand, is based on a convenience sample of people already participating in intervention trials. As such, a selection bias is possible in that people with a previous diagnosis of depression or anxiety disorders might not have participated in the original study. This notwithstanding, approximately half of the cohort showed single depressive symptoms at baseline. Comparing these to the pandemic results of the two PHQ-2 items “loss of interest or pleasure in activities” and “sadness” (in intersec-CM participants), our results imply that the pandemic did not impact symptoms of depression. (Only one participant showed a higher frequency than at baseline.) In fact, descriptively depressive symptoms before the baseline were shown to be more frequent than during the lockdown. Again, this may not reflect a real shift as all intersec-CM participants were initially recruited during an acute hospital stay. Summarizing, we can

show that depressive symptoms are present in our cohort, but they do not seem to change by the Corona pandemic and the restrictions imposed.

Impact of the lockdown on frequency of social contacts, social activities and quality of health care provision

The level of social isolation in the general population of elderly in Germany has been reported to be around 13%, in the age group of 60+ up to 20% [45]. Due to limitations of the telephone interview setting, we were only able to assess instrumental social support as a dimension of social isolation. On average, our cohort received medium to high levels of instrumental support even under lockdown-conditions and therefore cannot be considered isolated. We asked whether the frequency of pursuit in various social activities had changed due to the pandemic. On average the frequency did not change in six activities, one was pursued more and seven less often. (On average participants signaled that 10 of the 24 activities to choose from did not apply in their individual situation.) We can conclude that the level of social activities in the elderly in our cohort did not change significantly; restrictions in activities that are important to a given individual (especially those related to “staying in contact”) were, in fact, compensated during the lockdown.

Activities pursued less frequently after the lockdown are associated with activities that are difficult to perform under the conditions of social distancing. Participants met with other people less frequently, went out less frequently and visited social gatherings less frequently. However, the pandemic had a “positive effect” by increasing the frequency of talking to friends and relatives by phone, gardening or similar activities with less interpersonal contact. Thus, the elderly seem to find various ways to maintain social relations and stay in contact with their relatives and friends. Due to this, pandemic-related restrictions might not have direct and short-term consequences on measures of loneliness and social isolation. The elderly studied here – notable that they all suffered from impaired cognition– seem to retain sufficient resources to keep themselves active and socially connected, irrespective of the pandemic.

The strongest impact of the restrictions imposed on the elderly in our cohort were those referring to the provision and utilization of health care services. Our participants visited or were visited less by general physicians or other medical providers. They rated the provision of ambulatory services, day clinics and prescribed therapies as worse than before due to the pandemic. Especially the provision of services aiming at relief for caregivers was perceived to be worse. These results need further attention as a decrease in health care services may lead to significant long-term consequences.

We do not have any systematic information about the specific reasons participants rated the change in activity pursuit the way they did. From the semi-structured interviews, however, there is some indication that causes may be inter-individually quite diverse. For example, one participant explained that his prescription for a sleep apnea device was stopped “due to” Corona. Others reported that ambulatory care services reduced their services because of fear of infecting their clients or of becoming infected themselves. This had substantial consequences especially among participants with mobility restrictions

(defined as needing help when shopping or simply aid when walking). A more in-depth qualitative assessment is needed here.

The health of elderly people, who can not utilize health care services for a certain period of time may be put at risk. This might be devastating for the German health care system. A decreased number of patients admitted to the hospital, GPs or specialists within the Corona-pandemic leads to lower income for health care providers in private practice and, in the short-term to lower health care expenditures. However, as a result of the non-use of health care services, there will probably be significantly more serious cases in the clinics in the future and more patients with severe pre-existing conditions. This will likely increase long-term treatment costs and can put considerable strain on the health care system. Only in a few years, the full economic effects and consequences of the pandemic (i.e. high treatment costs and life-threatening diagnoses) can be assessed exactly.

Our data show that more attention must be focused on the needs of the caregivers. It is well documented that the burden of caregivers for people with cognitive impairment and dementia is often high and then is a risk factor for institutionalization [46-48]. Services aiming at relief for caregivers have been established to support caregivers and ease the situation for people living at home. With hindering and/ or closing these services over the pandemic the burden on caregivers will have increased. This is especially the case when caregivers need to compensate for professional services that are restricted or no longer provided during the pandemic. Thus, it is possible that caregiver health outcomes will be detrimentally affected. There is an urgent need to monitor this potential development and improve caregiver support wherever possible to alleviate the consequences.

Limitations

There are clear limitations that restrict the generalizability of these results.

Our results are limited to cognitively impaired people in primary care living at home. The pandemic on i.e mortality of institutionalized elderly has been reported to be severe; the outbreak was associated with a very high mortality rate and high rates of positively tested residents of nursing homes [49]. Isolating nursing home residents in their rooms is associated with morbidity and raises patient safety and staffing issues [50]. The SARS-CoV-2 pandemic could also have lasting psychological impacts on care home staff [50]. It is important that the necessary and appropriate support is provided. Nursing home care itself has been described as being “in crisis” because of the pandemic [51].

Results may look differently in regions with a higher infection rate than in the regions under investigation here. Participants were interviewed in the federal state of Mecklenburg-Western Pomerania, a rural state with the lowest infection rate in Germany, and in North-Rhine-Westphalia, a federal state with high infection rate and a higher population density. Thus, our results are more than regional results. However, samples from areas with high prevalence of Corona infection could be different. Further studies need to be conducted and compared to our results to get a more comprehensive picture of the impact and associated factors.

Our conclusions are based on data from a selective convenience sample from two ongoing intervention trials. The strengths of this sample are, that it was already recruited prior to the pandemic and direct contact was accessible during the pandemic. The main advantage was, that prior information on patient characteristics, cognitive and psychological status was readily available. Furthermore, the participants had previously – due to the ongoing trials and contact therein – established a relationship with the interviewers, which might have increased the validity of the data and might have decreased social desirability in the answers. This is important, because there can be reluctance to report attitudes and to rate services. This is illustrated by a comment of one participant: “Is [this] a check-up call to make sure we stay at home and adhere to the rules? Do you cooperate with the police? [Ist das ein Kontrollanruf, ob wir zu Hause sind und die Maßnahmen einhalten. Arbeiten Sie mit der Polizei zusammen?]”.

To our best knowledge, we are not aware of a comparable sample of people with cognitive impairments living at home examined during lockdown in Germany.

In this study, we focus on a generic list of health care services and a generic list of (social) activities, which makes it difficult to compare across studies. Nonetheless, we deemed the items relevant for this particular population and chose them for inclusion in our questionnaire based on expert opinions and other studies. Interviewing elderly people with cognitive impairments by phone results in limitations. Alertness and attention over time will be more prone to deterioration than in other cohorts. Therefore, we had to choose feasibility over comprehensiveness.

Conclusion

In summary, our study reflects the situation of people with cognitive impairment living at home during social distancing due to the Corona pandemic in Germany. It shows an only small impact of the pandemic and related measures on psychological variables like depression, anxiety and loneliness. People are well informed but not especially concerned about themselves or others. There is a decrease in social activities as expected and as intended by the restrictions imposed on the population. The impact on health care provision is significant. While this is a cross-sectional analysis, there is a need for longitudinal studies to assess mid-term and long-term effects on health-outcomes, especially for caregivers. There is also a need for comparative studies in areas with higher infection rates, in different health care systems and different countries to analyze factors associated with outcome heterogeneity.

Declarations

Trial registration and ethics

Intersec-CM is registered as clinical trial (ClinicalTrials.gov, NCT03359408) and Ethical approval has been obtained from the ethics committee of Greifswald Medical School (registry number: BB 159/17) and the ethics committee of the Chamber of Physicians of Westfalen-Lippe (registry number: 2017–688-b-S).

Ethical approval for the DCM:Impact study was obtained by the ethics committee of the University Medicine Greifswald (BB 01/2019).

Acknowledgment:

We do especially thank Ines Abraham, Stefanie Feldmann, Ulrike Kempe, Agnes Meyer, Andrea Pooch und Esther-Sarah Whittaker for recruiting the participants, conducting the interviews and listening attentive and patiently to the personal stories and worries of the participants. We thank Bernhard Holle for valuable nursing science input.

Funding: The study is funded by the German Federal Ministry of Education and Research (BMBF). The funding codes (FKZ) are 01GL1701A (German Center for Neurodegenerative Diseases / site Rostock/Greifswald, Greifswald); 01GL1701B (Evangelisches Klinikum Bethel, Division of Geriatric Psychiatry, Bielefeld); 01GL1701C (Institute of General Practice and Family Medicine, Faculty of Medicine, Ruhr-University Bochum (RUB)); 01GL1701D (Institute of Community Medicine, Department of Epidemiology and Community Health, Greifswald). The ministry had no role in the design of this study and will not have any role during implementation, analyses, interpretation of the data or publication of results.

Competing interests: The authors declare that they have no competing interests.

Authors' roles: Each author has made substantial contributions to this work. All have approved the submitted version and agreed to be personally accountable for the author's own contributions. All authors ensure that questions related to the accuracy or integrity of any part of the work, even those in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature.

Informed consent: Participants gave written informed consent to participate in the studies.

References

1. Chen Q, Zheng Z, Zhang C, Zhang X, Wu H, et al. (2020) Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. Infection.
2. Ji M, Yuan L, Shen W, Lv J, Li Y, et al. (2020) A predictive model for disease progression in non-severe illness patients with Corona Virus Disease 2019. European Respiratory Journal.
3. Lian J, Jin X, Hao S, Cai H, Zhang S, et al. (2020) Analysis of Epidemiological and Clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan. Clinical Infectious Diseases.
4. Chen Q, Zheng Z, Zhang C, Zhang X, Wu H, et al. (2020) Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. Infection: 1-9.

5. Lithander FE, Neumann S, Tenison E, Lloyd K, Welsh TJ, et al. (2020) COVID-19 in older people: a rapid clinical review. Age and Aging <https://doi.org/10.1093/ageing/afaa093>.
6. Pan A, Liu L, Wang C, Guo H, Hao X, et al. (2020) Association of Public Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. JAMA.
7. Glass RJ, Glass LM, Beyeler WE, Min HJ (2006) Targeted social distancing designs for pandemic influenza. Emerging infectious diseases 12: 1671.
8. Wilder-Smith A, Freedman DO (2020) Isolation, quarantine, social distancing and community containment: pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. Journal of travel medicine 27: taaa020.
9. Sarma R (2020) AN ANTHROPOLOGICAL PERSPECTIVE ON CORONA VIRUS, HEALTH AND SOCIETY. Purakala with ISSN 0971-2143 is an UGC CARE Journal 31: 1285-1292.
10. Schmitt F-J (2020) A simplified model for expected development of the SARS-CoV-2 (Corona) spread in Germany and US after social distancing. arXiv preprint arXiv:200310891.
11. Abel T, McQueen D (2020) The COVID-19 pandemic calls for spatial distancing and social closeness: not for social distancing! International journal of public health 65: 231.
12. Williams SN, Armitage CJ, Tampe T, Dienes K (2020) Public perceptions and experiences of social distancing and social isolation during the COVID-19 pandemic: A UK-based focus group study. medRxiv.
13. Ren S-Y, Gao R-D, Chen Y-L (2020) Fear can be more harmful than the severe acute respiratory syndrome coronavirus 2 in controlling the corona virus disease 2019 epidemic. World journal of clinical cases 8: 652.
14. Halek M, Reuther S, Schmidt J (2020) Herausforderungen für die pflegerische Versorgung in der stationären Altenhilfe: Corona-Pandemie 2020. Mmw Fortschritte Der Medizin 162: 51.
15. Brook SK, Webster RK, Smith LE, Woodland L, Wessely PS, et al. (2020) The psychological impact of quarantine and how to reduce it: rapid review of the evidence. The Lancet 395: P912-920.
16. Röhr S, Müller F, Jung F, Apfelbacher C, Seidler A, et al. (2020) Psychosocial Impact of Quarantine Measures During Serious Coronavirus Outbreaks: A Rapid Review [Psychosoziale Folgen von Quarantänemaßnahmen bei schwerwiegenden Coronavirus-Ausbrüchen: ein Rapid Review]. Psychiatr Prax 47.
17. Berkman LF, Melchior M, Chastang JF, Niedhammer I, Leclerc A, et al. (2004) Social integration and mortality: a prospective study of French employees of Electricity of France-Gas of France: the GAZEL Cohort. Am J Epidemiol 159: 167-174.
18. Barth J, Schneider S, von Kanel R (2010) Lack of social support in the etiology and the prognosis of coronary heart disease: a systematic review and meta-analysis. Psychosom Med 72: 229-238.
19. Kuiper JS, Zuidersma M, Oude Voshaar RC, Zuidema SU, van den Heuvel ER, et al. (2015) Social relationships and risk of dementia: A systematic review and meta-analysis of longitudinal cohort studies. Ageing Res Rev 22: 39-57.

20. Kroenke CH, Kubzansky LD, Schernhammer ES, Holmes MD, Kawachi I (2006) Social networks, social support, and survival after breast cancer diagnosis. *J Clin Oncol* 24: 1105-1111.
21. Wethington E, Kessler RC (1986) Perceived support, received support, and adjustment to stressful life events. *Journal of Health and Social behavior*: 78-89.
22. Berkman LF, Glass T (2000) Social integration, social networks, social support, and health. *Social epidemiology* 1: 137–173.
23. Uchino BN (2006) Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *J Behav Med* 29: 377-387.
24. Scotti M-AL, Carlton ED, Demas GE, Grippo AJ (2015) Social isolation disrupts innate immune responses in both male and female prairie voles and enhances agonistic behavior in female prairie voles (*Microtus ochrogaster*). *Hormones and behavior* 70: 7-13.
25. Sargin D, Oliver DK, Lambe EK (2016) Chronic social isolation reduces 5-HT neuronal activity via upregulated SK3 calcium-activated potassium channels. *Elife* 5.
26. Rodriguez FS, Pabst A, Luck T, König HH, Angermeyer MC, et al. (2018) Social Network Types in Old Age and Incident Dementia. *J Geriatr Psychiatry Neurol* 31: 163-170.
27. Armitage R, Nellums LB (2020) COVID-19 and the consequences of isolating the elderly. *Lancet Public Health* 5: e256.
28. Kessler E-M, Strumpfen S, Kricheldorf C, Franke A, Pantel J, et al. (2020) [Partizipation und soziale Teilhabe älterer Menschen trotz Corona-Pandemie ermöglichen].
29. Nicol GE, Piccirillo JF, Mulsant BH, Lenze EJ (2020) Action at a Distance: Geriatric Research during a Pandemic. *J Am Geriatr Soc* 68: 922-925.
30. Nikelski A, Keller A, Schumacher-Schonert F, Dehl T, Laufer J, et al. (2019) Supporting elderly people with cognitive impairment during and after hospital stays with intersectoral care management: study protocol for a randomized controlled trial. *Trials* 20: 543.
31. Thyrian JR, Fiss T, Dreier A, Bowing G, Angelow A, et al. (2012) Life- and person-centred help in Mecklenburg-Western Pomerania, Germany (DelpHi): study protocol for a randomised controlled trial. *Trials* 13: 56.
32. Thyrian JR, Hertel J, Wucherer D, Eichler T, Michalowsky B, et al. (2017) Effectiveness and Safety of Dementia Care Management in Primary Care: A Randomized Clinical Trial. *JAMA Psychiatry* 74: 996-1004.
33. Michalowsky B, Xie F, Eichler T, Hertel J, Kaczynski A, et al. (2019) Cost-effectiveness of a collaborative dementia care management-Results of a cluster-randomized controlled trial. *Alzheimers Dement* 15: 1296-1308.
34. van der Gaag M, Snijders TAB (2020) The Resource Generator: Social capital quantification with concrete items. *Social Networks* 27: 1-29.
35. De Jong Gierveld J, Van Tilburg T (2006) A 6-Item Scale for Overall, Emotional, and Social Loneliness: Confirmatory Tests on Survey Data. *Research on Aging* 28: 582-598.

36. Huxhold. O, Engstler. H, Hoffmann. E (2019) Entwicklung der Einsamkeit bei Menschen im Alter von 45 bis 84 Jahren im Zeitraum von 2008 bis 2017. DZA Fact Sheet. . Berlin: Deutsches Zentrum für Altersfragen.
37. Hughes ME, Waite LJ, Hawkley LC, Cacioppo JT (2004) A Short Scale for Measuring Loneliness in Large Surveys: Results From Two Population-Based Studies. *Res Aging* 26: 655-672.
38. Spitzer RL, Kroenke K, Williams JB, Lowe B (2006) A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 166: 1092-1097.
39. Lowe B, Decker O, Muller S, Brahler E, Schellberg D, et al. (2008) Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care* 46: 266-274.
40. Spitzer RL, Kroenke K, Williams JB (1999) Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. *JAMA* 282: 1737-1744.
41. Löwe. B, Spitzer. RL, Zipfel. S, Herzog. W (2002) PHQ-D Gesundheitsfragebogen für Patienten. Heidelberg: Pfizer.
42. Wancata J, Freidl M, Fabrian F (2011) Epidemiologie der Angststörungen. *Journal für Neurologie, Neurochirurgie und Psychiatrie* 12: 332-335.
43. Luppä M, Sikorski C, Luck T, Weyerer S, Villringer A, et al. (2012) Prevalence and risk factors of depressive symptoms in latest life—results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). *International Journal of Geriatric Psychiatry* 27: 286-295.
44. Luck T, Then FS, Engel C, Loeffler M, Thiery J, et al. (2017) [The Prevalence of Current Depressive Symptoms in an Urban Adult Population]. *Psychiatr Prax* 44: 148-153.
45. Rodriguez FS, Schroeter ML, Witte AV, Engel C, Löffler M, et al. (2017) Could High Mental Demands at Work Offset the Adverse Association Between Social Isolation and Cognitive Functioning? Results of the Population-Based LIFE-Adult-Study. *Am J Geriatr Psychiatry* 25: 1258-1269.
46. Laporte Uribe F, Heinrich S, Wolf-Ostermann K, Schmidt S, Thyrian JR, et al. (2017) Caregiver burden assessed in dementia care networks in Germany: findings from the DemNet-D study baseline. *Aging Ment Health* 21: 926-937.
47. Thyrian JR, Winter P, Eichler T, Reimann M, Wucherer D, et al. (2017) Relatives' burden of caring for people screened positive for dementia in primary care : Results of the Delphi study. *Z Gerontol Geriatr* 50: 4-13.
48. Seidel D, Thyrian JR (2019) Burden of caring for people with dementia - comparing family caregivers and professional caregivers. A descriptive study. *J Multidiscip Healthc* 12: 655-663.
49. Graham NS, Junghans C, Downes R, Sendall C, Lai H, et al. (2020) SARS-CoV-2 infection, clinical features and outcome of COVID-19 in United Kingdom nursing home.
50. Gordon AL, Goodman C, Achterberg W, Barker RO, Burns E, et al. (2020) Commentary: COVID in Care Homes-Challenges and Dilemmas in Healthcare Delivery. *Age Ageing*.
51. Grabowski DC, Mor V (2020) Nursing Home Care in Crisis in the Wake of COVID-19. *JAMA*.

Tables

Table 1: Sample (n=141)

Variable	Value	range	Male (N=51)		Female (N=82)		Total	
			n/m	%/SD	n/m	%/SD	n/m	%/SD
Sex	Female						82	61.7%
Age	Years	64-98	80.16	6.67	82.34	6.09	81.52	6.38
Living situation	Living alone (yes)		23	45.1%	22	27.2%	46	34.1%
Cognitive status	MMSE-score (0-30)	7-29	23.02	3.11	23.17	3.75	23.12	3.52
Instrumental social support	Total (0-12)	2-12	9.29	2.47	9.82	2.14	9.60	2.29
	Support in repairing something (0-3)	0-3	2.0	0.89	2.22	0.87	2.15	.88
	Shopping support (0-3)	1-3	2.64	0.63	2.58	0.67	2.60	0.65
	Support in filling out forms (0-3)	0-3	2.26	0.63	2.31	0.81	2.29	0.80
	Support in dealing with administration (0-3)	0-3	2.28	0.78	2.37	0.75	2.33	0.76
Loneliness	Yes							
	De Jong Gierveld-score (6-24)	6-20	10.73	3.64	11.35	3.06	11.12	3.24
	Hughes-score (3-9)	3-9	4.34	1.71	4.62	1.96	4.45	1.80
	Considered lonely (DZA)		4	10.3%	7	10.3%	11	10.2%
Anxiety symptoms	GAD-7 score (0-20)	0-18	3.64	3.67	3.79	2.95	3.73	3.22
	Minimal symptoms		34	72.3%	50	62.5%	84	65.1%
	Low symptoms		8	17.0%	24	30.0%	32	25.2%
	Medium symptoms		5	10.6%	6	7.5%	11	8.7%
	Severe symptoms							
Anxiety disorder	Possible (>10)		5	9.8%	2	2.4%	7	5.3%
	Probable (>15)		0	0	0	0	00	0
One main symptom of depression according to DSM-V	Yes		5	9.8%	10	12.20%	15	11.37%
Sadness (n=135)	Never		33	70.2%	43	53.8%	76	59.8%
	Single days		13	27.7%	33	41.3%	46	36.2%
	More than half the days		1	2.1%	3	3.8%	4	3.1%
	nearly daily		0	0	1	1.3%	1	0.8%
Loss of interest (n=135)	Never		22	47.8%	43	53.8%	65	51.6%
	Single days		19	41.3%	28	35.0%	47	37.3%
	More than half the days		5	10.9%	7	8.8%	12	9.5%
	nearly daily		0	0%	2	2.5%	2	1.6%
Change in sadness (n=102)	Sadness more often		0	0%	0	0%	0	0%
	Sadness equally often		0	0%	2	3.0%	2	2.0%
	Sadness less often		35	100.0%	65	97.0%	100	98.0%
Change in loss of interest (n=102)	Loss of interest more often		0	0%	1	1.5%	1	1.0%
	Loss of interest equally often		1	2.9%	5	7.5%	6	5.9%
	Loss of interest less often		34	97.1%	61	91.0%	95	93.1%
(Social) activities (24)	With a perceived increase in frequency	0-7	1.07	1.04	1.37	1.38	1.26	1.27
	With a perceived decrease in frequency	0-16	6.12	3.92	7.27	3.79	6.83	3.97
	With no perceived change	0-15	6.67	3.60	5.27	2.89	5.80	3.24
	n/a	0-24	9.84	4.74	10.4	4.57	9.96	4.56
Healthcare-related services (11)	With a perceived negative impact	0-5	0.86	1.02	1.12	1.27	1.02	1.18
	With no perceived change	0-11	3.75	2.50	4.02	2.24	3.92	2.34
	With a perceived positive impact	0-2	0.20	0.49	0.18	0.45	0.19	0.46
	n/a	0-10	6.18	2.35	5.59	2.31	5.81	2.34

Table 2: Attitudes, knowledge about and perceived personal impact of the pandemic in a sample of n=141 elderly with prior cognitive impairment

Statement	response	n	%
Are you sufficiently informed about Corona?	Yes	119	85.0%
Do you know anyone infected with Corona?	Yes	4	2.9%%
Do you think the measures against the pandemic are appropriate? (n=135)	Yes	90	64.3%
	No, too strict	15	10.7%
	No, not strict enough	17	12.1%%
Do you feel worried for yourself because of Corona?	Strongly	16	11.4%
	Moderately	74	52.9%
	Not at all	50	35.7%
How stressed out do you feel because of Corona?	Strongly	6	4.3%
	Moderately	44	31.7%
	Not at all	89	64.0%
How concerned are you about your own health because of Corona?	Strongly	16	11.4%
	Moderately	62	44.3%
	Not at all	62	44.3%
How concerned are you about your family's and friends' health because of Corona?	Strongly	21	15.1%
	Moderately	76	54.7%
	Not at all	42	30.2%
How much do you fear to carry over the infection to a relative?	Strongly	7	5.0%
	Moderately	24	17.3%
	Not at all	108	77.7%
Do you worry about Corona in general?	Strongly	10	7.2%
	Moderately	69	49.6%
	Not at all	60	43.2%
Has your everyday life changed by the pandemic?	Strongly	15	10.7%
	Moderately	44	31.4%
	Not at all	79	56.4%
Have the visiting restrictions for nursing homes affected your life?	Strongly	13	9.3%
	Moderately	7	5.0%
	Not at all	120	85.7%

Table 3: Perceived change in frequency of social activities during the lockdown in Germany by a sample of n= 141 elderly with prior cognitive impairment

(Social) activity	More frequently		Less frequently		Unchanged		Not applicable	
	n	%	n	%	n	%	n	%
Being visited by family	9	6.4%	65	46.4%	49	35.0%	17	12.1%
Being visited by neighbors	0	0%	73	52.1%	29	20.7%	38	27.1%
Being visited by a friend	3	2.1%	84	60.0%	20	14.3%	33	23.6%
Meet relatives	0	0%	86	61.4%	18	12.9%	36	25.7%
Meet friend	0	0%	86	61.4%	13	9.3%	41	29.3%
Visit birthdays/ festivities	1	0.7%	88	62.9%	11	7.9%	40	28.6%
Visit hairdresser	0	0%	87	62.1%	15	10.7%	38	27.1%
Make music	1	0.7%	9	6.4%	7	5.0%	123	87.9%
Go dancing	0	0%	11	7.9%	0	0%	129	92.1%
Go for coffee	1	0.7%	81	57.9%	5	3.6%	53	37.9%
Spend time outside	20	14.3%	49	35.0%	61	43.6%	10	7.1%
Go shopping	2	1.4%	69	49.3%	35	25.0%	34	24.3%
Talk with friends and relatives by phone	53	37.9%	1	0.7%	83	59.3%	3	2.1%
Watch TV	31	22.1%	10	7.1%	98	70.0%	1	0.7%
Clean the house	8	5.7%	8	5.7%	98	70.0%	1	0.7%
Work in the garden	13	9.2%	5	3.5%	31	22.0%	92	65.2%
Home improvement	5	3.5%	3	2.1%	22	15.6%	111	78.7%
Knitting, other crafts	7	5.0%	1	0.7%	28	19.9%	105	74.5%
Visit the GP	3	2.1%	46	32.9%	58	41.1%	33	23.6%
Visted by the GP	3	2.1%	17	12.1%	21	14.9%	100	70.9%
Visit medical specialist	4	2.8%	41	29.1%	64	45.4%	32	22.7%
Visit public institutions	0	0%	19	13.5%	32	22.7%	90	63.8%
Use computer/ tablet	7	5.0%	1	0.7%	30	21.3%	103	73.0%
Utilize voluntary services	1	0.7%	6	4.4%	4	2.9%	125	91.9%

Footnote:

Table 4: perceived impact of the lockdown on health care provision by a sample of n=141 elderly people with cognitive impairment in Germany

Health care services	Worse		Unchanged		Better		Not applicable	
	n	%	n	%	n	%	n	%
Ambulatory care service	8	5.7%	54	38.3%	5	3.5%	74	52.2%
Physiotherapy	10	7.1%	32	22.7%	1	0.7%	98	69.5%
Podiatry	30	21.3%	48	34.0%	0	0%	63	44.7%
Prescribed therapies	20	14.2%	36	25.5%	2	1.4%	82	58.2%
Hospital treatments	9	6.4%	44	31.2%	2	1.4%	84	59.6%
Day clinics/ services	8	5.7%	4	2.8%	1	0.7%	127	90.1%
Provision of medication	3	2.1%	129	91.5%	4	2.8%	4	2.8%
Utilities	6	4.3%	68	48.2%	2	1.4%	64	45.4%
Food service	1	0.7%	21	14.9%	1	0.7%	11	83.0%
Mobility	28	19.9%	92	65.2%	1	0.7%	20	14.2%
Relief services	16	11.3%	19	13.5%	6	4.3%	99	70.2%

Figures




Figure 1

Change in frequency of (social) activities during the Corona pandemic in Germany in a sample of n= 141 elderly with prior cognitive impairment. Footnote: the bars represent the distribution of answers by participants who pursued the respective activity before the pandemic. Participants who indicated “not applicable” were excluded.




Figure 2

Perceived impact of the lockdown on health care provision by a sample of n=141 elderly with cognitive impairment. Footnote: the bars represent the distribution of answers by participants who used the respective services before the pandemic. Participants who indicated “not applicable” were excluded.