

Relationship Between Foot Length And Gestational Age In Pakistan

Shiyam Sunder Tikmani (✉ shiyam.sunder@aku.edu)

Aga Khan University

Sana Roujani

Aga Khan University

Syed Iqbal Azam

Aga Khan University

Haleema Yasmin

Jinnah Sindh Medical University

Khadija Bano

Jinnah Sindh Medical University

Saleem Jessani

Aga Khan University

Sayyeda Reza

Aga Khan University

Elizabeth M McClure

RTI International

Robert L Goldenberg

Columbia University

Sarah Saleem

Aga Khan University

Research

Keywords: Gestational age, last menstrual period, Ballard scoring, foot length

Posted Date: July 1st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-38713/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Global Pediatric Health on January 1st, 2020. See the published version at <https://doi.org/10.1177/2333794X20974206>.

Abstract

Background: Neonatal mortality in preterm live births is high. Therefore, knowledge of the gestational age (GA) at birth is crucial for the management and predicting prognosis of a newborn. Common methods for estimating GA such as the last menstrual period, ultrasonography, and post-natal Ballard scoring have some limitations. This study aimed to determine the relationship between foot length and GA and to develop a regression equation for predicting GA of the neonates born in Pakistan.

Methods: Data for this analysis were extracted from the hospital files of eligible women by trained study midwives. Midwives were also trained in performing the Ballard examination and taking foot length using a disposable measuring tape within an hour of the birth. The measurement was taken from the mid-point of the heel to the end of the longest toe. The GA was calculated using an android-based GA calculator, which used a report of last menstrual period, ultrasound examination, and Ballard examination. The data were analyzed using SPSS version 19. Scatter plots were constructed to assess the linearity and coefficient of determination and the correlation was calculated. Simple and Multiple Linear regression was used to construct predicting equations for GA.

Results: Both foot length and GA were available for 1542 cases and were analyzed. The median GA was 34.5 weeks with an interquartile range (IQR 4.7) and the median foot length was 7 cm (IQR 1.4). There was a positive linear relationship between foot length and GA (r^2 81.7%, p -value <0.001). Stratified analysis showed an r^2 of 81.7% for males and 81.6% for females. The r^2 for stillbirths was 84.1% and, 82.3% for live births. The r^2 for macerated stillbirths was 88.6% and 90.6% for fresh stillbirths.

Conclusion: In resource poor settings, the use of foot length can estimate GA in both live births and stillbirths and can easily identify preterm infants to initiate early management.

Plain English Summary

Assessment of gestational age (GA) at the time of birth is crucial for the management and forecasting the complications of a preterm newborn. GA was commonly estimated by women's last menstrual period or ultrasonography in the first trimester. Other methods of estimating GA are post-natal Ballard scoring and body part measurements. Each method of estimating GA has some limitations. Considering the limitation of each method, this study was conducted to develop an equation to forecast GA of Pakistani neonates by measuring post-natal foot length. Data for this analysis were extracted from the patients' file by trained study midwives. In addition, within an hour of delivery, midwives did the Ballard examination and measured foot length of the baby using a disposable measuring tape. The measurement was taken from the mid-point of the heel to the end of the longest toe. The GA was calculated using an android-based GA calculator which used a report of last menstrual period, ultrasound examination, and Ballard examination. In this study, a positive linear relationship between foot length and GA was found. This relationship between foot length and GA is similar by sex of the baby and birth status (stillbirth vs. live

birth). In resource poor settings, the use of foot length can estimate GA in both live births and stillbirths and can identify preterm infants to initiate early management.

Introduction

Pakistan has the highest neonatal mortality across the globe with a neonatal mortality rate (NMR) of 43/1000 live births (LB). (1, 2) In Pakistan, 37% of deliveries occur at home and 42% of deliveries occurred at home in the rural areas of Sindh province alone. (2) Most of these home deliveries are attended by unskilled birth attendants. (2) According to the recent Pakistan Demographic & Health Survey (PDHS-2018), the most common causes of neonatal deaths are infections and prematurity, (3) and the majority of neonatal deaths occur in preterm newborns.

Knowledge about the gestational age (GA) at birth is crucial for the management and predicting the prognosis of a newborn, especially in preterm births. (4) There are different methods of estimating GA. During the antenatal period, GA is estimated using the first day of the last menstrual period (LMP) and ultrasonography. (5) LMP is a common method to assess GA if the menstrual cycle is regular. However, in the case of an irregular cycle, LMP may not be an accurate method to estimate the GA. Ultrasound (US) examination is considered to be a gold standard for dating, (5) however, dating using US is considered accurate when it is carried out in the first trimester with a variation of ± 10 days. (6)

If the antenatal GA is unknown, then the post-natal assessment may provide a reasonable estimation of GA. Postnatal assessment of GA can be assessed by using different techniques, for example, through Dubowitz scoring (7), New Ballard Scoring (8), Eregie model (9) and measurement of different parts of the body, such as chest circumference, leg length, foot length, etc. (10, 11). Nevertheless, both antenatal and postnatal estimations of GA have some limitations.

In low-middle-income countries like Pakistan where female illiteracy is high, women generally are unable to recall their LMP (10) and a good quality first-trimester ultrasound report is not commonly available. In such cases, Ballard scoring is the most commonly used post-natal method to determine the GA of the baby. Ballard scoring consists of two components; 1) physical and 2) neuromuscular maturity assessment. However, Ballard scoring is technical and subjective for health care workers to use and estimate the GA accurately. (10) In resource-poor settings where the US is not available and Ballard scoring is challenging, one of the simple ways to estimate GA is the measurement of post-natal foot length. Health care workers can measure foot length with minimal training and can estimate GA.

The Global Action Report on Preterm Birth, "Born too soon", emphasized using simple approaches of estimating GA for early identification of preterm infants and their management. (12) Recently, a study from Ethiopia reported that foot length could be used as a diagnostic tool for predicting LBW and prematurity. (10) However, in Pakistan, there is no standardized foot length chart available to assess GA. Therefore, this study aimed to determine the relationship between GA and foot length to construct a simple regression equation for GA assessment for Pakistani newborns.

Methods

Study design, study site and population

Data for this analysis were extracted from a large prospective, observational cohort study carried out in Pakistan with the primary objective to determine the cause of death in stillbirth and preterm live-born infants. (13).

The primary study was conducted in the Department of Obstetrics and Gynecology, Jinnah Postgraduate Medical Center (JPMC), Karachi, Pakistan. The Department of Obstetrics and Gynecology has 135 beds and around 15000 annual deliveries with approximately 15% preterm births and 3.5% stillbirths. The data for the study were collected from September 3rd, 2018 to December 31st, 2019.

Pregnant women ≥ 18 years of age admitted with an imminent preterm delivery based on clinical indications (20–36,6 weeks gestation) or women with a known stillbirth (intrauterine fetal death ≥ 20 weeks) were included in the study. Induced abortions (< 20 weeks), live births with unknown GA, lethal congenital malformations, neonates with limb deformities, and missing foot length measurements were excluded.

Data collection procedure

GA was calculated through an Android-based GA calculator with a predefined algorithm using the hierarchy of methods established by the American College of Obstetrics and Gynecology, (14) using the report of reliable LMP, ultrasound examination and Ballard examination. The LMP and ultrasound data were extracted from the woman's hospital file. The trained study midwives carried out Ballard examinations. Later, all the information was entered into the GA calculator (developed by RTI) i.e. the date of delivery, LMP, date of her first or only available ultrasound, and estimated GA by the US. For live births, the date of the Ballard examination, and the total score of the physical and neuromuscular examinations was collected and entered. Based on this information, the GA calculator then provided the best GA estimate. If the baby was sick or the neuromuscular examination could not be completed, study midwives doubled the score of the physical examination and entered it into the GA calculator.

Foot length of the right foot was measured by trained midwives using a disposable measuring tape within an hour of birth from the mid-point of the heel to the end of the longest toe. To minimize the effect of the plantar grasp reflex on the length of the foot, midwives were trained to hold the ankle with a finger placed on the dorsum of the foot to keep the foot straight. An average of three readings of foot length in centimeters was used for the final analysis.

Study midwives were trained for data extraction from the women's hospital files, carrying out a Ballard examination on livebirths, and measuring the foot length of the baby. They were also trained in operating the GA calculators. Random checks were carried out by the study supervisor nurse to verify the measurement of the foot length and appropriateness of the use of the GA calculator by the midwives.

Statistical analysis

Data were analyzed using SPSS version 19 (SPSS Inc., Chicago, IL, USA). The normality of GA and foot length was checked through the Kolmogorov-Smirnov test. For GA and foot length, median and interquartile ranges (IQR) are reported. Frequencies and percentages were calculated for categorical variables. For comparison of characteristics of the participants, student t-tests for continuous variables and chi-square tests for categorical variables were used.

A Scatter plot was made to assess the linearity between GA and foot length. Co-efficient of determination (r^2) and correlation (r) were calculated. Separate scatter plots were made to assess linearity by sex, livebirth, stillbirth and presence of maceration. Linear regression modeling was done for estimating predictive values of foot length for GA. For the assumption of the normality of residuals, a normality plot of residuals and predicted GA was constructed using PP plots and assessing heteroscedasticity. Cook's distance and Df Beta were estimated to assess influential values.

Multiple linear regression equation was made which is adjusted for sex and birth status. The equation for predicting GA is expressed below:

$$\hat{Y} = \beta_0 + \beta_1 (x) + \beta_2 (y) + \beta_3 (z)$$

Where \hat{Y} is the predicted GA), β_0 is the slope, and β_1 is the regression coefficient for foot length (x) cm, β_2 is regression coefficient for male sex (y) and β_3 is regression coefficient for livebirth (z).

Cross-validation of the model was done by randomly dividing the data set into two subsamples (1/3 & 2/3). The statistical model was generated with 2/3rd of the data. The model was then used to predict GA in the 1/3rd subsample. The standardized error term was calculated based on the predicted GA using the formula: $(\text{observed GA} - \text{predicted GA})^2 / (\text{predicted GA})$.

For validation, the foot length of Pakistani neonates for estimating GA was assessed using regression equations reported by different studies. The mean difference between observed and predicted GA was compared using a paired t-test. Mean differences and 95% CI were plotted in a graph.

Ethical approvals

The primary study was approved by the ethical review committees of Aga Khan University, Jinnah Post-graduate medical center, National Institute of Child Health and National Bioethics Committee of Pakistan. Informed consent was taken from each participant of the study.

Results

A total of 20,169 women delivered during the study period. Of these births, 1066 (6.6%) were stillbirths and 1844 (9.14%) were preterm live births. A total of 2,696 women were screened for imminent preterm birth or stillbirth and 2905 babies were born. Out of 2548 eligible mother-infant pairs, 2044 provided

consent and were enrolled in this study. Of these, 607 were stillbirths and 1437 were preterm live births. GA was available for 1436 preterm neonates and 554 stillbirths. Both foot length and GA were available for 1542 cases that were analyzed to determine this relationship, 767 (49.7%) were male, 773 (50.1%) were female, and in 2 (0.1%) cases, information on sex was not known. Of 1542 infants, 1132(73.4%) were live births and 410 (26.6%) were stillbirths (Table 1).

The Shapiro-Wilk test ($p > 0.05$) and a visual inspection of histograms, normal Q-Q plots, and box plots of maternal age, GA and foot length, showed that these variables were not normally distributed. The median maternal age was 26 (IQR 7) years, the median GA was 34.5 (IQR 4.7) weeks; median foot length was 7 (IQR 1.4) cm (Table 1).

$$\hat{Y} = 10.359 + (3.447 * FL) + (0.179 * Male) + (-0.357 * livebirth)$$

The mean difference of observed and predicted GA with CIs, using different equations are shown in Fig. 3.. The difference of observed and predicted GA of this study is consistent with the difference in the observed value of this study and predicted value of the equation of Manjunatha et al (12). However, all other equations (Table 2) appear to underestimate the GA compared to mean difference of the observed and predicted value of this study (Fig. 3).

Discussion

In this study, it was found that there is a positive linear relationship between GA and foot length ($r^2 = 81.7\%$). Streeter et al in 1920, first observed that the foot length can be used as a proxy for estimation of GA. (15) Many studies since then have shown a positive linear relationship between GA and foot length. (16–21) However, the coefficient of determination (r^2) of foot length and GA is different in different studies. Singhal et al found a positive relationship between foot length and GA with an r^2 of 93% (18). Mhaskar et al reported that foot length correlated very well with the GA with an r^2 of 85%. (20) Rakkapan et al reported a lower r^2 (65%) between foot length and GA. (17) In this study, the correlation between foot length and GA was 82.3% in live births and 84.1% in stillbirths, 88.6% macerated stillbirths and 90.6% for fresh stillbirths.

There are wide variations in the methods used across the studies. These variations include sample size, the GA of infants included in the analysis, method of measurement of GA, different tools to measure foot length, and different units used for the foot length measurement. However, most of the studies reported that the landmark for measuring foot length was similar to that used in this study. Several studies measured GA using different methods. Some used LMP (18), some used ultrasound examination (20), some used clinical examination(19), and some used the Ballard score for assessing GA. (16) Similarly, foot length was measured using different measuring equipment. The most commonly used equipment was a Vernier caliper (16, 17, 19, 20), followed by transparent rigid scales and steel or flexible tapes. (10, 18, 21) The unit of measurement was also different across studies. Some used centimeters (16–18) and some millimeters. (19–21) However, almost all studies measured foot length from the mid-point of the

heel to the longest toe. In this study, we determined GA using an algorithm and foot length by a disposable flexible paper tape in centimeters from mid-point of the heel to the longest toe.

Due to these variations, we compared the estimated GA using foot length from this study in the regression equations constructed by different studies (Table 2). Most of the prediction equations underestimate the GA compared to the predicted model developed in this study except for the model by Manjunatha et al. However, the mean difference of observed and predicted GA using different equations (Table 1) shows the minimal difference with precise confidence limits.

To the best of our knowledge, this is the first study in Pakistan for determining the relationship between foot length and GA and has some strengths. The data were prospectively collected. The GA was calculated using an algorithm through an Android application. The relationship of foot length and GA was compared by gender, stillbirth and live birth, macerated, and fresh stillbirth, which have not been reported in any single study so far. Foot length was measured by well-trained midwives with standardization methods. We excluded infants with congenital malformations and limb deformities from this study as this might have affected the GA estimation by 2–3 weeks. (22)

However, there are some limitations to the study. A flexible disposable measuring tape was used to measure the foot length. However, we believe that standardization exercises and taking an average of three readings should have minimized the error. We did not perform in-depth intra- or inter-observer variation analysis; however, study midwives were trained with quarterly refresher trainings and were monitored by a senior study nurse, thus minimizing the inter-observer variation. In conclusion, a simple measurement of foot length can estimate GA and can identify preterm infants to initiate treatment to reduce mortality in preterm neonates in a resource-limited setting.

Abbreviations

Gestational age (GA); Interquartile range (IQR); Jinnah Postgraduate Medical Center (JPMC); Last menstrual period (LMP); Research Triangle Institute (RTI); Ultrasound (US)

Declarations

Ethical approvals: The primary study was approved by the ethical review committees of Aga Khan University, Jinnah Post-graduate medical center, National Institute of Child Health and National Bioethics Committee of Pakistan. Informed consent was taken from each participant of the study.

Consent for publication: Not applicable

Availability of data and material: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interest: The authors declare that they have no competing interests.

Funding:

The primary study was funded by the Bill and Malinda Gates Foundation via the Research Triangle Institute (RTI).

Author's contribution: SST, EM, SS RLG conceptualized and designed the study. SST conducted data analysis, interpretation and prepared the first draft of the manuscript. SR did literature search and oversaw data collection. SR and SRe monitored the data quality. SJ helped in data analysis and interpretation, IA supervise data analysis and interpretation of data. SST, SR, IA, HY, KB, EM RLG, and SS reviewed the manuscript and provided critical feedback. All authors (SST, SR, IA, HY, KB, SJ, SRe, EM, RLG, SS) read and approved the final version of the manuscript.

Acknowledgements: The authors recognize the contributions of the midwives who collected data, Kanwal Ilyas and Neelofar Sami for monitoring the data collection and Zaheer Habib and team for data management. We also acknowledge the parents of the enrolled infants who voluntarily participated in the study.

Dedication: To my Father Late Prof. Pirbhulal Tikmani and mother Late Asha Devi (Revti).

References

1. UNICEF
UNICEF. For, Every, Child. Neonatal mortality. Unicef.org; 2019.
2. National, Institute, of, Population, Studies, (NIPS), et al. Pakistan Demographic and Health Survey 2017-18. Islamabad, Pakistan, Rockville, Maryland, USA, NIPS and ICF; 2019.
3. Hussain S. Neonatal morbidity and mortality pattern in a Tertiary Care Neonatal Unit of a Teaching Hospital. *Ann Pak Inst Med Sci.* 2014;10(1):7–11.
4. Lee AC, Mullany LC, Ladhani K, Uddin J, Mitra D, Ahmed P, et al. Validity of newborn clinical assessment to determine gestational age in Bangladesh. *Pediatrics.* 2016;138(1):e20153303.
5. Unger H, Thriemer K, Ley B, Tinto H, Traoré M, Valea I, et al. The assessment of gestational age: a comparison of different methods from a malaria pregnancy cohort in sub-Saharan Africa. *BMC Pregnancy Childbirth.* 2019;19(1):1–9.
6. McClure EM, Nathan RO, Saleem S, Esamai F, Garces A, Chomba E, et al. First look: a cluster-randomized trial of ultrasound to improve pregnancy outcomes in low income country settings. *BMC Pregnancy Childbirth.* 2014;14(1):73.
7. Dubowitz LM, Dubowitz V, Goldberg C. Clinical assessment of gestational age in the newborn infant. *J Pediatr.* 1970;77(1):1–10.
8. Marín MG, Martín JM, Lliteras GF, Delgado SG, Pallás CA, Pérez EE. Assessment of the new Ballard score to estimate gestational age. *Anales de pediatria (Barcelona, Spain: 2003).* 2006;64(2):140-5.

9. Eregie CO. Assessment of gestational age: modification of a simplified method. *Developmental Medicine Child Neurology*. 1991;33(7):596–600.
10. Gidi NW, Berhane M, Girma T, Abdissa A, Lim R, Lee K, et al. Anthropometric measures that identify premature and low birth weight newborns in Ethiopia: a cross-sectional study with community follow-up. *Arch Dis Child*. 2020;105(4):326–31.
11. Thawani R, Dewan P, Faridi M, Arora SK, Kumar R. Estimation of gestational age, using neonatal anthropometry: a cross-sectional study in India. *J Health Popul Nutr*. 2013;31(4):523.
12. Howson CP, Kinney MV, McDougall L, Lawn JE. Born too soon: preterm birth matters. *Reproductive health*. 2013;10(S1):1.
13. McClure EM, Saleem S, Goudar SS, Dhaded S, Guruprasad G, Kumar Y, et al. The project to understand and research preterm pregnancy outcomes and stillbirths in South Asia (PURPOSE): a protocol of a prospective, cohort study of causes of mortality among preterm births and stillbirths. *Reproductive health*. 2018;15(1):89.
14. Obstetricians, ACo. Gynecologists. Committee opinion no 611: method for estimating due date. *Obstet Gynecol*. 2014;124(4):863–6.
15. Streeter GL. Weight, sitting height, head size, foot length and menstrual age of the human embryo. *Contrib Embryol*. 1920;11:143–70.
16. Tenali ASL, Tenali RK. Study of foot length as an alternate measurement for assessment of gestational maturity in neonate. *Int J Contemp Pediatr*. 2019;6(2):477–81.
17. Rakkappan I, Kuppusamy N. Newborn foot length measurement to identify high-risk neonate. *Int J Sci Stud*. 2016;4:13–9.
18. Singhal S, Tomar A, Masand R, Purohit A. A simple tool for assessment of gestational age in newborns using foot length. *J Evol Med Dent Sci*. 2014;3:6424–9.
19. Manjunatha B, Nithin M, Sameer S. Cross sectional study to determine gestational age by metrical measurements of foot length. *Egyptian Journal of Forensic Sciences*. 2012;2(1):11–7.
20. Mhaskar R, Agarwal N, Takkar D, Buckshee K, Deorari A. Fetal foot length—a new parameter for assessment of gestational age. *International Journal of Gynecology Obstetrics*. 1989;29(1):35–8.
21. Daga S, Daga A, Patole S, Kadam S, Mukadam Y. Foot length measurement from foot print for identifying a newborn at risk. *J Trop Pediatr*. 1988;34(1):16–9.
22. Sherwood RJ, Meindl R, Robinson H, May R. Fetal age: methods of estimation and effects of pathology. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*. 2000;113(3):305–15.

Tables

Table 1
Characteristics of infants

Characteristics	Livebirth (n-1130)	Stillbirth (n-410)	Total (n-1542)
Gender			
Male	560 (49.6%)	207 (50.5%)	767 (49.8%)
Female	570 (50.4%)	203 (49.5%)	773 (50.2%)
Gestational age (weeks) Median (IQR)	34.6 (3.9)	33.3 (7.4)	34.5 (4.7)
Foot length (cm) Median (IQR)	7.1 (1.2)	6.5 (1.9)	7.0 (1.4)

Table 2
Simple Linear Regression equations constructed by different studies

Author & year	Regression equation	R-Square
Tenali ASL et al. (2019)	$GA = 21.84 + (2.09 * FL)$	0.9
Rakkappan I, et al. (2016)	$GA = 15.343 + (3.183 * FL)$	0.65
Singhal S, et al. (2014)	$GA = 6.278 + (4.15 * FL)$	0.86
Munjunatha B, et al. (2012)	$GA = 5.60 + (4.11 * FL)$	0.97
Mhaskar R, et al. (1989)	$GA = 12.79 + (3.36 * FL)$	0.85
Daga SR, et al. (1988)	$GA = 11.55 + (3.41 * FL)$	0.84
FL-foot length, GA-gestational age		

Figures

Figure 1: Scatter plot showing a linear relationship between foot length and gestational age (A-D).

A. Overall relationship between gestational age and foot length with correlation coefficient of 0.903 and coefficient of determination 0.817 (n-1542)





Figure 1


Scatter plot showing a linear relationship between foot length and gestational age (A-D). A. Overall relationship between gestational age and foot length with correlation coefficient of 0.903 and coefficient of determination 0.817 (n-1542)

A. Relationship between foot length and gestational age by sex

Scatter plot (Foot length and gestational age-**Male**), (n = 766)
 $r = 0.902$, $r^2 = 0.817$




Scatter plot (Foot length and gestational age-**Female**), (n = 772)
 $r = 0.903$, $r^2 = 0.816$



B. Relationship of foot length and gestational age by livebirth and stillbirth

Scatter plot (Foot length and gestational age-**Stillbirth**), (n = 410)
 $r = 0.917$, $r^2 = 0.841$



Scatter plot (Foot length and gestational age-**Livebirth**), (n = 1132)
 $r = 0.907$, $r^2 = 0.823$




Figure 2

Scatter plot showing a linear relationship between foot length and gestational age (A-D). B. Relationship between foot length and gestational age by sex C. Relationship of foot length and gestational age by livebirth and stillbirth

A. Relationship between foot length and gestational age by presence and absence of maceration among stillbirth

Scatter plot (Foot length and gestational age-Macerated Stillbirth), (n = 228)
 $r=0.941$, $r^2 = 0.886$

Scatter plot (Foot length and gestational age-Fresh Stillbirth), (n = 181)
 $r=0.951$, $r^2 = 0.906$




Figure 3

Scatter plot showing a linear relationship between foot length and gestational age (A-D). D. Relationship between foot length and gestational age by presence and absence of maceration among stillbirth

Figure 2: Graph showing distribution regression standardized residuals




Figure 4

Graph showing distribution regression standardized residuals

Figure 3: Mean difference and 95% CI of observed and predicted values using different prediction equations




Figure 5

Mean difference and 95% CI of observed and predicted values using different prediction equations

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CoverLetter.docx](#)