

Intrathoracic Fracture-dislocation of the Humeral Head:A Case Report and Review of the Literature

Ruifeng Yang

Lishui Central Hospital and Fifth Affiliated Hospital of Wenzhou Medical College

chong wang (✉ 252000966@qq.com)

Lishui Central Hospital and Fifth Affiliated Hospital of Wenzhou Medical College

<https://orcid.org/0000-0002-2418-804X>

Research Article

Keywords: Arthroplasty, humeral fractures, intrathoracic dislocation, thoracic injuries, high-energy trauma

Posted Date: April 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-390864/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Intrathoracic displacement of a humeral head fracture is extremely rare. Only slightly more than 30 cases have been reported. Because few cases have been reported, there is no consensus on how to treat this injury. The etiology, injury mechanism, related lesions and treatment of the injury are diverse.

Case presentation

A 73-year-old woman presented with multiple fractures of the left ribs, bilateral lung contusions, extensive emphysema of the anterior and posterior chest wall, massive left hemopneumothorax, fracture-dislocation of the proximal humerus and intrathoracic displacement of the humeral head. The patient was sent to the operating room for emergency thoracotomy surgery. The head of the humerus was confirmed to be completely removed from the thoracic cavity during the operation. After discussion with the orthopedic surgeon, the humeral head was discarded considering avascular necrosis, and open reduction and internal fixation were not performed. Three weeks later, the orthopedic surgery team performed reverse shoulder arthroplasty. During follow-up, the patient's shoulder was free from pain, and its range of movement (ROM) included 110° flexion, 70° abduction, 35° external rotation and 50° internal rotation.

Conclusions

Intrathoracic displacement of the humeral head due to proximal humeral fracture is a very rare and serious trauma that requires multidisciplinary treatment. Considering the extremely high risk of humeral head necrosis, actively removing broken bone fragments of the humeral head in the early stage is recommended, and we advocate for shoulder arthroplasty for elderly patients. Detailed preoperative evaluations and individualized operation plans should be made to achieve the best effect.

Background

Intrathoracic fracture-humeral head dislocation has been rarely reported and requires a combination of high-energy trauma and specific mechanisms. Enumerating the mechanism of injury, most of which are high-energy injuries. We can review including 18 falls from height [1–18], 7 motor vehicle accidents [19–25], and 2 were knocked down by motor vehicles [26, 27], 2 cases of falling from standing height [28, 29], and 1 case of falling from a bicycle [30].

The purpose of this study is to review published cases to examine the etiology, mechanism, related injuries and treatment of this rare injury. This study described a patient who suffered an intrathoracic fracture-dislocation of the humeral head after a high-energy traffic accident.

Case Presentation

A 73-year-old woman presented to the emergency department after a high-speed motor vehicle accident with the chief complaint of left chest pain accompanied by dyskinesia with dyspnea. The patient's left shoulder was injured, with local tenderness and loss of function. She denied loss of consciousness. The patient was evaluated and treated according to the concepts of advanced trauma life support (ATLS) and damage control surgery (DCS). The Glasgow coma scale was 15. Admission vital signs included body temperature of 36.7°C, blood pressure of 104/68 mmHg, pulse 87 of times/min, respiratory rate of 25 times/min, and blood oxygen saturation of 90%. Physical examination of the left upper limb revealed left shoulder swelling, deformity, severe instability, dyskinesia, ecchymosis, strong pulsation of the radial artery, and no obvious hypoesthesia. There was obvious subcutaneous emphysema in the left chest, tenderness in the front of the left ribs, and decreased respiratory movement and respiratory sounds. Radiography and computed tomography (CT) showed multiple fractures of the left ribs, bilateral lung contusions, extensive emphysema of the anterior and posterior chest wall, massive left hemopneumothorax, fracture-dislocation of the proximal humerus and intrathoracic displacement of the humeral head(Fig. 1–3).

After stabilizing the patient with the ATLS regimen, the patient was sent to the operating room for emergency thoracotomy surgery. A thoracic drainage tube was inserted and fixed in the sixth intercostal space of the left axillary midline after complete hemostasis. The third and fourth ribs were fixed with a rib embracing fixator. The operation was successful, and the patient was transfused with 370 mL plasma and 5 units of packed red cells. Subsequently, the patient was further treated in the intensive care unit.

The head of the humerus was confirmed to be completely removed from the thoracic cavity during the operation(Fig. 4). After discussion with the orthopedic surgeon, the humeral head was discarded considering avascular necrosis, and open reduction and internal fixation were not performed. Three weeks later, electromyography (EMG) showed normal axillary nerve. Thus, the orthopedic surgery team performed reverse shoulder arthroplasty(Fig. 5). The patient is being well after reoperation.

Postoperative rehabilitation protocol: 0 ~ 6w after surgery: Patients were educated to make them understand the significance of postoperative rehabilitation. It includes boxing pump, assisted flexion and extension, pendulum movement, active flexion and extension of elbow and wrist, shoulder abduction pillow and shrug. Oral celebrex was given to patients within 3 weeks after surgery to reduce pain and prevent ectopic ossification. 7 ~ 12w after operation: Assisted external rotation, deltoid isometric contraction, table cleaning exercises and wall climbing exercises. At the final follow-up, the patient's shoulder was free from pain, and its range of movement (ROM) included 110° flexion, 70° abduction, 35° external rotation and 50° internal rotation.

Discussion And Conclusions

Humeral head fracture-dislocation and penetration into the chest are injuries that have been rarely reported in the literature. To date, only 30 cases have been recorded in the literature. No treatment

guidelines are currently available, there is a high variability in the management of these patients, even diametrically opposed treatment options. Treatment has to be individualized for each patient.

The literature on this subject is limited, so the mechanism of the injury has not been elucidated. Hardcastle et al.[31] believed that the occurrence of injury can be divided into two stages. First, a drop in forced abduction with external rotation causing dislocation and moves the humeral head toward the chest, and then a sharp adduction leads to fracture of the humeral head. Glessner[32] and Patel et al[15] proposed a similar mechanism, in which a forced abduction dislocated the shoulder into the chest cavity, and in the process of impact against the ribs would be when the fracture took place. Karr et al.[7] proposed a two-stage mechanism of injury. When the elbow was in the abduction position, the humeral head was fractured and dislocated, and then the shoulder hit the ground, causing the rib to fracture and the humeral head to be embedded in the chest cavity. As all these authors have concluded, a single traumatic force is unlikely to cause this type of injury[21]. The injury mechanism of our case was associated with high-energy trauma of abduction and external rotation of the shoulder, and there was no need for a reduction operation. Spontaneous dislocation reduction occurred in most cases, and no reduction operation was needed.

This mechanism often leads to fracture-dislocation of four parts of the proximal end of the humerus. As an exception, two cases of two-part fracture-dislocation were reported[1, 26], and one case of three-part fracture-dislocation[31] and separation fracture of proximal humerus epiphysis[30] were reported in the literature. Humeral head fracture-dislocation and chest penetration often occur due to high-energy injuries, mostly accompanied by related injuries. Pneumothorax, hemothorax and other lung injuries almost always coexist. It is often associated with thoracic and limb injuries. Fractures of the scapula, humeral shaft, elbow, wrist, pelvis and tibial plateau have also been reported. Vascular injuries, including aortic compression and subclavian artery and costal neck trunk injuries, were also reported. Cases of rupture of the main bronchus have also been reported in the literature[24]. In addition to humeral head displacement to the ipsilateral thoracic cavity, there are other special cases, such as 1 case of the humeral head in the retroperitoneal space[33] and 2 cases of movement to the contralateral thoracic cavity[4, 5].

When a nerve injury occurs, the axillary nerves generally tend to be involved, which is very difficult to evaluate in the initial physical examination. There have also been reports of median nerve, ulnar nerve and secondary brachial plexus trunk injuries. Most patients spontaneously recover without surgery, although other patients do not fully recover[3, 4, 6, 27, 34]. In this paper, EMG showed normal axillary nerve.

Treatment of the humeral head differed from previous reports. There is no agreement on whether to remove the humeral head, most of which is removed to prevent potential complications. In early and subsequent individual cases, fragments of the humeral head remained in the chest[1, 7, 10, 15, 27]. Some scholars have suggested that it is necessary to remove fragments only when there are cardiopulmonary complications. The reason for removal of bone fragments from the humeral head is that there may be

related complications, such as infection or foreign body reactions[18]. It has been reported that the retained humeral head was removed 6 weeks late, and was embedded in the pulmonary parenchyma with extensive adhesion, requiring extensive resection[35]. We believe that fragments of the humeral head should be actively removed at an early stage. Although humeral head fragments are sterile, bone fragments may become a potential focus of infection, complicating treatment. Extraction can be performed by thoracotomy or thoracoscopy. Ideally, minimally invasive methods, such as video-assisted thoracoscopy, should be adopted to remove the humeral head. After removing it, subsequent techniques for shoulder reconstruction will depend on specific conditions of the patient.

In terms of treatment of fracture of the proximal humerus, a surgeon must consider the risk of humeral head necrosis when determining the surgical plan, and vitality of the humeral head determines treatment of the proximal humerus. In three- or four-part displaced fractures, there is high risk of ischemic necrosis or nonunion of the head. Humeral head necrosis increased from approximately 25% of three-part fractures to nearly 60% of four-part fractures[34, 36]. In the orthopedic literature, there is still a great deal of controversy about which surgical option is the most appropriate. Open reduction and internal fixation are possible in younger patients[5, 22, 26, 30, 31]. In published articles, the majority of cases were treated with hemiarthroplasty of the glenohumeral joint[2–6, 8, 9, 11–13, 18–21, 23, 27–29], and one patient was treated with reverse shoulder arthroplasty[12].

Intrathoracic displacement of the humeral head due to proximal humeral fracture is a very rare and serious trauma that requires multidisciplinary treatment. Considering the extremely high risk of humeral head necrosis, actively removing broken bone fragments of the humeral head in the early stage is recommended, and we advocate for shoulder arthroplasty for elderly patients. Detailed preoperative evaluations and individualized operation plans should be made to achieve the best effect.

Abbreviations

ROM:Range Of Movement;ATLSAdvanced Trauma Life Support (ATLS) ;DCS:Damage Control Surgery ; CT: Computed Tomography; EMG:electromyography

Declarations

Acknowledgements:Not applicable.

Ethics approval and consent to participate: The experimental protocol was established, according to the ethical guidelines of the Helsinki Declaration and was approved by the Human Ethics Committee of Lishui Central Hospital,Written informed consent was obtained from individual or guardian participants.

Consent for publication:We obtained written consent from the participants before publishing this information.

Authors Contributions: WC collected the data of the case and consulted relevant literature. YRF was a major contributor in writing the manuscript. All authors read and approved the final manuscript.

Availability of data and materials:All the data needed to achieve the conclusion are presented in the paper.

Funding: Not applicable

Competing Interests: The authors declare that they have no competing interests.

References

- [1] WEST E. F. (1949). Intrathoracic dislocation of the humerus. *The Journal of bone and joint surgery. British volume*, 31B(1), 61.
- [2] Boyer, P., Alsac, J. M., Ettori, M. A., Lesèche, G., & Hutten, D. (2007). Four-part fracture after intrathoracic displacement of the humeral head: a case report and review of the literature. *Archives of orthopaedic and trauma surgery*, 127(8), 651–654.
- [3] Wiesler, E. R., Smith, A. M., & Shilt, J. S. (2004). Humeral head fracture-dislocation into the thoracic outlet: case report and review of the literature. *Journal of shoulder and elbow surgery*, 13(5), 576–579.
- [4] Ebersson, C. P., Ng, T., & Green, A. (2000). Contralateral intrathoracic displacement of the humeral head. A case report. *The Journal of bone and joint surgery. American volume*, 82(1), 105–108.
- [5] Abbott, G. F., Gaissert, H., & Faberman, R. S. (1999). Contralateral intrathoracic migration of humeral head fracture dislocation. *AJR. American journal of roentgenology*, 172(5), 1403–1404.
- [6] al-Kaar, M., Garcia, J., Khan, H. G., & Manueddu, C. (1998). Fracture-luxation intrathoracique de la tête humérale [Intrathoracic dislocation fracture of the head of the humerus]. *Journal de radiologie*, 79(1), 57–59.
- [7] Kaar, T. K., Rice, J. J., & Mullan, G. B. (1995). Fracture-dislocation of the shoulder with intrathoracic displacement of the humeral head. *Injury*, 26(9), 638–639.
- [8] Jun-Song, W., Jing-Yu, D., Zhi-Qiang, W., Fang, G., & Xiang-Jin, L. (2012). Intrathoracic displacement of the humeral head in a trauma patient. *Indian journal of orthopaedics*, 46(5), 596–598.
- [9] Hawkes, T., Parra, M., & Shrock, K. (2014). Intrathoracic fracture-dislocation of the proximal humerus treated with rib fixation and shoulder arthroplasty. *American journal of orthopedics (Belle Mead, N.J.)*, 43(4), E74–E78.
- [10] Du Plessis, N., Rajendran, S., Naqi, S. A., & Cahill, R. (2012). Intrathoracic fracture-dislocation of the humeral head. Is removal of the humeral head necessary?. *BMJ case reports*, 2012, bcr2012007478.

- [11] Daffner, S. D., Cipolle, M. D., & Phillips, T. G. (2010). Fracture of the humeral neck with intrathoracic dislocation of the humeral head. *The Journal of emergency medicine*, 38(4), 439–443.
- [12] Maroney, S. S., & Devinney, D. S. (2009). Intrathoracic fracture-dislocation of the proximal humerus treated with reverse total shoulder arthroplasty. *Orthopedics*, 32(12), 924.
- [13] Chen, J., Yan, J., Wang, S., Zhong, H., & Zhou, H. (2015). Intrathoracic dislocation of the humeral head accompanied by polytrauma: How to treat it?. *Ulusal travma ve acil cerrahi dergisi = Turkish journal of trauma & emergency surgery : TJTES*, 21(2), 149–151.
- [14] Schoffl, H., Froschauer, S. M., & Huemer, G. M. (2009). Intrathoracic displacement of the humeral head after fracture. *Wiener klinische Wochenschrift*, 121(7-8), 246.
- [15] PATEL, M. R., PARDEE, M. L., & SINGERMAN, R. C. (1963). INTRATHORACIC DISLOCATION OF THE HEAD OF THE HUMERUS. *The Journal of bone and joint surgery. American volume*, 45, 1712–1714.
- [16] Brogdon, B. G., Crotty, J. M., MacFeely, L., McCann, S. B., & Fitzgerald, M. (1995). Intrathoracic fracture-dislocation of the humerus. *Skeletal radiology*, 24(5), 383–385.
- [17] Rupperecht, H., Ghidau, M., & Gaab, K. (2017). Severe Thoracic Trauma Due to an Intrathoracic Dislocation of a Fractured Humeral Head in an Aged Patient; A Case Report. *Bulletin of emergency and trauma*, 5(3), 212–214.
- [18] Fernández, A. I., & Fernández, A. G. (2018). Intrathoracic fracture-Dislocation of the shoulder. *Journal of clinical orthopaedics and trauma*, 9(Suppl 2), S54–S57.
- [19] Kocer, B., Gulbahar, G., Aktekin, C. N., Gunal, N., Birinci, B., Dural, K., & Sakinci, U. (2007). Intrathoracic humeral head fracture-dislocation: is removal of the humeral head necessary?. *The Annals of thoracic surgery*, 84(4), 1371–1372.
- [20] Harman, B. D., Miller, N. G., & Probe, R. A. (2004). Intrathoracic humeral head fracture-dislocation. *Journal of orthopaedic trauma*, 18(2), 112–115.
- [21] Liu, H. H., Lee, Y. H., Yang, S. W., & Wong, C. Y. (2007). Fracture of the proximal humerus with intrathoracic dislocation of the humeral head. *The Journal of trauma*, 63(2), E37–E39.
- [22] Anderson, M., Rose, P., Jacofsky, D. J., Torchia, M. E., & Dahm, D. L. (2007). Intrathoracic fracture-dislocation of the proximal humerus: a case report and report of a new surgical technique. *The Journal of trauma*, 63(4), 920–923.
- [23] Tsai, Y. M., Shen, P. H., Chang, H., Tzao, C., Lee, S. C., & Cheng, Y. L. (2014). Intrathoracic displacement of a fractured humeral head due to blunt trauma. *Annals of thoracic and cardiovascular surgery : official journal of the Association of Thoracic and Cardiovascular Surgeons of Asia*, 20 Suppl, 592–594.

- [24] Hayes, N., White, J., Lillie, P., Bennetts, J. S., Tu, C. G., & Bain, G. I. (2020). Intrathoracic shoulder dislocation causing rupture of the right main bronchus. *Archives of orthopaedic and trauma surgery*, 140(4), 443–447.
- [25] Sola Junior, W. C., & Santos, P. S. (2017). Intrathoracic fracture-dislocation of the humerus - case report and literature review. *Revista brasileira de ortopedia*, 52(2), 215–219.
- [26] Simpson, N. S., Schwappach, J. R., & Toby, E. B. (1998). Fracture-dislocation of the humerus with intrathoracic displacement of the humeral head. A case report. *The Journal of bone and joint surgery. American volume*, 80(6), 889–891.
- [27] Abellan, J. F., Melendreras, E., Gimenez, D. J., Carrillo, F. J., Ruano, L., & Rivkin, J. (2010). Intrathoracic fracture-dislocation of the humeral head: a case report. *Journal of orthopaedic surgery (Hong Kong)*, 18(2), 254–257.
- [28] Salhiyyah, K., Potter, D., & Sarkar, P. K. (2012). Fracture-dislocation of humeral head with intrathoracic displacement. *Asian cardiovascular & thoracic annals*, 20(2), 196–198.
- [29] Griffin, N. C., Temes, R. T., Gill, I. S., & Rice, T. W. (2007). Intrathoracic displacement of a fractured humeral head. *The Annals of thoracic surgery*, 84(4), 1400.
- [30] Guiotti Filho, J. , Borges, Antônio Carlos Wall, & Leite, M. D. C. . (2008). Fratura epifisiolise da extremidade proximal do úmero com luxação intratorácica: relato de caso. *Revista Brasileira De Ortopedia*, 43(1-2), 48-52.
- [31] Hardcastle, P. H., & Fisher, T. R. (1981). Intrathoracic displacement of the humeral head with fracture of the surgical neck. *Injury*, 12(4), 313–315.
- [32] Glessner, J. R. . (1961). Intrathoracic dislocation of the humeral head. *The Journal of Bone and Joint Surgery*, 43(3), 383-385.
- [33] Wirth, M. A., Jensen, K. L., Agarwal, A., Curtis, R. J., & Rockwood, C. A., Jr (1997). Fracture-dislocation of the proximal part of the humerus with retroperitoneal displacement of the humeral head. A case report. *The Journal of bone and joint surgery. American volume*, 79(5), 763–766.
- [34] Hawkes, T., Parra, M., & Shrock, K. (2014). Intrathoracic fracture-dislocation of the proximal humerus treated with rib fixation and shoulder arthroplasty. *American journal of orthopedics (Belle Mead, N.J.)*, 43(4), E74–E78.
- [35] Griffin, N. C., Temes, R. T., Gill, I. S., & Rice, T. W. (2007). Intrathoracic displacement of a fractured humeral head. *The Annals of thoracic surgery*, 84(4), 1400.
- [36] Besch, L., Daniels-Wredenhagen, M., Mueller, M., Varoga, D., Hilgert, R. E., & Seekamp, A. (2009). Hemiarthroplasty of the shoulder after four-part fracture of the humeral head: a long-term analysis of 34

Figures




Figure 1

CT-scan showing the intrathoracic localization of the humeral head.




Figure 2

CT-scan showing the intrathoracic localization of the humeral head.




Figure 3

Three-dimensional reconstruction CT showed intrathoracic displacement of humeral head and ribs fracture.



Figure 4

Intra-operative picture demonstrating the fractured humeral head.



Figure 5

Postoperative radiographs of the reverse shoulder arthroplasty