

Complementary feeding patterns of Filipino infants and toddlers lack diversity, especially among children from poor households

Emma F Jacquier

Nestle Research Center

Imelda Angeles-Agdeppa

Food and Nutrition Research Institute

Yvonne Lenighan (✉ yvonne.lenighan@rd.nestle.com)

Nestle Research Center <https://orcid.org/0000-0001-9179-6072>

Marvin B Toledo

Food and Nutrition Research Institute

Mario V Capanzana

Food and Nutrition Research Institute

Research article

Keywords: Complementary Feeding, Infants, Children, Philippines

Posted Date: August 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-39100/v3>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on October 26th, 2020. See the published version at <https://doi.org/10.1186/s40795-020-00376-1>.

Abstract

Consumption of nutritionally adequate complementary foods is essential for optimal growth and development of infants and toddlers, particularly in developing countries. The aim of this study was to describe percentages of consumption, energy contribution (6-23.9mo), and usual nutrient intakes (12-23.9mo) from complementary foods among Filipino infants and toddlers stratified by household wealth. Data from 1087 infants and toddlers from the 2013 National Nutrition Survey were included. Dietary intake data was assessed using a 24Hr recall and population food intakes were stratified into pre-defined wealth categories. Breast milk, infant formula, powdered milk and rice were the most commonly consumed foods and beverages across the age groups. Several differences in complementary feeding by wealth status were observed. Infants and toddlers from poor households reported significantly greater consumption of human milk, in comparison to those from the rich households who reported a significantly greater consumption of infant/toddler formula ($P>0.05$). A higher percentage of toddlers from rich households consumed protein-containing foods, cookies and cakes. There was no difference in fruit and vegetable consumption between wealth groups in 6-11.9mo and 18-23.9mo children. Human milk and formula were the top contributors to energy in 6.11.9mo and 12-17.9mo children, while rice was the top energy contributor in 18-23.9mo children. Intakes of protein, fat, vitamin C, B vitamins, vitamin D, vitamin E, iron and calcium were higher in 12-23.9mo children from rich households. Interventions are required to enable caregivers of young Filipino children to provide complementary foods of high nutritional quality, particularly among children from the poor households.

Key Messages

A greater number of mothers from the poor households complied with the WHO recommendation to breastfeed, and children from the poor households were breastfed for longer than the rich households.

Milk consumption among older toddlers in the poor and middle wealth households decreased with age compared to toddlers from the rich households.

Consumption of nutrient poor foods is more common among infants and toddlers from the low wealth households compared to those in the rich households.

Background

The nutritional quality of appropriate complementary foods is an important factor in the healthy growth and development of infants and young children [1]. During this period of early childhood, dietary habits and preferences begin to form, and caregivers must pay special attention to the choices and quality of foods offered to meet nutritional needs of the growing child. However, access to a variety of affordable and nutritious complementary foods may be challenging for households of limited means. Socioeconomic status has been recognised as a determinant of health [2] and individuals with higher socioeconomic status are more likely to have access to, and consume, healthier diets [3]. Hence, guidelines for complementary feeding aimed at low- and middle-income countries encourage feeding practices that promote consumption of nutrient dense foods [1].

Despite the economic growth of the Philippines over recent years, the National Nutrition Survey (NNS) reported that among infants aged 1 year old about 36.2% of children are stunted and that 20.8% are underweight [4]. There is also a high prevalence of anaemia (40.5%) in infants aged 6 months to less than one year [4]. Indeed, studies from the Philippines found that inadequate nutrient intakes in Filipino schoolchildren were more marked among those from poor families. Infants and young children in the Philippines also had a high prevalence of inadequate nutrient intakes [5].

At present, there has been no comprehensive assessment of socioeconomic differences in complementary feeding patterns in a nationally representative sample of Filipino infants and toddlers. Regional studies conducted to date have focused on sub-populations in Cebu and urban areas [6-10] and report sub-optimal nutrient intakes during the complementary feeding period. Furthermore, an association between complementary feeding practices and anaemia, stunting, iron and vitamin A deficiencies have also been identified [7]. The aim of this study was to describe the food and beverage consumption patterns among 6-23.9 month old Filipino infants and toddlers, and the associated impact on nutrient intakes, by household wealth status. This study provides novel results on the role of household wealth in determining nutrient adequacy in this vulnerable population.

Methods

Participants

The National Nutrition Survey (NNS) is a nationally representative, cross-sectional, epidemiological survey of the health and nutritional status of the Filipino population. The survey covers all 17 regions and 80 provinces of the country and applies a three-stage sampling system which enables full geographical coverage in both rural and urban areas. Survey response rate was high (87.7%) among the 35, 825 households surveyed. This study includes 1087 infants and young children aged 6-23.9 months who participated in the 2013 NNS in the Philippines. The protocol and survey instruments were approved by the Ethics committee of the Food and Nutrition Research Institute (FNRI) and informed consent was obtained from all households that participated. Anthropometric data (height, length and weight) were measured in the participant's home, and the World Health Organization-Child Growth Standards were used to assess the nutritional status of the children, based on weight and height measurements. The socioeconomic status of the household was determined based on the possession of certain items e.g. vehicles, gadgets and household appliances. This data was then applied to classify households into wealth categories [4]. For the purpose of the food group analysis, the population was split into three categories; poor, middle and rich, however, for the in-depth analysis on usual nutrient intakes the poor and rich categories were further characterised into poor, poorest, rich and richest.

Dietary Intake Data collection

A full description of the methods of the Filipino 2013 NNS has been previously described [4, 5]. All dietary interviews were carried out in-person with the primary caregiver of the child. Trained dietitians collected data on all foods and beverages consumed using a structured 24Hr dietary recall. A single 24 hour recall is not reflective of habitual intakes of a person, however evidence suggests that two 24 hr recalls in a proportion of the population can provide an approximate measure of habitual/long-term diet. Therefore, a second 24Hr recall was collected on a non-consecutive day from 50% of the households and was used to estimate usual nutrient intakes. Usual intakes were calculated using SAS software and the validated National Cancer Institute (NCI) method to calculate usual intakes was applied to calculate means and standard deviations [11, 12]. This method has been validated [13, 14] and is considered to be reflective of long term intakes within a given time period e.g. childhood. Amounts of foods and beverages were estimated using common household measures, and were converted into grams using a portion-to-weight list developed by FNRI for the survey.

The Individual Dietary Evaluation System (IDES), developed by FNRI, was used to process the dietary records and to estimate nutrient intakes from the 24Hr recalls [5]. Quality control of the data collected occurred at both the food level (amounts reported and coding to appropriate food groups) and at nutrient level to exclude implausible intakes [15]. Breast milk consumption was estimated using the child's age and assigned volumes as in other national nutrition surveys of the infant population [16]. Foods were assigned to a food grouping system consisting of 85 food groups

similar to those used in other dietary intake surveys of the infant and toddler population but adapted to include foods common in the local food culture [16].

Statistical methods

The percentage of infants and young children consuming each food group was calculated by wealth category, as previously described [5]. The percent contribution of energy from food groups was calculated, at the population level, by summing the energy provided by each food group and dividing by the total energy intake by age and wealth category. Chi-squared tests were applied to examine the relationships between age, household wealth category and food group consumption. Usual nutrient intakes were estimated using the PC-SIDE software (Version 1.0) developed by Iowa State University (Ames, IA, USA) and data were analysed using Stata Version 13 (StataCorp, College Station, TX, USA). Since dietary variety increases at 12 months of age when children are eating more foods of the family table, the decision was taken to examine nutrient intakes starting from 12 months and also to examine, more closely, usual nutrient intakes by household wealth status.

Results

The characteristics of the study population are presented in Table 1. The population was split almost equally by gender and age. In terms of wealth status the population was divided as follows; 50% from poor households; 20% from middle-class households and 30% from rich households. Additionally, approximately 54% of the children were from urban areas, and 46% resided from a rural setting. Approximately 5% of the children were underweight, 5-11% stunted, 2-4% wasted, and 4% overweight.

Milk types by age and household wealth category.

The percent of consumers by milk types and by wealth category is presented in Table 2.

Among 6-11.9 month old infants, there were significant differences by wealth status. Breast milk was consumed by 69% of infants from the poor households, compared to 42% in the rich households ($P<0.001$). In contrast, infant formula was consumed by 22% of 6-11.9 month old infants from the poor households and 56% of children from rich households ($P<0.001$). Breast milk was consumed by 54% of 12-17.9 month olds from the poor households, compared to 24% from the rich households ($P<0.001$). Powdered milk was consumed by 34% of children from the poor households in this age group, compared to 15% from the rich households ($P=0.001$). Toddler/preschooler formula was consumed by 10% of 12-17.9 month olds from the poor households, and 47% from the rich households ($P<0.001$). In the older age group (18-23.9 month olds), the percentage of milk (all types) consumers in the poor and middle households were lower than those from rich households ($P<0.001$). There was a wide variation in breast milk consumption in 18-23.9 month old children, ranging from 35% in children from the poor households to 19% from the rich households. Similar to the younger group, toddler/preschooler formula was consumed by 7% of 18-23.9 month olds from the poor households, compared to 43% from the rich households ($P<0.001$).

Food groups by age and household wealth category.

There were notable differences in food group consumption by wealth status among 6-11.9 month old infants (Table 3). Infant cereal (grain-based cereal with nutrient fortification) was consumed by 7% of 6-11.9 month old infants from the poor households, compared to 23% of infants from the middle-class households and 19% from the rich households ($P=0.001$). Among 12-17.9 month old children, consumption of eggs was greater in middle and rich households ($P=0.015$). Consumption of cookies and cakes was higher in 12-17.9 month old children from rich households, while sugar sweetened beverage (SSBs) consumption was higher in infants from the poor households ($P<0.05$). There was no

significant difference in rice consumption between the age or wealth groups, however consumption of bread and noodles were greater among 12-17.9 month old toddlers from the rich households, compared to the poor households ($P<0.05$). Protein-rich foods were more likely to be consumed by 12-17.9 month old toddlers from the rich households; pork (13% rich, 2% poor), sausages/luncheon meat (18% rich, 6% poor) and eggs (27% rich, 13% poor) ($P<0.05$). Vegetable consumption was significantly greater among toddlers from the rich households ($P=0.021$). While, percentage consumption of SSBs was greater among 12-17.9 month olds from the poor households (24%) compared to the rich households (13%) ($P=0.002$). Similar consumption patterns of rice and grits were observed among the 12-17.9 month old and 18-23.9 month old toddlers. A greater percentage of 18-23.9 month old toddlers from richer households consumed sausages/luncheon meat compared to those from middle-income and poorer households ($P=0.002$). There was an apparent difference in consumption of cookies ($P=0.017$) and cakes ($P=0.028$) also, with 38% of 18-23.9 month old toddlers from middle households consuming cookies and 18% of toddlers from the rich households consuming cakes.

Total daily energy intakes by age and household wealth category.

The contribution of food to daily energy intakes by wealth status are presented in Figure 1. Among 6-11.9 month olds, breast milk was the greatest contributor to energy intakes in infants from the poor households (44%). In contrast, infant formula provided the greatest energy contribution in 6-11.9 month old infants from the rich households (50%). Rice provided 16% of energy intakes in poor households, compared to 8% in infants from rich households, while infant cereal provided greater contributions in 6-11.9 month old infants from rich households than the poorer households.

Similar to the 6-11.9 month old infants, breast milk (27%) and rice (23%) were the greatest contributors to energy intakes in 12-17.9 month old toddlers from the poor households, while toddler/preschooler formula (26%) and rice (16%) were the greatest contributors to energy intakes in toddlers from rich households. Powdered milk contributed to 16% of energy intakes in 12-17.9 month old toddlers in the poor households, compared to 8% of intakes in the rich households, and infant formula contributed to 11% of energy in the rich households, compared to 3% in the poor households. Rice provided the greatest energy contributions in 18-23.9 month old toddlers from the poor households (32%), with toddler formula providing the greatest contributions in the rich households (19%). Breast milk (16%) and powdered milks (11%) were the second and third highest contributors in the poor households, compared to rice (18%) and powdered milks (15%) in the rich households. Bread and noodles contributed to energy intakes across the three wealth groups.

Usual nutrient intakes in 12-23.9mo toddlers by household wealth category.

The usual nutrient intakes of 12-23.9mo toddlers are presented in Table 4. Protein and total fat intakes were higher in toddlers from the rich and richest households, both in grams/day and as a percentage of total energy. Intakes of vitamin C, riboflavin, niacin, vitamin B₆, folate, vitamin B₁₂, vitamin D and vitamin E were higher among toddlers from the wealthiest households. Similarly, intakes of calcium, iron, magnesium and potassium were higher in toddlers from the richest households.

Discussion

The objective of this analysis was to describe the food and beverage consumption patterns among 6-23.9 month old Filipino infants and toddlers, and the associated impact on nutrient intakes, by household wealth status. The analysis highlights differences in food group consumption patterns, according to wealth status and provides insight into complementary feeding behaviours that may drive inequities in nutrient intake in infants and toddlers in the Philippines.

The prevalence of breastmilk consumption in this population has been previously reported by Denney *et al.* as 60% in 6-11.9 month old infants and 37% of 12-23.9 month old toddlers [5]. This analysis showed that there was higher breastmilk consumption among infants and toddlers from poorer households (Infants; Poor: 69%, Rich: 42%), and as a result breastmilk had a greater contribution to energy among these infants and toddlers (Infants; Poor: 44% Rich: 21%). Overall, the prevalence of breastfeeding in Filipino infants is sub-optimal and does not adhere to the guidelines of continued, on-demand breastfeeding until 2 years of age and beyond [1]. Nonetheless, the situation is better than that reported in national surveys from other geographies e.g. China (44% 0-5.9mo) [17] and Mexico (15% 0-3.9mo) [18]. Public health strategies, including educational campaigns and the availability of lactation consultants, are required to further increase the rates of breastfeeding among Filipino mothers.

In the current study, breast milk contributes 7-40% of energy intakes of infants and toddlers between 6-24 months, depending on the child's stage and wealth status, with infant/toddler formula providing from 1-50% of energy intakes. Additionally, rice contributed from 8-23% of energy intakes in this population. However, breast milk and rice are low in minerals such as iron and zinc, and infant need complementary foods, in particular, those that provide iron, zinc and calcium, from the age of 6 months [19, 20]. Infants typically consume relatively small amounts of complementary foods between 6-11.9 months, therefore, the nutrient density of these foods needs to be high [21]. The current analysis suggests that consumption of nutrient dense food at this age is low, therefore educational strategies are needed to inform mothers of the importance of consuming nutrient dense foods at this early life stage. An extensive study examining Filipino infant and child feeding (6-23 months of age) in urban areas reported that children who are breastfed were more likely to be anaemic or iron deficient after 6 months of age than those receiving fortified foods [7]. This is similar to results reported in other countries including India, China and Japan [22-24]. Fortified young child beverages provide a source of iron among Filipino infants (6-11.9mo) and toddlers (12-23.9mo), respectively [5]. However, consumption of these products was more prevalent in infants and toddlers from higher income households. This may, in part, explain some of the differences in nutrient intakes according to household wealth category.

Rice is the most commonly consumed (94.8%) food in the Philippines, and is also the primary source of energy, protein, iron, thiamin and niacin [4]. In the current study the percentage contribution of rice to total daily energy, was consistently higher among children from the poorest households in infants and toddlers. This was reflected by poorer intakes of vitamins and minerals in toddlers from poorer backgrounds. However, a recent paper by Denney *et al.* in the same population, identified inadequate of protein and B vitamins in Filipino infants and toddlers at population level [5]. Ideally, the mandatory rice with vitamins and minerals in the Philippines could provide a public health opportunity to improve nutrient intakes, particularly among children from poorer households. For example, a study among school kids revealed significant reduction in anaemia after a 120 day supervised fortified lunch feeding program [25]. Unfortified complementary foods that are plant-based generally do not provide sufficient key nutrients such as iron, zinc and calcium in adequate amounts for very young children in developing countries [26]. The current analysis identified low intakes of iron, zinc and calcium in young Filipino children, particularly in those from poor households. Therefore, initiatives to support fortification of commonly consumed foods could help to close this gap in nutrient intakes.

In general, infant cereal consumption is low in the Philippines, with the exception of a sub-population from the rich households. In the current study, iron-fortified infant cereal, as recommended in complementary feeding by the American Academy of Paediatricians [27], was lacking from the diet, except among 6% of 6-11.9mo infants from the middle and rich households. The current study identified a very limited diversity of foods in the diets of Filipino infants and children, with less than 5 foods making up 80% of the diet in 6-11.9 month olds and 60% of the diet in 18-23.9 month olds. Nutrient-dense protein foods such as eggs, pork and chicken were more likely to be consumed by children from richest households. Whereas, fish was more likely to be consumed by children from low wealth households. Improving access to nutritious or fortified foods for children, particularly breast-fed infants such as infant cereal, could

help to improve micronutrient intakes. However, it must be noted, that even in the U.S., iron and zinc were identified as problem nutrients [28] despite the widespread availability of fortified foods for infants. Therefore, ensuring adequate nutrient supply in complementary feeding is clearly complex and requires further detailed study in order to examine *how* young children can meet their nutrient needs in these challenging circumstances.

In the current study, SSBs were more likely to be consumed by infants and young toddlers from the poorest households. However, SSBs were consumed at approximately equal prevalence across wealth categories in 18-23.9 month olds. Since the nutrient needs, according to body weight, of infants and young children are very high [27] there is typically no room in the diet for nutrient-poor foods such as SSBs. In the current study, infants were more likely to consume fruit juice and powdered chocolate milk than other types of SSBs. In order to help combat the nutritional inadequacies in the Philippines, fruit juice and chocolate milk are often fortified and have been shown to be among the top sources of micronutrients in the diets of Filipino children [5]. Fortified fruit juice has been effective at reducing the basal level of iron deficient anaemia in Filipino school children from 100% to 13% [29, 30]. However, it must be noted that unpublished data from the most recent Philippines National Nutrition Survey shows that the prevalence of overweight has increased in 3-5 year old children between 2015 and 2018, with a similar trend observed in school-aged children (Preliminary Results National Nutrition Survey 2018, Food and Nutrition Research Institute). Evidence suggests that excess weight gain in infancy, of which consumption of a poor diet (low in fruit and vegetables and high in sugar) or a high calorie diet, in early childhood could be a contributing factor, has been associated with increased risk of developing obesity and associated comorbidities in later life [31-33]. The current study demonstrated that the diet of Filipino infants is rich in carbohydrates, with limited protein intakes and low adherence to the local Filipino *Pinggang Pinoy* recommendation for frequent consumption of fruit and vegetables. This may be due to a number of factors including high cost, limited access to fresh fruit and vegetables, poor nutrition knowledge and family members' food preferences [34]. Therefore, the importance of fruit and vegetables, meat, poultry, fish and eggs, should be emphasised in Filipino households, regardless of wealth status. Public health authorities may wish to investigate methods to improve accessibility to such foods, particularly among low-income households.

Conclusion

Strategies aiming to make fortified foods affordable and accessible to low-income populations, and the widespread use of vitamin and mineral supplements of young children and their breastfeeding mothers could help to improve nutrient intakes. This study provided a detailed assessment of nutrient intakes in this population according to wealth category, but additional assessments of foods available in the local food supply would be required to determine if food fortification or supplementation would be the most appropriate strategy for meeting nutrient gaps. In addition, environmental and conservational issues need to be considered when evaluating nutrient sources for low-income and vulnerable populations in the Philippines.

In light of the high rates of anaemia and inadequate nutrient intakes among Filipino infants, future research should focus on determining how best to meet the nutritional needs of young Filipino children. Further work is needed to identify potential targets for food fortification, and to clarify the impact of fortification (or supplementation) with single or multiple nutrients is needed. Diet modelling studies on the impact of fortification of commonly consumed foods or supplementation may shed light on the feasibility of potential strategies aimed at improving nutrient intakes, particularly among infants and toddlers from poor households.

Declarations

Ethics approval and consent to participate: The protocol and survey instruments were approved by the Ethics committee of the Food and Nutrition Research Institute (FNRI) and written informed consent was obtained from all

households that participated (through parent / guardian for children under 10 years old). Written informed consent was obtained from a parent/guardian for participants under 16 years old.

Consent for publication: NA

Availability of data and materials: The datasets supporting the conclusions of the study are included in the article. Any additional data will be available on request. The datasets used and/or analyzed during the current study are available from the FNRI on reasonable request.

Competing interests: E.F.J. and Y.M.L. are employees of Nestec. Ltd. (Nestlé Institute of Health Sciences), Lausanne, Switzerland. The opinions expressed in the article are those of the authors alone and do not necessarily reflect the views and recommendations of their affiliations.

Funding: The National Nutrition Survey including was funded by the Government of the Philippines. This included design of the study and collection and analysis of the data. This analysis of food consumption by wealth status, interpretation of the data and writing of the manuscript was funded by the Nestlé Institute of Health Sciences, Lausanne, Switzerland.

Authors' contributions E.F.J designed the analysis, interpreted the data, wrote the manuscript and has the final responsibility for the contents. M.B.T conducted the analysis. Y.M.L and I.A.A, contributed to the data analysis, data interpretation and manuscript editing. M.V.C. contributed to the critical review and editing of the manuscript. All authors read and finally approved this manuscript for submission.

Acknowledgements: We acknowledge the contribution of Royce Ann Octavio, Kristine Biona and Regina Rodriguez for building food composition tables and food groups, along with Mark Lester Cayadong, and Glen Melvin Gironella for work in data management and programming. We would also like to thank the field workers who assisted in data collection and the caregivers and children who participated in the study. We also thank Liya Denney for her guidance.

Abbreviations

24Hr: 24 hour, NNS: National Nutrition Survey, FNRI: Food and Nutrition Research Institute, IDES: Individual Dietary Evaluation System, SSBs; sugar-sweetened beverages

References

1. World Health Organisation. Guiding principles for complementary feeding of the breastfed child. 2004. https://www.who.int/nutrition/publications/guiding_principles_compfeeding_breastfed.pdf?ua=1
2. Mayen AL, Marques-Vidal P, Paccaud F, Bovet P, Stringhini S. Socioeconomic determinants of dietary patterns in low- and middle-income countries: a systematic review. *Am J Clin Nutr.* 2014;100(6):1520-31.
3. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr.* 2008;87(5):1107-17.
4. Food and Nutrition Institute. Philippine Nutrition Facts and Figures 2013: 8th National Nutrition Survey Overview. Food and Nutrition Institute, Metro Manila, Philippines; 2015.
5. Denney L, Angeles-Agdeppa I, Capanzana M, Toledo M, Donohue J, Carriquiry A. Nutrient Intakes and Food Sources of Filipino Infants, Toddlers and Young Children are Inadequate: Findings from the National Nutrition Survey 2013. *Nutrients.* 2018;10(11):1730.
6. Perlas LA, Gibson RS, Adair LS. Macronutrient and selected vitamin intakes from complementary foods of infants and toddlers from Cebu, Philippines. *Int J Food Sci Nutr.* 2004; 55(1):1-15.

7. Rohner F, Bradley AW, Grant JA, Elizabeth AY, Lebanan MAO, Rayco-Solon P, et al. Infant and young child feeding practices in urban Philippines and their associations with stunting, anemia, and deficiencies of iron and vitamin A. *Food and Nutrition Bulletin*. 2013;34(2_suppl1):S17-S34.
8. Wright M, Bentley M, Mendez M, Adair L. The interactive association of dietary diversity scores and breast-feeding status with weight and length in Filipino infants aged 6–24 months. *Public Health Nutr*. 2015;10(1762-1773).
9. Kennedy GL, Pedro MR, Seghieri C, Brouwer I, Nantel G. Dietary Diversity Score Is a Useful Indicator of Micronutrient Intake in Non-Breast-Feeding Filipino Children. *J Nutr*. 2007;137(2):472-7.
10. Tengco LW, Rayco-Solon P, Solon JA, Sarol Jr JN, Solon FS. Determinants of anemia among preschool children in the Philippines. *J Am Coll Nutr*. 2008;27(2):229-43.
11. Toozé JA, Midthune D, Dodd KW, Freedman LS, Krebs-Smith SM, Subar AF, et al. A new statistical method for estimating the usual intake of episodically consumed foods with application to their distribution. *J Am Diet Assoc*. 2006;106(10):1575-87.
12. Dodd KW, Guenther PM, Freedman LS, Subar AF, Kipnis V, Midthune D, et al. Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. *J Am Diet Assoc*. 2006;106(10):1640-50.
13. Toozé JA, Kipnis V, Buckman DW, Carroll RJ, Freedman LS, Guenther PM, et al. A mixed-effects model approach for estimating the distribution of usual intake of nutrients: the NCI method. *Stat Med*. 2010;29(27):2857-68.
14. Freedman LS, Guenther PM, Dodd KW, Krebs-Smith SM, Midthune D. The population distribution of ratios of usual intakes of dietary components that are consumed every day can be estimated from repeated 24-hour recalls. *J Nutr*. 2010;140(1):111-6.
15. López-Olmedo N, Carriquiry AL, Rodríguez-Ramírez S, Ramírez-Silva I, Espinosa-Montero J, Hernández-Barrera L, et al. Usual Intake of Added Sugars and Saturated Fats Is High while Dietary Fiber Is Low in the Mexican Population—4. *J Nutr*. 2016;146(9):1856S-65S.
16. Roess AA, Jacquier EF, Catellier DJ, Carvalho R, Lutes AC, Anater AS, et al. Food consumption patterns of infants and toddlers: findings from the Feeding Infants and Toddlers Study (FITS) 2016. *J Nutr*. 2018;148(suppl_3):1525S-35S.
17. Hong Qin LZ, Lingling Zhang, Wei Zhang, Li Li, Xin Deng, Danping Tian, Jing Deng, Guoqing Hu. Prevalence of Breastfeeding: Findings from the First Health Service Household Interview in Hunan Province, China. *Int J Environ Res Public Health*. 2017;14:150.
18. Deming DM, Afeiche MC, Reidy KC, Eldridge AL, Villalpando-Carrión S. Early feeding patterns among Mexican babies: findings from the 2012 National Health and Nutrition Survey and implications for health and obesity prevention. *BMC Nutrition*. 2015;1(1):40.
19. World Health Organization/United Nation Children's Fund. *Global Strategy for Infant and Young Child Feeding*. Geneva: WHO Press 2003. <https://apps.who.int/iris/bitstream/handle/10665/42590/9241562218.pdf?sequence=1>
20. Dewey KG. Nutrition, growth, and complementary feeding of the breastfed infant. *Pediatric Clinics of North America*. 2001;48(1):87-104.
21. United Nations Children's Fund. *Programming Guide: Infant and Young Child Feeding*. New York UNICEF; 2011. https://www.unicef.org/nutrition/files/Final_IYCF_programming_guide_2011.pdf
22. Chandyo RK, Henjum S, Ulak M, Thorne-Lyman AL, Ulvik RJ, Shrestha PS, et al. The prevalence of anemia and iron deficiency is more common in breastfed infants than their mothers in Bhaktapur, Nepal. *Eur J Clin Nutr*. 2016;70(4):456-62.
23. Clark KM, Li M, Zhu B, Liang F, Shao J, Zhang Y, et al. Breastfeeding, mixed, or formula feeding at 9 months of age and the prevalence of iron deficiency and iron deficiency anemia in two cohorts of infants in China. *J Pediatrics*. 2017;181:56-61.

24. Amano I, Murakami A. Prevalence of infant and maternal anemia during the lactation period in Japan. *Pediatrics Intl.* 2019;61(5):495-503.
25. Angeles-Agdeppa, Capanzana, Barba, Florentino, Takanashi. Efficacy of iron-fortified rice in reducing anemia among schoolchildren in the Philippines. *Int J Vitam Nutr Res.* 2008;78(2):74-86.
26. Motuma Adimasu Abeshu AL, Bekesho Geleta. Complementary Feeding: Review of Recommendations, Feeding Practices, and Adequacy of Homemade Complementary Food Preparations in Developing Countries – Lessons from ethiopia. *Front Nutr* 2016;3:41.
27. American Academy of Pediatrics. *Pediatric Nutrition.*, American Academy of Pediatrics: Itasca I, USA, ; 2014.
28. Roess AA, Jacquier EF, Catellier DJ, Carvalho R, Lutes AC, Anater AS, et al. Food consumption patterns of Infants and Toddlers: Findings from the Feeding Infants and Toddlers Study (FITS) 2016. *J Nutr.* 2018;152:5S.
29. FAO. The State of Food Security and Nutrition. 2019 <http://www.fao.org/3/ca5162en/ca5162en.pdf>
30. Imelda Angeles-Agdeppa CRM, Mario V Capanzana Fortified juice drink improved iron and zinc status of schoolchildren. *Asia Pac J Clin Nutr.* 2011;20(4):535-43.
31. Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, et al. Early life risk factors for obesity in childhood: cohort study. *BMJ.* 2005;330(7504):1357.
32. Rooney BL, Mathiason MA, Schauburger CW. Predictors of obesity in childhood, adolescence, and adulthood in a birth cohort. *Mater Child Health J.* 2011;15(8):1166-75.
33. Baidal JAW, Locks LM, Cheng ER, Blake-Lamb TL, Perkins ME, Taveras EM. Risk factors for childhood obesity in the first 1,000 days: a systematic review. *Am J Preventive Med.* 2016;50(6):761-79.
34. Marilou M. Lopez-Madrid CCSA, Consuelo L. Orense, Charmaine A. Duante, Robby Carlo A. Tan, Mario V. Capanzana. Awareness of and Adherence to the Food Based Dietary Guidelines Among Household Meal Planners in the Philippines. *Philippine J Sci.* 2018;147(3):523-35.

Tables

Table 1. Sample demographics of the 6-23.9 month Filipino study population

		n	%
Gender (all)	Boy	544	50.1
	Girl	543	49.9
Age	6-11.9 months old	362	33.0
	12-17.9 months old	360	33.0
	18-23.9 months old	373	34.0
Wealth categories (all)	Poor	528	50
	Middle	213	20.1
	Rich	320	29.9
Region (all)	Urban	592	54.0
	Rural	504	45.9
Underweight	6-11.9 months old	54	5.0
	12-17.9 months old	65	6.1
	18-23.9 months old	63	5.8
Stunting	6-11.9 months old	52	4.9
	12-17.9 months old	81	7.6
	18-23.9 months old	117	11.0
Wasting	6-11.9 months old	43	4.0
	12-17.9 months old	24	2.2
	18-23.9 months old	26	2.4
Normal weight	6-11.9 months old	199	55.0
	12-17.9 months old	177	49.2
	18-23.9 months old	151	40.5
Overweight	6-11.9 months old	14	1.3
	12-17.9 months old	13	1.2
	18-23.9 months old	16	1.5

Table 2. Milk consumption (expressed as percent consuming) of Filipino 6-23 month olds, by wealth category.

	6-11.9mo			P [†]	12-17.9			p [†]	18-23.9			p [†]
	Poorest	Middle	Richest		Poorest	Middle	Richest		Poorest	Middle	Richest	
	n=170	n=71	n=105		n=183	n=68	n=101		n=175	n=74	n=114	
All milk	96	100	98	0.162	91	93	97	0.204	70	81	95	<0.001
Breast milk	69	63	42	<0.001	54	44	24	<0.001	35	27	19	0.061
Powdered milks	17	24	18	0.449	34	19	15	0.001	34	42	29	0.242
Formula	25	31	62	<0.001	13	40	61	<0.001	7	23	48	<0.001
Infant	22	31	56	<0.001	3	9	15	0.001	0	1	5	0.076
Toddler/preschooler	4	0	7	0.088	10	31	47	<0.001	7	22	43	<0.001

[†]Chi-square test for independence between wealth groups, P<0.05 considered significant

Table 3. Food consumption (expressed as percent consuming) of Filipino 6-23.9 month olds, by wealth category.

	6-11.9mo				12-17.9mo				18-23.9mo			
	Poor	Middle	Rich	p†	Poor	Middle	Rich	p†	Poor	Middle	Rich	p†
	n=170	n=71	n=105		n=183	n=68	n=101		n=175	n=74	n=114	
Rice	72	68	66	0.548	89	85	91	0.545	86	93	92	0.272
Grain-based mixed dishes	11	11	6	0.279	9	19	10	0.065	4	10	9	0.403
Corn grits	7	3	2	0.102	8	4	2	0.096	11	3	2	0.014
Infant cereal	7	23	19	0.001	0	2	4	0.022	0	0	0	-
Bread	7	4	12	0.118	11	18	25	0.011	23	30	27	0.639
Noodles	7	3	5	0.388	11	21	27	0.011	26	23	18	0.469
Vegetables	11	9	13	0.603	17	27	31	0.021	34	31	27	0.630
Fruits	9	7	2	0.071	14	10	15	0.621	16	14	18	0.741
Fish	12	9	8	0.484	29	34	21	0.146	45	41	33	0.247
Eggs	11	6	11	0.372	13	22	27	0.015	18	16	25	0.314
Sausages/luncheon meats	4	4	4	0.988	6	13	18	0.009	6	12	25	0.002
Pork	4	1	6	0.363	2	7	13	0.001	8	12	18	0.206
Chicken	2	1	2	0.969	6	7	12	0.172	7	10	16	0.186
Cookies	21	21	35	0.022	24	32	24	0.360	18	38	22	0.017
Sugar sweetened beverages (SSBs)	9	3	5	0.159	24	6	13	0.002	20	24	23	0.813
Table sugar	2	1	2	0.969	4	4	2	0.661	5	8	6	0.824
Crackers	7	17	6	0.020	18	19	21	0.871	18	19	14	0.589
Cakes	2	6	10	0.007	9	4	12	0.209	5	8	18	0.028

†Chisquare test for independence between wealth groups, P<0.05 considered significant

Table 4. Usual nutrient intakes in a subpopulation of Filipino 12-23.9 month old children, by wealth category.

	Poorest n=208	Poor n=141	Middle n=139	Rich n=116	Richest n=92
Carbohydrate (g/d)	108.2 ± 3.3	109.2 ± 4.1	112.4 ± 4.2	114 ± 4.3	116 ± 4.4
Total sugars (g/d)	31.3 ± 1.3	33 ± 1.6	30.8 ± 1.5	33.7 ± 2.6	29.3 ± 2.5
Dietary fiber (g/d)	2.3 ± 0.1	2.2 ± 0.1	2.3 ± 0.1	2.4 ± 0.1	2.4 ± 0.2
Protein (g/d)	19 ± 0.6	22.8 ± 1	21.8 ± 0.8	27.5 ± 1.3	27.7 ± 1.4
Total fat (g/d)	19.7 ± 0.6	25 ± 1	24 ± 0.9	30.5 ± 1.5	31.6 ± 1.5
SFA (g)	8.7 ± 2.6	8 ± 0.4	8 ± 0.4	9.4 ± 0.5	8.1 ± 0.6
MUFA (g)	5.6 ± 0.2	8.4 ± 0.6	6.4 ± 0.4	9.7 ± 0.8	6.9 ± 0.6
PUFA (g)	1.7 ± 0.1	2.7 ± 0.2	2.1 ± 0.1	2.9 ± 0.2	2.2 ± 0.2
As percentage of total energy					
Carbohydrate (%)	62.5 ± 0.7	58 ± 0.7	59.8 ± 0.6	55.2 ± 0.9	54.5 ± 0.9
Protein (%)	11.2 ± 0.2	11.8 ± 0.2	11.4 ± 0.2	12.8 ± 0.3	12.6 ± 0.3
Total Fat (%)	26.2 ± 0.7	29.9 ± 0.6	29.7 ± 0.6	31.9 ± 0.8	32.6 ± 0.7
Vitamins					
Vitamin A (µgRE/d)	195.7 ± 12.2	315 ± 31.1	197.1 ± 18.6	335.4 ± 40	159.8 ± 27.2
Vitamin C (mg/d)	21.2 ± 1	23.7 ± 1.8	32.2 ± 2.9	37.7 ± 3	55.7 ± 4.4
Thiamine (mg/d)	0.3 ± 0.02	0.4 ± 0.02	0.4 ± 0.03	0.5 ± 0.03	0.7 ± 0.05
Riboflavin (mg/d)	0.5 ± 0.03	0.8 ± 0.07	0.7 ± 0.1	1.1 ± 0.1	1.2 ± 0.1
Niacin (mg/d)	4.2 ± 0.1	4.4 ± 0.2	5.3 ± 0.3	5.8 ± 0.3	7.7 ± 0.5
Vitamin B6 (mg)	0.3 ± 0.02	0.6 ± 0.1	0.5 ± 0.03	0.6 ± 0.05	1.1 ± 0.1
Folate (DFE µg)	92.9 ± 6.2	85.4 ± 4.7	118.6 ± 9.3	107.8 ± 8.2	161.9 ± 14
Vitamin B12 (mg)	0.9 ± 0.04	0.8 ± 0.1	1.1 ± 0.1	1.2 ± 0.1	1.6 ± 0.1
Vitamin D (mg)	1.2 ± 0.1	1.4 ± 0.2	2.5 ± 0.2	2.7 ± 0.3	5.2 ± 0.7
Vitamin E (mg)	1.5 ± 0.1	1.5 ± 0.1	2.5 ± 0.3	2.1 ± 0.2	3.6 ± 0.4
Minerals					
Calcium (mg/d)	291.4 ± 16.5	447.3 ± 35.1	413.8 ± 30	632.9 ± 52.7	768.2 ± 62.7
Phosphorus (mg/d)	311.7 ± 13.4	453.2 ± 30.5	389.8 ± 21.8	544.1 ± 40.1	491.6 ± 33.1
Iron (mg/d)	3.2 ± 0.1	3.5 ± 0.2	4.6 ± 0.3	5.7 ± 0.4	9 ± 0.8
Sodium (mg/d)	469.5 ± 23.7	504.6 ± 21.2	474.2 ± 27	576.7 ± 32.4	555.7 ± 33.9
Zinc (mg)	3.1 ± 0.2	5.3 ± 1	3.8 ± 0.2	3.7 ± 0.2	5.2 ± 0.4
Magnesium (mg)	49 ± 1.9	49.2 ± 2.3	57.5 ± 3	63.5 ± 4.2	73.4 ± 4.8
Potassium (mg)	456.5 ± 153	441.8 ± 16.1	562.7 ± 25.7	606.3 ± 45.6	693.1 ± 47
Selenium (mg)	31.2 ± 1.3	27.3 ± 1.3	32.5 ± 1.7	31.8 ± 1.6	30 ± 1.9

Figures




Figure 1

Top 5 contributors to energy intakes in Filipino infants and toddlers