

Propofol attenuated the effect of TNF- α on occludin expression by inhibiting HIF-1 α /VEGF/VEGFR-2/ERK signaling pathway in hCMEC/D3 cells

Yue Zhang

Fudan University Shanghai Cancer Center

Xiaowei Ding

Fudan University Shanghai Cancer Center

Changhong Miao

Fudan University Shanghai Cancer Center

Jiawei Chen (✉ jiawei_chen@hotmail.com)

Fudan University Shanghai Cancer Center <https://orcid.org/0000-0003-3726-9823>

Research article

Keywords: TNF- α , propofol, occludin, hCMEC/D3 cells

Posted Date: June 11th, 2019

DOI: <https://doi.org/10.21203/rs.2.409/v5>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 9th, 2019. See the published version at <https://doi.org/10.1186/s12871-019-0788-5>.

Abstract

Purpose: The levels of tight junction proteins (TJs), especially occludin, correlate with blood-brain barrier (BBB) disruption caused by inflammation in central nervous system (CNS). It has been reported that propofol, the most commonly used anesthetic, could inhibit inflammation response in CNS. In this study, we investigated the effects of tumor necrosis factor- α (TNF- α) and propofol on occludin expression in human cerebral microvascular endothelial cell line, D3 clone (hCMEC/D3 cells), and explored the underlying mechanisms. **Methods:** The hCMEC/D3 cells were treated with propofol, followed by TNF- α . The expression and phosphorylation of Hif-1 α , VEGF, VEGFR-2, ERK, p38MAPK and occludin were measured by Western blot analysis. The cell viability of hCMEC/D3 cells was measured by cell counting kit-8. **Results:** TNF- α (10 ng/ml, 4 h) significantly decreased the expression of occludin, which was attenuated by propofol (25 μ M). TNF- α induced Hif-1 α /VEGF/VEGFR-2/ERK signaling pathway, while propofol could inhibit it. TNF- α induced the phosphorylation of p38MAPK, while propofol had no effect on it. In addition, the inhibitors of Hif-1 α , VEGF, VEGFR-2, and ERK could reduce the effect of TNF- α on occludin expression. **Conclusion:** TNF- α could decrease the expression of occludin via activating Hif-1 α /VEGF/VEGFR-2/ERK signaling pathway, which was attenuated by propofol.

Background

Blood-brain barrier (BBB) is a multicellular structure and mainly consists of microvascular endothelial cells, astrocytes, pericytes, intercellular tight junctions and basal lamina [1]. It is accepted that BBB can protect central nervous system (CNS) from harmful molecules and pathogens in the peripheral blood circulation. A dysfunctional or impaired BBB allows augmented permeation of macrophages, leukocytes, endotoxins and bacteria into the CNS. Studies have revealed that the disruption of BBB may be involved in neuropathological status, such as postoperative cognitive dysfunction [2], Alzheimer disease [3], and dementia [4].

Cerebral microvascular endothelial cells, one of the most important components of BBB, are linked together through multiple tight junction proteins including claudins, occludin and zonula occludens(ZOs) [5]. The tight junctions play a crucial role in the biological functions of BBB. It has been reported that the decreased expression of tight junction proteins, especially occludin, correlates with the disruption of BBB [6, 7]. A study also revealed that truncation of occludin decreased transendothelial electrical resistance, suggesting a key role of occludin in the barrier function of BBB [8].

It has been reported that BBB disruption could be induced by many factors, including neuroinflammation, oxidative stress, and hypoxia [9-11]. Recent studies showed that inflammatory reactions may be the primary cause for BBB disruption [12]. Tumor necrosis factor- α (TNF- α) is one of the most important inflammation factors and plays an essential role in neuroinflammation. It has been reported that TNF- α could decrease the expression of occludin in endothelial cells [13]. However, the underlying mechanism of TNF- α -mediated occludin expression in microvascular endothelial cells has not been fully elucidated. Hypoxia inducible factor-1 α (Hif-1 α) has been reported to be activated by TNF- α and was correlated with

occludin expression [14, 15]. However, the exact mechanism is unclear. As a direct target of Hif-1 α , vascular endothelial growth factor(VEGF) can regulate paracellular permeability via the activation of extracellular regulated protein kinases(ERK) [16]. Therefore, we sought to investigate and verify these molecules in TNF- α -mediated occludin expression.

Propofol, a widely used intravenous anesthetic agent, has been reported to own anti-inflammation effects. Studies have revealed that propofol could protect hypoxia-induced inflammation in BV2 microglial cells [17]. It was also demonstrated that propofol inhibited sevoflurane-induced inflammation in human neuroglioma cells [18]. In this study, we sought to explore whether propofol could attenuate the effect of TNF- α on occludin expression and identify the underlying mechanisms.

Materials And Methods

Cell culture and reagents

Human cerebral microvascular endothelial cell (hCMEC) line, D3 clone, was purchased from Shanghai GuanDao biological engineering company and maintained in Dulbecco's modified Eagle's medium (DMEM; Hyclone, Australia); all culture media were supplemented with 10% fetal bovine serum (Gibco, Australia), 100 mg/mL streptomycin, and 100 U/mL penicillin. The hCMEC/D3 cells were cultured in a humidified 5% CO₂ atmosphere at 37°C. The cells were passaged and cultured on reaching 80% confluence and the cell medium was changed every other day.

TNF- α powder was obtained from PeproTech China and dissolved in serum-free DMEM. Propofol (Sigma, St. Louis, MO, USA) was dissolved in dimethyl sulfoxide (DMSO; Sigma-Aldrich). The ERK inhibitor LY3214996 was purchased from Sigma Aldrich. SU5416, an inhibitor of VEGFR-2, and KC7F2, an inhibitor of Hif-1 α , were purchased from MedChemExpress (Shanghai, China). KC7F2, SU5416, LY3214996 were dissolved in DMSO, respectively. The final concentration of DMSO was adjusted to 0.01% for each solution to avoid possible non-specific effects.

Western blot analysis

After designed treatments, the cell extracts were collected and lysed with RIPA lysis buffer (Beyotime Biotechnology, Suzhou, China) supplemented with protease and phosphatase inhibitors (Roche, Rotkreuz, Switzerland). The lysates were mixed with 5 \times loading buffer and boiled for 5 min at 100°C. Equivalent amounts of protein in each sample were separated by 10% or 12% SDS-polyacrylamide gel and transferred to a polyvinylidene fluoride membrane (Millipore, Billerica, MA) using a semidry electroblotting system. The membranes were blocked with 5% skim milk in phosphate-buffered saline-Tween 20 (PBST) for 1 h at room temperature, and then incubation with appropriate primary antibody at dilution of 1:1000 at 4°C overnight. The primary antibodies included: antibodies against occludin(Proteintech, Wuhan, China), p-ERK(CST, Massachusetts, USA), ERK(CST, Massachusetts, USA), p-p38MAPK(CST, Massachusetts, USA), p38MAPK(CST, Massachusetts, USA), p-VEGFR-2(CST, Massachusetts, USA), VEGF(CST,

Massachusetts, USA), Hif-1 α (CST, Massachusetts, USA) (Abcam, Shanghai, China), β -actin(Proteintech, Wuhan, China). After that, the membranes were washed with PBST and incubated with secondary antibodies conjugated with horseradish peroxidase (HRP) for 1 h at room temperature. After washing, the signals were detected using a LAS-4000 mini CCD camera (GE Healthcare). β -actin was used as internal control and the intensity of each protein band was normalized with that of β -actin. Each assay was performed in five replicates.

Cell viability assay

The cell viability of hCMEC/D3 cells was measured by cell counting kit-8 (CCK8) (Beyotime Institute of Biotechnology, Shanghai, China). In this study, the hCMEC/D3 cells were seeded on 96-multiwell plates at a density of 5×10^3 cells/well and cultured for 24 h. After designed treatment, the culture medium was replaced by 20 ml CCK8 along with 180 ml fresh culture medium and incubated in 37°C for 2 h. The absorbance at a 450 nm wavelength was measured by a microplate reader (Synergy H4, Bio-Tek) and the cell viability curve was determined by calculating the mean value and standard deviation (SD) of the optical density for every 6 wells.

Statistical analysis

Data were expressed as mean \pm SD, and results were obtained from at least 5 separately performed experiments. Differences between groups were determined by one-way analysis of variance (ANOVA) followed by the Newman–Keuls test. InStat statistical program (GraphPad Software, San Diego, CA, USA) was used for the statistical analyses. All results in this study were considered statistically significant at a value of $p < 0.05$.

Results

TNF- α decreased the expression of occludin in hCMEC/D3 cells.

As shown in Fig 1A, we treated different concentrations of TNF- α (1, 10, 25, 50, 100ng/ml) for 8h, and observed that 1ng/ml TNF- α had no effect on occludin expression, while 10, 25, 50, 100 ng/ml TNF- α all significantly decreased occludin expression. It is noted that 10ng/ml was the minimal dose of TNF- α to exert significant effect on occludin expression. Then we treated cells with 10ng/ml TNF- α for different times (2, 4, 8, 12, 24 h), and found that 2h treatment of TNF- α had no effect on occludin expression, while 4, 8, 12, 24 h treatment of TNF- α greatly decreased occludin expression (Fig. 1B). It is noted that 4 h was the minimal treatment time of TNF- α to exert significant effect on occludin expression. As shown in Fig. 1C, the cell viability was not changed by TNF- α at different concentrations (1, 10, 25, 50, 100ng/ml) or

different times (2, 4, 8, 12, 24 h). Then, we treated cells with 10 ng/ml TNF- α for 4 h in follow-up experiments to explore the underlying mechanism.

Propofol inhibited the effect of TNF- α on occludin in hCMEC/D3 cells.

As shown in Fig. 2A, we pretreated hCMEC/D3 cells with different concentrations of propofol (5, 10, 25, 50 μ M) for 2 h and followed by TNF- α treatment (10 ng/ml, 4 h). We found 5 or 10 μ M propofol pretreatment had no effect on TNF- α -modulated occludin expression, while 25 and 50 μ M propofol pretreatment significantly attenuated the TNF- α -modulated occludin expression. Please noted that 25 μ M was the minimal dose of propofol to show significant effect. As shown in Fig. 2B, we detected that DMSO (0.01%), the solvent of propofol, had no effect on TNF- α -modulated occludin expression or basal occludin expression. In addition, we found that 25 μ M propofol pretreatment alone had no effect on basal occludin expression. Therefore, 25 μ M propofol pretreatment for 2 h was used in the following experiments to investigate the mechanisms of propofol.

TNF- α decreased the expression of ZO-1, while propofol had no effect on it.

As shown in supplemental figure, we treated hCMEC/D3 cells with different concentration of TNF- α (10, 50, 100ng/ml) for 8h, and observed that TNF- α had no effect on claudin-5 expression. In addition, we found that although TNF- α (10ng, 4h) decreased the expression of ZO-1, while 25 or 50 μ M propofol could not repress the effect of TNF- α . Thus, we decided not to pursue further with claudin-5 and ZO-1 in this study.

ERK was involved in the TNF- α and propofol-mediated occludin expression in hCMEC/D3 cells.

It has been reported that the phosphorylation of ERK and p38MAPK may be involved in the expression of occludin [19]. As shown in Fig. 3A, TNF- α (10ng, 4h) induced the phosphorylation of ERK, which was attenuated by 25 μ M propofol in hCMEC/D3 cells. TNF- α and propofol had no effect on the expression of ERK. Then, we found that the phosphorylation of p38MAPK was induced by TNF- α , while propofol had no effect on p38MAPK phosphorylation (Fig. 3A). To further confirm the role of ERK, we treated cells with the inhibitor of ERK. We found that 5 μ M LY3214996, the inhibitor of ERK, significantly attenuated the effect of TNF- α on ERK phosphorylation and occludin expression. And these effects were similar to those of propofol (Fig. 3B).

VEGF/ VEGFR-2 signaling pathway was involved in the effects of propofol and TNF- α on ERK and occludin in hCMEC/D3 cells.

It has been reported that VEGF could activate ERK pathway via the activation of VEGF receptor-2(VEGFR-2) [20]. As shown in Fig. 4A, TNF- α (10ng, 4h) induced the expression of VEGF and the phosphorylation of VEGFR-2 in hCMEC/D3 cells. Moreover, these effects were inhibited by 25 μ M propofol. To further confirm the role of VEGF/VEGFR-2 signaling pathway, we treated cells with the inhibitor of VEGFR-2. We found that 10 μ M SU5416, the inhibitor of VEGFR-2, markedly inhibited the effect of TNF- α on VEGFR-2 phosphorylation, ERK phosphorylation, and occludin expression in hCMEC/D3 cells. And these effects were similar to those of propofol (Fig. 4B).

Hif-1 α was involved in the effects of propofol and TNF- α on VEGF, VEGFR-2, ERK and occludin in hCMEC/D3 cells.

It has been reported that VEGF could be regulated by Hif-1 α [21]. As shown in Fig. 5A, TNF- α (10ng, 4h) induced the expression of Hif-1 α , which was inhibited by 25 μ M propofol. To further confirm the role of Hif-1 α , we treated cells with the inhibitor of Hif-1 α . We found that 10 μ M KC7F2, the inhibitor of Hif-1 α , significantly attenuated the effects of TNF- α on Hif-1 α expression, VEGF expression, VEGFR-2 phosphorylation, ERK phosphorylation, and occludin expression in hCMEC/D3 cells. And these effects were similar to those of propofol (Fig. 5B).

The confirmation of Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway that induced by TNF- α .

To confirmed the sequential activation of Hif-1 α , VEGF, VEGFR-2 and ERK, we treated hCMEC/D3 cells with ERK inhibitor (LY3214996), TNF- α (10ng, 4h), and 25 μ M propofol. We found that 5 μ M LY3214996, the inhibitor of ERK, had no effect on TNF- α -induced Hif-1 α expression, VEGF expression and VEGFR-2 phosphorylation (Fig. 6A). Then, we treated cells with VEGFR-2 inhibitor (SU5416), TNF- α (10ng, 4h), and 25 μ M propofol. We found that 10 μ M SU5416, the inhibitor of VEGFR-2, had no effect on TNF- α -induced Hif-1 α expression (Fig. 6B). Accordingly, we inferred that TNF- α decreased occludin expression via sequentially activating Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway in hCMEC/D3 cells.

Discussion

In this study, we observed that propofol inhibited TNF- α -modulated occludin expression. More importantly, our data implied that the Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway was involved in this process.

It is known that BBB plays a crucial role in maintaining a stable environment of the CNS. Brain microvascular endothelial cells, one of the major components of BBB, are linked by intercellular tight junction protein complexes. Tight junctions among endothelial cells limit BBB permeability [22]. The tight junction proteins mainly include claudin-5, occludin and ZO-1 [23]. It has been reported inflammation plays a key role in tight junction disruption and many inflammation factors can modulate the expression of tight junction proteins [24]. For example, TNF- α was shown to decrease occludin expression in glomerular endothelial cells of mice [25]. It has also been revealed that TNF- α induced rat pulmonary microvascular endothelial cells injury via downregulating the expression of ZO-1 [26]. A study revealed that TNF- α reduced occludin and ZO-1 expression in pulmonary microvascular endothelial cells [27]. Consistently, we found that TNF- α significantly decreased the expression of occludin and ZO-1 in hCMEC/D3 cells. Interestingly, TNF- α was shown to reduce claudin-5 expression in pulmonary microvascular endothelial cells [26, 27], while we found TNF- α had no effect on claudin-5 expression in hCEMC/D3 cells. We inferred that TNF- α may exert different effects on occludin expression in different organs. To rule out the possibility that TNF- α affected cell viability. We treated hCMEC/D3 cells with TNF- α at different concentrations (1, 10, 25, 50, 100ng/ml) and different times (2, 4, 8, 12, 24 h). And we found the cell viability was not changed by TNF- α (Fig. 1C). Accordingly, we inferred that the TNF- α -modulated occludin expression was not due to its effect on cell viability.

Propofol has been reported to own anti-inflammatory effects. It was revealed that propofol reversed TNF- α -induced human vascular endothelial cells apoptosis [28]. It has also been reported that propofol exerted anti-inflammatory effects in rats [29]. In this study, propofol inhibited lipopolysaccharide-induced acute lung injury by inhibition of cluster of differentiation (CD) 14 and Toll-like receptor (TLR) 4 expression [29]. In recent years, the anti-inflammatory effects of propofol in CNS attract widespread concerns. For example, it has been reported that propofol inhibited TNF- α -induced apoptosis in mouse hippocampal HT22 cells [30]. A study also revealed that propofol inhibited the inflammation and apoptosis of mouse microglia BV2 cells [17]. Here, we wondered if propofol attenuated the effect of TNF- α on the expression of occludin and ZO-1. As shown in figure 2, we found that the effect of TNF- α on occludin expression was markedly attenuated by 25 μ M propofol, which is within clinical plasma concentration(5-50 μ M) during the induction and maintenance of general anesthesia [31]. In addition, the supplementary data showed that propofol had no effect on TNF- α -modulated ZO-1 expression. Although it has been reported that propofol attenuated hypoxia-modulated ZO-1 expression in mouse brain microvascular endothelial cells [27, 32]. We reasoned that hypoxia and inflammation may reduce ZO-1 expression through different intracellular signaling pathway, and that propofol may have different effect on these pathways.

Hif-1 α has been reported to be involved in occludin expression [33]. In addition, the role of Hif-1 α in tight junction damage has recently become evident [14]. Although Hif-1 α was usually induced by hypoxia, studies showed many inflammation mediators induced the expression of Hif-1 α [15, 34]. Consistently, we found TNF- α increased Hif-1 α expression in hCMEC/D3 cells, while propofol repressed it. In addition, we found that the inhibitor of Hif-1 α repressed TNF- α -modulated occludin expression as propofol, suggesting Hif-1 α may play a key role in the effect of propofol and TNF- α on occludin expression.

VEGF is a homodimeric 45-kDa glycoprotein and is secreted by a variety of cells including cerebral microvascular endothelial cells [35]. It is accepted that VEGF can affect endothelial cell survival and function. It has been revealed that VEGF was involved in occludin expression [36]. The study revealed that the level of occludin was lowered by VEGF treatment and that VEGF increased BBB permeability in brain microvascular endothelial cells monolayer cultures [36]. It has also been reported that VEGF was activated by Hif-1 α in endothelial progenitor cell [37]. VEGF exerted its physiological function via interacting with receptors on vascular endothelial cells [38]. The receptors of VEGF(VEGFR) are receptor tyrosine kinases, and include VEGFR-1, VEGFR-2, and VEGFR-3 [39]. Although these receptors are highly similar in amino acid sequence, different VEGFRs have disparate functions and play key role in different pathological processes and signal transductions [40]. Previous studies indicated that VEGFR-1 was mainly involved in angiogenesis [41] and VEGFR-3 was involved in migration and invasion in cancer cells [42], while VEGFR-2 was closely involved with inflammation response [43]. It has been reported that VEGF activated VEGFR-2 to promote its auto-phosphorylation to generate p-VEGFR-2 [44], which exerted specific effects on intercellular permeability [45]. As shown in figure 4A, the expression of VEGF and the phosphorylation of VEGFR-2 could be induced by TNF- α , while propofol attenuated the effects of TNF- α . Then, we found that the inhibitor of VEGFR-2 attenuated TNF- α -modulated occludin expression. Furthermore, our data also showed that the inhibitor of Hif-1 α markedly decreased VEGF expression and VEGFR-2 phosphorylation as propofol. Our finding demonstrated that TNF- α modulated occludin expression via Hif-1 α /VEGF/VEGFR-2 signaling pathway, which was also involved in the effect of propofol.

Studies have revealed that the breakdown of BBB was due to the activation of ERK signaling pathway [46]. As one of the downstream molecules of VEGF/VEGFR-2 signaling pathway [47], ERK has been reported to be involved in the expression of occludin [48]. In our study, we found that TNF- α induced the phosphorylation of ERK, which was attenuated by propofol. And we also demonstrated that the inhibitor of ERK significantly alleviated the effect of TNF- α on occludin expression. In addition, we found the phosphorylation of ERK was repressed by the inhibitor of Hif-1 α and the inhibitor of VEGFR-2, suggesting the role of Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway in propofol- and TNF- α -modulated occludin expression.

It has been reported that p38MAPK was activated by TNF- α and played a crucial role in occludin expression [13]. As shown in Fig 3A, we found TNF- α induced the phosphorylation of p38MAPK, while propofol had no effect on it. In addition, the inhibitor of ERK almost completely blocked TNF- α -modulated occludin expression in hCMEC/D3 cells. Thus, we inferred that compared with ERK, p38MAPK plays a minor role in TNF- α -modulated occludin expression.

In conclusion, our data strongly implied that propofol exerts protective effect on TNF- α -modulated occludin expression via inhibiting Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway. We have realized this needs to be further confirmed in the animal study. Actually, we are planning to examine the protective effect of propofol on TNF- α -modulated occludin expression and BBB impairment in the mouse model.

Declarations

Ethics approval and consent to participate

Not applicable

Consent to publish

All authors read and approved the manuscript.

Availability of data and materials

All the data were presented within the manuscript and additional supporting files.

Competing interests

No conflict of interests exists.

Funding

This work was supported by Shanghai Shenkang hospital development center clinical science and technology innovation project (SHDC12018105) and by National Key R&D Program of China (No. 2018YFC2001900-04).

Authors' Contributions

Yue Zhang performed the experiments and wrote the paper.

Xiaowei Ding performed the experiments and wrote the paper.

Changhong Miao designed the study and modified the manuscript.

Jiawei Chen designed the study and modified the manuscript.

Acknowledgements

We thank Zhen Wang (Fudan University Shanghai Cancer Center, Shanghai, China) for her assistance with technical aspects of performing the experiments.

References

1. Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ: Structure and function of the blood-brain barrier. *Neurobiol Dis* 2010, 37(1):13-25.
2. Li Z, Mo N, Li L, Cao Y, Wang W, Liang Y, Deng H, Xing R, Yang L, Ni C *et al*: Surgery-Induced Hippocampal Angiotensin II Elevation Causes Blood-Brain Barrier Disruption via MMP/TIMP in Aged Rats.

Front Cell Neurosci 2016, 10:105.

3. Sweeney MD, Sagare AP, Zlokovic BV: Blood-brain barrier breakdown in Alzheimer disease and other neurodegenerative disorders. *Nat Rev Neurol* 2018, 14(3):133-150.
4. Raja R, Rosenberg GA, Caprihan A: MRI measurements of Blood-Brain Barrier function in dementia: A review of recent studies. *Neuropharmacology* 2018, 134(Pt B):259-271.
5. Luo D, Zhao J, Rong J: Plant-derived triterpene celastrol ameliorates oxygen glucose deprivation-induced disruption of endothelial barrier assembly via inducing tight junction proteins. *Phytomedicine* 2016, 23(13):1621-1628.
6. Chen JT, Lin YL, Chen TL, Tai YT, Chen CY, Chen RM: Ketamine alleviates bradykinin-induced disruption of the mouse cerebrovascular endothelial cell-constructed tight junction barrier via a calcium-mediated redistribution of occludin polymerization. *Toxicology* 2016, 368-369:142-151.
7. Tian J, Shi R, Liu T, She R, Wu Q, An J, Hao W, Soomro MH: Brain Infection by Hepatitis E Virus Probably via Damage of the Blood-Brain Barrier Due to Alterations of Tight Junction Proteins. *Frontiers in cellular and infection microbiology* 2019, 9:52.
8. Maherally Z, Fillmore HL, Tan SL, Tan SF, Jassam SA, Quack FI, Hatherell KE, Pilkington GJ: Real-time acquisition of transendothelial electrical resistance in an all-human, in vitro, 3-dimensional, blood-brain barrier model exemplifies tight-junction integrity. *FASEB J* 2018, 32(1):168-182.
9. Nathoo N, Jalal H, Natah SS, Zhang Q, Wu Y, Dunn JF: Hypoxia and Inflammation-Induced Disruptions of the Blood-Brain and Blood-Cerebrospinal Fluid Barriers Assessed Using a Novel T1-Based MRI Method. *Acta Neurochir Suppl* 2016, 121:23-28.
10. Banks WA, Gray AM, Erickson MA, Salameh TS, Damodarasamy M, Sheibani N, Meabon JS, Wing EE, Morofuji Y, Cook DG *et al*: Lipopolysaccharide-induced blood-brain barrier disruption: roles of cyclooxygenase, oxidative stress, neuroinflammation, and elements of the neurovascular unit. *J Neuroinflammation* 2015, 12:223.
11. Vutukuri R, Brunkhorst R, Kestner RI, Hansen L, Bouzas NF, Pfeilschifter J, Devraj K, Pfeilschifter W: Alteration of sphingolipid metabolism as a putative mechanism underlying LPS-induced BBB disruption. 2018, 144(2):172-185.
12. Terrando N, Eriksson LI, Ryu JK, Yang T, Monaco C, Feldmann M, Jonsson Fagerlund M, Charo IF, Akassoglou K, Maze M: Resolving postoperative neuroinflammation and cognitive decline. *Ann Neurol* 2011, 70(6):986-995.
13. Ni Y, Teng T, Li R, Simonyi A, Sun GY, Lee JC: TNFalpha alters occludin and cerebral endothelial permeability: Role of p38MAPK. *PLoS One* 2017, 12(2):e0170346.

14. Engelhardt S, Al-Ahmad AJ, Gassmann M, Ogunshola OO: Hypoxia selectively disrupts brain microvascular endothelial tight junction complexes through a hypoxia-inducible factor-1 (HIF-1) dependent mechanism. *J Cell Physiol* 2014, 229(8):1096-1105.
15. Kim KW, Lee SJ, Kim JC: TNF-alpha upregulates HIF-1alpha expression in pterygium fibroblasts and enhances their susceptibility to VEGF independent of hypoxia. *Exp Eye Res* 2017, 164:74-81.
16. Miao Z, Dong Y, Fang W, Shang D, Liu D, Zhang K, Li B, Chen YH: VEGF increases paracellular permeability in brain endothelial cells via upregulation of EphA2. *Anatomical record (Hoboken, NJ : 2007)* 2014, 297(5):964-972.
17. Lu Y, Gu Y, Ding X, Wang J, Chen J: Intracellular Ca²⁺ homeostasis and JAK1/STAT3 pathway are involved in the protective effect of propofol on BV2 microglia against hypoxia-induced inflammation and apoptosis. 2017, 12(5):e0178098.
18. Tian Y, Guo S, Guo Y, Jian L: Anesthetic Propofol Attenuates Apoptosis, Abeta Accumulation, and Inflammation Induced by Sevoflurane Through NF-kappaB Pathway in Human Neuroglioma Cells. *Cell Mol Neurobiol* 2015, 35(6):891-898.
19. Yang L, Liu G, Lian K, Qiao Y, Zhang B, Zhu X, Luo Y, Shang Y, Gu XL: Dietary leonurine hydrochloride supplementation attenuates lipopolysaccharide challenge-induced intestinal inflammation and barrier dysfunction by inhibiting the NF-kappaB/MAPK signaling pathway in broilers. *J Anim Sci* 2019, 97(4):1679-1692.
20. Park W, Baek YY, Kim J, Jo DH, Choi S, Kim JH, Kim T, Kim S, Park M, Kim JY *et al*: Arg-Leu-Tyr-Glu Suppresses Retinal Endothelial Permeability and Choroidal Neovascularization by Inhibiting the VEGF Receptor 2 Signaling Pathway. *Biomol Ther (Seoul)* 2019.
21. Dewangan J, Srivastava S, Mishra S, Divakar A, Kumar S, Rath SK: Salinomycin inhibits breast cancer progression via targeting HIF-1alpha/VEGF mediated tumor angiogenesis in vitro and in vivo. *Biochem Pharmacol* 2019, 164:326-335.
22. Liu WY, Wang ZB, Zhang LC, Wei X, Li L: Tight junction in blood-brain barrier: an overview of structure, regulation, and regulator substances. *CNS Neurosci Ther* 2012, 18(8):609-615.
23. Keaney J, Campbell M: The dynamic blood-brain barrier. *Febs j* 2015, 282(21):4067-4079.
24. Shen Y, Zhou M, Yan J, Gong Z, Xiao Y, Zhang C, Du P, Chen Y: miR-200b inhibits TNF-alpha-induced IL-8 secretion and tight junction disruption of intestinal epithelial cells in vitro. *Am J Physiol Gastrointest Liver Physiol* 2017, 312(2):G123-g132.
25. Sun Z, Xie Q, Pan J, Niu N: Cadmium regulates von Willebrand factor and occludin expression in glomerular endothelial cells of mice in a TNF-alpha-dependent manner. *Ren Fail* 2019, 41(1):354-362.

26. Zhou W, Shi G, Bai J, Ma S, Liu Q, Ma X: Colquhounia Root Tablet Protects Rat Pulmonary Microvascular Endothelial Cells against TNF-alpha-Induced Injury by Upregulating the Expression of Tight Junction Proteins Claudin-5 and ZO-1. 2018, 2018:1024634.
27. Fang M, Zhong WH, Song WL, Deng YY, Yang DM, Xiong B, Zeng HK, Wang HD: Ulinastatin Ameliorates Pulmonary Capillary Endothelial Permeability Induced by Sepsis Through Protection of Tight Junctions via Inhibition of TNF-alpha and Related Pathways. *Front Pharmacol* 2018, 9:823.
28. Luo T, Xia Z: A small dose of hydrogen peroxide enhances tumor necrosis factor-alpha toxicity in inducing human vascular endothelial cell apoptosis: reversal with propofol. *Anesth Analg* 2006, 103(1):110-116, table of contents.
29. Ma L, Wu XY, Zhang LH, Chen WM, Uchiyama A, Mashimo T, Fujino Y: Propofol exerts anti-inflammatory effects in rats with lipopolysaccharide-induced acute lung injury by inhibition of CD14 and TLR4 expression. *Braz J Med Biol Res* 2013, 46(3):299-305.
30. Xu Z, Lu Y, Wang J, Ding X, Chen J, Miao C: The protective effect of propofol against TNF-alpha-induced apoptosis was mediated via inhibiting iNOS/NO production and maintaining intracellular Ca(2+) homeostasis in mouse hippocampal HT22 cells. *Biomed Pharmacother* 2017, 91:664-672.
31. Mathy-Hartert M, Deby-Dupont G, Hans P, Deby C, Lamy M: Protective activity of propofol, Diprivan and intralipid against active oxygen species. *Mediators Inflamm* 1998, 7(5):327-333.
32. Chen W, Ju XZ, Lu Y, Ding XW, Miao CH, Chen JW: Propofol improved hypoxia-impaired integrity of blood-brain barrier via modulating the expression and phosphorylation of zonula occludens-1. 2019, 25(6):704-713.
33. Zhan CY, Chen D, Luo JL, Shi YH, Zhang YP: Protective role of down-regulated microRNA-31 on intestinal barrier dysfunction through inhibition of NF-kappaB/HIF-1alpha pathway by binding to HMOX1 in rats with sepsis. *Mol Med* 2018, 24(1):55.
34. Palazon A, Goldrath AW, Nizet V, Johnson RS: HIF transcription factors, inflammation, and immunity. *Immunity* 2014, 41(4):518-528.
35. Jacob A, Potin S, Saubamea B, Crete D, Scherrmann JM, Curis E, Peyssonnaud C, Decleves X: Hypoxia interferes with aryl hydrocarbon receptor pathway in hCMEC/D3 human cerebral microvascular endothelial cells. *J Neurochem* 2015, 132(4):373-383.
36. Wang W, Dentler WL, Borchardt RT: VEGF increases BMEC monolayer permeability by affecting occludin expression and tight junction assembly. *Am J Physiol Heart Circ Physiol* 2001, 280(1):H434-440.
37. Hsu SL, Yin TC, Shao PL, Chen KH, Wu RW, Chen CC, Lin PY, Chung SY, Sheu JJ, Sung PH *et al*: Hyperbaric oxygen facilitates the effect of endothelial progenitor cell therapy on improving outcome of rat critical limb ischemia. *American journal of translational research* 2019, 11(4):1948-1964.

38. Heinolainen K, Karaman S, D'Amico G, Tammela T, Sormunen R, Eklund L, Alitalo K, Zarkada G: VEGFR3 Modulates Vascular Permeability by Controlling VEGF/VEGFR2 Signaling. *Circ Res* 2017, 120(9):1414-1425.
39. Nasir A, Reising LO, Nedderman DM, Fulford AD, Uhlik MT, Benjamin LE, Schade AE, Holzer TR: Heterogeneity of Vascular Endothelial Growth Factor Receptors 1, 2, 3 in Primary Human Colorectal Carcinoma. *Anticancer Res* 2016, 36(6):2683-2696.
40. Ferrara N: Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev* 2004, 25(4):581-611.
41. Park K, Amano H, Ito Y, Kashiwagi S, Yamazaki Y, Takeda A, Shibuya M, Kitasato H, Majima M: Vascular endothelial growth factor receptor-1 (VEGFR-1) signaling enhances angiogenesis in a surgical sponge model. *Biomed Pharmacother* 2016, 78:140-149.
42. Zhu G, Huang Q, Zheng W, Huang Y, Hua J, Yang S, Zhuang J, Wang J, Chang J, Xu J *et al*: LPS Upregulated VEGFR-3 Expression Promote Migration and Invasion in Colorectal Cancer via a Mechanism of Increased NF-kappaB Binding to the Promoter of VEGFR-3. *Cell Physiol Biochem* 2016, 39(5):1665-1678.
43. Allred EN, Dammann O, Fichorova RN, Hooper SR, Hunter SJ, Joseph RM, Kuban K, Leviton A, O'Shea TM, Scott MN: Systemic Inflammation during the First Postnatal Month and the Risk of Attention Deficit Hyperactivity Disorder Characteristics among 10 year-old Children Born Extremely Preterm. *J Neuroimmune Pharmacol* 2017, 12(3):531-543.
44. Kurihara T, Westenskow PD, Bravo S, Aguilar E, Friedlander M: Targeted deletion of Vegfa in adult mice induces vision loss. *J Clin Invest* 2012, 122(11):4213-4217.
45. Li X, Padhan N, Sjostrom EO, Roche FP, Testini C, Honkura N, Sainz-Jaspeado M, Gordon E, Bentley K, Philippides A *et al*: VEGFR2 pY949 signalling regulates adherens junction integrity and metastatic spread. 2016, 7:11017.
46. Lei S, Cheng T, Guo Y, Li C, Zhang W, Zhi F: Somatostatin ameliorates lipopolysaccharide-induced tight junction damage via the ERK-MAPK pathway in Caco2 cells. *Eur J Cell Biol* 2014, 93(7):299-307.
47. Almalki SG, Agrawal DK: ERK signaling is required for VEGF-A/VEGFR2-induced differentiation of porcine adipose-derived mesenchymal stem cells into endothelial cells. *Stem Cell Res Ther* 2017, 8(1):113.
48. Cao C, Dai L, Mu J, Wang X, Hong Y, Zhu C, Jin L, Li S: S1PR2 antagonist alleviates oxidative stress-enhanced brain endothelial permeability by attenuating p38 and Erk1/2-dependent cPLA2 phosphorylation. *Cell Signal* 2019, 53:151-161.

Figures




Figure 1

TNF- α could decrease the expression of occludin in hCMEC/D3 cells. (A) The hCMEC/D3 cells were treated with different concentrations of TNF- α (1, 10, 25, 50 and 100 ng/ml) for 8 h. Left side shows the image of a representative Western blot for occludin; right side is the plot of normalized ratios of optical densities. (B) The hCMEC/D3 cells were incubated with TNF- α (10 ng/ml) for different hours (2, 4, 8, 12, 24 h). Left side is the image of a representative Western blot for occludin; right side is the plot of normalized ratios of optical densities. (C) The cell viability of hCMEC/D3 cells after TNF- α treatment. β -actin was served as internal loading control. Data were represented as the mean \pm SD. *p < 0.05, compared with the control group. Each assay was performed in five replicates.




Figure 2

Propofol could inhibit TNF- α -modulated occludin expression in hCMEC/D3 cells. (A) The hCMEC/D3 cells were pretreated with different concentrations of propofol (5, 10, 25, 50 μ M) for 2 h, followed by TNF- α treatment (10 ng/ml, 4 h). Left side is the image of a representative Western blot for occludin. Right side is the normalized ratios of optical densities. (B) DMSO served as solvent for propofol in this study. Left side is the protein expression of occludin in hCMEC/D3 cells that treated with TNF- α , propofol and DMSO (0.01%). Right side is the normalized ratios of optical densities. β -actin was served as internal loading control. Data were represented as the mean \pm SD. * $p < 0.05$, compared with the control group. # $p < 0.05$, compared with TNF- α treatment group. Each assay was performed in five replicates.




Figure 3

The phosphorylation of ERK was involved in the effects of propofol and TNF- α on occludin expression in hCMEC/D3 cells. (A) Left side is the protein expression of p-ERK, p-p38MAPK, ERK and p38MAPK in hCMEC/D3 cells that treated with TNF- α and propofol. Right side is the plot of normalized ratios of optical densities. (B) The hCMEC/D3 cells were treated with TNF- α , propofol, and ERK inhibitor (LY3214996, 5 μ M). Left side is the protein expression of p-ERK, ERK, and occludin. Right side is the plot of normalized ratios of optical densities. β -actin was served as internal loading control. Data were represented as the mean \pm SD. * $p < 0.05$, compared with the control group. # $p < 0.05$, compared with TNF- α treatment group. Each assay was performed in five replicates.




Figure 4

The VEGF/VEGFR-2 signaling pathway was involved in the effects of propofol and TNF- α on occludin expression in hCMEC/D3 cells. (A) Left side shows the image of a representative Western blot for VEGF and p-VEGFR-2 in hCMEC/D3 cells that treated with TNF- α and propofol. Right side is the plot of normalized ratios of optical densities. (B) The hCMEC/D3 cells were incubated with TNF- α , propofol, and VEGFR-2 inhibitor (SU5416, 10 μ M). Left side shows the image of a representative Western blot for VEGF, p-VEGFR-2, p-ERK, ERK and occludin in hCMEC/D3 cells. Right side is the plot of normalized ratios of optical densities. β -actin was served as internal loading control. Data were represented as the mean \pm SD. *p < 0.05, compared with the control group. #p < 0.05, compared with TNF- α treatment group. Each assay was performed in five replicates.




Figure 5

The expression of Hif-1 α was involved in the effects of propofol and TNF- α on occludin expression in hCMEC/D3 cells. (A) Left side shows the image of a representative Western blot for Hif-1 α in hCMEC/D3 cells that treated with TNF- α and propofol. Right side is the plot of normalized ratios of optical densities. (B) The hCMEC/D3 cells were treated with TNF- α , propofol, and Hif-1 α inhibitor (KC7F2, 10 μ M). Left side shows the image of a representative Western blot for Hif-1 α , VEGF, p-VEGFR-2, p-ERK, ERK and occludin. Right side is the plot of normalized ratios of optical densities. β -actin was served as internal loading control. Data were represented as the mean \pm SD. *p < 0.05, compared with the control group. #p < 0.05, compared with TNF- α treatment group. Each assay was performed in five replicates.




Figure 6

Propofol attenuated TNF- α -modulated occludin expression via inhibiting Hif-1 α / VEGF/ VEGFR-2/ ERK signaling pathway in hCMEC/D3 cells. (A) Left side shows the image of a representative Western blot for Hif-1 α , VEGF, p-VEGF in hCMEC/D3 cells that treated with TNF- α , ERK inhibitor (LY3214996, 5 μ M) and propofol. Right side is the plot of normalized ratios of optical densities. (B) Left side shows the image of a representative Western blot for Hif-1 α in hCMEC/D3 cells that treated with TNF- α , VEGFR-2 inhibitor (SU5416, 10 μ M) and propofol. Right side is the plot of normalized ratios of optical densities. β -actin was served as internal loading control. Data were represented as the mean \pm SD. *p < 0.05, compared with the control group. #p < 0.05, compared with TNF- α treatment group. Each assay was performed in five replicates.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement2.docx](#)
- [supplement2.jpg](#)
- [supplement3.tif](#)