

Association of *VEGF* Haplotypes with Breast Cancer Risk in North-West Indians

Vasudha Sambyal

Guru Nanak Dev University

Kamlesh Guleria (✉ guleria_k@yahoo.com)

Guru Nanak Dev University <https://orcid.org/0000-0002-1626-601X>

Ruhi Kapahi

Guru Nanak Dev University

Mridu Manjari

Sri Guru Ram Das Institute of Medical Sciences and Research

Meena Sudan

Sri Guru Ram Das Institute of Medical Sciences and Research

Manjit Singh Uppal

Sri Guru Ram Das Institute of Medical Sciences and Research

Neeti Rajan Singh

Sri Guru Ram Das Institute of Medical Sciences and Research

Research article

Keywords: VEGF, Polymorphism, Haplotype, Breast cancer

Posted Date: October 5th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42841/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Medical Genomics on August 24th, 2021. See the published version at <https://doi.org/10.1186/s12920-021-01060-4>.

Abstract

Background: Angiogenesis is a complex and coordinated process regulated by different growth factors and is one of the hallmark features of cancer. VEGF is one of the most important endothelial cell mitogen and has a critical role in normal physiological and tumor angiogenesis. The objective of this study was to investigate the potential association of haplotypes of six *VEGF* polymorphisms with breast cancer risk in North-West Indians.

Methods: Samples of 250 breast cancer patients and 250 age and gender matched controls were genotyped for *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms. Haplotypes were generated to determine the better contribution of *VEGF* polymorphisms to breast cancer risk.

Results: Haplotypes CDTCCC (OR = 0.56, 95%CI, 0.38-0.81; $p = 0.003$) and CDTGCC (OR = 0.63, 95%CI, 0.44-0.92; $p = 0.018$) of *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms were significantly associated with decreased risk of breast cancer. CDTCCC haplotype was also significantly associated with reduced risk of breast cancer in pre and post menopausal as well as both obese and non obese patients. Haplotype CDTGCC was marginally associated ($p = 0.07$) with reduced risk of breast cancer in non-obese patients as compared with non-obese controls where as haplotype AICGTC was marginally associated ($p = 0.09$) with reduced risk of breast cancer in obese patients when compared with non-obese patients. The CDTGCC haplotype was significantly associated with increased risk of breast cancer in premenopausal obese patients (OR = 1.98, 95%CI, 1.10-3.56; $p = 0.02$).

Conclusions: Our data indicated that CDTCCC and CDTGCC haplotypes of *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms were significantly associated with breast cancer risk in North-West Indians. Further studies on multiethnic groups with larger sample size are required to confirm our results.

Background

Angiogenesis is one of the hallmark features of cancer [1]. It is a complex and coordinated process regulated by different growth factors like platelet derived growth factor, transforming growth factor and angiopoietins among which vascular endothelial growth factors (VEGF) play a crucial role [2-4]. VEGF is one of the most powerful endothelial cell mitogen and has a very critical role in normal physiological and tumor angiogenesis [5-7]. It enhances tumor vessel permeability and endothelial cell proliferation, migration, differentiation, capillary formation and also has proinflammatory actions [8-12].

The *VEGFA* also known as *VEGF* is located at 6p21.3 and it comprises eight exons and seven introns [13]. It is highly polymorphic with several polymorphisms in the promoter, 5'-untranslated region (5'-UTR) and 3'-UTR [14, 15]. Polymorphisms in the promoter and UTRs have been reported to regulate VEGF expression via alternative initiation of transcription and internal initiation of translation [16, 17]. Functional genetic polymorphisms which alter the regulation of gene expression are predicted to have a

significant impact on disease pathogenesis [18]. *VEGF* -2578C/A, -2549I/D, -460T/C, -116G/A, +405C/G and +936C/T polymorphisms have been associated with differential expression of VEGF [14, 15, 19-22]. The importance of *VEGFA* in breast cancer has been described in several studies [23, 24]. Increased expression of VEGF has been documented in invasive and non invasive breast cancer tissue [25, 26]. Polymorphisms in promoter, 5'-UTR and 3'-UTR of *VEGF* have been reported to affect translation efficiency, circulating plasma concentrations and tumor tissue expression of VEGF [19, 27]. It has been documented that *VEGF* polymorphisms influencing VEGF expression in normal cells might have an impact on tumorigenesis, tumor progression, and response to anti-VEGF agents [22, 28-30].

Haplotype analysis could be a better predictive approach rather than investigating individual polymorphism. It estimates more specific risk and reduces the dimension of association tests and increase statistical power [31]. Due to the important role of *VEGF* in carcinogenesis, the present study aimed to investigate the association of haplotypes of *VEGF* -2578C/A (-1540C/A), -2549I/D (-1511I/D), -460T/C (-1498T/C), +405C/G (-634C/G), -7C/T (+1032C/T) and +936C/T polymorphisms with breast cancer risk in North-West Indians. So far there is no combined report on these six *VEGF* polymorphisms in breast cancer. To the best of our knowledge, this is the first study evaluating the potential association of haplotypes of *VEGF* -2578C/, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphism with breast cancer risk.

Methods

Subjects

The study was performed according to Declaration of Helsinki and was approved by the Ethics Committee of Guru Nanak Dev University, Amritsar, Punjab, India. All the subjects gave a written informed consent with a signature. A total of 500 subjects (250 breast cancer patients and 250 healthy controls) were analyzed in this study. The patients were investigated at Sri Guru Ram Das Institute of Medical Sciences and Research, Vallah, Amritsar, Punjab (India). The selection criteria of patients and controls have been described in our previous study [32]. All the subjects gave 5 ml blood samples for genetic analyses.

Genotyping of *VEGF* Polymorphisms and Analyses of Data

The DNA was extracted from EDTA-anti-coagulated blood samples using organic method [33] with few modifications. Three promoter (*VEGF* -2578C/A, -2549I/D, -460T/C), two 5'-UTR (+405C/G, -7C/T), and one 3'-UTR (+936C/T) polymorphisms were analyzed in this study. The genotyping techniques of six *VEGF* polymorphisms and analysis of genotype data have been described in our published studies [34, 35]. To determine the better contribution of *VEGF* polymorphisms to breast cancer risk, haplotypes of six *VEGF* polymorphisms were generated using the online SNPStats software [36]. Further we predicted the possible influence of studied *VEGF* polymorphisms on the transcription factor binding sites using online software TFSEARCH (<http://www.cbrc.jp/research/db/TFSEARCH.html>).

Results

Characteristics of study Participants

The demographic characteristics of study participants were presented in Table 1. The mean age of patients was 49.38 ± 11.87 years and of controls was 49.34 ± 11.85 years. Of the 250 breast cancer patients, 234 (93%) had infiltrating ductal carcinoma, 4 (2%) had infiltrating lobular carcinoma and 12 (5%) had other types of cancer like medullary carcinoma, mucinous carcinoma, Paget's disease and phyllodes tumor. In the present study, 127 (51%) cases had tumor in left breast, 112 (45%) in right breast and 11 (4%) cases had tumor in both breasts. Of the 250 breast cancer patients, 65 (26%) had stage I, 119 (48%) had stage II, 48 (19%) had stage III, and 18 (7%) had stage IV tumor.

Association of *VEGF* Polymorphisms with Breast Cancer

The results of association of individual *VEGF* polymorphism were summarized in (Supplementary Table S 1). The AA genotype and A allele of *VEGF*-2578C/A, II genotype and I allele of *VEGF*-2549I/D, CC genotype and C allele of *VEGF*-460T/C, GG genotype and G allele of *VEGF*+405C/G polymorphism was significantly associated with increased risk of breast cancer. No association of *VEGF*-7C/T and +936C/T polymorphism with breast cancer risk was observed. We analyzed haplotypes of *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms to determine if there is any difference in *VEGF* haplotypes between breast cancer patients and healthy controls. The most common haplotype in the present study was AICGCC, with the frequencies of 31.3% in breast cancer patients and 23.8% in healthy control individuals. We observed that CDTCCC (OR = 0.56, 95%CI, 0.38-0.81; $p = 0.003$) and CDTGCC (OR = 0.63, 95%CI, 0.44-0.92; $p = 0.018$) haplotypes were significantly associated with decreased risk of breast cancer (Table 2). CDTCCC haplotype was significantly associated with reduced risk of breast cancer in pre and post menopausal patients (Tables 3 and 4). None of the *VEGF* haplotype was associated with breast cancer risk in pre menopausal patients when compared with post menopausal patients (Table 5). The CDTCCC haplotype was also significantly associated with decreased risk of breast cancer in both obese and non obese patients (Tables 6 and 7). Haplotype CDTGCC was marginally associated ($p = 0.07$) with reduced risk of breast cancer in non-obese patients as compared with non-obese controls (Table 7) where as haplotype AICGTC was marginally associated ($p = 0.09$) with reduced risk of breast cancer in obese patients when compared with non-obese patients (Table 8). Further we compared pre menopausal obese patients with post menopausal obese patients and observed that CDTGCC haplotype was significantly associated ($p = 0.02$) with increased risk of breast cancer in premenopausal obese patients (Table 9).

The TFSEARCH software was used to predict the functional significance of *VEGF* polymorphisms. Based on the difference in TFSEARCH TFBS scores, *VEGF*-2578C/A and +405C/G polymorphisms were predicted to alter a transcription factor binding site. *VEGF*-2578A allele abolish the binding site of GATA-2 transcription factor where as *VEGF*+405G allele created the binding site of MZF1 (Myeloid zinc finger 1) transcription factor.

Discussion

In the present study we investigated the potential association of *VEGF* haplotypes based on six polymorphisms with breast cancer risk. In previous reported studies, by using the single/double or triple polymorphism approach, *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms have been analyzed to evaluate their potential association with breast cancer risk in different ethnic groups and results are conflicting (**Supplementary Table S 2**). The ethnicity difference and inadequate sample size could be the potential cause of inconsistent results.

In the present study, we observed that CDTCCC (OR = 0.56, 95%CI, 0.38-0.81; p = 0.003) and CDTGCC (OR = 0.63, 95%CI, 0.44-0.92; p = 0.018) haplotypes of *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms were significantly associated with reduced risk of breast cancer. In none of the previous studies, these six polymorphisms have been reported together. In Caucasian subjects, -460T/+405C/-7C/, +936C haplotype was associated with reduced risk of breast cancer [46]. Significant association of *VEGF*-2578A/-1154A/+405G haplotype with decreased risk of invasive breast cancer has been reported in American population [44]. Haplotype *VEGF*-1154A/-2578A/-634G/-460C was associated with decreased risk of breast cancer in Moroccan population [39]. The -2578A/-1154G/+405G haplotype was associated with decreased risk whereas haplotype -2578C/-1154G/+405G was associated with increased risk of breast cancer recurrence in Caucasian women [58]. Association of -2578C/+405C haplotype with tumor size and higher histological grade has been documented in breast cancer patients [45]. None of the haplotype of *VEGF*-2578C/A, -2549I/D, -460T/C, +405C/G, and +936C/T polymorphisms was associated with breast cancer risk in Iranian population [38].

VEGF-460C/+405G/+936T haplotype was associated with decreased risk of lung cancer in Koreans [59] and increased risk of esophageal adenocarcinoma in Caucasian [60]. The TGC haplotype of *VEGF*-460C/T, +405C/G and +936C/T polymorphism was significantly associated with decreased risk of adenocarcinoma among male non-small cell lung cancer patients [61]. In Turkish population, *VEGF*-2578A/+936T/-460T haplotype has been reported to be associated with increased risk of colorectal cancer [62]. In Tunisians, CIC haplotype of *VEGF*-2578C/A, -2549I/D and +936C/T polymorphisms was associated with increased risk of urothelial bladder cancer [63].

There are some studies from India on different diseases showing association of VEGF haplotypes with disease risk. The CTIG haplotype of VEGF-2578C/A,-7C/T, -2549I/D, and -1001G/C polymorphisms was associated with increased risk of bladder cancer [64] whereas TACI haplotype of *VEGF*+936C/T, -1154G/A, -2578C/A and -2549I/D polymorphisms was associated with increased risk of end stage renal disease [65]. Haplotypes CGCC and CGGC of *VEGF*-460T/C, -1154G/A, +405C/G, and +936C/T polymorphism were associated with aggressiveness of disease in epithelial ovarian cancer patients [66]. No association of *VEGF*+405C/G and +936C/T haplotypes with lung cancer risk has been reported in Kashmiri patients [67].

In the present study, CDTCCC haplotype was significantly associated with reduced risk of breast cancer in pre menopausal as well as in post menopausal patients when compared with pre and post menopausal

controls. The breast cancer risk has also been reported to be modulated by menopause [68]. Estrogen exposure has been described as an important risk factor for breast cancer development and progression [69]. It has been documented that estrogen modulates angiogenesis via effects on endothelial cells under both physiologic and pathologic conditions [70]. Association of *VEGF*-460T/+405G/+936T haplotype with reduced risk of breast cancer has been reported in Chinese premenopausal women [47]. Among postmenopausal breast cancer patients, CCCC haplotype of *VEGF*-2578C/A, -2489C/T, -460T/C, +405C/G and -7C/T polymorphisms was associated with reduced risk of distant metastases [71].

The CDTCCC haplotype was significantly associated with decrease risk of breast cancer in obese as well as in non obese patients compared to obese and non obese controls where as CDTGCC haplotype was significantly associated with increased risk of breast cancer in premenopausal obese patients. About 75.2% of patients and 72% of controls were obese in the present study. It has been hypothesized that hormonal mechanisms and metabolic factors are involved in the link between obesity and breast cancer. Insulin resistance and hyperinsulinemia have been reported to be associated with increased breast cancer risk and with worst prognosis in both pre and post menopausal women [72-74]. In mouse model, it has been demonstrated that over expression of VEGFA in adipose tissue provide protection against high fat diet induced obesity and insulin sensitivity [75, 76]. It has been documented that angiogenesis plays an important role in the regulation of adipogenesis [77]. *VEGF* has been described as an important angiogenic factor in adipose tissue and it regulates the development of new vessels required for the expansion of adipose tissue [76, 78]. It has been reported that adiponectin, a regulator of insulin resistance block angiogenesis by increasing the expression of *TP53* and decreasing the expression of *VEGF* [79].

In the present study we predicted that *VEGF*-2578A allele of *VEGF*-2578C/A polymorphism abolished the binding site of GATA-2 transcription factor. The GATA family of transcription factors is regulator of gene expression in hematopoietic cells [80, 81]. Correlation of reduced GATA binding promoter activity has been documented with attenuation of VEGF mediated signaling [82]. G allele of *VEGF*+405C/G polymorphism created the binding site of MZF1 transcription factor. MZF1 transcription factor has been reported to be involved in transcriptional regulation during myelopoiesis [83]. Disruption of MZF1 transcription factor binding site by *VEGF*-634C (+405C) allele has also been reported in peripheral blood mononuclear cells [15]. It has been reported that substitution of C by G at +405 position in 5'-UTR may affect internal ribosome entry site (IRES) and increases the transcription of large isoform of VEGFA [84].

Polymorphisms of *VEGFA* have been reported to be associated with efficacy and toxicity of anti - VEGF agents [41, 85, 86]. Haplotype -460T/+405C/+936C haplotype was associated with better survival among Chinese breast cancer patients [87]. *VEGF*-2578A/-1154G/+405G haplotype was associated with marginally improved prognosis whereas haplotype -2578C/-1154G/+405G was significantly associated with adverse prognosis in HER2 positive breast cancer patients [88]. Apart from breast cancer, correlation of *VEGF* haplotypes with therapy response has also been documented in other cancer types. The CACC haplotype of *VEGF*-460T/C, -116G/A, +405C/G, and +936C/T polymorphism was significantly associated with worse survival in Korean gastric cancer patients [89]. In esophageal cancer, CGC haplotype of *VEGF*

-460T/C, +405C/G and +936C/T polymorphism was associated with poorer outcome as compared to other haplotypes [90]. The AGCGC haplotype of *VEGF* -2578 C/A, -1154 G/A, -460 T/C, +405 G/C and +936C/T polymorphisms was found to be associated with improved progression-free survival in epithelial ovarian cancer patients [91]. Haplotype -2578C/-460T/+405C/+936C and -2578C/-460T/+405C/+936T was associated with inferior response rate in metastatic colorectal cancer patients to first line XELOX treatment [92]. Assessment of haplotypes of *VEGF* polymorphisms can be used for identification of cancer risk, aggressiveness and selection of patients suitable for anti-VEGF therapy.

Conclusions

We report for the first time that CDTCCC and CDTGCC haplotypes of *VEGF* -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T polymorphisms were significantly associated with breast cancer risk in North-West Indians. Further studies on multiethnic groups with larger sample size are required to confirm our results.

List Of Abbreviations

VEGF: Vascular endothelial growth factor; UTR: untranslated region; OR: odds ratio

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of Guru Nanak Dev University, Amritsar, Punjab, India. All the subjects gave a written informed consent with a signature or thumb impression.

Consent for publication: Not applicable

Availability of data and materials

All data generated in this study is included in supplementary file.

Competing interests

The authors declare that they have no competing interests.

Funding: This study was supported by the DBT grant BT/PR 13252/GBD/27/236/2009 and DST-PURSE grant sanctioned to VS and KG. The funders had no role in the designing of the study or in the collection of samples, analysis of data or preparation of the manuscript.

Authors' contributions

KG and VS conceptualized and designed the study. RK and KG performed the experiments. KG and VS analyzed the results and prepared the manuscript. MM, MS, MSU and NRS did clinical diagnosis and

histopathological classification of patients and also helped in collection of blood samples of breast cancer patients. All authors read and approved the final manuscript.

Acknowledgements

We thank the patients and the healthy control individuals for their participation.

References

1. Hanahan D, Weinberg RA. The hallmarks of cancer. *cell*. 2000 Jan 7;100(1):57-70.
2. Hanahan D, Folkman J. Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *cell*. 1996 Aug 9;86(3):353-64.
3. Ferrara N, Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocrine reviews*. 1997 Feb 1;18(1):4-25.
4. Carmeliet P. Mechanisms of angiogenesis and arteriogenesis. *Nature medicine*. 2000 Apr;6(4):389-95.
5. Ferrara N. Vascular endothelial growth factor: molecular and biological aspects. *Curr Top Microbiol Immunol*. 1999;237:1-30.
6. Yoshiji H, Kuriyama S, Yoshii J, Ikenaka Y, Noguchi R, Hicklin DJ, Wu Y, Yanase K, Namisaki T, Kitade M, Yamazaki M. Halting the interaction between vascular endothelial growth factor and its receptors attenuates liver carcinogenesis in mice. *Hepatology*. 2004 Jun;39(6):1517-24.
7. Roy H, Bhardwaj S, Ylä-Herttuala S. Biology of vascular endothelial growth factors. *FEBS letters*. 2006 May 22;580(12):2879-87.
8. Dvorak HF, Brown LF, Detmar M, Dvorak AM. Vascular permeability factor/vascular endothelial growth factor, microvascular hyperpermeability, and angiogenesis. *The American journal of pathology*. 1995 May;146(5):1029.
9. Kim I, Moon SO, Kim SH, Kim HJ, Koh YS, Koh GY. Vascular endothelial growth factor expression of intercellular adhesion molecule 1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), and E-selectin through nuclear factor- κ B activation in endothelial cells. *Journal of Biological Chemistry*. 2001 Mar 9;276(10):7614-20.
10. Reinders ME, Fang JC, Wong W, Ganz P, Briscoe DM. Expression patterns of vascular endothelial growth factor in human cardiac allografts: association with rejection. *Transplantation*. 2003 Jul 15;76(1):224-30.
11. Ferrara N. Vascular endothelial growth factor: basic science and clinical progress. *Endocrine reviews*. 2004 Aug 1;25(4):581-611.
12. Kimura Y, Morohashi S, Yoshizawa T, Suzuki T, Morohashi H, Sakamoto Y, Koyama M, Murata A, Kijima H, Hakamada K. Clinicopathological significance of vascular endothelial growth factor, thymidine phosphorylase and microvessel density in colorectal cancer. *Molecular medicine reports*. 2016 Feb 1;13(2):1551-7.

13. Vincenti V, Cassano C, Rocchi M, Persico MG. Assignment of the vascular endothelial growth factor gene to human chromosome 6p21. 3. Circulation. 1996 Apr 15;93(8):1493-5.
14. Brogan IJ, Khan N, Isaac K, Hutchinson JA, Pravica V, Hutchinson IV. Novel polymorphisms in the promoter and 5' UTR regions of the human vascular endothelial growth factor gene. Human immunology. 1999 Dec 1;60(12):1245-9.
15. Watson CJ, Webb NJ, Bottomley MJ, Brenchley PE. Identification of polymorphisms within the vascular endothelial growth factor (VEGF) gene: correlation with variation in VEGF protein production. Cytokine. 2000 Aug 1;12(8):1232-5.
16. Akiri G, Nahari D, Finkelstein Y, Le SY, Elroy-Stein O, Levi BZ. Regulation of vascular endothelial growth factor (VEGF) expression is mediated by internal initiation of translation and alternative initiation of transcription. Oncogene. 1998 Jul;17(2):227-36.
17. Huez I, Créancier L, Audigier S, Gensac MC, Prats AC, Prats H. Two independent internal ribosome entry sites are involved in translation initiation of vascular endothelial growth factor mRNA. Molecular and cellular biology. 1998 Nov 1;18(11):6178-90.
18. Taylor JG, Choi EH, Foster CB, Chanock SJ. Using genetic variation to study human disease. Trends in molecular medicine. 2001 Nov 1;7(11):507-12.
19. Renner W, Kotschan S, Hoffmann C, Obermayer-Pietsch B, Pilger E. A common 936 C/T mutation in the gene for vascular endothelial growth factor is associated with vascular endothelial growth factor plasma levels. Journal of vascular research. 2000;37(6):443-8.
20. Awata T, Inoue K, Kurihara S, Ohkubo T, Watanabe M, Inukai K, Inoue I, Katayama S. A common polymorphism in the 5'-untranslated region of the VEGF gene is associated with diabetic retinopathy in type 2 diabetes. Diabetes. 2002 May 1;51(5):1635-9.
21. Mohammadi M, Ollier WE, Hutchinson IV. A functional association study of VEGF gene promoter polymorphisms with VEGF expression by stimulated pbm cells. Human Immunology. 2003;10(64):S125.
22. Stevens A, Soden J, Brenchley PE, Ralph S, Ray DW. Haplotype analysis of the polymorphic human vascular endothelial growth factor gene promoter. Cancer research. 2003 Feb 15;63(4):812-6.
23. Adams J, Carder PJ, Downey S, Forbes MA, MacLennan K, Allgar V, Kaufman S, Hallam S, Bicknell R, Walker JJ, Cairnduff F. Vascular endothelial growth factor (VEGF) in breast cancer: comparison of plasma, serum, and tissue VEGF and microvessel density and effects of tamoxifen. Cancer research. 2000 Jun 1;60(11):2898-905.
24. Gasparini G. Prognostic value of vascular endothelial growth factor in breast cancer. The oncologist. 2000 Apr;5:37-44.
25. Yoshiji H, Gomez DE, Shibuya M, Thorgeirsson UP. Expression of vascular endothelial growth factor, its receptor, and other angiogenic factors in human breast cancer. Cancer Research. 1996 May 1;56(9):2013-6.
26. Guidi AJ, Schnitt SJ, Fischer L, Tognazzi K, Harris JR, Dvorak HF, Brown LF. Vascular permeability factor (vascular endothelial growth factor) expression and angiogenesis in patients with ductal

- carcinoma in situ of the breast. *Cancer*. 1997 Nov 15;80(10):1945-53.
27. Koukourakis MI, Papazoglou D, Giatromanolaki A, Bougioukas G, Maltezos E, Sivoridis E. VEGF gene sequence variation defines VEGF gene expression status and angiogenic activity in non-small cell lung cancer. *Lung cancer*. 2004 Dec 1;46(3):293-8.
 28. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nature medicine*. 2003 Jun;9(6):669-76.
 29. Willett CG, Boucher Y, Di Tomaso E, Duda DG, Munn LL, Tong RT, Chung DC, Sahani DV, Kalva SP, Kozin SV, Mino M. Direct evidence that the VEGF-specific antibody bevacizumab has antivasculature effects in human rectal cancer. *Nature medicine*. 2004 Feb;10(2):145-7.
 30. Jain RK. Antiangiogenic therapy for cancer: current and emerging concepts. *Oncology (Williston Park, NY)*. 2005 Apr;19(4 Suppl 3):7-16.
 31. Clark AG. The role of haplotypes in candidate gene studies. *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society*. 2004 Dec;27(4):321-33.
 32. Kapahi R, Manjari M, Uppal MS, Singh NR, Sambyal V, Guleria K. Association of -2549 insertion/deletion polymorphism of vascular endothelial growth factor with breast cancer in North Indian patients. *Genetic testing and molecular biomarkers*. 2013 Mar 1;17(3):242-8.
 33. Adeli K, Ogbonna G. Rapid purification of human DNA from whole blood for potential application in clinical chemistry laboratories. *Clinical chemistry*. 1990 Feb 1;36(2):261-4.
 34. Kapahi R, Manjari M, Sudan M, Uppal MS, Singh NR, Sambyal V, Guleria K. Association of +405C>G and +936C>T polymorphisms of the vascular endothelial growth factor gene with sporadic breast cancer in North Indians. *Asian Pac J Cancer Prev*. 2014 Jan 1;15(1):257-63.
 35. Kapahi R, Guleria K, Sambyal V, Manjari M, Sudan M, Uppal MS, Singh NR. Association of VEGF and VEGFR1 polymorphisms with breast cancer risk in North Indians. *Tumor Biology*. 2015 Jun 1;36(6):4223-34.
 36. Solé X, Guinó E, Valls J, Iniesta R, Moreno V. SNPStats: a web tool for the analysis of association studies. *Bioinformatics*. 2006 Aug 1;22(15):1928-9.
 37. Al Balawi IA, Mir R, Abu-Duhier FM. Potential impact of vascular endothelial growth factor gene variation (-2578C>A) on breast cancer susceptibility in Saudi Arabia: a Case-Control Study. *Asian Pacific journal of cancer prevention: APJCP*. 2018;19(4):1135.
 38. Rezaei M, Hashemi M, Sanaei S, Mashhadi MA, Taheri M. Association Between Vascular Endothelial Growth Factor Gene Polymorphisms with Breast Cancer Risk in an Iranian Population. *Breast Cancer (Auckl)*. 2016 Jul 4;10:85-91.
 39. Rahoui J, Laraqui A, Sbitti Y, Touil N, Ibrahim A, Ghrab B, Al Bouzidi A, Rahali DM, Dehayni M, Ichou M, Zaoui F. Investigating the association of vascular endothelial growth factor polymorphisms with breast cancer: a Moroccan case-control study. *Medical Oncology*. 2014 Sep 1;31(9):193.
 40. Sa-Nguanraksa D, Chuangsuwanich T, Pongpruttipan T, Kummalue T, Rojananin S, Ratanawichitrasin A, Prasartong-Osoth P, Chuthatisith S, Pisarturakit P, Aumrithacharoenchok W, Rushatamukayanunt P. Vascular endothelial growth factor-634G/C polymorphism is associated with

- increased breast cancer risk and aggressiveness. *Molecular medicine reports*. 2013 Oct 1;8(4):1242-50.
41. Schneider BP, Radovich M, Sledge GW, Robarge JD, Li L, Storniolo AM, Lemler S, Nguyen AT, Hancock BA, Stout M, Skaar T. Association of polymorphisms of angiogenesis genes with breast cancer. *Breast cancer research and treatment*. 2008 Sep 1;111(1):157-63.
 42. Langsenlehner T, Langsenlehner U, Renner W, Krippel P, Mayer R, Wascher TC, Kapp KS. Single nucleotide polymorphisms and haplotypes in the gene for vascular endothelial growth factor and risk of prostate cancer. *European journal of cancer*. 2008 Jul 1;44(11):1572-6.
 43. Pharoah PD, Antoniou AC, Easton DF, Ponder BA. Polygenes, risk prediction, and targeted prevention of breast cancer. *New England Journal of Medicine*. 2008 Jun 26;358(26):2796-803.
 44. Jacobs EJ, Feigelson HS, Bain EB, Brady KA, Rodriguez C, Stevens VL, Patel AV, Thun MJ, Calle EE. Polymorphisms in the vascular endothelial growth factor gene and breast cancer in the Cancer Prevention Study II cohort. *Breast Cancer Research*. 2006 Apr 1;8(2):R22.
 45. Jin Q, Hemminki K, Enquist K, Lenner P, Grzybowska E, Klaes R, Henriksson R, Chen B, Pamula J, Pekala W, Zientek H. Vascular endothelial growth factor polymorphisms in relation to breast cancer development and prognosis. *Clinical cancer research*. 2005 May 15;11(10):3647-53.
 46. Balasubramanian SP, Cox A, Cross SS, Higham SE, Brown NJ, Reed MW. Influence of VEGF-A gene variation and protein levels in breast cancer susceptibility and severity. *International journal of cancer*. 2007 Sep 1;121(5):1009-16.
 47. Kataoka N, Cai Q, Wen W, Shu XO, Jin F, Gao YT, Zheng W. Population-based case-control study of VEGF gene polymorphisms and breast cancer risk among Chinese women. *Cancer Epidemiology and Prevention Biomarkers*. 2006 Jun 1;15(6):1148-52.
 48. James R, Ramesh G, Krishnamoorthy L, Bhagat R, Chadaga S, Deshmane V, Ramaswamy G. Prevalence of +405G>C, -1154G>A vascular endothelial growth factor polymorphism in breast cancer. *Indian Journal of Clinical Biochemistry*. 2014 Jan 1;29(1):21-8.
 49. Rani J, Rahul B, Ramesh G, Krishnamoorthy L, Shilpa C, Ramaswamy G, Deshmane V. Association of vascular endothelial growth factor single nucleotide polymorphisms on the prognosis of breast cancer patients. *Indian Journal of Cancer*. 2014 Oct 1;51(4):512.
 50. Luo T, Chen L, He P, Hu QC, Zhong XR, Sun Y, Yang YF, Tian TL, Zheng H. Vascular endothelial growth factor (VEGF) gene polymorphisms and breast cancer risk in a Chinese population. *Asian Pac J Cancer Prev*. 2013 Jan 1;14(4):2433-7.
 51. Oliveira C, Lourenço GJ, Silva PM, Cardoso-Filho C, Favarelli MH, Gonçalves NS, Gurgel MS, Lima CS. Polymorphisms in the 5'-and 3'-untranslated region of the VEGF gene and sporadic breast cancer risk and clinicopathologic characteristics. *Tumor Biology*. 2011 Apr 1;32(2):295-300.
 52. Absenger G, Szkandera J, Stotz M, Pichler M, Winder T, Langsenlehner T, Langsenlehner U, Samonigg H, Renner W, Gerger A. A common and functional gene variant in the vascular endothelial growth factor predicts clinical outcome in early-stage breast cancer. *Molecular carcinogenesis*. 2013 Nov;52(S1):96-102.

53. Rodrigues P, Furriol J, Tormo E, Ballester S, Lluch A, Eroles P. The single-nucleotide polymorphisms+ 936 C/T VEGF and- 710 C/T VEGFR1 are associated with breast cancer protection in a Spanish population. *Breast cancer research and treatment*. 2012 Jun 1;133(2):769-78.
54. Lin GT, Tseng HF, Yang CH, Hou MF, Chuang LY, Tai HT, Tai MH, Cheng YH, Wen CH, Liu CS, Huang CJ. Combinational polymorphisms of seven CXCL12-related genes are protective against breast cancer in Taiwan. *OMICS A Journal of Integrative Biology*. 2009 Apr 1;13(2):165-72.
55. Jakubowska A, Gronwald J, Menkiszak J, Górski B, Huzarski T, Byrski T, Edler L, Lubiński J, Scott RJ, Hamann U. The VEGF_936_C> T 3' UTR polymorphism reduces BRCA1-associated breast cancer risk in Polish women. *Cancer letters*. 2008 Apr 8;262(1):71-6.
56. Eroğlu A, Öztürk A, Cam R, Akar N. Vascular endothelial growth factor gene 936 C/T polymorphism in breast cancer patients. *Medical Oncology*. 2008 Mar 1;25(1):54-5.
57. Krippel P, Langsenlehner U, Renner W, Yazdani-Biuki B, Wolf G, Wascher TC, Paulweber B, Haas J, Samonigg H. A common 936 C/T gene polymorphism of vascular endothelial growth factor is associated with decreased breast cancer risk. *International journal of cancer*. 2003 Sep 10;106(4):468-71.
58. Kidd LR, Brock GN, VanCleave TT, Benford ML, Lavender NA, Kruer TL, Wittliff JL. Angiogenesis-associated sequence variants relative to breast cancer recurrence and survival. *Cancer Causes & Control*. 2010 Oct 1;21(10):1545-57.
59. Lee SJ, Lee SY, Jeon HS, Park SH, Jang JS, Lee GY, Son JW, Kim CH, Lee WK, Kam S, Park RW. Vascular endothelial growth factor gene polymorphisms and risk of primary lung cancer. *Cancer Epidemiology and Prevention Biomarkers*. 2005 Mar 1;14(3):571-5.
60. Zhai R, Liu G, Asomaning K, Su L, Kulke MH, Heist RS, Nishioka NS, Lynch TJ, Wain JC, Lin X, Christiani DC. Genetic polymorphisms of VEGF, interactions with cigarette smoking exposure and esophageal adenocarcinoma risk. *Carcinogenesis*. 2008 Dec 1;29(12):2330-4.
61. Zhai R, Liu G, Zhou W, Su L, Heist RS, Lynch TJ, Wain JC, Asomaning K, Lin X, Christiani DC. Vascular endothelial growth factor genotypes, haplotypes, gender, and the risk of non-small cell lung cancer. *Clinical cancer research*. 2008 Jan 15;14(2):612-7.
62. Jannuzzi AT, Özhan G, Yanar HT, Alpertunga B. VEGF gene polymorphisms and susceptibility to colorectal cancer. *Genetic testing and molecular biomarkers*. 2015 Mar 1;19(3):133-7.
63. Ben Wafi S, Kallel A, Ben Fradj MK, Sallemi A, Ben Rhouma S, Ben Halima M, Sanhaji H, Nouria Y, Jemaa R, Feki M. Haplotype-based association of Vascular Endothelial Growth Factor gene polymorphisms with urothelial bladder cancer risk in Tunisian population. *Journal of Clinical Laboratory Analysis*. 2018 Nov;32(9):e22610.
64. Jaiswal PK, Tripathi N, Shukla A, Mittal RD. Association of single nucleotide polymorphisms in vascular endothelial growth factor gene with bladder cancer risk. *Medical Oncology*. 2013 Jun 1;30(2):509.
65. Prakash S, Prasad N, Sharma RK, Faridi RM, Agrawal S. Vascular endothelial growth factor gene polymorphisms in North Indian patients with end stage renal disease. *Cytokine*. 2012 May

1;58(2):261-6.

66. Janardhan B, Vaderhobli S, Bhagat R, Chennagiri Srinivasamurthy P, Venketeshiah Reddihalli P, Gawari R, Krishnamoorthy L. Investigating impact of vascular endothelial growth factor polymorphisms in epithelial ovarian cancers: a study in the Indian population. *PloS one*. 2015 Jul 9;10(7):e0131190.
67. Naikoo NA, Afroze D, Rasool R, Shah S, Ahangar AG, Bhat IA, Qasim I, Siddiqi MA, Shah ZA. SNP and haplotype analysis of vascular endothelial growth factor (VEGF) gene in lung cancer patients of Kashmir. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2017;18(7):1799.
68. Mohanty SS, Mohanty PK. Obesity as potential breast cancer risk factor for postmenopausal women. *Genes & Diseases*. 2019 Sep 10.
69. Lapidus RG, Nass SJ, Davidson NE. The loss of estrogen and progesterone receptor gene expression in human breast cancer. *Journal of mammary gland biology and neoplasia*. 1998 Jan 1;3(1):85-94.
70. Losordo DW, Isner JM. Estrogen and angiogenesis: a review. *Arteriosclerosis, thrombosis, and vascular biology*. 2001 Jan;21(1):6-12.
71. Langsenlehner U, Hofmann G, Renner W, Gerger A, Krenn-Pilko S, Thurner EM, Krippel P, Langsenlehner T. Association of vascular endothelial growth factor–a gene polymorphisms and haplotypes with breast cancer metastases. *Acta Oncologica*. 2015 Mar 16;54(3):368-76.
72. Goodwin PJ, Ennis M, Pritchard KI, Trudeau ME, Koo J, Madarnas Y, Hartwick W, Hoffman B, Hood N. Fasting insulin and outcome in early-stage breast cancer: results of a prospective cohort study. *Journal of clinical oncology*. 2002 Jan 1;20(1):42-51.
73. Morimoto LM, White E, Chen Z, Chlebowski RT, Hays J, Kuller L, Lopez AM, Manson J, Margolis KL, Muti PC, Stefanick ML. Obesity, body size, and risk of postmenopausal breast cancer: the Women's Health Initiative (United States). *Cancer Causes & Control*. 2002 Oct 1;13(8):741-51.
74. Schairer C, Hill D, Sturgeon SR, Fears T, Pollak M, Mies C, Ziegler RG, Hoover RN, Sherman ME. Serum concentrations of IGF-I, IGFBP-3 and c-peptide and risk of hyperplasia and cancer of the breast in postmenopausal women. *International Journal of Cancer*. 2004 Feb 20;108(5):773-9.
75. Elias I, Franckhauser S, Ferré T, Vilà L, Tafuro S, Muñoz S, Roca C, Ramos D, Pujol A, Riu E, Ruberte J. Adipose tissue overexpression of vascular endothelial growth factor protects against diet-induced obesity and insulin resistance. *Diabetes*. 2012 Jul 1;61(7):1801-13.
76. Sun K, Asterholm IW, Kusminski CM, Bueno AC, Wang ZV, Pollard JW, Brekken RA, Scherer PE. Dichotomous effects of VEGF-A on adipose tissue dysfunction. *Proceedings of the National Academy of Sciences*. 2012 Apr 10;109(15):5874-9.
77. Xue Y, Petrovic N, Cao R, Larsson O, Lim S, Chen S, Feldmann HM, Liang Z, Zhu Z, Nedergaard J, Cannon B. Hypoxia-independent angiogenesis in adipose tissues during cold acclimation. *Cell metabolism*. 2009 Jan 7;9(1):99-109.
78. Cao Y. Adipose tissue angiogenesis as a therapeutic target for obesity and metabolic diseases. *Nature reviews Drug discovery*. 2010 Feb;9(2):107-15.

79. Khan S, Shukla S, Sinha S, Meeran SM. Role of adipokines and cytokines in obesity-associated breast cancer: therapeutic targets. *Cytokine & growth factor reviews*. 2013 Dec 1;24(6):503-13.
80. Tsai FY, Keller G, Kuo FC, Weiss M, Chen J, Rosenblatt M, Alt FW, Orkin SH. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature*. 1994 Sep 15;371(6494):221-6.
81. Shimizu R, Yamamoto M. Gene expression regulation and domain function of hematopoietic GATA factors. In *Seminars in cell & developmental biology* 2005 Feb 1 (Vol. 16, No. 1, pp. 129-136). Academic Press.
82. Minami T, Murakami T, Horiuchi K, Miura M, Noguchi T, Miyazaki JI, Hamakubo T, Aird WC, Kodama T. Interaction between hex and GATA transcription factors in vascular endothelial cells inhibits flk-1/KDR-mediated vascular endothelial growth factor signaling. *Journal of Biological Chemistry*. 2004 May 14;279(20):20626-35.
83. Lenny N, Westendorf JJ, Hiebert SW. Transcriptional regulation during myelopoiesis. *Molecular biology reports*. 1997 Aug 1;24(3):157.
84. Huez I, Bornes S, Bresson D, Créancier L, Prats H. New vascular endothelial growth factor isoform generated by internal ribosome entry site-driven CUG translation initiation. *Molecular Endocrinology*. 2001 Dec 1;15(12):2197-210.
85. Etienne-Grimaldi MC, Formento P, Degeorges A, Pierga JY, Delva R, Pivot X, Dalenc F, Espié M, Veyret C, Formento JL, Francoual M. Prospective analysis of the impact of VEGF-A gene polymorphisms on the pharmacodynamics of bevacizumab-based therapy in metastatic breast cancer patients. *British journal of clinical pharmacology*. 2011 Jun;71(6):921-8.
86. Formica V, Palmirotta R, Del Monte G, Savonarola A, Ludovici G, De Marchis ML, Grenga I, Schirru M, Guadagni F, Roselli M. Predictive value of VEGF gene polymorphisms for metastatic colorectal cancer patients receiving first-line treatment including fluorouracil, irinotecan, and bevacizumab. *International journal of colorectal disease*. 2011 Feb 1;26(2):143-51.
87. Lu H, Shu XO, Cui Y, Kataoka N, Wen W, Cai Q, Ruan ZX, Gao YT, Zheng W. Association of genetic polymorphisms in the VEGF gene with breast cancer survival. *Cancer research*. 2005 Jun 15;65(12):5015-9.
88. Maae E, Andersen RF, Steffensen KD, Jakobsen EH, Brandslund I, Sørensen FB, Jakobsen A. Prognostic impact of VEGFA germline polymorphisms in patients with HER2-positive primary breast cancer. *Anticancer research*. 2012 Sep 1;32(9):3619-27.
89. Kim JG, Sohn SK, Chae YS, Cho YY, Bae HI, Yan G, Park JY, Lee MH, Chung HY, Yu W. Vascular endothelial growth factor gene polymorphisms associated with prognosis for patients with gastric cancer. *Annals of oncology*. 2007 Jun 1;18(6):1030-6.
90. Bradbury PA, Zhai R, Ma C, Xu W, Hopkins J, Kulke MJ, Asomaning K, Wang Z, Su L, Heist RS, Lynch TJ. Vascular endothelial growth factor polymorphisms and esophageal cancer prognosis. *Clinical cancer research*. 2009 Jul 15;15(14):4680-5.
91. Steffensen KD, Waldstrøm M, Brandslund I, Jakobsen A. The relationship of VEGF polymorphisms with serum VEGF levels and progression-free survival in patients with epithelial ovarian cancer.

Gynecologic oncology. 2010 Apr 1;117(1):109-16.

92. Hansen TF, Spindler KG, Andersen RF, Lindebjerg J, Brandslund I, Jakobsen A. The predictive value of genetic variations in the vascular endothelial growth factor A gene in metastatic colorectal cancer. The pharmacogenomics journal. 2011 Feb;11(1):53-60.

Tables

Table 1. Characteristics of Breast cancer patients and healthy controls

Variable	Patients	%	Controls	%
	n		n	
Total No. of subjects	250		250	
Gender				
Males	7	2.8	7	2.8
Females	243	97.2	243	97.2
Age at diagnosis (Years)				
≤40	66	26.4	66	26.4
>40	184	73.6	184	49.34±11.85
<i>Mean age</i>	49.38±11.87		25-85	
<i>Range</i>	25-85			
Habitat				
Rural	186	74.4	186	74.4
Urban	64	25.6	64	25.6
Diet				
Vegetarian	147	58.8	154	61.6
Non vegetarian	103	41.2	96	38.4
Obesity				
Non obese	62	24.8	70	28.0
Obese	188	75.2	180	72.0
Menstrual status				
Premenopausal	114	46.91	150	61.73
Postmenopausal	129	53.09	93	38.27
Mean age at menarche	14.74±1.78		14.75±1.47	
Mean age at first child birth	22.70±4.08		22.45±3.43	
Breastfeeding				
Yes	215	88.48	224	92.18
No	28	11.52	19	7.82
Oral Contraceptives				
Yes	21	8.64	16	6.58

Table 2. Association between *VEGF* haplotypes and breast cancer risk

Haplotype*	Cases (%)	Controls (%)	OR(95%CI)	p value
AICGCC	31.3	23.8	1(Reference)	
CDTCCC	23.7	28.9	0.56(0.38-0.81)	0.003
CDTGCC	19.6	25.6	0.63(0.44-0.92)	0.018
AICGTC	9.0	9.3	0.75(0.43-1.30)	0.31
AICGCT	3.1	1.8	1.22(0.40-3.74)	0.73
AICGTT	1.8	2.0	0.76(0.25-2.33)	0.64
CDTCCT	1.7	1.2	1.25(0.30-5.19)	0.76
CDTGCT	1.4	1.5	0.65(0.13-3.35)	0.61

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 3. *VEGF* haplotypes and breast cancer risk in premenopausal subjects

Haplotype*	Pre menopausal Cases (%) n=114	Pre menopausal Controls (%) n=150	OR(95%CI)	p value
AICGCC	29.0	24.5	1(Reference)	
CDTCCC	20.6	28.5	0.65(0.43-0.99)	0.04
CDTGCC	25.0	26.3	0.95(0.64-1.42)	0.81
AICGTC	9.0	9.9	0.90(0.50-1.64)	0.73
AICGCT	3.1	1.4	2.23(0.67-7.43)	0.18
AICGTT	3.2	1.5	2.27(0.69-7.51)	0.16
CDTCCT	3.3	2.1	1.87(0.46-7.56)	0.38

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 4. *VEGF* haplotype and breast cancer risk in post menopausal subjects

Haplotype*	Post menopausal Cases (%) n=129	Post menopausal Controls (%) n=93	OR(95%CI)	p value
AICGCC	31.9	23.4	1 (Reference)	
CDTCCC	24.8	33.3	0.52(0.30-0.92)	0.02
CDTGCC	17.6	22.4	0.59(0.31-1.12)	0.11
AICGTC	9.2	7.9	0.82(0.34-1.99)	0.66
AICGCT	3.3	2.4	1.05(0.16-6.83)	0.96
AICGTT	0.7	3.3	0.27(0.04-1.94)	0.19

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 5. Association of *VEGF* haplotypes with breast cancer risk in pre menopausal and post menopausal patients

Haplotype*	Pre menopausal (%) n=114	Post menopausal (%) n=129	OR(95%CI)	p value
AICGCC	29.0	31.9	1(Reference)	
CDTCCC	20.6	24.8	0.95(0.54-1.67)	0.86
CDTGCC	25.0	17.6	1.42(0.82-2.47)	0.22
AICGTC	9.0	9.2	1.05(0.48-2.33)	0.90
AICGCT	3.1	3.3	0.70(0.12-3.90)	0.68
CDTCCT	3.8	1.1	3.01(0.55-16.65)	0.21
AITGTC	2.8	1.8	0.24(0.03-1.97)	0.18
AITGCC	1.5	1.2	1.31(0.29-5.96)	0.73

OR-odds ratio; CI-confidence interval

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 6. Association of *VEGF* haplotypes with breast cancer risk in obese subjects

Haplotype*	Obese Patients (%) n=188	Obese Controls (%) n=180	OR(95%CI)	p value
AICGCC	31.2	24.6	1(Reference)	
CDTCCC	25.2	28.9	0.63(0.41-0.99)	0.04
CDTGCC	19.3	25.0	0.76 (0.49-1.19)	0.23
AICGTC	7.0	9.3	0.70(0.35-1.39)	0.31
AICGTT	3.0	1.9	1.27(0.39-4.12)	0.69
CDTGCT	2.4	1.7	1.23(0.29-5.22)	0.78
AICGCT	1.8	1.7	0.77(0.14-4.26)	0.76
CDTCCT	1.8	1.5	1.17(0.26-5.18)	0.84

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 7. Association of *VEGF* haplotypes with breast cancer risk in non obese subjects

Haplotype*	Non-obese Patients (%) n=62	Non-obese Controls (%) n=70	OR(95%CI)	p value
AICGCC	34.1	23.7	1(Reference)	
CDTCCC	20.9	31.2	0.44(0.21-0.93)	0.03
CDTGCC	18.2	22.7	0.47 (0.21-1.05)	0.07
AICGTC	13.5	9.5	1.11(0.38-3.21)	0.85
CDTGTC	1.2	1.4	0.46(0.02-12.99)	0.65

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 8. Association of *VEGF* haplotypes with breast cancer risk in obese and non obese patients

Haplotype*	Obese (%) n=188	Non-obese (%) n=62	OR(95%CI)	p value
AICGCC	31.2	34.1	1(Reference)	
CDTCCC	25.1	20.9	1.28(0.70-2.34)	0.42
CDTGCC	19.3	18.2	1.11(0.61-2.03)	0.73
AICGTC	7.0	13.5	0.48(0.21-1.12)	0.09
CDTGTC	3.1	1.2	2.74(0.33-22.40)	0.35
AICGCT	1.8	4.8	0.27(0.04-1.91)	0.19

OR-odds ratio; CI-confidence interval

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Table 9. Association of *VEGF* haplotypes with breast cancer risk in pre menopausal obese and post menopausal obese patients

Haplotype*	Pre menopausal Obese (%) n=85	Post menopausal obese (%) n=102	OR(95%CI)	p value
AICGCC	25.0	35.1	1 (Reference)	
CDTCCC	21.8	25.5	1.13(0.60-2.13)	0.7
CDTGCC	27.6	15.0	1.98(1.10-3.56)	0.02
CDTGCT	1.8	2.7	1.45(0.27-7.71)	0.66
AICGTC	8.6	7.9	1.27(0.50-3.24)	0.62

OR-odds ratio; CI-confidence interval; Significant p values are shown in bold

*In order of -2578C/A, -2549I/D, -460T/C, +405C/G, -7C/T and +936C/T

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTable.docx](#)