

Association Between Recent Overnight Travel And Use Of Long-Lasting Insecticidal Nets In Rural Uganda: A Prospective Cohort Study In Tororo

Emmanuel Arinaitwe (✉ emmanuel.arinaitwe@yahoo.com)

Infectious Diseases Research Collaboration

Joaniter I. Nankabirwa

Infectious Diseases Research Collaboration, Kampala; Department of Medicine, Makerere University, Kampala

Paul Krezanoski

Department of Medicine, University of California, San Francisco, California

John Rek

Infectious Diseases Research Collaboration, Kampala

Victor Kamyra

Infectious Diseases Research Collaboration, Kampala

Adrienne Epstein

Department of Medicine, University of California, San Francisco, California

Philip J. Rosenthal

Department of Medicine, University of California, San Francisco, California

Chris Drakeley

London School of Hygiene and Tropical Medicine Faculty of Infectious and Tropical Diseases

Moses R. Kamyra

Department of Medicine, Makerere University, Kampala; Infectious Diseases Research Collaboration, Kampala

Grant Dorsey

Department of Medicine, University of California, San Francisco, California

Sarah G. Staedke

London School of Hygiene and Tropical Medicine Faculty of Infectious and Tropical Diseases

Research

Keywords: Human behaviour, Recent overnight travel, Malaria risk

Posted Date: July 21st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-45494/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 11th, 2020. See the published version at <https://doi.org/10.1186/s12936-020-03475-3>.

Abstract

Background: The burden of malaria in Uganda remains high, but has become increasingly heterogenous following intensified malaria control. Travel within Uganda is recognised as a risk factor for malaria, but behaviours associated with travel are not well-understood. To address this knowledge gap, malaria-relevant behaviours of cohort participants were assessed during travel and at home in Uganda.

Methods: Residents from 80 randomly selected households in Nagongera sub-county, Tororo district were enrolled into a cohort to study malaria in rural Uganda. All participants were given long-lasting insecticidal nets (LLINs) at enrolment and were evaluated every 4 weeks at the study clinic. Participants were asked if they had travelled overnight from their home, and if so, a questionnaire was administered to capture information on travel details and behaviours. Behaviour while at home was assessed using a similar questionnaire during two-weekly home visits. Behaviours while travelling vs at home were compared using generalized estimating equations adjusting for repeated measures in the same individual.

Results: Between October 2017 and October 2019, 527 participants were enrolled and assessed for travel. Of these, 123 (23.2%) reported taking 211 overnight trips; 149 (70.6%) trips were within Tororo. Participants were less likely to use LLINs when travelling than when at home (41.0% vs. 56.2%, relative risk [RR] 0.73, 95% CI: 0.60-0.89, $p=0.002$); this difference was noted for women (38.8% vs 59.2%, RR 0.66, 95% CI 0.52-0.83, $p=0.001$) but not men (48.3% vs 46.6%, RR 0.96, 95% CI 0.67-1.40, $p=0.85$). In an adjusted analysis, factors associated with LLIN use when travelling included destination (travelling to districts not receiving indoor residual spraying [IRS] 65.8% vs Tororo district 32.2%, RR 1.80, 95% CI 1.31-2.46, $p<0.001$) and duration of travel (>7 nights 60.3% vs one night 24.4%, RR 1.97, 95% CI 1.07-3.64, $p=0.03$).

Conclusions: Travellers, particularly women, were less likely to use LLINs when travelling than when at home. LLIN adherence was higher among those who travelled to non-IRS districts and for more than one week, suggesting that perceived malaria risk influences LLIN use. Strategies are needed to raise awareness of the importance of using LLINs while travelling.

Introduction

Malaria control interventions have been scaled-up globally, resulting in significant declines in malaria burden [1–3]. Despite these achievements, malaria morbidity remains high worldwide; in 2018, 228 million malaria cases, 93% from Africa, were reported [1]. In Uganda, key malaria control strategies include prompt treatment with artemisinin-based combination therapy (ACTs), universal distribution of long-lasting insecticidal nets (LLINs), and targeted indoor residual spraying of insecticides (IRS) [2, 4–6]. Although intervention coverage has expanded remarkably in Uganda over the past decade, progress on malaria control has been uneven [7]. According to the 2018-19 Malaria Indicator Survey, parasite prevalence, measured by microscopy in children under-five, ranged from 0.2% in Kampala to 34.3% in the

Karamoja region [7]. Factors contributing to the heterogeneity of malaria in Uganda include geographical variation in transmission intensity, increasing urbanisation, and delivery of IRS to a limited number of districts[8, 9].

Travel is a well-recognised risk factor for malaria [10–15]. Studies from Uganda and elsewhere in Africa have shown that overnight travel is associated with an increased risk of malaria, especially when individuals travel from areas of lower transmission intensity to higher risk areas [16, 17]. Although travellers may be at increased risk of malaria due to exposure to higher malaria transmission [18, 19], changes in behaviour while away from home may also contribute. Some studies have suggested that individuals who travel within malaria-endemic areas may take part in outdoor activities, go to bed late, and be less likely to use LLINs, all behaviours that increase exposure to mosquitoes and risk of malaria infection [20–23]. To further explore associations between overnight travel and behaviours that might modify the risk of malaria infection, data collected over a two-year period from a cohort of individuals living in Tororo, Uganda, were analysed.

Methods

Study site

The study was conducted in Nagongera sub-county in Tororo district, Uganda. Details about the site have been described elsewhere [16, 24, 25]. Briefly, Nagongera is a rural area with very high malaria transmission, which is now under intensive malaria control. In 2012, the entomological inoculation rate (EIR) in Tororo was 310 infectious bites per person per year [26]. Because of its high malaria burden, Tororo was selected to receive IRS starting in 2015, and to date the district has received seven rounds of IRS (three rounds of the carbamate Bendiocarb, followed by four rounds of the organophosphate pirimiphos-methyl [Actellic]). In addition to IRS, LLINs were distributed to all households in Tororo through national campaigns in 2013 and 2017. The interventions have been associated with a drastic reduction in key malaria indicators, including malaria incidence, parasite prevalence and EIR which was <1% in 2018 [5, 27, 28].

Study design and participant enrolment

The cohort study has been described in detail elsewhere [29]. Briefly, all households in the study area were enumerated, and 80 households were randomly selected for participation. Households were included if they met the following selection criteria: 1) at least two household members under 5 years of age, 2) no more than 7 permanent residents, 3) no intention for the household to move from Nagongera sub-county during the study period, and 4) willingness to participate in study follow up activities.

All members of the enrolled households were screened and enrolled in the cohort study if they met the following selection criteria: 1) full-time resident of the selected household, 2) agreement to come to the study clinic for any illness and scheduled follow up, and 3) provision of written informed consent. Participants were followed up for 2 years, and the cohort was dynamic; any residents that were born into

or joined the household were screened for enrolment during the course of the study. Participants were withdrawn from the study if they met the following criteria: 1) permanent movement out of Nagongera sub-county, 2) unable to be located for > 120 days, 3) withdrawal of informed consent, or 4) unable to comply with the study schedule and procedures. All enrolled participants were given LLINs at enrolment and were encouraged to come to the study clinic for all of their medical care.

Study participant follow up and data collection

Participants were seen at the study clinic monthly for routine follow up. At these visits, participants were asked whether they had travelled overnight since the last visit. A detailed questionnaire was administered to those who travelled to capture data on destination and duration of travel, behavioural factors such as time spent outdoors in the evening, time to bed, and use of LLINs during travel. Every two weeks, participants were visited at home and the same questionnaire was administered to collect data on behavioural factors while at home.

Statistical analysis

Data were collected by trained study staff using standardised case record forms and double-entered using Microsoft Access (Microsoft Corporation, Redmond, Washington, USA). All the analyses were performed using Stata, version 14 (Stata Corporation, College Station, Texas, USA). This analysis included data on any overnight travel and behavioural factors collected between October 2017 and October 2019. Overnight travel was defined as travel out of the sub-county of residence and spending at least one night away.

Behavioural factors including adherence to reported LLIN usage the prior night and time to bed were evaluated when study participants were at home and during overnight travel. LLINs use during travel was dichotomised into use most of the time during travel and no use, and time to bed was dichotomised into going to bed before 9pm most of the time and going to bed at 9pm or later mostly. Each overnight trip was paired with the most recent assessment at home as a comparison. For each pair, comparisons between behavioural factors during travel and while at home were made using generalised estimating equations adjusting for repeated measures in the same individual, and estimates were reported as relative risks (RR). In addition, an analysis of factors associated with LLIN adherence, such as destination and duration of travel, time to bed during travel, gender and age at time of travel, were also assessed using generalised estimating equations and expressed as relative risks. A p-value <0.05 was considered statistically significant.

Results

Characteristics of study participants

From October 2017 through October 2019, 531 study participants from 80 randomly selected households were enrolled (Fig. 1). Of these, 527 were assessed for overnight travel. Overall, 123 (23.2%) participants

reported at least one overnight trip and were included in the analysis (Table 1). Of these participants who travelled, 65.9% were female. Adults were more likely to travel than children, but school-aged children (5–15 years) were more likely to take longer trips (17 days) than younger children (7 days) or adults (3 days). Most participants travelled short distances (< 30 km) and generally stayed within Tororo district (70.6%). Travel destinations outside Tororo included Bugiri (65 km away) and Butaleja (25 km away), both districts receiving IRS (Fig. 2). Other participants travelled to non-IRS districts (18.0%), and to Kampala (8.1%). The main reason older children and adults travelled was to visit relatives, while children under-five mainly accompanied their parents or guardians. Travellers most commonly stayed with their relatives while away.

Table 1
 Characteristics of study participants and individual overnight trips

Characteristic	Age categories			
	All ages	< 5 years	5–15 years	> 15 years
Characteristics of study participants*				
Participants with any overnight trip, n (% total participants)	123 (23.2%)	39 (22.0%)	21 (11.1%)	63 (39.1%)
Female gender, n (% among participants with any travel)	81 (65.9%)	20 (51.3%)	13 (61.9%)	48 (76.2%)
Characteristics of individual overnight trips**				
Number of overnight trips	211	53	34	124
Duration of each trip, median (range)	5 (1-115)	7(1–53)	17 (1–53)	3 (1-115)
Duration of travel categories, n (% total trips)				
1 night	41 (19.4%)	7 (13.2%)	1 (2.9%)	33 (26.6%)
2–3 nights	49 (23.2%)	9 (17.0%)	4 (11.8%)	36 (29.0%)
4–7 nights	43 (20.4%)	11 (20.8%)	6 (17.7%)	26 (21.0%)
More than 7 nights	78 (37.0%)	26 (49.1%)	23 (67.7%)	29 (23.4%)
Destination of travel, n (% total trips)				
Tororo (IRS district)	149 (70.6%)	37 (69.8%)	22 (64.7%)	90 (72.6%)
Other IRS districts	7 (3.3%)	1 (1.9%)	1 (2.9%)	5 (4.0%)
Kampala (no IRS)	17 (8.1%)	5 (9.4%)	7 (20.6%)	5 (4.0%)
Other non-IRS districts	38 (18.0%)	10 (18.9%)	4 (11.8%)	24 (19.4%)
Reason for travel, n (% total trips)				
Visiting relatives	106 (50.2%)	20 (37.7%)	27 (79.4%)	59 (47.6%)
* based on age at the time of study enrolment				
** based on age at the time of travel				

Characteristic	Age categories			
	All ages	< 5 years	5–15 years	> 15 years
Funeral rite	45 (21.3%)	1 (1.9%)	0	44 (35.5%)
Accompanying parents	39 (18.5%)	32 (60.4%)	7 (20.6%)	0
Caring for the sick	10 (4.7%)	0	0	10 (8.1%)
Business	8 (3.8%)	0	0	8 (6.5%)
Pleasure	2 (1.0%)	0	0	2 (1.6%)
Attending school	1 (0.5%)	0	0	1 (0.8%)
Where participant stayed, n (% total trips)				
Friend/relative's home	176 (83.4%)	49 (92.5%)	33 (97.1%)	94 (75.8%)
Hospital	13 (6.2%)	2 (3.8%)	1 (2.9%)	10 (8.1%)
Camp or Gardens	22 (10.4%)	2 (3.8%)	0	20 (16.1%)
* based on age at the time of study enrolment				
** based on age at the time of travel				

Differences in behaviour during overnight travel versus at home

Overall, LLIN use among cohort participants who travelled was low both at home and while travelling. Participants were significantly less likely to use LLINs when they travelled than when at home (41.0% travel vs. 56.2% home, relative risk (RR) 0.73, 95% CI: 0.60–0.89, $p = 0.002$) (Table 2). However, this difference was modified by gender and age. Women were less likely to use LLINs when they travelled than when at home (38.8% vs 59.2%; RR 0.66, 95% CI: 0.52–0.83, $p = 0.001$) but no difference in LLIN use was observed for men (46.6% vs 48.3%; RR 0.96, 95% CI: 0.67–1.40, $p = 0.85$). Stratifying by age, no differences in LLIN use during travel were observed for younger or school-aged children. However, older participants (> 15 years) were significantly less likely to use LLINs when travelling than at home (33.9% vs 61.3%; RR 0.55, 95% CI: 0.41–0.74, $p < 0.001$). Overall, there were no differences in the proportion of participants who went to bed after 9 pm when travelling versus at home.

Table 2
Comparison of behavioural factors at home of residence and during overnight travel

Behavioural factor	Groups		Number of paired observations	Number with a reported behavioural factor (%)		RR (95% CI)	p-value
				At home of residence	During overnight travel		
Sleeping under an LLIN	All		210	118 (56.2%)	86 (41.0%)	0.73 (0.60–0.89)	0.002
	Gender	Male	58	28 (48.3%)	27 (46.6%)	0.96 (0.67–1.40)	0.85
		Female	152	90 (59.2%)	59 (38.8%)	0.66 (0.52–0.83)	0.001
	Age	< 5 years	53	26 (49.1%)	24 (45.3%)	0.92 (0.62–1.38)	0.70
		5–15 years	33	16 (48.5%)	20 (60.6%)	1.25 (0.86–1.82)	0.25
		> 15 years	124	76 (61.3%)	42 (33.9%)	0.55 (0.41–0.74)	< 0.001
Going to bed after 9 pm	All		210	147 (70.0%)	149 (71.0%)	1.01 (0.91–1.13)	0.80
	Gender	Male	58	27 (46.6%)	31 (53.5%)	1.15 (0.86–1.53)	0.35
		Female	152	120 (79.0%)	118 (77.6%)	0.98 (0.88–1.10)	0.77
	Age	< 5 years	53	15 (28.3%)	20 (37.7%)	1.33 (0.77–2.31)	0.30
		5–15 years	33	23 (69.7%)	19 (57.6%)	0.83 (0.58–1.17)	0.28
		> 15 years	124	109 (87.9%)	110 (88.7%)	1.01 (0.92–1.11)	0.85

Factors associated with LLIN adherence during overnight travel

Participants who travelled to districts without an IRS program were more likely to sleep under LLINs than those who travelled within Tororo district (65.8% vs 32.2%; RR 1.80, 95% CI: 1.31–2.46, $p < 0.001$) (Table 3). There were no differences in LLIN use when travelling to other destinations (Kampala and other IRS districts). Participants who travelled for more than 7 nights were significantly more likely to use LLINs while travelling than those who travelled for only one night (60.3% vs 24.4%; RR 1.97, 95% CI: 1.07–3.64, $p = 0.03$). Other factors that were not associated with LLIN use during travel included time to bed, gender of the participant, and age at the time of travel.

Table 3
Factors associated with LLIN adherence during overnight travel

Factors	Categories	Proportion of trips adherent to LLINs (%)	Univariable analysis		Multivariable analysis	
			RR (95% CI)	p-value	RR (95% CI)	p-value
Destination of travel	Tororo district (IRS)	48/149 (32.2%)	Reference	-	Reference	-
	Other IRS districts	1/7 (14.3%)	0.44 (0.07–2.77)	0.39	0.39 (0.05–2.88)	0.35
	Kampala district	12/17 (70.6%)	2.19 (1.49–3.22)	< 0.001	1.49 (0.93–2.40)	0.10
	Other non-IRS districts	25/38 (65.8%)	2.04 (1.47–2.83)	< 0.001	1.80 (1.31–2.46)	< 0.001
Duration of travel	1 night	10/41 (24.4%)	Reference	-	Reference	-
	2–3 nights	11/49 (22.5%)	0.92 (0.43–1.95)	0.83	0.90 (0.43–1.90)	0.79
	4–7 nights	18/43 (41.9%)	1.72 (0.90–3.27)	0.10	1.44 (0.75–2.80)	0.27
	More than 7 nights	47/78 (60.3%)	2.47 (1.40–4.37)	0.002	1.97 (1.07–3.64)	0.03
Time to bed during travel	Before 9 pm	28/62 (45.2%)	Reference	-	Reference	-
	9 pm or later	58/149 (38.9%)	0.86 (0.61–1.21)	0.39	1.08 (0.72–1.62)	0.73
Gender	Male	27/58 (46.6%)	Reference	-	Reference	-
	Female	59/153 (38.6%)	0.83 (0.59–1.17)	0.28	1.00 (0.70–1.44)	0.99
Age at time of travel	< 5 years	24/53 (45.3%)	Reference	-	Reference	-
	5–15 years	20/34 (58.8%)	1.30 (0.86–1.96)	0.21	1.16 (0.78–1.73)	0.47
	> 15 years	42/124 (33.9%)	0.75 (0.51–1.10)	0.14	0.87 (0.56–1.37)	0.55

Discussion

To better understand behavioural factors that might modify the risk of malaria during travel, a cohort of individuals living in Tororo under highly effective malaria control were assessed. Overall, LLIN use in travellers was low, and participants were less likely to use LLINs when they travelled than when at home. However, this finding was true only for women, and adults. Factors associated with higher LLIN use while travelling included travel to non-IRS districts, and travelling for more than one week, suggesting that perceived risk of malaria may influence the decision to sleep under an LLIN while away from home.

Travel has been identified as a risk factor for malaria across Africa [11, 15, 19, 30, 31]. In Uganda, a study conducted at three sites of varied malaria transmission demonstrated that the incidence of malaria in travellers was over three times higher in the 60 days after overnight travel compared to the 60 days before travelling [16]. Another study in western Uganda found that travelling within the previous 4 weeks from highland areas with low malaria transmission to higher transmission areas was strongly associated with increased malaria risk [17]. Two studies conducted on Bioko Island, Equatorial Guinea, demonstrated that island residents who travelled to the mainland were at increased risk of malaria infection [19, 30]. A survey of Bioko island travellers found that malaria prevalence was significantly higher in passengers returning to the island from the mainland compared to those departing the island [30]. The odds of malaria among Bioko Island residents who travelled was significantly higher than in non-travellers, suggesting that imported malaria cases contributed to the sustained transmission of malaria on the island [19]. Similarly, a study in northern Ethiopia found that travel from high-altitude (low transmission) villages to other areas within the previous month was associated with increased odds of malaria [31]. This evidence suggests that travel within Africa is a risk factor for malaria infection. However, behavioural factors associated with travel that might increase exposure to mosquito vectors, and thus malaria infection, have been less well-explored.

There are several potential reasons why people may be at increased risk of malaria during travel. In this study, gender differences in LLIN use while travelling were observed. Women were less likely to use LLINs when travelling than at home, but this was not true for men. Interestingly, women reported using LLINs more often than men when at home; however, when travelling, the opposite was true. This suggests that at home, women may be more aware of the importance of sleeping under LLINs to protect against malaria, perhaps reflecting routine distribution of LLINs at antenatal clinics and targeted campaigns to increase LLIN use among pregnant women [32–34]. When travelling, women may either lack LLINs or the agency to use them, particularly when visiting the home of a friend or relative.

Individuals who travelled to non-IRS districts and those who travelled for more than 7 days were more likely to use LLINs. These findings suggest that the decision to use LLINs may be influenced by destination or duration of travel and the individuals' perceptions of malaria risk. Indeed, perceptions of malaria risk have been shown to influence the use of LLINs when people travel [20]. In south-eastern Tanzania, in-depth interviews were used to assess perceptions of malaria risk during outdoor and indoor activities. In this study, participants believed that outdoor activities such as fishing in the river late at night, travelling to farms overnight, and attending parties and funerals held at night, all increased their risk of malaria infection. For situations where use of LLINs was not feasible, participants believed that alternative malaria prevention approaches, including use of mosquito repellents and chemoprophylaxis, were needed.

LLINs are known to reduce malaria morbidity and mortality and are widely used for vector control in Africa [35], but achieving high adherence to LLINs, even at home, is challenging. In this study, just over half of cohort participants who travelled slept under LLINs when at home, despite universal access. Many barriers to LLIN use have been described, including many household members [36, 37], lack of space to

hang LLINs [38], lower socioeconomic status, and time since the last LLINs distribution [39]. In this study setting, where malaria transmission dropped substantially, individuals may have felt that it was no longer necessary to use their LLINs [5, 28]. During travel, a possible barrier to LLIN adherence is limited availability of LLINs to use away from home. Mass distribution of LLINs in Uganda follows WHO guidelines, which recommend distributing one LLIN for every two household residents [40]. This may leave no spare LLINs for visitors, or for carrying during travel. In this study, other factors that may have contributed to limited use of LLINs during travel include social barriers, such as attending a funeral or wedding where individuals are expected to stay outdoors all night, or fear of appearing rude or disrespectful during communal gatherings [22]. These factors should be considered when designing strategies to increase LLIN adherence in travellers.

A strength of this study is that behaviours at home and during travel within the same individuals were prospectively compared, minimizing the potential for confounding. Similar studies have only assessed malaria-relevant behaviours while travelling, or at home, but not both [23, 41, 42]. A study conducted in south-eastern Tanzania evaluated human behaviour of participants at home [20]. The study found that a high proportion of participants (75%) stayed outdoors in the evenings (between 6 pm and 9 pm), resulting in exposure to malaria vectors before going to bed. Another study carried out in the Kilombero Valley of Tanzania from November 2015 to March 2016, assessed patterns of behaviour only when travelling, and demonstrated that when individuals travelled for religious, cultural and social gatherings, they stayed outdoors at night till dawn [41]. Previous studies in Uganda that have assessed travel and malaria risk also examined behavioural factors during travel, such as use of LLINs [16, 17]. However, differences in behaviour while travelling versus at home were not explored. The findings from this study suggest that a better understanding of circumstances leading to lower use of LLINs when travelling may be important in guiding malaria prevention measures.

This study had several limitations. First, data on behavioural factors during travel could have been subject to recall bias. However, questionnaires were administered within 4 weeks following travel, and adherence to LLINs at home was assessed every two weeks by home visits, to closely evaluate the relationship between behaviours at home and when travelling. Second, the study was conducted in rural Tororo, and few individuals travelled outside of the district. Thus, results may not be generalisable to other settings. Lastly, intensive malaria control with IRS and LLINs resulted in few malaria cases in Tororo. Thus, it was not possible to directly measure the association between behaviours and malaria risk.

Conclusion

Travel is an important individual risk factor for malaria, and individuals who travel may also threaten malaria control gains, especially in areas on a pathway to elimination. Results from this study suggest that individuals were less likely to use LLINs when travelling. Strategies to increase awareness about the importance of LLIN adherence, particularly in travellers, should be developed and deployed by the National Malaria Control Division of the Ministry of Health, or other stakeholders. Use of LLIN during

travel, especially during the holiday season when most people are likely to visit family and friends, should be emphasised. Travellers should be encouraged to carry an extra LLIN when travelling, especially when visiting rural areas or those without ongoing IRS. Further research on innovative approaches to prevent malaria in travellers including portable LLINs, effective ways to influence behaviour and increase LLIN use, and acceptability of other malaria prevention measures such as mosquito repellents and chemoprophylaxis, should be encouraged.

Declarations

Acknowledgement

We thank the study participants for their involvement in this study, and grateful to Infectious Diseases Research Collaboration for administrative support.

Author's contributions

EA, JIN, JR, PJR, CD, MRK, GD, and SGS conceived and designed the study. EA, JIN, JR, VK, and AE participated in the data collection. EA, JIN, PK, VK, AE, MRK, GD, CD, and SGS participated in the management and analysis of data. All authors participated in writing of the manuscript, and approved the final manuscript.

Funding

This work was supported by the Fogarty International Centers of the National Institutes of Health under Award Number D43TW010526 and D43TW7375. JIN is supported by the Fogarty International Center (Emerging Global Leader Award grant number K43TW010365). The main PRISM study was funded by the National Institute of Allergy and Infectious Diseases (NIAID) as part of the International Centers of Excellence in Malaria Research (ICEMR) Program (U19AI089674).

Disclaimer. The content of this report is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders of this study did not participate in the study design, data collection, data analysis, and data interpretation or writing of findings.

Availability of data and study materials

Data and study material are available on request by an e-mail to the corresponding author.

Ethics approval and consent to participate

Study approvals were obtained from the Makerere University School of Medicine Research and Ethics Committee (ref. 2017-099), the Uganda National Council for Science and Technology (ref. HS-119ES), the London School of Hygiene & Tropical Medicine Ethics Committee (ref. 14266), and the University of California, San Francisco Committee on Human Research (ref. 17-22544).

All study participants provided written informed consent (with assent from minors) before enrolment. All study staff received training on procedures for maintaining confidentiality.

Consent for publication

All authors held final responsibility for the decision to submit and consented to this publication.

Conflict of interest

All authors declare no conflict of interest.

References

1. World Health Organization: World malaria report 2019. 2019.
2. Bhatt S, Weiss DJ, Cameron E, Bisanzio D, Mappin B, Dalrymple U, Battle KE, Moyes CL, Henry A, Eckhoff PA, et al: The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* 2015, 526:207-211.
3. Weiss DJ, Lucas TCD, Nguyen M, Nandi AK, Bisanzio D, Battle KE, Cameron E, Twohig KA, Pfeffer DA, Rozier JA, et al: Mapping the global prevalence, incidence, and mortality of *Plasmodium falciparum*, 2000-17: a spatial and temporal modelling study. *Lancet* 2019, 394:322-331.
4. Uganda Bureau of Statistics (UBOS): Uganda Malaria Indicator Survey 2014-2015. 2015.
5. Zinszer K, Charland K, Vahey S, Jahagirdar D, Rek JC, Arinaitwe E, Nankabirwa J, Morrison K, Sadoine ML, Tutt-Guerette MA, et al: The impact of multiple rounds of indoor residual spraying on malaria incidence and haemoglobin levels in a high transmission setting. *J Infect Dis* 2019.
6. Oguttu DW, Matovu JKB, Okumu DC, Ario AR, Okullo AE, Opigo J, Nankabirwa V: Rapid reduction of malaria following introduction of vector control interventions in Tororo District, Uganda: a descriptive study. *Malar J* 2017, 16:227.
7. Uganda Bureau of Statistics (UBOS): Uganda Malaria Indicator Survey 2018-2019. 2019.
8. The PMI VectorLink project, Uganda [<https://pmivectorlink.org/where-we-work/uganda/>]
9. Kigozi SP, Pindolia DK, Smith DL, Arinaitwe E, Katureebe A, Kilama M, Nankabirwa J, Lindsay SW, Staedke SG, Dorsey G, et al: Associations between urbanicity and malaria at local scales in Uganda. *Malar J* 2015, 14:374.
10. Le Menach A, Tatem AJ, Cohen JM, Hay SI, Randell H, Patil AP, Smith DL: Travel risk, malaria importation and malaria transmission in Zanzibar. *Sci Rep* 2011, 1:93.
11. Marshall JM, Toure M, Ouedraogo AL, Ndhlovu M, Kiware SS, Rezai A, Nkhama E, Griffin JT, Hollingsworth TD, Doumbia S, et al: Key traveller groups of relevance to spatial malaria transmission: a survey of movement patterns in four sub-Saharan African countries. *Malar J* 2016, 15:200.
12. Shanks GD, Biomndo K, Guyatt HL, Snow RW: Travel as a risk factor for uncomplicated *Plasmodium falciparum* malaria in the highlands of western Kenya. *Trans R Soc Trop Med Hyg* 2005, 99:71-74.

13. Yukich JO, Taylor C, Eisele TP, Reithinger R, Nauhassenay H, Berhane Y, Keating J: Travel history and malaria infection risk in a low-transmission setting in Ethiopia: a case control study. *Malaria Journal* 2013, 12.
14. Malede A, Alemu K, Aemero M, Robele S, Kloos H: Travel to farms in the lowlands and inadequate malaria information significantly predict malaria in villages around Lake Tana, northwest Ethiopia: a matched case-control study. *Malar J* 2018, 17:290.
15. Lowa M, Sitali L, Siame M, Musonda P: Human mobility and factors associated with malaria importation in Lusaka district, Zambia: a descriptive cross sectional study. *Malar J* 2018, 17:404.
16. Arinaitwe E, Dorsey G, Nankabirwa JI, Kigozi SP, Katureebe A, Kakande E, Rek J, Rosenthal PJ, Drakeley C, Kanya MR, Staedke SG: Association Between Recent Overnight Travel and Risk of Malaria: A Prospective Cohort Study at 3 Sites in Uganda. *Clin Infect Dis* 2019, 68:313-320.
17. Lynch CA, Bruce J, Bhasin A, Roper C, Cox J, Abeku TA: Association between recent internal travel and malaria in Ugandan highland and highland fringe areas. *Trop Med Int Health* 2015, 20:773-780.
18. Smith JL, Auala J, Haindongo E, Uusiku P, Gosling R, Kleinschmidt I, Mumbengegwi D, Sturrock HJ: Malaria risk in young male travellers but local transmission persists: a case-control study in low transmission Namibia. *Malar J* 2017, 16:70.
19. Guerra CA, Kang SY, Citron DT, Hergott DEB, Perry M, Smith J, Phiri WP, Osa Nfumu JO, Mba Eyono JN, Battle KE, et al: Human mobility patterns and malaria importation on Bioko Island. *Nat Commun* 2019, 10:2332.
20. Finda MF, Moshi IR, Monroe A, Limwagu AJ, Nyoni AP, Swai JK, Ngowo HS, Minja EG, Toe LP, Kaindoa EW, et al: Linking human behaviours and malaria vector biting risk in south-eastern Tanzania. *PLoS One* 2019, 14:e0217414.
21. Koenker HM, Loll D, Rweyemamu D, Ali AS: A good night's sleep and the habit of net use: perceptions of risk and reasons for bed net use in Bukoba and Zanzibar. *Malar J* 2013, 12:203.
22. Monroe A, Harvey SA, Lam Y, Muhangi D, Loll D, Kabali AT, Weber R: "People will say that I am proud": a qualitative study of barriers to bed net use away from home in four Ugandan districts. *Malar J* 2014, 13:82.
23. Monroe A, Mihayo K, Okumu F, Finda M, Moore S, Koenker H, Lynch M, Haji K, Abbas F, Ali A, et al: Human behaviour and residual malaria transmission in Zanzibar: findings from in-depth interviews and direct observation of community events. *Malar J* 2019, 18:220.
24. Nankabirwa JI, Briggs J, Rek J, Arinaitwe E, Nayebare P, Katrak S, Staedke SG, Rosenthal PJ, Rodriguez-Barraquer I, Kanya MR, et al: Persistent Parasitemia Despite Dramatic Reduction in Malaria Incidence After 3 Rounds of Indoor Residual Spraying in Tororo, Uganda. *J Infect Dis* 2019, 219:1104-1111.
25. Kanya MR, Arinaitwe E, Wanzira H, Katureebe A, Barusya C, Kigozi SP, Kilama M, Tatem AJ, Rosenthal PJ, Drakeley C, et al: Malaria transmission, infection, and disease at three sites with varied transmission intensity in Uganda: implications for malaria control. *Am J Trop Med Hyg* 2015, 92:903-912.

26. Kilama M, Smith DL, Hutchinson R, Kigozi R, Yeka A, Lavoy G, Kanya MR, Staedke SG, Donnelly MJ, Drakeley C, et al: Estimating the annual entomological inoculation rate for Plasmodium falciparum transmitted by Anopheles gambiae s.l. using three sampling methods in three sites in Uganda. *Malar J* 2014, 13:111.
27. Katureebe A, Zinszer K, Arinaitwe E, Rek J, Kakande E, Charland K, Kigozi R, Kilama M, Nankabirwa J, Yeka A, et al: Measures of Malaria Burden after Long-Lasting Insecticidal Net Distribution and Indoor Residual Spraying at Three Sites in Uganda: A Prospective Observational Study. *PLoS Med* 2016, 13:e1002167.
28. Musiime AK, Smith DL, Kilama M, Rek J, Arinaitwe E, Nankabirwa JI, Kanya MR, Conrad MD, Dorsey G, Akol AM, et al: Impact of vector control interventions on malaria transmission intensity, outdoor vector biting rates and Anopheles mosquito species composition in Tororo, Uganda. *Malar J* 2019, 18:445.
29. Nankabirwa JI, Arinaitwe E, Rek J, Kilama M, Kizza T, Staedke SG, Rosenthal PJ, Rodriguez-Barraquer, Briggs J, Greenhouse B, et al: Malaria Transmission, Infection and Disease Following Sustained Indoor Residual Spraying of Insecticide in Tororo, Uganda. *Am J Trop Med Hyg* 2020.
30. Bradley J, Monti F, Rehman AM, Schwabe C, Vargas D, Garcia G, Hergott D, Riloha M, Kleinschmidt I: Infection importation: a key challenge to malaria elimination on Bioko Island, Equatorial Guinea. *Malar J* 2015, 14:46.
31. Haile M, Lemma H, Weldu Y: Population Movement as a Risk Factor for Malaria Infection in High-Altitude Villages of Tahtay-Maychew District, Tigray, Northern Ethiopia: A Case-Control Study. *Am J Trop Med Hyg* 2017, 97:726-732.
32. Hill J, Hoyt J, van Eijk AM, ter Kuile FO, Webster J, Steketee RW: Prioritizing pregnant women for long-lasting insecticide treated nets through antenatal care clinics. *PLoS Med* 2014, 11:e1001717.
33. Muhumuza E, Namuhani N, Balugaba BE, Namata J, Ekirapa Kiracho E: Factors associated with use of malaria control interventions by pregnant women in Buwunga subcounty, Bugiri District. *Malar J* 2016, 15:342.
34. Theiss-Nyland K, Lynch M, Lines J: Assessing the availability of LLINs for continuous distribution through routine antenatal care and the Expanded Programme on Immunizations in sub-Saharan Africa. *Malar J* 2016, 15:255.
35. Lengeler C: Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2004:CD000363.
36. Buchwald AG, Walldorf JA, Cohee LM, Coalson JE, Chimbiya N, Bauleni A, Nkanaunena K, Ngwira A, Kapito-Tembo A, Mathanga DP, et al: Bed net use among school-aged children after a universal bed net campaign in Malawi. *Malar J* 2016, 15:127.
37. Wanzira H, Katamba H, Rubahika D: Use of long-lasting insecticide-treated bed nets in a population with universal coverage following a mass distribution campaign in Uganda. *Malar J* 2016, 15:311.
38. Tassew A, Hopkins R, Deressa W: Factors influencing the ownership and utilization of long-lasting insecticidal nets for malaria prevention in Ethiopia. *Malar J* 2017, 16:262.

39. Gonahasa S, Maiteki-Sebuguzi C, Rugnao S, Dorsey G, Opigo J, Yeka A, Katureebe A, Kyohere M, Lynd A, Hemingway J, et al: LLIN Evaluation in Uganda Project (LLINEUP): factors associated with ownership and use of long-lasting insecticidal nets in Uganda: a cross-sectional survey of 48 districts. *Malar J* 2018, 17:421.
40. World Health Organization: Achieving and maintaining universal coverage with long-lasting insecticidal nets for malaria control 2017.
41. Moshi IR, Manderson L, Ngowo HS, Mlacha YP, Okumu FO, Mnyone LL: Outdoor malaria transmission risks and social life: a qualitative study in South-Eastern Tanzania. *Malar J* 2018, 17:397.
42. Monroe A, Asamoah O, Lam Y, Koenker H, Psychas P, Lynch M, Ricotta E, Hornston S, Berman A, Harvey SA: Outdoor-sleeping and other night-time activities in northern Ghana: implications for residual transmission and malaria prevention. *Malar J* 2015, 14:35.

Figures




Figure 1

Study profile.




Figure 2

Map of Uganda showing travel destination of study participants to the districts level.