

Morphological, Biochemical and Molecular Characterization of Indigenous *Aromatic* Rice of Assam

Dibosh Bordoloi

Assam Agricultural University Faculty of Agriculture

Debojit Sarma (✉ debojit.sarma@aaau.ac.in)

Assam Agricultural University <https://orcid.org/0000-0002-3808-7231>

Nagendra Sarma Barua

Assam Agricultural University Faculty of Agriculture

Bikram Kishore Das

Bhabha Atomic Research Centre

Research Article

Keywords: Joha rice, DUS characteristics, genetic variability, divergence, fatty acid profile, molecular diversity

Posted Date: May 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-455040/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

We carried out the morphological, biochemical and molecular characterization of 20 indigenous *Joha* (aromatic) rice cultivars of Assam. Distinctiveness, Uniformity and Stability (DUS) characterization of the cultivars revealed polymorphism in thirty-seven traits, establishing distinctiveness for their utilization in breeding programmes. Unweighted Neighbour Joining (UNJ) clustering based on usual Euclidean distances for the polymorphic markers grouped the cultivars into three multi-genotypic clusters. The *Joha* rice cultivars showed highly significant differences for all the quantitative traits except for panicle length. The genotypic and phenotypic coefficients of variability (GCV & PCV) were high for grain yield ha^{-1} (24.62 & 24.85%) and filled grains panicle $^{-1}$ (23.69 & 25.02%). All the traits except days to flowering and maturity, flag leaf breadth and spikelet fertility exhibited high heritability along with high to moderate genetic advance, indicating the predominant role of additive gene action. Mahalanobis D^2 analysis revealed three multi-genotypic and four mono-genotypic clusters of the cultivars. The cultivars' average polyunsaturated fatty acids were 37.9% oleic acid, 39.22% linoleic acid and 0.5% linolenic acid. The fatty acid profile of *Local Joha* was superior to the other cultivars as it showed a high level of linoleic and linolenic acid and low saturated fatty acid content. *Kon Joha 4* and *Ronga Joha* contained the highest iron (82.88 mg kg^{-1}) and zinc (47.39 mg kg^{-1}), respectively, while protein content of *Kon Joha-1* and amylose content of *Harinarayan* were the highest. *Joha-Bihpuria* showed the highest gel consistency of 140.50 mm. *Kalijeera*, *Kunkuni Joha*, *Kon Joha-5*, *Manimuni Joha* and *Kon Joha-2* accorded a strong aroma. PCR amplified 174 alleles with a mean value of 2.64 across the 66 polymorphic SSR markers. PIC values ranged from 0.091 to 0.698, with an average of 0.326. The highly informative (PIC > 0.50) markers were RM316, RM283, RM585, RM1388, RM3562, RM171, R1M30, RM118, RM11 and RM29 for identification of the twenty aromatic rice cultivars. The UNJ clustering based on Jaccard's coefficients classified the 20 cultivars into three distinct clusters with eight, ten and two entries.

Introduction

Globally rice is planted to about 162 million ha and 755 million tonnes of produce harvested annually (FAO, 2020). Ninety per cent of the production and consumption of the world's rice occur in Asia. India has the world's largest area under rice with 44.0 million ha and is the second-largest producer (96.0 million tonnes in 2010), next only to China. It contributes 21.5 per cent of global rice production. The FAO (2009) estimated a requirement of 70 per cent more food for over nine billion people expected to inhabit planet earth by 2050. As the staple food for most Indians, rice's future demand would increase with the growing population, projected at 1.378 billion by 2030 (Goyal and Singh, 2002). Assam has a wide range of variation of cultivated rice along with several new wild varieties. In Assam, rice cultivation occupies about 70% of the total cropped area that dominates the state's agriculture scenario (Singh et al., 2009). Rice, being the single primary source of agricultural GDP, plays a significant role in the state economy.

Aromatic rice is a significantly small but significant sub-group of rice gained in general popularity for consumers. The fragrance of *Joha* rice is the crucial feature, which increases its popularity in the international market (Nayak et al., 2002). Aromatic rice constitutes a particular group, which is considered the best in quality and fetches a much higher price than non-aromatic rice (Hien et al., 2007). A unique class of aromatic rice grown as winter rice in Assam is very popular and highly valued due to its quality, popularly known as *Joha* rice in Assam. The traditional fragrant *Joha* rice is best known for its superfine kernel, unique aroma, better cooking properties, as well as excellent palatability (Das et al., 2010).

Despite domestic aromatic rice, there are large numbers of wild varieties, which are still unexplored. The agricultural productivity of *Joha rice* is the lowest with the demand for production, and the gap is increasing over time. Agro-morphological characterization of rice germplasm is fundamental to provide information for plant breeding programmes (Lin, 1991). There are several agro-morphological markers used in the description and study of rice germplasm diversity. The nutritional status of rice is becoming increasingly important among consumers because of deficiency disorders. There is limited knowledge regarding the nutritional composition of different rice varieties available in Assam. However, the characterization of the rice cultivars in terms of composition and physicochemical aspects is relatively meagre. Rice is a leading food crop that possesses a different nutritional status that helps alleviate poverty (Otegbayo et al., 2001). The proximate analysis is an essential parameter for the routine description of foodstuffs. This analysis consists of six fractions viz., amylose, crude protein, iron, zinc and fatty acid content. The proximate composition, sensory, gelatinization temperature, and alkali digestion analysis of rice are physicochemical and biochemical analysis, which help generate nutritionally complete rice varieties (Dong et al., 2007).

Rice showed a genetically greater extent of diversity. DNA-based molecular markers play essential roles in assessing genetic variation and the elucidation of gene relationships within and among species. Molecular markers have been extensively used in crop improvement research throughout the globe as an appropriate and effective tool addressing biological parameters in agricultural productivity (Jones et al., 1997). Among PCR-based markers, SSR or microsatellites markers are excellent due to their locus of identity, high polymorphisms, and multi-allelic nature. Moreover, SSR markers are tandemly interspersed repeats throughout the genome that get amplified through primers that flank these regions. Studies to characterize and improve the aromatic rice of Assam are limited, even though this region of the country stores enormous rice diversity. The present study aimed at characterizing the aromatic rice of Assam using morphological, biochemical and molecular markers for their usage in breeding manipulation.

Material And Methods

The experiments were carried out during *the Sali* season of 2018 and 2019 at Instruction-cum-Research (ICR) Farm, Assam Agricultural University. The site is located at 26°45' north latitude and 94°12' east longitude having an elevation of 86.6 m above the mean sea level. The experimental area was relatively uniform in respect of topography and fertility. All the molecular works, including DNA extraction, PCR and gel electrophoresis, were done in Mutation Breeding Section-I Laboratory of Nuclear Agriculture and Biotechnology Division (NA&BTD), Bhabha Atomic Research Centre, Trombay. The twenty aromatic *Joha* rice cultivars (Table 1) were grown in randomized complete block design with three replications.

The seedlings' age was 30 days at transplanting in the main field. Each genotype constituted ten rows of 2.5 m long spaced 20 cm apart with one seedling per hill. Gap filling was done in the first week of transplanting to maintain a uniform plant population. A fertilizer dose of 60 kg N, 20 kg P_2O_5 , and 40 kg K_2O was

applied as per Sali rice recommendation for Assam. The entire quantity of P_2O_5 and K_2O and a half dose of N were applied as basal dose at the time of final field preparation; the remaining nitrogen applied in two equal splits at maximum tillering and booting stage. The standard agronomic practices recommended for the state of Assam were adopted in both experiments. Observations were recorded according to the National Test Guidelines for DUS test in rice developed by the Directorate of Rice Research, Hyderabad (Shobha Rani et al., 2004). The yield attributing traits were based on five random plants per replication, while days to flowering and maturity recorded on a plot basis. The characteristics observed were days to first flowering, days to 50% flowering, days to maturity, number of productive tillers plant⁻¹, plant height (cm), panicle length (cm), spikelet fertility (%), 1000-grain weights (g), grain yield plant⁻¹ (g), biological yield plant⁻¹ (g), harvest index (%), protein content (Kjeldahl's method), iron (Fe) and zinc (Zn) content (AAS), fatty acid profiling in rice bran.

Genomic DNA extraction

Seeds of cultivars were germinated and grown in a growth chamber, maintaining a temperature of 30°C, 10 hours of light and 85% relative humidity. At the three-leaf stages, the leaves were harvested by liquid nitrogen and taken for DNA isolation. The genomic DNA then was isolated by the CTAB (Cetyl Trimethyl Ammonium Bromide) method (Murray and Thompson, 1980).

DNA quantification

The concentration and quality of genomic DNA samples were estimated on a 260, 280 nm, and 230 nm spectrophotometer, the samples with a 260/280 ratio exceeding 1.8 were considered good quality DNA. Quality of DNA fragment was also confirmed by 0.8% agarose gel electrophoresis using 1XTBE buffer at 100V for 90 min.

Molecular marker selection

Seventy-one SSR markers were used for the genotyping (**Supplementary Table S1**). The markers were selected from various published literatures as well as from Gramene database (www.gramene.org) maintaining the genome wide distribution of markers.

PCR amplification using SSR primers

PCR reaction was performed in 25 µL mixture containing 1 µL (25 ng/ µL) template DNA, 2.5 µL of 10x PCR buffer with 25 mM $MgCl_2$, 1.0 µL 5 mM of each forward and reverse SSR primers, 1.0 µL of 10 mM dNTPs and 0.2 µL of *Taq* DNA polymerase. PCR reactions were performed in a thermal-cycler (Eppendorf, Hamburg, Germany). The amplification profile consisted of initial denaturation for 2 min at 95°C; 35–40 cycles of denaturation at 95°C, annealing at 50–60°C and extension at 72°C. After that, final extension was carried out at 72°C for 7 min.

Gel electrophoresis

PCR products of SSR markers were resolved on capillary electrophoresis system (Qiagen Pvt. Ltd., Hamburg, Germany).

Scoring of SSR data

The molecular weight of the PCR products for each SSR primer was determined from a ladder of known molecular weights. During band scoring, faint bands and bands with smeared background were avoided, and only intense bands were scored on the basis of their product size. The presence of a product in a certain genotype was designated as '1', and the absence was designated as '0'. Only the specific PCR products showing consistency in the successive amplifications were selected to minimize the possibility of mis-scoring markers.

Statistical analysis

A pooled ANOVA for the traits over the two years was done considering replication, genotype and environment as fixed effects (Singh and Chaudhary, 1985) in MS Excel 2007. Genetic parameters were estimated by the formula given by Burton (1952) for GCV and PCV, Hanson, Robinson and Comstock (1956) for heritability and Allard (1960) for expected genetic advance in MS Excel 2007. Mahalanobis D^2 analysis was done in Windostat version 9.2 (<http://www.windostat.org>). Usual Euclidian distances between the cultivars were worked out from the standardized data matrix in DARwin version 6.0.021 (Perrier and Jacquemoud-Collet, 2006) and represented through cluster analysis using the algorithm of Unweighted Neighbour-joining (UNJ) method by feeding the distance matrix as input data. Genetic relatedness among the genotypes was computed by using the Jaccard's coefficient of similarity (Jaccard, 1908) and a dendrogram was constructed illustrating the genetic relationship among the rice genotypes using UNJ method as proposed by Gascuel (1997), which uses a criterion of weighted average, in DARwin 6 (Perrier and Jacquemoud-Collet, 2006). The number of different alleles per locus (N_a), major allele frequency (MAF), number of effective allele (N_e), Shannon's information index (I), observed heterozygosity (H_o), expected heterozygosity (H_e) and polymorphism information content (PIC) values were calculated using GenAlEx version 6.5 (Peakall and Smouse, 2012).

Results And Discussion

Morphological characterization of the cultivars

A total of sixty-two qualitative and quantitative traits characterized the distinctiveness of the twenty *Joha* rice cultivars. Qualitative characteristics are considered morphological markers for identifying rice landraces because environmental changes least influence these traits (Raut, 2003). The stable morphological traits serve as reliable morphological markers for the identification of a cultivar. Each cultivar must have certain novel diagnostic features which will distinguish a variety from the others. Such diagnostic characters should be uniformly present in the population and inherit in the next generation, and then only the character is supposed to be stable and can be used as morphological marker traits to distinguish that variety. Among the traits, the spectrum of variability in the twenty cultivars revealed 22 monomorphic and 37 polymorphic characteristics (**Tables 2 & 3**), suggesting their potential for cultivar

characterization and distinctiveness. Earlier studies of Bisne and Sarawgi (2008), Mathure et al. (2011), Subudhi et al. (2012), Subba Rao et al. (2013), Sarawgi et al. (2014), Parikh et al. (2012) and Sinha et al. (2015) corroborated the above findings.

Cluster analysis of the cultivars based on polymorphic DUS characteristics

Clustering by unweighted neighbour-joining (UNJ) of usual Euclidean distance based on thirty-seven polymorphic traits as dissimilarity measure grouped the twenty indigenous aromatic rice cultivars into three multi-genotypic clusters viz., *G1*, *G2* and *G3* (Fig. 1). The clusters further subdivided into sub-clusters and had an unequal distribution of the cultivars. The cultivars viz., *Kon Joha-Bongaigaon*, *Kon Joha 1*, *Joha-Golaghat*, *Manimuni Joha*, *Kola Joha*, *Joha-Bihpuria*, *Soru Joha-Tinsukia*, *Kon Joha 4* and *Kalijeera* belonged to *G1*. At the same time, *G2* had *Kon Joha 5*, *Kon Joha-Moran*, *Harinarayan*, *Jeera Joha*, *Local Joha*, *Keteki Joha*, *Kon Joha 3*, *Kon Joha 2* and *Ronga Joha*. These clusters had the most significant number of nine cultivars, followed by *G3*, including *Kon Joha-Teok* and *Kunkuni Joha*. The group constellation pattern proved the existence of a substantial amount of variability among the indigenous *Joha* rice cultivars. The present findings agree with Mondal et al. (2014) and Sripunitha and Sivasubramaniam (2014).

Pooled analysis of variance

The pooled ANOVA over the two years (Supplementary Table S2) revealed that the mean squares due to years were significant for fifteen traits viz., days to first flowering, flag leaf length, flag leaf breadth, flag leaf area, days to 50% flowering, days to maturity, plant height, productive tillers plant⁻¹, filled grains panicle⁻¹, spikelet fertility, 1000-grain weights, biological yield plant⁻¹, grain yield plant⁻¹, harvest index and grain yield kg ha⁻¹, suggesting a significant influence of the years on the phenotypic expression of these traits. The yearly variation was mainly due to the difference in the 2018 and 2019 crops' sowing time. The crops were sown on 11th July and 17th June in 2018 and 2019, respectively. The June planted crop in 2019 exhibited higher mean performances for most of the traits mentioned above. Delaying the sowing time decreased the number of days to flowering and maturity for most of the cultivars. A similar observation was reported by Song et al. (1996) for days to heading reduced in different rice cultivars due to delayed sowing. Nahar et al. (2009) observed a significant decrease in filled grain production consequent upon delayed transplanting, attributed to low temperature at anthesis and spikelet primordial formation. Khalifa (2009) noted the early date of sowing the best time for maximizing morpho-physiological traits such as tillering, panicle initiation, chlorophyll content, leaf area index, sink capacity, panicle length, panicle number and grain yield. Delayed sowing significantly reduces the number of filled grains, panicles and test weight, finally lowering rice cultivars' grain yield (Patel et al., 2019). The cultivar differences for all the traits except for panicle length were also evident from highly significant mean squares. The Years x Cultivars interaction component was substantial for days to first flowering, days to 50% flowering, days to maturity, filled grains per panicle and spikelet fertility, suggesting differential behaviour of the cultivars in the early and late sown crops. The findings of Ganesan et al. (1995), Karim et al. (2007), Padmaja et al. (2008), Sahidullah et al. (2009), Singh et al. (2010), Khan et al. (2012), Gangashetty et al. (2013) and Verma et al. (2014) further supported the presence of significant cultivar differences. Mahapatra et al. (1996), Reddy et al. (1998) and Panwar et al. (2008) also reported substantial Years x Cultivars interaction for days to 50% flowering and days to maturity.

Mean performance of the cultivars

The variation among the various traits under study revealed free variability in different cultivars populations, reflecting the unforeseen impact of potential variability on yield. The cultivar mean performance for the observed traits (Supplementary Table S3) identified the top-ranking cultivars *Kon Joha 2* for stem thickness; *Joha-Bihpuria*, *Joha-Golaghat* and *Kola Joha* for the earliest days to first flowering; *Kon Joha-Teok* for flag leaf length; *Joha-Bihpuria*, *Ronga Joha* and *Soru Joha-Tinsukia* for flag leaf breadth; *Ronga Joha* for flag leaf area; *Kola Joha* for the earliest days to 50% flowering and maturity; *Keteki Joha* for the shortest height and the highest number of productive tillers; *Manimuni Joha*, *Kon Joha-Moran*, *Kon Joha 2*, *Jeera Joha* and *Kon Joha-Teok* for filled grains per panicle; *Jeera Joha* and *Kon Joha-Teok* for spikelet fertility; *Kola Joha* for thousand-grain weights; *Soru Joha-Tinsukia* for the most extended rice length and the highest rice length/breadth ratio; *Kon Joha 2* for the broadest rice breadth; *Ronga Joha* and *Kola Joha* for biological yield; *Ronga Joha* for grain yield per plant; *Joha-Bihpuria* for harvest index; and *Soru Joha-Tinsukia* for grain yield per hectare (Table 4). The majority of the *Joha* rice cultivars showed low to medium tillering, whereas none of the cultivars showed high tillering ability, as also supported by Ogunbayo et al. (2005), Singh et al. (2010), Mathure et al. (2011) and Sarawgi et al. (2014). The presence of considerable genetic variability for yield components observed in the present study was also similar to the findings of Marchenzan et al. (2005) and Singh et al. (2010). The variations in the grain characteristics of the *Joha* rice cultivars was in tune with the findings of Singh et al. (2010), Singh et al. (2012), Pachauri et al. (2013), Sarawgi et al. (2014), Semwal et al. (2014) and Sarhadi et al. (2015). The findings of Koutroubas et al. (2004), Vanaja and Babu (2006), Bajpai and Singh (2010) and Srivastava and Jaiswal (2013) further corroborated the present results on grain physical quality characteristics.

Genetic variability parameters

Grain yield per hectare showed the highest range of variation followed by filled grains per panicle, thousand-grain weights, grain length/breadth ratio, grain yield per plant, productive tillers per plant, rice length, stem thickness and biological yield per plant. Flag leaf breadth showed minor variability among the cultivars.

The magnitude of the genotypic and phenotypic coefficient of variation (Table 5) was the highest for grain yield ha⁻¹ (24.62 & 24.85%), followed by filled grains per panicle (23.69 & 25.02%). The genotypic and phenotypic coefficient of variation ranked moderate for rice length/breadth ratio (19.61 & 19.63%) followed by thousand-grain weights (19.41 & 19.47%), grain length/breadth ratio (18.86 & 18.90%), rice length (16.99 & 17.01%), grain length (16.55 & 16.56%), productive tillers per plant (13.15 & 17.27%), flag leaf area (10.62 & 10.66%) and stem thickness (10.23 & 10.37%). The phenotypic variations for all the above traits except productive tillers per plant were determined largely by the genotypes. Therefore, phenotype-based selection would be effective for these traits. These findings were in tune with Singh and Choudhary (1996), DebChoudhary and Das (1998), Kavitha and Reddy (2002), Karim et al. (2007), Osman et al. (2012), Babu et al. (2012) and Karupaiyan et al. (2013). High GCV and PCV for the number of filled grains per panicle agreed with Pandey and Anurag (2010) and Mazid et al. (2013). Kavitha and Reddy (2002) and Gangashetty et al. (2013) also reported moderate GCV and PCV for grain breadth and panicle length.

Heritability estimates ranged from 37.36 per cent for grain yield per plant to 99.99 per cent for flag leaf length (**Table 5**). Most of the traits exhibited high heritability values (82.64 to 99.99%); the exceptions were productive tillers (58.00%), biological yield (41.19%) and harvest index (45.26%) with average estimates and grain yield were low heritable. The genetic advances as per cent of mean were high (> 20%) for grain yield kg per ha (50.24%), filled grain per panicle (46.22%), rice length/breadth ratio (40.33%), thousand-grain weights (39.85%), grain length/breadth ratio (38.77%), rice length (34.98%), grain length (34.06%), flag leaf area (21.79%), stem thickness (20.79% and productive tillers per plant (20.64%). The traits excluding days to first/50 per cent flowering and maturity, flag leaf breadth and spikelet fertility exhibited high heritability in conjunction with high to moderate genetic advance, indicating the most likely role of additive gene action and effectiveness of simple selection for the traits. Similar results on days to 50 per cent flowering were in agreement with Bihari et al. (2004), Sankar et al. (2006) and Karthikeyan et al. (2009). High heritability and low genetic advance for days to 50% flowering agreed with Chaurasia et al. (2012). Plant height registered high heritability coupled with the moderate genetic advancement in conformity with the findings of Mishra and Verma (2002) and Chaurasia et al. (2012). Expression of moderate heritability coupled with high genetic advance for productive tillers plant⁻¹ was in tune with Mishra and Verma (2002), Kumari *et al.* (2003), Jaiswal et al. (2007) and Nandan *et al.* (2010). For filled grains panicle⁻¹, a high heritability concomitant with high genetic advance was in agreement with the results of Hasib et al. (2004) and Panwar (2005). High heritability in concurrence with high genetic advance for 1000-grain weights was in accord with the findings of Reddy et al. (1997), Murthy et al. (1999), Rao (2000), Mishra and Verma (2002) and Nandan *et al.* (2010). The grain quality traits *viz.*, grain length, breadth and length-breadth ratio, rice length, breadth, and length-breadth ratio registered high heritability coupled with the high genetic advance in consonance with the findings of Jaiswal et al. (2007), Reddy et al. (2013), Fatema et al. (2011) for grain length; Murthy et al. (1999) and Jaiswal et al. (2007) for grain breadth; and Jaiswal et al. (2007) and Fatema et al. (2011) for length-breadth ratio. A low heritability coupled with moderate genetic advance for grain yield plant⁻¹ was in agreement with Adjah et al. (2020). Similarly, Islam et al. (2015) observed high estimates of both heritability and genetic advance for grain yield kg ha⁻¹ as obtained in the present study.

Genetic divergence among the twenty *Joha* rice cultivars

The twenty aromatic *Joha* rice cultivars were assessed for the nature and magnitude of genetic divergence (Singh and Chaudhary, 1985) based on the twenty-three quantitative traits following Mahalanobis D^2 statistics. The V statistics and the analysis of dispersion (**Supplementary Table S4**) showed that the mean differences for the pooled effect of p characters between the cultivars were highly significant. Based on the D^2 values, the twenty *Joha* rice cultivars belonged to seven clusters, which showed three multi-genotypic and four mono-genotypic clusters. **Table 6** summarizes the cluster-wise distribution of the cultivars. The average intra- and inter-cluster distances (**Table 7**) showed cluster I to have the maximum intra-cluster distance (331.08) followed by cluster IV (307.84) and cluster II (223.91). The rest clusters were mono-genotypic, consisting of one cultivar each. The inter-cluster D^2 values were the least between the clusters V and VI (353.94), while it was the maximum between clusters IV and VI (7303.31). Cluster I, the largest group comprising eight cultivars, was distantly apart from cluster IV and cluster VI by 2281.86 and 1815.30 units, respectively, and was nearest to cluster III (490.87), followed by cluster V (854.97). Cluster II, with three cultivars, had the lowest inter-cluster distance from cluster III (564.34), followed by cluster V (892.56), while the highest inter-cluster distance was from cluster IV (4709.44). Cluster III consisted of one aromatic *Joha* rice cultivar and was nearest to cluster VII (434.10) whereas distantly placed from cluster IV (2323.71). Cluster IV was a multi-genotypic cluster having four aromatic *Joha* rice cultivars and was nearest to cluster VII (1256.85) while most distantly related to cluster VI (7303.31). Mono-genotypic cluster V was closest to cluster VI (353.94) and separated from cluster IV by 5187.67 units. Cluster VII having one aromatic *Joha* rice cultivar, was distantly placed from cluster VI (4073.37). Hybridization between parents from the widely divergent cluster pairs *viz.*, IV-VI, IV-V, II-IV, VI-VII, V-VI and III-HV would be likely to produce a broad spectrum of variability and transgressive segregations with high heterotic effects as also suggested by Shahidullah et al. (2009), Patil et al. (2012), Ali et al. (2012), Allam et al. (2014) and Patel et al. (2015).

The cluster mean performances for the various traits showed a wide range of variations among the clusters (**Supplementary Table S5**). Cluster III registered the earliest days to first and 50 per cent flowering (108 & 114) and maturity (147) and the highest mean performance for flag leaf breadth (0.84 cm), grain yield per plant (14.83 g) and harvest index (36.40%). Cluster VII showed superiority for spikelet fertility (90.92%), thousand-grain weights (22.50 g), grain and rice length (9.04 & 6.97 mm), rice length-breadth ratio (3.32) and grain yield per hectare (3012 kg). The highest mean performances for flag leaf length (54.21 cm), flag leaf area (32.42 cm²), panicle length (27.26 cm) and biological yield per plant (36.73 g) characterized cluster IV while cluster VI exhibited superiority for stem thickness (4.62 mm), filled grains per panicle (252), grain and rice breadth (2.82 & 2.33 mm). Cluster II was superior to other clusters in having the highest grain length-breadth ratio (3.89) and the shortest plant height (111 cm).

Of the twenty-three traits observed, only eight contributed to the genetic divergence. The contribution towards the total variation was the maximum for flag leaf length (72.11%) followed by rice length (13.68%), grain length (6.84%), rice breadth (3.16%), grain yield per hectare (1.58%), thousand-grain weights (1.05%), grain breadth (1.05%) and rice length-breadth ratio (0.53%). These results agreed with the findings of Allam et al. (2014) for rice length, Nayak et al. (2004), Chandra et al. (2007), Singh et al. (2008) and Shahidullah et al. (2009) for grain length.

The clustering pattern of the twenty *Joha* rice cultivars also confirmed the quantum of diversity present in the indigenous aromatic rice of Assam and offer scope for its exploitation through breeding for yield improvement. Previous studies reported different numbers of clusters in fragrant rice, *e.g.*, two clusters by Das et al. (2012), four clusters by Kole (2000), five clusters by Patel et al. (2015), six clusters by Singh *et al.* (1996) and Awasthi et al. (2005) and Shahidullah et al. (2009), Patil et al. (2012) and Allam et al. (2014), ten clusters by Pradhan and Mani (2005) and Ali et al. (2012).

Biochemical characterization of the *Joha* rice cultivars

In **Table 8** showing the biochemical characterization of the twenty *Joha* rice cultivars based on fatty acid profile, Fe and Zn content, crude protein, amylose, gel consistency and aroma, a cultivar's mean value was considered desirable for all other biochemical traits except for polyunsaturated fatty acids when it exceeded the cultivars' mean plus the standard deviation. A low mean less than the cultivars' mean minus the standard deviation was desirable for polyunsaturated fatty acids.

Fatty acid profile

Palmitic acid content ranged from 17.98% (*Local Joha*) to 20.57% (*Kon Joha-Teok*), with an average of 19.01%. Stearic acid content was the lowest (0.90%) in *Harinarayan* and the highest (1.86%) in *Kalijeera*; the average was 1.40 per cent. The range of oleic acid was from 33.53% in *Kunkuni Joha* to 41.32% in *Joha-Golaghat*, having an average of 37.90 per cent. The linoleic acid content varied from 36.02% in *Joha-Golaghat* to 44.61% in *Local Joha*, with an average of 39.22 per cent. *Local Joha* recorded the highest linolenic acid content (2.14%), whereas the *Kola Joha* had the lowest estimate (1.06%). Arachidic acid content ranged from 0.28% (*Local Joha*) to 0.61% (*Kon Joha 3*), with an average of 0.48 per cent. Fatty acids are vital components of food and human health. Fatty acids are the major constituents of cell membrane structure and play important biological, structural and functional roles in the human body (Nagy and Tiuca 2017). They act as modulators of gene transcription, cytokine precursors, and energy sources in complex interconnected systems (Glick and Fischer 2013) by producing a vast ATP quantity during their metabolism (Nagy and Tiuca 2017). The role of dietary fatty acids in human health is strongly evident for their influence on cardiovascular disease and mental health (Glick and Fischer 2013). Besides, rice is a dietary consumption; rice fats have unique health benefits (Jennings and Akoh, 2009). In the present investigation, the contents of oleic, linoleic and palmitic acids were the primary fatty acids, and those of stearic, linolenic and arachidic acids were minor in the aromatic rice cultivars. Palmitic, stearic and arachidic acids were the saturated fatty acids present in rice bran, increasing the health risk such as atherosclerosis, a disease associated with a heart attack (Oluremi et al. 2013). Linoleic acid gets absorbed as a predominant unsaturated fatty acid, followed by oleic and linolenic acid. The high contents of polyunsaturated fatty acids are desirable for human health as their consumption minimizes the risk of heart-related diseases (Law, 2000). The mean polyunsaturated fatty acid (PUFA) contents of the aromatic cultivars were 37.9% for oleic acid, 39.22% for linoleic acid and 0.5% for linolenic acid, whereas the contents of saturated fatty acids (SFA) accorded 1.40% for stearic acid and 19.01% for palmitic acid. These estimates were comparable or even better than the values of 38.4% oleic acid, 34.4% linoleic acid and 2.2% α -linolenic acid of PUFA; 2.9% stearic acid and 21.5% palmitic acid of SFA (Sayre and Saunders, 1990). The present results were also comparable with those reported by Resurreccion and Juliano (1975) and Taira and Itani (1988).

Similarly, the variations in the fatty acid profile of the present study proved better in having lower maximum limits for SFA and higher maximum limits for linoleic and linolenic acid than those reported by Goffman et al. (2003), who obtained 13.9–22.1% for palmitic, 1.5–2.7% for stearic, 35.9–49.2% for oleic, 27.3–41.0% for linoleic and 1.0–1.9% for linolenic acid in rice bran. Stearic acid and arachidic acid were present in trace amount in all the studied aromatic rice cultivars. Comparatively, the fatty acid profile of *Local Joha* was better than the other cultivars as it showed a high level of linoleic and linolenic acid and low saturated fatty acid content. In general, the *Joha* rice cultivars' fatty acid profile qualified for extraction of quality bran oil for consumption.

Iron and Zn content (mg kg⁻¹) of the cultivars

The cultivars' iron content varied from 21.33 (*Keteki Joha*) to 82.88 mg kg⁻¹ in *Kon Joha 4* with an average of 43.57 mg kg⁻¹. The zinc content ranged from 12.26 (*Keteki Joha*) to 47.39 mg kg⁻¹ in *Ronga Joha*, showing an average of 28.43 mg kg⁻¹. Iron and Zinc deficiency is severe nutritional problems for humans and is particularly prevalent among children and pregnant women, especially in developing countries. The *Joha* rice cultivars with very high iron and zinc contents in brown rice could play an essential role in the developing world's nutritional upliftment. Therefore, increasing the iron and zinc content and bioavailability in rice grains is an essential target for breeders and offers potential benefits to a large proportion of the human population. Substantial variations in iron and zinc content of brown rice agreed with Banerjee et al. (2010), Nachimuthu et al. (2014) and Chowdhury et al. (2016).

Protein content (%), amylose (%), gel consistency (mm) and aroma score

The range of variation in the cultivars' protein content was from 7.51 per cent in *Ronga Joha* to 10.32 per cent in *Kon Joha 1* with an average of 9.09 per cent. The variation in protein content was in agreement with Banerjee et al. (2011), who reported 4.91 to 12.08% protein in 258 diverse rice landraces with a mean of 6.63 per cent. The amylose content of the cultivars varied from 15.20% in *Jeera Joha* to 24.40% in *Harinarayan*, with an average of 19.86%. The cultivars exhibited two classes of amylose content - medium (20–25%) and low (10–20%), as also noted by Bajpai and Singh (2010). The lowest and the highest gel consistency in the cultivars were 61.50 mm in *Kon Joha 5* and 140.50 mm in *Joha-Bihpuria*, respectively, with an average of 100.25 mm. *Joha-Golaghat* (129.50 mm), *Kunkuni Joha* (123.00 mm) and *Manimuni Joha* (121.50 mm) followed *Joha-Bihpuria*. The cultivars viz., *Kalijeera*, *Kunkuni Joha*, *Kon Joha 5*, *Manimuni Joha* & *Kon Joha 2* showed strong aroma, *Joha-Bihpuria*, *Keteki Joha*, *Kon Joha 3*, *Kon Joha-Moran*, *Joha-Golaghat*, *Kola Joha*, *Jeera Joha*, *Kon Joha-Bongaigaon*, *Local Joha*, *Harinarayan*, *Kon Joha 1*, *Soru Joha-Tinsukia* & *Kon Joha 4* had a mild aroma, and *Ronga Joha* and *Kon Joha-Teok* possessed light aroma. In aromatic rice, Bajpai and Singh (2010), Mia et al. (2010), Golam et al. (2010), Mathure et al. (2011), Pachauri et al. (2013), Semwal et al. (2014) and Sarhadi et al. (2015) also observed variation in aroma and accordingly classified the genotypes.

Molecular characterization

Among these 71 SSR markers, 66 markers showed polymorphism. The analysis excluded the markers with monomorphic banding patterns. **Tables 9** summarize the results obtained on the analysis of 20 aromatic rice cultivars using the polymorphic SSR loci. Figure 1 depicts representative gel pictures of the PCR products.

The 66 polymorphic SSR loci amplified a total of 174 alleles. The allelic richness per locus ranged from 2 to 4 alleles, with an average of 2.64 alleles. Among the polymorphic markers, 30 markers produced two alleles each, 30 produced three alleles each, and 6 generated four alleles. The markers RM 283, RM 118, RM316, RM29, RM 585 and RM 26063 amplified the maximum number of alleles. The result revealed that all the markers showed distinct polymorphisms among the cultivars studied, indicating the robust nature of microsatellites revealing polymorphism. The number of alleles per locus (2.64) obtained in the present study was comparable with the earlier reports by Joshi and Behera (2007), Shah et al. (2013), and Venkatesan and Bhat (2015), who reported 2.6, 2.75 and 2.3 alleles per locus, respectively. The mean allele number (2.64) obtained in the present study was higher than the result of Meti et al. (2013), who detected 2.08 alleles per locus using 48 traditional indigenous aromatic rice germplasm grown under the Eastern part of India through 12 polymorphic SSR loci. In contrast, the mean alleles (2.64) detected was markedly lower than the average number of alleles reported in previous diversity studies by Pervaiz et al. (2010) and Rahman et al. (2012), who reported an average of 4.4 and 4.18 alleles per locus, respectively. The variability in the number of alleles detected per locus might be due to the use of diverse genotypes and the selection of different SSR primers with scorable alleles. Among the sixty-six markers, the highest major allele frequency was 0.950 in 4 markers (RM 237, RM 215, RM6641, RM3481 & RM21), followed by 0.900 in 9 markers and 0.875 in 2 markers. However,

RM 316 recorded the lowest major allele frequency of 0.300. The mean value of major allele frequency among the markers was 0.727, and it ranged from 0.300 to 0.950 among the sixty-six markers scored against a set of twenty aromatic *Joha* rice cultivars. Similarly, Sajib et al. (2012) reported major allele frequency ranging from 0.41 to 0.91; Shah et al. (2013) noted a range of 0.425 to 0.975 with an average of 0.647, and Kumar et al. (2015) observed it to vary from 0.510 to 0.970 averaging 0.74. The number of effective alleles was from 1.105 (RM 237) to 3.922 (RM 316), with a mean value of 1.759. A greater number of alleles generated by SSR markers suggest this marker system's usefulness for detecting genetic polymorphisms. Aljumaili et al. (2018) detected 1.48 effective alleles per SSR locus among 53 rice cultivars. On the contrary, the effective allele number detected in the present study was lower than the average number of effective alleles (5.51) reported by Yelome et al. (2018) among West African rice accessions. Shannon's information index ranged from 0.199 (RM 237) to 1.376 (RM 316), with a mean value of 0.640. Aljumaili et al. (2018) reported a similar Shannon's informative index by evaluating fifty-three aromatic rice accession using 32 SSR markers, and they reported a mean value of 0.580. The high value of Shannon's information index indicated the presence of high genetic diversity in the rice germplasm under consideration (Kibria et al. 2008). The gene diversity varied from 0.095 to 0.745, with a mean value of 0.377. RM 316 showed the highest gene diversity (0.745), followed by RM 283 (0.715) and RM 1388 (0.645). In contrast, Shah et al. (2013) recorded an average gene diversity of 0.448, ranging from 0.049 to 0.664, whereas Kumar et al. (2015) reported gene diversity ranging from 0.045 to 0.588 with a mean of 0.340. A set of four SSRs viz., RM 271, RM 279, RM 7434 and RM501 represented heterozygosity among sixty-six SSRs. All four markers showed the same value, 0.050, with a mean value of 0.003. The rest of the markers showed no heterozygosity. Similarly, Choudhury et al. (2014), Nachimuthu et al. (2014) and Yelome et al. (2018) reported low observed heterozygosity. The autogamous mode of reproduction in rice could be the reason for the low level of observed heterozygosity. The polymorphism information content (PIC) values, a reflection of allele diversity and frequency among the genotypes, also varied from one locus to another. PIC values varied from 0.091 to 0.698, with an average of 0.326, indicating that not all the SSR markers used were highly informative. The highest PIC value 0.698 was obtained for RM 316 followed by RM 283 (0.665), RM 585 (0.573), RM 1388 (0.572), RM 3562 (0.559), RM 171 (0.559), RM130 (0.534), RM118 (0.531), RM11 (0.513) and RM29 (0.509), suggesting that these ten markers were highly informative (PIC > 0.50) for identification of the twenty aromatic rice cultivars. Among the remaining polymorphic SSR markers, 30 markers were informative (0.50 < PIC < 0.25) and 26 markers were slightly informative (PIC < 0.25). The level of polymorphism detected in the present study was consistent with the reported mean PIC values in previous works (Pal et al. 2004; Hossain et al. 2007; Sajib et al. 2012). However, Nadia et al. (2014) reported an average PIC value of 0.84, markedly higher than the present average PIC value. The presence of sufficient polymorphism by the 66 SSR markers among the twenty indigenous *Joha* rice cultivars justifies their proper classification and use in the genetic improvement programme, based on the extent of genetic variation for desirable alleles.

Molecular diversity among the aromatic rice cultivars

An unweighted neighbour-joining (UNJ) cluster analysis based on Jaccard's similarity coefficients resolved the phylogenetic relationships among the aromatic rice cultivars collected from different parts of Assam. The UNJ cluster diagram showed three major clusters (G1, G2 & G3) with additional sub-clusters (Fig. 2). This dendrogram revealed that the cultivars derived from a genetically similar type clustered together. Cluster I comprised eight cultivars, whereas cluster II had ten cultivars, forming the most significant cluster. Cluster III had only two cultivars (*Local Joha* & *Kunkuni Joha*). Cluster I included *Kon Joha 3*, *Jeera Joha*, *Kon Joha 4*, *Keteki Joha*, *Joha-Bihpuria*, *Joha-Golaghat*, *Kon Joha 5* and *Kon Joha 2*. *Kon Joha-Bongaigaon*, *Kon Joha-Teok*, *Soru Joha-Tinsukia*, *Ronga Joha*, *Kola Joha*, *Kon Joha-Moran*, *Kalijeera*, *Kon Joha1*, *Manimuni Joha* and *Harinarayan* belonged to cluster II. The cluster I sub-divided into two clusters, IA and IB, consisting of six and two cultivars. Cluster II had two sub-clusters, IIA and IIB, consisting of nine and one cultivars, respectively. Similar to the present investigation's clustering pattern, Meti et al. (2013) obtained two major clusters for 48 aromatic rice genotypes from Odisha using 12 SSR markers at 49 per cent genetic similarity. Shah et al. (2013) effectively differentiated the basmati cultivars from non-basmati ones based on the cluster analysis using 24 microsatellite loci, classifying 40 rice cultivars into three groups. Pervaiz et al. (2010) reported four significant clusters of 75 rice landraces using 35 SSR markers at 0.40 similarity coefficients. Thus, SSR markers provided an adequate resolution to discriminate between aromatic rice accessions, and it could serve as a potential tool in the identification and characterization of genetically distant accessions from various sources. The microsatellite assay generated genotype-specific alleles in some of the cultivars evaluated for DNA fingerprints for cultivar identification and differentiation of aromatic rice. DNA fingerprints would be enormous assistance for establishing and defending the proprietary rights and maintaining the cultivar purity.

Conclusion

The present investigation on morpho-molecular and biochemical profiling of a sample of popular indigenous *Joha* cultivars of Assam has been a step forward for exploiting variability in this unique rice class to improve its inherently low yield potential through breeding. In light of the results on diversity analyses, it is evident that the *Joha* rice cultivars are highly diverse regarding yield and quality traits and utilization of these diverse traits specific genotypes to develop crop varieties with a broad genetic base would prove beneficial for aromatic *Joha* rice improvement programme. The low to a high degree of dissimilarity among the aromatic rice accessions observed in the present study exemplified the high level of diversity at the molecular level among the aromatic rice cultivars used and indicated their possible utilization in breeding programmes targeted at developing elite aromatic rice varieties. Besides, marker-based identification and differentiation of aromatic rice cultivars might help maintain this high-quality product's integrity to benefit both farmers and consumers. The *Joha* rice cultivar *Soru Joha* (*Tinsukia*) with the highest yield (3012 kg ha⁻¹), high spikelet fertility (90.9%) and high Fe content (61.09 mg kg⁻¹) could be considered for an immediate recommendation in the state of Assam. The *Joha* rice cultivars' fatty acid profile qualified for extraction of quality bran oil for consumption.

Declarations

ACKNOWLEDGEMENTS

The first author sincerely acknowledges the financial grant for his PhD research work from the Board of Research in Nuclear Sciences (BRNS), Bhabha Atomic Research Centre (BARC), Department of Atomic Energy (DAE), Government of India, Trombay under the project "Breeding for higher productivity of indigenous *Joha* rice of Assam through induced mutations for morpho-agronomic traits" (Sanction No. 35/14/17/2016-BRNS Dated 04/06/2016). The authors are thankful to the Assam Agricultural University for providing land and laboratory facilities in the department of Plant Breeding and Genetics. *Sincere*

thanks are due to Dr Vikash Kumar, Dr Gautam Vishawkarma, Dr Abhijit Shinde, Dr Suvendu Mondal, Miss. Anjali Chauhan and Mr Sudhakar Mali for their support during the first author's laboratory work at BARC, Trombay.

CONFLICT OF INTEREST

The authors declare no competing interest.

Funding: The research was supported by the financial grant from the Board of Research in Nuclear Sciences (BRNS), Bhabha Atomic Research Centre (BARC), Department of Atomic Energy (DAE), Government of India, Trombay, Mumbai, Maharashtra 400085 (Sanction No. 35/14/17/2016-BRNS Dated 04/06/2016).

Availability of data and material: Supplementary file 1

References

- Adjah KL, Abe A, Adetimirin VO, Asante M D (2020) Genetic variability, heritability and correlations for milling and grain appearance qualities in some accessions of rice (*Oryza sativa* L.). *Physiology and molecular biology of plants* 26(6):1309-1317.
- Ali M, Mian MAK, Rasul MG, Aziz Miah MA, Alam MZ, Hossain MA (2012) Genetic diversity in local aromatic rice (*Oryza sativa* L.) genotypes. *Bangladesh Journal of Plant Breeding and Genetics* 25(2): 33-40.
- Aljumaili SJ, Rafii MY, Latif MA, Sakimin SZ, Arolu IW, Miah G (2018) Genetic diversity of aromatic rice germplasm revealed by SSR markers. *BioMed research international* 2018: 1-11. DOI: 10.1155/2018/7658032
- Allam CR, Jaiswal HK, Qamar A (2014) Divergence analysis for yield and quality traits in some indigenous basmati rice genotypes (*Oryza sativa* L.). *International Journal of Applied Biology and Pharmaceutical Technology* 5(4): 257-263.
- Allard RW (1960) *Principles of Plant Breeding*. 1st Edn. John Wiley and Sons, Inc. New York.
- Awasthi, L. P., Mishra, C. H. and Pandey, V. K. (2005). Genetic divergence in Indian aromatic rice. *Crop Research*. 3(2): 199-201.
- Babu VR, Shreya K, Dangi KS, Usharani G, Nagesh P (2012) Genetic variability studies for qualitative and quantitative traits in popular rice (*Oryza sativa* L.) hybrids of India. *Int J Sci & Res Publ* 2(6): 1-5.
- Bajpai A, Singh Y (2010) Study of quality characteristics of some small and medium grained aromatic rices of Uttar Pradesh and Uttarakhand. *Agric Sci Digest* 30(4): 241-245.
- Banerjee S, Chandel G, Mandal N, Meena M, Saluja T (2011) Assessment of nutritive value in milled rice grain of some Indian rice landraces and their molecular characterization. *Bangladesh J Agril Res* 36(3): 369-380.
- Banerjee S, Sharma DJ, Verulkar SB, Chandel G (2010) Use of in silico and semiquantitative RT-PCR approaches to develop nutrient rich rice (*Oryza sativa* L.). *Ind J Biotechnol* 9(2): 203-212.
- Bihari PK, Richharia AK, Sahu RS (2004) Genetic advance for yield attributes in aromatic rice. *J Appl Biol* 14: 1-5.
- Bisne R, Sarawgi AK (2008) Morphological and quality characterization of Badsabhog group from aromatic rice germplasm of Chhattisgarh. *Bangladesh J Agric Res* 33(4): 479-492.
- Burton GV (1952) Quantitative inheritance in pearl millet. *Agron J* 50:503.
- Chandra R, Pradhan SK, Singh S, Bose LK, Singh ON (2007) Multivariate analysis in upland rice genotypes. *World Journal of Agriculture Science*. 3(3): 295–300.
- Chaurasia AK, Rai PK, Kumar A (2012) Estimation of genetic variability, heritability and genetic advance in aromatic fine grain rice. *Romanian journal of biology-Plant biology* 57(1): 71-76.
- Choudhury DR, Singh N, Singh AK, Kumar S, Srinivasan K, Tyagi RK (2014) Analysis of genetic diversity and population structure of rice germplasm from North-eastern region of India and development of a core germplasm set. *PLoS One* 9(11):e113094.
- Chowdhury BD, Nath A, Dasgupta T (2016) Evaluation of some popular rice genotypes with special emphasis on zinc, iron and protein content. *Int J Sci & Res Publ* 6(7): 609-619.
- Das, A., Kesari, V., and Rangan, L. (2010). Aromatic joha rice of Assam-A review. *Agricultural Reviews*. 31(1): 1-10.
- Das B, Sengupta S, Ghosh M, Ghose TK (2012) Assessment of diversity amongst a set of aromatic rice genotypes from India. *International Journal of Biodiversity and Conservation* 4(5): 206-218.
- Debchoudhary PK, Das PK (1998) Genetic variability, correlation and path coefficient analysis in deep water rice. *Annals of Agricultural Research* 19: 120-124.

- Dong MH, Sang DZ, Wang P, Wang XM, Yang JC (2007) Changes in cooking and nutrition qualities of grains at different positions in a rice panicle under different nitrogen levels. *Rice Science* 14: 141-148.
- FAO (2009) Food Security and agricultural mitigation in developing countries: Options for capturing synergies. Rome, Italy.
www.fao.org/docrep/012/i1318e/i1318e00.pdf
- FAO (2020) Production/Yield quantities of rice, paddy in World + (Total). 22/12/2020. Visited on 05/01/2021.
<http://www.fao.org/faostat/en/#data/QC/visualize>
- Fatema K, Rasul MG, Mian MAK, Rahman MM (2011) Genetic variability for grain quality traits in aromatic rice. *Bangladesh Journal of Plant Breeding and Genetics* 24(2): 19-24.
- Ganesan KW, Manuel W, Sundaram T (1995) Analysis of yield and yield components in rice. *IRRN* 20: 1-4.
- Gangashetty PI, Salimath PM, Hanamaratti NG (2013) Genetic variability studies in genetically diverse non-basmati local aromatic genotypes of rice (*Oryza sativa* L.). *Rice Genomics and Genetics* 4(2): 4-8.
- Gascuel O (1997) BIONJ: an improved version of the NJ algorithm based on a simple model of sequence data. *Molecular biology and evolution* 14(7): 685-695.
- Glick NR, Fischer MH (2013) The Role of Essential Fatty Acids in Human Health. *Journal of Evidence-Based Complementary & Alternative Medicine* 18(4) 268-289 DOI: 10.1177/2156587213488788
- Goffman FD, Pinson S, Bergman C (2003) Genetic diversity for lipid content and fatty acid profile in rice bran. *J Am Oil Chem Soc* 80(5): 485-490
- Golam F, Norzulaani K, Jennifer AH, Subha B, Zulqarnain M, Osman M, Nazia AM, Mohammad O (2010) Evaluation of kernel elongation ratio and aroma association in global popular aromatic rice cultivars in tropical environment. *Afr J Agric Res* 5(12): 1515-1522.
- Goyal SK, Singh JP (2002) Demand versus supply of food grains in India: Implications to food security. 13th International Farm Management Congress, Wageningen, The Netherlands, July 7-12, 2002 (No. 1026-2016-82014).
- Hanson GH, Robinson HF, Comstock RE (1956) Biometric studies in segregating population of Korean lespedesa. *Agron J* 48: 268-272.
- Hasib KM, Ganguli PK, Kole PC (2004) Evaluation of the performance of advanced generation lines of mutant x Basmati crosses of scented rice. *Journal of Interacademia* 8(1): 7-10.
- Hien NL, Ahmad WS, Oikawa Y, Hirata Y (2007) Genetic Diversity of Morphological Responses and the Relationships among Asia Aromatic Rice (*Oryza sativa* L.) Cultivar. *Tropics* 16(4): 343-355.
- Hossain MZ, Rasul MG, Ali MS, Iftekharuddaula KM, Mian MAK (2007) Molecular characterization and genetic diversity in fine grain and aromatic landraces of rice using microsatellite markers. *Bangladesh J Genet Plant Breed* 20(2): 1-10.
- Islam MA, Raffi SA, Hossain MA, Hasan AK (2015) Character association and path coefficient analysis of grain yield and yield related traits in some promising early to medium duration rice advanced lines. *Int J Expt Agric* 5(1): 8-12.
- Jaccard P (1980) Nouvelle recherches sur la distribution florale. *Bull Soc Vaud Sci Nat* 44: 223-70.
- Jaiswal HK, Shrivastava AK, Dey A (2007) Variability and association studies in indigenous aromatic rice. *Oryza* 44(4): 351-353.
- Jennings BH, Akoh CA (2009) Effectiveness of natural versus synthetic antioxidants in a rice bran oil-based structured lipid. *Food Chem* 114(4): 1456-1461.
- Jones N, Ougham H, Thomas H (1997) Genome mapping, molecular markers and marker-assisted selection in crop plants. *New Phytologist*. 137: 165-177.
- Joshi RK, Behera L (2007) Identification and differentiation of indigenous non-Basmati aromatic rice genotypes of India using microsatellite markers. *Afr. J. Biotechnol* 6(4): 348-354.
- Karim D, Sarkar U, Siddique MNA, Khaleque MMA, Hasant MZ (2007) Variability and genetic parameter analysis in aromatic rice. *International Journal of Sustainable Crop Production* 2(5):15-18.
- Karthikeyan P, Anbuselvam Y, Elangaimannan R, Venkatesan M (2009) Variability and heritability studies in rice (*Oryza sativa* L.) under coastal salinity. *Electronic Journal of Plant Breeding* 1(2): 196-198.
- Karuppaiyan R, Kapoor C, Gopi R (2013) Variability, heritability and genetic divergence in lowland rice genotypes under the midhills of Sikkim. *Oryza* 50: 81-84.
- Kavitha S, Reddy SR (2002) Variability, heritability and genetic advance of some important traits in rice (*Oryza sativa* L.). *Andhra Agric J* 49(34): 222-224
- Khalifa AABA (2009) Physiological evaluation of some hybrid rice varieties under different sowing dates. *Australian Journal of Crop Science* 3(3): 178-183.

- Khan MA, Khan AS, Khan SHU, Ahmad R (2012) Association of various morphological traits with yield and genetic divergence in rice (*Oryza sativa* L.). International Journal of Agriculture and Biology 14(1): 55-62.
- Kibria K, Islam MM, Begum SN (2008) Screening of aromatic rice lines by phenotypic and molecular markers. Bangladesh Journal of Botany 37(2): 141-147.
- Kole PC (2000) Genetic divergence in aromatic rice involving induced mutants. Oryza 37 (2): 40-42.
- Koutroubas SD, Mazzini F, Pons B, Ntanos DA (2004) Grain quality variation and relationships with morpho-physiological traits in rice (*Oryza sativa* L.) genetic resources of Europe. Field Crops Res 86: 115-130.
- Kumar V, Kumar S, Charabarty SK, Mohapatra T, Dadlani M (2015) Molecular characterization of farmers' varieties of rice (*Oryza sativa*). Indian Journal of Agricultural Sciences 85(1): 118-124.
- Kumari S, Kewat RN, Singh RP, Singh P (2013) Studies of quality characteristics in short grain scented rice (*Oryza sativa* L.) varieties accessions. Trends in Biosciences 6(2):177-179.
- Law M (2000) Dietary fat and adult diseases and the implications for childhood nutrition: An Epidemiologic approach. Am J Clin Nutr 72 (5): 1291-1296.
- Lin MS (1991) Genetic base of japonica rice varieties released in Taiwan. Euphytica 56(1): 43-46.
- Mahapatra KC, Mishra CHP, Singh B (1996) Assessment of adaptability in early rice varieties. Indian J Genet 56: 477-85.
- Marchezan E, Martin TN, Santos FMD, Camargo ER (2005) Path coefficient analysis of rice yield components. Ciencia Rural 35(5): 1027- 1033.
- Mathure S, Shaikh A, Renuka N, Wakte K, Jawali N, Thengane R, Nadaf A (2011) Characterization of aromatic rice (*Oryza sativa* L.) germplasm and correlation between their agronomic and quality traits. Euphytica 179: 237-249.
- Mazid MS, Rafii MY, Hanafi MM, Rahim HA, Latif MA (2013) Genetic variation, heritability, divergence and biomass accumulation of rice genotypes resistant to bacterial blight revealed by quantitative traits and ISSR markers. Physiologia Plantarum 149(3): 432–447.
- Meti N, Samal KC, Bastia DN, Rout GR (2013) Genetic diversity analysis in aromatic rice genotypes using microsatellite based simple sequence repeats (SSR) marker. Afr J Biotechnol 12(27): 4238-4250.
- Mia MF, Begum SN, Islam MM, Manidas AC, Halder J (2010). Identification and differentiation of aromatic rice genotypes using SSR markers. Int J Bio Res 2(9): 7-12.
- Mishra LK and Verma RK (2002) Correlation and path coefficient analysis for morphological and quality traits in rice (*Oryza sativa* L.). Plant Archives 2(2): 275-284.
- Mondal B, Singh SP, Joshi DC (2014) DUS characterization of rice (*Oryza sativa* L.) using morphological descriptors and quality parameters. Outlook on Agriculture 43(2): 131-137.
- Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. Nucleic Acids Res 8: 4321-4325.
- Murthy N, Kulkarni RS, Udaykumar M, Murthy N (1999) Genetic variability, heritability and genetic advance for morphological traits in rice. Oryza 36(2): 159-160.
- Nachimuthu VV, Robin S, Sudhakar D, Rajeswari S, Raveendran M, Subramanian KS, Tannidi S, Pandian BA (2014) Genotypic variation for micronutrient content in traditional and improved rice lines and its role in biofortification programme. Indian J Sci Technol 7(9): 1414–1425.
- Nadia I, Mohiuddin AKM, Sultana S, Ferdous J (2014) Diversity analysis of indica rice accessions using morphological and SSR markers. Ann Biol Res 5(11): 20-31.
- Nagy K, Tiuca ID (2017) Importance of Fatty Acids in Physiopathology of Human Body. Fatty Acids DOI: 10.5772/67407
- Nahar K, Hasanuzzaman M, Majumder RR (2009) Effect of low temperature stress in transplanted aman rice varieties mediated by different transplanting dates. Academic Journal of Plant Sciences 2(3): 132-138.
- Nayak AR, Chaudhury D, Reddy JN (2002) Genetic variability, heritability and genetic advance in scented rice. Indian Journal of Agricultural Science 46(12): 45-47.
- Nayak AR, Chaudhury D, Reddy JN (2004) Genetic divergence in scented rice. Oryza 41(3-4): 79-82.
- Ogunbayo SA, Ojo DK, Guei RG, Oyelakin OO, Sanni KA (2005) Phylogenetic diversity and relationships among 40 rice accessions using morphological and RAPDs techniques. Afr. J. Biotechnol 4(11): 1234- 1244.

- Oluremi, OI, Solomon AO, Saheed AA (2013) Fatty acids, metal composition and physico-chemical parameters of Igbemo Ekiti rice bran oil. *J Environ Chem Ecot* 5(3): 39-46.
- Osman KA, Mustafa AM, Ali F, Yonglain Z, Fazhan Q (2012) Genetic variability for yield and related attributes of upland rice genotypes in semi arid zone (Sudan). *African Journal of Agricultural Research* 7(33): 4613–4619.
- Otegbayo BO, Osamuel F, Fashakin JB (2001) Effect of parboiling on physicochemical qualities of two local rice varieties in Nigeria. *Journal of Food Technology in Africa* 6(4): 130-132.
- Pachauri V, Nilay T, Vikram P, Singh NK, Singh S (2013) Molecular and morphological characterization of Indian farmer rice varieties. *Australian Journal of Crop Science* 7(7): 923-932.
- Padmaja D, Rao KR, Subba LV, Padma V (2008) Studies on variability, heritability and genetic advance for quantitative characters in rice (*Oryza sativa* L.). *Indian Journal of Plant Genetic Resources* 21(3): 71-84.
- Pal S, Jain S, Saini N, Jain RK (2004) Identification of microsatellite markers for differentiating some Basmati and non-Basmati rice varieties. *Indian J Biotechnol* 3: 519-526.
- Pandey P, Anurag PJ (2010) Estimation of genetic parameters in indigenous rice. *Advances in Agriculture & Botany* 2(1): 79-84
- Panwar LL (2005) Genetic variability, heritability and genetic advance for panicle characters in transplanted rice. *Research on Crops* 6(3): 505-508.
- Panwar LL, Joshi VN, Ali M (2008) Genotype x environment interaction in scented rice. *Oryza* 45(2):103-109.
- Parikh M, Motiramani NK, Rastogi NK, Sharma B (2012) Agromorphological characterization and assessment of variability in aromatic rice germplasm. *Bangladesh J Agric Res* 37(1): 1-8.
- Patel AR, Patel ML, Patel RK, Mote BM (2019) Effect of different sowing date on phenology, growth and yield of rice – a review. *Plant Archives* 19(1):12-16
- Patel NB, Shrivastava R, Kumar V, Dhirhi N (2015) Genetic diversity among some traditional aromatic Rice (*Oryza sativa* L.) landraces of Chhattisgarh. *Electronic Journal of Plant Breeding* 6(1): 204-209.
- Patil KG, Nagesh P, Veeresh G, Kulkarni RS, Shashidhar HE (2012) Diversity studies among basmati and non-basmati type aromatic rice (*Oryza sativa* L.) genotypes for grain yield and its attributing characters. *Annals of Biological Research* 3(12):5766-5771.
- Peakall ROD, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular ecology notes* 6(1): 288-295.
- Perrier X, Jacquemoud-Collet JP (2006) DARwin software: Dissimilarity analysis and representation for windows. <http://darwin.cirad.fr/Darwin>.
- Pervaiz ZH, Rabbani MA, Khaliq I, Pearce SR, Malik SA (2010). Genetic diversity associated with agronomic traits using microsatellite markers in Pakistani rice landraces. *Electron J Biotechnol* 13(3): 717-745.
- Pradhan SK, Mani, SC (2005) Genetic diversity in basmati rice. *Oryza* 42(2): 150- 152.
- Rahman MM, Rasaul MG, Hossain MA, Iftekharuddaula KM, Hasegawa H (2012). Molecular characterization and genetic diversity analysis of rice using SSR markers. *J Crop Improv* 26: 244-257.
- Rao SS (2000) Estimation of grain yield and inter-relationship with yield components in upland rice. *Mysore Journal of Agriculture Science* 34(2): 142-146
- Raut VM (2003) Qualitative genetics of Soyabean-a review. *Soybean Res* 1: 1-28.
- Reddy JN, Pani D, Roy JK (1998) Genotype × environment interaction for grain yield in low land rice cultivars. *Indian J Genet* 58: 209-13.
- Reddy JN, Pani D, Roy JK (1997) Variability and character association in lowland rice. *Indian Agriculturist* 41(3): 159-65.
- Reddy RM, Raju S, Reddy DT, Reddy NS (2013) Study of genetic parameters in F2 generation involving aromatic rice genotypes (*Oryza sativa* L.). *International Journal of Applied Biology and Pharmaceutical Technology* 4(4): 238-241.
- Resurreccion AP, Juliano BO (1975) Fatty acid composition of rice oils. *J Sci Food Agric* 26(4): 437-439.
- Sajib AM, Hossain MM, Mosnaz ATMJ, Islam MM, Ali MS, Prodhan SM (2012) SSR marker-based molecular characterization and genetic diversity analysis of aromatic landraces of rice (*Oryza sativa* L.). *J Bio Sci Biotech* 1(2): 107-116.
- Sankar PD, Sheeba A, Anbumalaramathi J (2006) Variability and character association studies in rice (*Oryza sativa* L.). *Agric Sci Digest* 26(3): 182-184.
- Sarawgi AK, Parikh M, Sharma B, Sharma D (2014) Phenotypic divergence for agro-morphological traits among dwarf and medium duration rice germplasm and interrelationships between their quantitative traits. *The Bioscan* 9(4): 1677-1681.

- Sarhadi WA, Shams S, Bahram GM, Sadeghi MB (2015) Comparison of yield and yield components among different varieties of rice. *Global Journal of Biology, Agriculture and Health Sciences* 4(1):158-163.
- Sayre RN, Saunders RM (1990) Rice bran and rice bran oil. *Lipid Technol* 2:72
- Semwal DP, Pandey A, Bhandari DC, Dhariwal OP, Sharma SK (2014) Variability study in seed morphology and uses of indigenous rice landraces (*Oryza sativa* L.) collected from West Bengal, India. *AJCS* 8(3): 460-467.
- Shah SM, Naveed SA, Arif M (2013) Genetic diversity in Basmati and non-Basmati rice varieties based on microsatellite markers. *Pak J Bot* 45(51): 423-431.
- Shahidullah SM, Hanafi MM, Ashrafuzzaman M, Razi IM, Salam MA (2009) Phenological characters and genetic divergence in aromatic rice. *African Journal of Biotechnology* 8(14): 3199-3207.
- Shobha Rani N, Shobha Rao LV, Viraktamath BC, Mishra B (2004) National Guidelines for the Conduct of Tests for Distinctiveness, Uniformity and Stability. Directorate of Rice Research pp 6-13.
- Singh AK, Singh PK, Nandan R, Rao M (2012) Grain quality and cooking properties of rice germplasm. *Ann Pl Soil Res* 14(1): 52-57.
- Singh B, Mishra MK, Naik RK (2010) Genetic diversity among some traditional aromatic rice (*Oryza sativa* L.) varieties of Orissa. *Ind J Agric Res* 44(2): 141-145.
- Singh RK, Chaudhary, BD (1985) Biometrical methods in quantitative genetic analysis. Kalyani Publishers, New Delhi.
- Singh S, Choudhary BS (1996) Variability, heritability and genetic advance in cultivars of rice (*Oryza sativa* L.). *Crop Research Hisar* 12(2): 165-167.
- Singh SP, Mishra NC, Rir-Sima-Ah J, Campen M, Kurup V, Razani-Boroujerdi S, Sopori M L (2009) Maternal exposure to secondhand cigarette smoke primes the lung for induction of phosphodiesterase-4D5 isozyme and exacerbated Th2 responses: rolipram attenuates the airway hyperreactivity and muscarinic receptor expression but not lung inflammation and atopy. *Journal of Immunology* 183(3): 2115-2121.
- Singh Y, Pani DR, Pradhan SK, Bajpai A, Singh US (2008) Divergence analysis for quality traits in some indigenous Basmati rice genotypes. *Crop Improvement* 45(4): 263–267.
- Sinha AK, Mallick GK, Mishra PK (2015) Diversity of grain morphology on traditional rice varieties of lateritic regions of West Bengal. *World J Agril Sci* 1(1): 48-54.
- Song Y, Oh N, Hwang C (1996) Growth and response of yield characters of dry seeded rice at different seeding dates in plain area. RDA. *Journal of Agricultural Sciences* 38(2): 20-31.
- Srivastava AK, Jaiswal HK (2013) Grain characteristics and cooking quality of indigenous aromatic and non-aromatic genotypes of rice (*Oryza sativa* L.). *Int J Scientific Res Reviews* 2(1): 36-41.
- Subba Rao LV, Shiva Prasad G, Chiranjivi M, Chaitanya U, Surendhar R (2013) DUS characterization of farmer varieties of rice. *IOSR J Agric Vet Sci (IOSR-JAVS)* 4(5): 35-43.
- Subudhi HN, Samantaray S, Swain D, Singh ON (2012) Collection and agro-morphological characterization of aromatic short grain rice in eastern India. *Afr J Agric Res* 7(36): 5060-5068.
- Taira H, Itani T (1988) Lipid content and fatty acid composition of brown rice of cultivars of the United States. *J Agric Food Chem* 36(3): 460-462.
- Vanaja T, Babu LC (2006) Variability in grain quality attributes of high yielding rice varieties (*Oryza sativa* L.) of diverse origin. *J Trop Agric* 44 (1-2): 61-63.
- Verma PK, Chaurasia AK, Bara BM, Deepshikha M (2014) Evaluation of aromatic short grain rice cultivars and elite lines for yield and quality parameters. *International Journal of Research in Engineering and Technology* 03(06): 103-106.
- Yelome OI, Audenaert K, Landschoot S, Vanhove ADW, Silue D, Van Damme P, Haesaert G (2018) Analysis of population structure and genetic diversity reveals gene flow and geographic patterns in cultivated rice (*O. sativa* and *O. glaberrima*) in West Africa. *Euphytica* 11: 214-215. DOI: 10.1007/s10681-018-2285-1.

Tables

Table 1 List of indigenous *Joha* rice cultivars used in the investigation

S No	Cultivar name	Pedigree	Origin
1	Joha (Bihpuria)	Landrace	Assam
2	Kalijeera	Landrace	Assam
3	Ronga Joha	Landrace	Assam
4	Joha (Golaghat)	Landrace	Assam
5	Manimuni Joha	Landrace	Assam
6	Kon Joha (Moran)	Landrace	Assam
7	Keteki Joha	Savitri/Badsabhog	Assam
8	Kon Joha-1	Landrace	Assam
9	Soru Joha (Tinsukia)	Landrace	Assam
10	Kon Joha-2	Landrace	Assam
11	Jeera Joha	Landrace	Assam
12	Kon Joha-3	Landrace	Assam
13	Kon Joha-4	Landrace	Assam
14	Kunkuni Joha	Landrace	Assam
15	Kon Joha-5	Landrace	Assam
16	Local Joha	Landrace	Assam
17	Harinarayan	Landrace	Assam
18	Kon Joha (Teok)	Landrace	Assam
19	Kola Joha	Landrace	Assam
20	Kon Joha (Bongaigaon)	Landrace	Assam

Table 2 Distribution of the *Joha* rice cultivars based on monomorphic characteristics

S No*	Characteristics	States	Notes
1	Coleoptile: colour	Green	2
3	Leaf: intensity of green colour	Medium	5
4	Leaf anthocyanin colouration	Present	9
6	Leaf sheath: anthocyanin colouration	Absent	9
9	Leaf: auricles	Present	9
10	Leaf: anthocyanin colouration of auricles	Colourless	1
11	Leaf: collar	Present	9
12	Leaf: anthocyanin colouration of collar	Absent	1
13	Leaf: ligule	Present	9
14	Leaf: shape of ligule	Split	3
15	Leaf: colour of ligule	White	1
17	Leaf: width of blade	Narrow	3
19	Culm: attitude	Erect	1
20	Time of heading (% of plants with panicles) (days)	Late	7
23	Male sterility	Absent	1
35	Panicle curvature of main axis	Dropping	7
43	Panicle: presence of secondary branch	Present	9
46	Panicle: exertion	Well exerted	7
57	Decorticated grain: colour	Light brown	2
58	Endosperm presence of amylose	Present	9
61	Gelatinization temperature through alkali spreading value	Low	1
62	Decorticated grain: aroma	Present	9

*Serial numbers are as per DUS descriptor.

Table 3 Distribution of the *Joha* rice cultivars based on polymorphic characteristics

Characteristics	States	%	Characteristics	States	%	
Basal leaf: sheath colour	Green (1)	95	Lemma and palea: colour	Straw (1)	5	
	Purple lines (3)	5		Brown spot on straw (3)	10	
Leaf: distribution of anthocyanin colouration	On margins only (1)	40		Brown furrows on straw (4)	10	
	On tips only (2)	60		Brown (tawny) (5)	5	
Leaf pubescence of blade surface	Weak (3)	85		Black (9)	45	
	Medium (5)	15	Panicle: awns		Absent (1)	70
Leaf: length of blade	Medium (5)	25		Present (9)	30	
	Long (7)	75	Panicle: colour of awns (late observation)	Yellowish white (1)	10	
Flag leaf: attitude of blade (early observation)	Semi-erect (3)	85		Brown (3)	5	
	Horizontal (5)	15		Purple (8)	5	
				Black (9)	10	
Spikelet: density of pubescence of lemma	Medium (5)	85	Panicle: length of the longest awn	Very short (1)	10	
	Strong (7)	15		Short (3)	10	
Lemma: anthocyanin colouration of keel	Absent (1)	30			Very long (9)	10
	Strong (7)	20	Panicle: distribution of awns	Tip only (1)	5	
	Very strong (9)	50		Upper half only (3)	10	
Lemma: anthocyanin colouration of area below apex	Absent (1)	30			Whole length (5)	15
	Strong (7)	20	Panicle: secondary branching	Strong (2)	80	
	Very strong (9)	50			Clustered (3)	20
Lemma: anthocyanin colouration of apex	Absent (1)	30	Panicle attitude of branches	Semi-erect (5)	75	
	Strong (7)	25			Semi-erect to spreading (7)	25
	Medium (5)	5	Time maturity (days)	Late (7)	85	
	Very strong (9)	40			Very late (9)	15
Spikelet: colour of stigma	White (1)	80	Leaf senescence	Early (3)	20	
	Light green (2)	10			Medium (5)	15
	Yellow (3)	5			Late (7)	65
	Purple (5)	5	Sterile lemma: colour	Straw (1)	45	
Stem thickness	Thin (3)	50			Purple (4)	55
	Medium (5)	50	Grain weight of 1000 fully developed grains	Very low (1)	5	
Stem: length (excluding panicle; excluding floating rice)	Very short (1)	10			Low (3)	75
	Short (3)	90			Medium (5)	15
Stem: anthocyanin colouration of nodes	Absent (1)	90			High (7)	5
	Present (9)	10	Grain: length	Short (3)	80	
Stem: intensity of anthocyanin colouration of nodes	Weak (3)	5			Medium (5)	20
	Medium (5)	5	Grain: width	Narrow (3)	70	
Stem: anthocyanin colouration of internodes	Absent (1)	90			Medium (5)	30
	Present (9)	10	Grain: Phenol reaction of lemma	Absent (1)	80	
Panicle: length of main axis	Medium (5)	40			Present (9)	20
	Long (7)	55	Decorticated grain: length	Short (1)	75	
	Very long (9)	5			Medium (3)	25
Flag leaf: attitude of blade (late observation)	Semi-erect (3)	40	Decorticated grain: width	Narrow (3)	10	
	Horizontal (5)	60			Medium (5)	85
Panicle: number per plant	Few (3)	65				

	Medium (5)	35		Broad (7)	5
Spikelet: colour of tip of lemma	Yellowish (2)	25	Decorticated grain: shape (in lateral view)	Short bold (2)	70
				Long bold (4)	10
	Brown (3)	15	Endosperm content of amylose	Long Slender (5)	15
				Low	50
	Black (6)	60		Medium (5)	50

Table 4 Top ranking cultivars with desirable characteristics

Cultivar	Desirable characteristics
Kola Joha	Days to 1st flowering*, days to 50% flowering*, days to maturity*, 1000-grain weights (g), biological yield (g Plant ⁻¹)
Soru Joha-Tinsukia	Flag leaf breath (cm), rice length (mm), rice length/breadth ratio, grain yield (kg ha ⁻¹)
Ronga Joha	Flag leaf area (cm ²), grain yield (g Plant ⁻¹)
Keteki Joha	Plant height (cm), productive tillers Plant ⁻¹
Kon Joha-5	Filled grains Panicle ⁻¹ , grain breadth (mm)
Kon Joha-2	Stem thickness (mm), rice breadth (mm)
Kon Joha-Teok	Flag leaf length (cm)
Jeera Joha	Spikelet fertility (%)
Kon Joha-3	Grain length (mm)
Local Joha	Grain length/breadth ratio
Joha-Bihpuria	Harvest index (%)

*Earliness is desirable.

Table 5 Genetic variability parameters for the traits of the twenty indigenous *Joha* rice cultivars of Assam evaluated during *Sali* season of 2018 and 2019

Character	Range	Mean ± SE	GCV (%)	PCV (%)	h ² _{bs} (%)	GA (5%) As % of mean
Stem thickness (mm)	3.00 - 4.96	4.09 ± 0.07	10.23	10.37	97.34	20.79
Days to 1st flowering	108.00 - 123.33	114.51 ± 0.52	3.83	3.86	98.62	7.84
Flag leaf length (cm)	40.84 - 54.82	47.96 ± 0.05	9.08	9.08	99.99	18.71
Flag leaf breath (cm)	0.75 - 0.84	0.79 ± 0.01	3.61	3.73	93.80	7.21
Flag leaf area (cm ²)	24.21 - 34.0	28.29 ± 0.26	10.62	10.66	99.23	21.79
Days to 50% flowering	112.50 - 128.50	119.80 ± 0.56	3.93	3.96	98.59	8.05
Days to maturity	144.50 - 162.03	152.16 ± 0.69	3.35	3.38	98.18	6.85
Plant height (cm)	98.90 - 142.90	127.80 ± 3.74	7.83	8.36	87.75	15.11
Productive tillers Plant ⁻¹	7.80 - 13.80	10.42 ± 1.17	13.15	17.27	58.00	20.64
Filled grains Panicle ⁻¹	101.73 - 252.07	177.21 ± 14.24	23.69	25.02	89.69	46.22
Spikelet fertility (%)	78.97 - 93.03	87.04 ± 1.88	4.72	5.19	82.64	8.83
1000-grain weights (g)	13.02 - 26.69	17.60 ± 0.27	19.41	19.47	99.38	39.85
Grain length (mm)	5.95 - 9.30	7.28 ± 0.05	16.55	16.56	99.84	34.06
Grain breadth (mm)	2.09 - 2.87	2.41 ± 0.02	9.68	9.73	99.00	19.84
Grain length/breadth ratio	2.08 - 4.30	3.04 ± 0.04	18.86	18.90	99.58	38.77
Rice length (mm)	4.13 - 6.97	5.18 ± 0.04	16.99	17.01	99.83	34.98
Rice breadth (mm)	1.89 - 2.57	2.20 ± 0.02	8.54	8.57	99.16	17.51
Rice length/breadth ratio	1.79 - 2.32	2.38 ± 0.03	19.61	19.63	99.70	40.33
Biological yield (g Plant ⁻¹)	28.98 - 44.97	33.56 ± 3.30	8.23	12.83	41.19	10.88
Grain yield (g Plant ⁻¹)	8.64 - 17.20	12.94 ± 1.65	9.83	16.09	37.36	12.38
Harvest index (%)	22.99 - 36.40	31.65 ± 3.16	9.08	13.49	45.26	12.58
Grain yield (kg ha ⁻¹)	1000.00 - 3011.67	2012.08 ± 68.16	24.62	24.85	98.14	50.24

Table 6 Composition of the Tocher's clusters based on Mahanobis *D*² analysis

Cluster identity	No. of cultivars	Name of the cultivars
I	8	<i>Kon Joha-Moran, Jeera Joha, Manimuni Joha, Kon Joha-1, Kon Joha-Bongaigaon, Kon Joha-4, Kalijeera, Kon Joha-2</i>
II	3	<i>Keteki Joha, Local Joha, Kon Joha-3</i>
III	1	<i>Joha-Bihpuria</i>
IV	5	<i>Ronga Joha, Kola Joha, Joha-Golaghat, Kon Joha-Teok, Harinarayan</i>
V	1	<i>Kunkuni Joha</i>
VI	1	<i>Kon Joha-5</i>
VII	1	<i>Soru Joha-Tinsukia</i>

Table 7 Intra- (bold) and inter-cluster distances

Cluster	I	II	III	IV	V	VI	VII
I	331.08	1293.16	490.87	2281.86	854.97	1815.30	1167.98
II	1293.16	223.91	564.34	4709.44	892.56	1187.69	1560.50
III	490.87	564.34	0.00	2323.71	1046.91	1930.59	434.10
IV	2281.86	4709.44	2323.71	307.84	5187.67	7303.31	1256.85
V	854.97	892.56	1046.91	5187.67	0.00	353.94	2733.52
VI	1815.30	1187.69	1930.59	7303.31	353.94	0.00	4073.37
VII	1167.98	1560.50	434.10	1256.85	2733.52	4073.37	0.00

Table 8 Biochemical characteristics of the twenty *Joha* rice cultivars

Cultivars	Palmitic acid (16:0)	Stearic acid (18:0)	Oleic acid (18:1)	Linoleic acid (18:2)	Linolenic acid (18:3)	Arachidic acid (20:0)	Fe (mg kg ⁻¹)	Zn (mg kg ⁻¹)	Protein content (%)	Amylose content (%)	Gel consistency (mm)	Aroma (KOH test score)
Joha-Bihpuria	19.45	1.29	37.93	37.91	1.09	0.38	23.41	26.73	8.94	20.10	140.50	2
Kalijeera	18.50	1.86	40.70	36.77	1.23	0.61	31.19	23.49	9.69	17.40	84.50	3
Ronga Joha	18.22	1.25	38.55	39.62	1.33	0.54	47.00	47.39	7.51	19.10	113.00	1
Joha-Golaghat	19.16	1.50	41.32	36.02	1.11	0.51	28.58	32.60	7.93	19.42	129.50	2
Manimuni Joha	18.97	1.59	40.56	36.51	1.36	0.57	36.23	26.71	9.37	23.20	121.50	3
Kon Joha-Moran	19.19	1.48	36.90	39.83	1.66	0.54	25.61	28.56	9.87	17.30	98.00	2
Keteki Joha	18.08	1.20	37.37	40.90	1.77	0.37	21.33	12.26	9.07	22.60	82.00	2
Kon Joha 1	18.19	1.70	38.66	38.92	1.50	0.61	61.00	17.50	10.32	16.20	105.50	2
Soru Joha-Tinsukia	18.78	1.59	40.23	37.32	1.09	0.59	61.09	22.98	8.20	19.50	94.50	2
Kon Joha-2	19.21	1.56	38.07	38.23	2.10	0.51	62.50	23.00	9.89	22.60	86.00	3
Jeera Joha	18.86	1.19	35.71	41.82	1.63	0.43	43.07	37.95	8.59	15.20	118.50	2
Kon Joha-3	19.18	1.40	40.19	36.30	1.86	0.61	24.05	45.32	8.65	21.40	79.50	2
Kon Joha-4	20.11	1.50	37.91	38.26	1.24	0.55	82.88	38.25	9.23	21.40	71.00	2
Kunkuni Joha	19.78	1.04	33.53	43.13	1.78	0.40	34.61	28.76	9.17	15.70	123.00	3
Kon Joha-5	18.94	1.72	39.84	36.72	1.93	0.50	34.91	24.45	8.41	18.92	61.50	3
Local Joha	17.98	1.08	33.66	44.61	2.14	0.28	51.91	20.13	9.38	23.84	85.50	2
Harinarayan	19.94	0.90	35.56	41.14	1.81	0.29	53.83	31.30	9.18	24.40	91.00	2
Kon Joha-Teok	20.57	1.10	34.92	41.41	1.34	0.34	63.66	31.80	8.66	18.00	113.50	1
Kola Joha	18.76	1.63	40.26	37.36	1.06	0.53	37.31	25.86	10.20	17.20	95.50	2
Kon Joha-Bongaigaon	18.38	1.48	36.10	41.58	1.63	0.49	47.21	23.62	9.65	23.80	111.00	2
Lowest mean	17.98	0.90	33.53	36.02	1.06	0.28	21.33	12.26	7.51	15.20	61.50	1.00
Highest mean	20.57	1.86	41.32	44.61	2.14	0.61	82.88	47.31	10.32	24.40	140.50	3.00
Mean	19.01	1.40	37.90	39.22	1.53	0.48	43.57	28.43	9.09	19.86	100.25	2.15
SE	0.16	0.06	0.53	0.56	0.08	0.02	3.75	1.96	0.35	0.56	3.23	0.13
CV (%)	3.71	18.49	6.30	6.35	22.53	22.01	38.51	30.76	8.22	14.62	20.72	27.30

Aroma score: 0 – No aroma; 1 - Slight aroma; 2 - Mild aroma; 3 - Strong aroma; 4 - Very strong aroma. Values > (Mean + SD) are considered as desirable for all the traits except PUFA; For PUFA, values < (Mean - SD) were desirable (bold figures).

Table 9 SSR markers with their number of different alleles amplified (*Na*), major allele frequency (*MAF*), number of effective alleles (*Ne*), Shannon's informative index (*I*), observed heterozygosity (*Ho*), expected heterozygosity (*He*) and polymorphic information content (PIC) in the twenty indigenous *Joha* rice cultivars.

Locus	Na	MAF	Ne	I	Ho	He	PIC	Locus	Na	MAF	Ne	I	Ho	He	PIC	Locus	Na	MAF	Ne	I	Ho	He	PIC
RM495	2	0.90	1.22	0.33	0.00	0.18	0.16	RM489	3	0.70	1.80	0.75	0.00	0.46	0.38	RM 20236	2	0.90	1.22	0.33	0.00	0.19	0.16
RM 283	4	0.40	3.51	1.32	0.00	0.72	0.67	RM55	3	0.85	1.36	0.52	0.00	0.27	0.25	RM 7434	3	0.78	1.58	0.65	0.05	0.38	0.30
RM 237	2	0.95	1.11	0.20	0.00	0.10	0.09	RM 510	2	0.55	1.98	0.69	0.00	0.51	0.37	RM2126	2	0.90	1.22	0.33	0.00	0.19	0.16
RM431	3	0.70	1.80	0.75	0.00	0.45	0.38	RM474	3	0.60	2.25	0.94	0.00	0.57	0.49	RM 253	3	0.55	2.15	0.85	0.00	0.55	0.44
RM154	3	0.70	1.80	0.75	0.00	0.45	0.38	RM 171	3	0.45	2.74	1.05	0.00	0.65	0.56	RM217	2	0.90	1.22	0.33	0.00	0.19	0.16
OSR 13	3	0.80	1.50	0.61	0.00	0.34	0.30	RM212	2	0.90	1.22	0.33	0.00	0.19	0.16	RM7434	3	0.85	1.36	0.52	0.00	0.27	0.25
RM338	2	0.65	1.84	0.65	0.00	0.46	0.35	RM23	3	0.60	2.06	0.82	0.00	0.53	0.42	RM 481	3	0.70	1.85	0.80	0.00	0.47	0.41
RM514	3	0.60	2.06	0.82	0.00	0.52	0.35	RM 229	2	0.70	1.72	0.61	0.00	0.43	0.33	RM11	3	0.55	2.41	0.98	0.00	0.60	0.51
RM124	2	0.90	1.22	0.33	0.00	0.18	0.16	RM1003	2	0.70	1.72	0.61	0.00	0.43	0.33	RM 505	3	0.65	1.94	0.79	0.00	0.50	0.41
RM 161	2	0.55	1.98	0.69	0.00	0.50	0.37	RM29	4	0.60	2.30	1.03	0.00	0.58	0.51	RM501	3	0.73	1.75	0.75	0.05	0.44	0.35
RM133	3	0.75	1.68	0.73	0.00	0.41	0.37	RM 221	2	0.80	1.47	0.50	0.00	0.33	0.27	RM25	3	0.85	1.36	0.52	0.00	0.27	0.25
RM 125	3	0.85	1.36	0.52	0.00	0.27	0.25	RM6641	2	0.95	1.11	0.20	0.00	0.10	0.09	RM407	3	0.70	1.85	0.80	0.00	0.47	0.41
RM 118	4	0.55	2.47	1.07	0.00	0.60	0.53	RM 279	2	0.88	1.28	0.38	0.05	0.22	0.09	RM3481	2	0.95	1.11	0.20	0.00	0.10	0.09
RM 152	2	0.85	1.34	0.42	0.00	0.26	0.35	RM3562	3	0.45	2.74	1.05	0.00	0.65	0.56	RM3395	3	0.90	1.23	0.39	0.00	0.19	0.18
RM284	3	0.85	1.36	0.52	0.00	0.27	0.25	RM60	2	0.65	1.84	0.65	0.00	0.47	0.35	RM228	3	0.85	1.36	0.52	0.00	0.27	0.25
RM316	4	0.30	3.92	1.38	0.00	0.75	0.70	RM1388	3	0.45	2.82	1.07	0.00	0.66	0.57	RM590	2	0.80	1.47	0.50	0.00	0.33	0.27
RM 215	2	0.95	1.11	0.20	0.00	0.10	0.09	RM471	2	0.85	1.34	0.42	0.00	0.26	0.22	RM591	3	0.70	1.85	0.80	0.00	0.47	0.41
RM 271	2	0.88	1.28	0.38	0.05	0.22	0.16	RM17467	3	0.70	1.87	0.82	0.00	0.48	0.42	RM 26063	4	0.60	2.38	1.09	0.00	0.60	0.44
RM484	2	0.85	1.34	0.42	0.00	0.26	0.35	RM 317	2	0.85	1.34	0.42	0.00	0.26	0.22	RM21	2	0.95	1.11	0.20	0.00	0.10	0.09
RM 536	2	0.50	2.00	0.69	0.00	0.50	0.38	RM 187	2	0.85	1.34	0.42	0.00	0.26	0.22	RM27601	3	0.90	1.23	0.39	0.00	0.19	0.18
RM 227	2	0.60	1.92	0.67	0.00	0.48	0.37	RM 3322	2	0.90	1.22	0.33	0.00	0.19	0.16	R1M7	2	0.85	1.34	0.42	0.00	0.26	0.22
RM 29	3	0.45	2.41	0.95	0.00	0.59	0.50	RM 585	4	0.50	2.74	1.14	0.00	0.65	0.57	R1M30	3	0.45	2.60	1.01	0.00	0.63	0.53
Mean	2.64	0.73	1.76	0.64	0.00	0.38	0.33	Mean	2.64	0.73	1.76	0.64	0.00	0.39	0.33	Mean	2.64	0.73	1.76	0.64	0.00	0.39	0.33
SE	0.08	0.02	0.07	0.04	0.00	0.02	0.02	SE	0.08	0.02	0.07	0.04	0.00	0.02	0.02	SE	0.08	0.02	0.07	0.04	0.00	0.02	0.02

Figures




Figure 1

Representative gel pictures showing the PCR products




Figure 2

Hierarchical horizontal clustering of the 20 Joha rice cultivars using Unweighted Neighbour-Joining (UNJ) method based on usual Euclidean distances estimated from 37 polymorphic morphological markers




Figure 3

Hierarchical horizontal clustering of the 20 Joha rice cultivars using Unweighted Neighbour-Joining (UNJ) method based on Jaccard's coefficients of similarity estimated from 66 polymorphic SSR markers

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryfile1.xlsx](#)