

Table 1: Patient demographic and cancer characteristics and their correlation with survival

<u>Characteristic</u>	<u>Value</u>	<u>p</u>	<u>OS in months</u> <u>(95% CI)/HR</u>
Gender [n (%)]		0.766	
male	607 (70.2%)		31.4 (27.0-35.8)
female	258 (29.8%)		27.9 (23.0-32.8)
Age (years)		<0.001	1.013 (1.006-1.020)
median (min, max)	64 (28; 92)		
Body mass index (BMI)		0.247	0.985 (0.961-1.010)
median (min, max)	24.8 (13.6; 50.0)		
Localization of cancer [n (%)]		0.005	
Gastric	345 (39.9%)		32.4 (25.5-39.3)
Gastroesophageal junction	281 (32.5%)		26.7 (18.6-34.8)
Esophageal	239 (27.6%)		27.3 (20.9-33.7)
Histological type [n (%)]		0.001	
Adenocarcinoma	681 (78.7%)		31.7 (27.1-36.3)
Squamous cell carcinoma	184 (21.3%)		24.2 (18.2-30.2)
Stage [n (%)]		<0.001	
Stage I	155 (17.9%)		87.6 (65.4-109.8)
Stage II	276 (31.9%)		38.6 (30.2-47.0)
Stage III	422 (48.8%)		20.1 (18.0-22.2)
Resectable, but unknown stage	12 (1.4%)		