

The Prevalence, Awareness, Management and Control of Hypertension in Men and Women in Benin, West Africa: the TAHES study.

ILEANA DESORMAIS (✉ ileana.desormais@orange.fr)

Hopital Dupuytren <https://orcid.org/0000-0003-3728-568X>

Salimanou Ariyoh Amidou

Faculty of Health Science, Cotonou

Yessito Corine Houehanou

Faculty of Health Science, Cotonou

Stephan Dismand Houinato

Faculty of Health Science, Cotonou

Gwladys Nadia Gbagoudi

Faculty of Health Science Cotonou

Pierre Marie Preux

INSEM 1094 Limoges

Victor Aboyans

Department of cardiology, Dupuytren University Hospital Limoges

Philippe Lacroix

Hopital Dupuytren

Research article

Keywords: Hypertension, general population, Benin

Posted Date: October 11th, 2019

DOI: <https://doi.org/10.21203/rs.2.14078/v2>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Cardiovascular Disorders on December 1st, 2019. See the published version at <https://doi.org/10.1186/s12872-019-01273-7>.

Abstract

Background Sub-Saharan Africa is facing a growing burden of non-communicable diseases, including cardiovascular diseases (CVDs), due to epidemiological transitions. Among their risk factors, hypertension is a major determinant of CVDs, but the prevalence and level of awareness and management of this condition are poorly studied in African populations. The aim of this study was to determine the prevalence of hypertension and identify its associated risk factors as well as the awareness and management of this condition in a community-dwelling cohort in Benin. **Methods** A cross-sectional door-to-door study was conducted in the population over the age of 25 years in Tanve, a rural setting in Benin. The questionnaire and anthropometric measurements of the World Health Organization STEPWISE survey were used. Blood pressure was measured using standard procedures. **Results** The sample included 1777 subjects (60.9% females, mean age was 42.5 ± 16.5 years). The prevalence of hypertension was 32.9%, similar in men (32.8%) and women (33.0%, $p=0.9342$). Age and obesity were significantly associated with hypertension. Less than half (42%) of hypertensive people were aware about their condition and only 46.3% of them were treated. Awareness ratios differed between men and women (respectively 32.9% vs. 47.5%; $p=0.0039$) and was not influenced by age, education, occupation, marital status or income. Female sex was the only factor associated with better controlled HTN, independent of socio-economic parameters. **Conclusion** This large population-based study confirms the high prevalence, low awareness, and low control of hypertension in men and women in sub-Saharan Africa. Only half of the populations with hypertension are aware of their hypertension, indicating a high burden of undiagnosed and un-controlled high blood pressure in these populations.

Background

Hypertension (HTN) is a major risk factor for cardiovascular diseases (CVDs), especially ischemic heart disease and stroke(1). As the global number of hypertensive people in 2015 was estimated at almost 1.13 billion(2) with a prevalence of HTN within the 30%-45% range in adults(3), the global burden of morbidity and mortality related to HTN becomes one of the major public health concerns worldwide. Given its increasing prevalence with advancing age(3) and the trends for population ageing and increased sedentary lifestyle and obesity, the prevalence of HTN is expected to increase even more. It has been predicted that, by 2025, the number of hypertensive individuals will increase by 15-20%, nearing one and a half billion(4). Although scarce evidence is available in developing countries, the high prevalence of HTN seems consistent across the world, irrespective of the income status (i.e. lower-income, middle-income and higher-income countries)(3). Few population-based studies indicate that hypertension is also a widespread problem in sub-Saharan Africa (SSA), with a prevalence up to 38% in some communities (5, 6). It is estimated that out of the approximately 650 million people in SSA, between 10 to 20 million may have hypertension. These estimations are based on scarce heterogeneous studies in the past and many countries in SSA still lack detailed recent basic data on the prevalence of HTN(7). Furthermore, few studies have reported on the proportion of awareness, treatment and good control of HTN in these populations with socio-economic particularities. In Benin, the study conducted in general population in 2008 showed an alarming prevalence of 28%(8). This condition was highly undiagnosed, as 78% of the subjects were unaware of their high blood pressure.

Our aim was to estimate, using an updated rigorous methodology in accordance to the recent guidelines, the prevalence of hypertension, to identify its associated risk factors and assess the level of awareness and management of hypertension in a rural area of Benin in West Africa.

Methods

Study population

This is an analysis of inclusion data collected in the TAnve HEalth Study (TAHES) a prospective population-based cohort study initiated in 2015 in Tanve, a rural setting 150 km north of Cotonou, the capital of Benin. The TAHES involved all adults above 25 years old living in Tanve(9). Its main objective is to assess the frequency of CVDs and their associated risk factors. Exclusion criteria were pregnant women and refusals to participate.

All participants and/or their families gave informed consent prior to inclusion in the study. Written consent was obtained whenever feasible. For illiterate people, the study's objectives were verbally explained and consent was obtained by thumbprint.

The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki and had prior approval of the Benin national health's research ethics committee and the "Comité de Protection des Personnes du Sud-Ouest et d'Outre-Mer 4 in France".

Data collection

An exhaustive sampling using a door-to-door approach was performed. Data were collected by 8 teams of 3 trained investigators, using a questionnaire adapted from WHO STEPS tools(10) in the households.

Sociodemographic data

Sociodemographic data included age, sex, marital status (never married, living with someone as a couple, widow/divorced/separated), education (none, primary education, higher), profession (employee/government employee, craftsman/ storekeeper, farmer/breeder/fisherman, homemade/retired, jobless), and the household income per month (low, middle, high) according to the World Bank indicators(11).

Cardiovascular risk factors

The cardiovascular risk factors were defined according to the WHO STEPS surveillance manual(12). Tobacco use was assessed and participants were classified as never users and current/former users (including cigarette, cigar, pipe or other modes of tobacco use including chewing tobacco).

Weight was measured to the nearest 100 g on mechanical scales (Seca, Hamburg, Germany) and height was measured to the nearest centimeter using a carpenter meter. While the stand upright position was impossible, height was estimated according to the knee height (KH) using Chumlea's formula for non-Hispanic Black people. Body mass index (BMI) was calculated as weight/height². Underweight was defined as BMI <18.5 kg/m², normal weight: BMI= 18.5-24.9 kg/m², overweight: BMI= 25-29.9 kg/m², and obese BMI ≥ 30 kg/m².

Diabetes was defined as currently taking antidiabetic drugs or having a fasting capillary whole blood glucose value³ 126 mg/dL(13).

Blood pressure measurements and HTN definition

Systolic (SBP) and diastolic (DBP) blood pressures were recorded, in seated position, after a rest of at least 15 minutes, using an electronic device (OMRON® M3, OMRON Corporation, Japan). Three measures were recorded, in both arms, at 5-minutes intervals. In accordance to the 2017 ESC Guidelines on hypertension the average of the last two measurements was used in the analyses and hypertensive subjects was defined by self-reporting ongoing treatment, or SBP ≥140 mm Hg and/or DBP ≥90 mm Hg (14). The hypertension was defined as controlled when SBP <140 mm Hg and DBP <90 mm Hg under pharmacological treatment.

Other Data

Nutritional variables included the dietary sodium intake using a food frequency questionnaire(15) and defined according to the WHO guidelines for sodium intake(16).

"Rare" salt intake was defined by low-salt food intake, seasoning less than once a day and consuming ready-made-dishes less than twice a week.

Low intake of fruit and vegetable was defined as consuming less than five total servings (400 g) of fruit and vegetables per day.

Harmful (moderate to heavy) use of alcohol was defined as consumption of >60 g of alcohol for men or 40 g for women in one occasion within the last 30 day. Consumption below these thresholds was considered as light.

Sedentary lifestyle was defined as <150 min of moderate-intensity activity (walk, bicycle) per week, or equivalent.

Data Analysis

Descriptive analyses were performed to compare the socio-demographic, cardiovascular risk factors and nutritional variables in men and women. The averages (\pm SD) and numbers (ratios) were compared with Fisher's exact test and the Chi-square test, as appropriate.

The association between variables and HTN was assessed by univariable and multivariable analyses. A multivariable logistic regression model was performed to identify associated factors for HTN within demographic variables and CVD risk factors when p -value <0.20 in univariable logistic regression. Interactions between independent variables in the final model were examined.

We performed several models. In these models, age and sex were forced systematically.

First, we adjusted for sociodemographic factors such as age, sex, country, area, marital status, and previous occupation (model 1). Second, cardiovascular risk factors – tobacco use, BMI, physical activity, diabetes – were additionally adjusted (Model 2).

Third, nutritional factors (salt consumption and alcohol consumption, fruits and vegetables) were also adjusted for (Model 3).

The level of significance was fixed at 0.05 for all analyses. Statistical analyses were carried out using Statview 5.0 software (SAS Institute, Cary, USA).

Results

Study Population

Among the 1779 subjects aged 25 years and older who were approached, data for hypertension were missing for 2 participants, resulting in a total sample size of 1777 participants, mean age 42.5 ± 16.5 years. Subjects who were 25–34 years old comprised the largest group (41.0%). Females accounted for more than 60% of this population (Table 1).

The global prevalence of hypertension was 32.9% (584 subjects), with the prevalence of hypertension being similar in men and women (32.8% (228) and 33.0% (356) respectively). General characteristics of our study population are detailed in Table 1. The prevalence of hypertension increased significantly with age (OR: 1.04, $p < 0.001$) (Table 2).

The univariable analysis indicated that increased age, the past or current use of tobacco, increased BMI, sedentary lifestyle, widowed or divorced status, previous occupation other than employee, lower education, animist religion, and increased salt consumption were associated with an increased likelihood of hypertension (Table 2). In the full model (model 3), increased age and BMI, homemade and retired status as well as animist religion remained significantly associated with an increased likelihood of hypertension, whereas the moderate alcohol consumption remained associated with a lower ratio of HTN (Table 2). No interactions between independent variables have been found.

Further analyses suggested a higher likelihood of hypertension in the case of high to very high salt consumption and sedentary lifestyle in women as compared to men who seemed less exposed to HTN by a moderate alcohol intake (Table 3). Age, increasing BMI and animist religion remain significant associated HTN factors in both men and women.

Awareness, Treatment, and Control ratios of hypertension

Among hypertensive participants, less than one-half (41.8%) were aware of their condition (Figure 1). Awareness ratios differed between men and women (32.9% vs. 47.5%; $p=0.0039$) (Figure 2) and was not influenced by age, education, profession, marital status or income. Female sex was the only associated factor with controlled HTN, independently of the education, occupation, income or marital status (Table 4).

Only 46.3% (113) of hypertensive participants declared receiving antihypertensive treatment at the time of the study (Figure 1). No difference in the prevalence of treatment was observed between men and women (37.3% vs. 50.3%, $p=0.0610$), nor in uncontrolled hypertensive (34.1% vs. 35.7%, $p=0.1520$). Treatment was not influenced by the socio-economic status.

Discussion

To our knowledge, this population-based study, conducted following the STEPS method recommended by the WHO for screening and monitoring risk factors of noncommunicable diseases, is the one of the largest investigations in the past five years on the prevalence of HTN and associated factors in rural population in West Africa. The study on hypertension, conducted in 2008, in Benin, showed an alarming prevalence in general population (27.9%)(8). This condition was highly undiagnosed, as 78% of the subjects were unaware of their high blood pressure. As hypothesized, our study underlines the highly increased HTN prevalence (32.9%) since 2008 (8). Previous studies also suggested a higher prevalence of HNT in urban areas compared to rural zones, explained by the fact that rural African populations were characterized by a traditional lifestyle associating cultural and dietary habits more in favour of HTN prevention (27.1% in rural areas vs 29.5% in urban zones in 2011) (17). Almost twenty years later, our study points out the sociocultural transition in rural areas with a prevalence of HNTN even higher than the prevalence described previously in urban population. The prevalence of HTN observed in the present study is comparable to the recently reported prevalence in Ghana, Nigeria, South Africa, Sudan and Tanzania (18-20). The progressive aging of the population with the increase in life expectancy may partially explain the progression of HTN prevalence in African countries.

The factors generally associated with hypertension include age, BMI, sedentary lifestyle, tobacco use, alcohol consumption. Most of these have previously been identified as important risk factors for hypertension in different studies conducted in SSA(21-26). Our results also confirm some of these observed associations such as increasing age, sedentary lifestyle and BMI. Our study observed that tobacco users, generally considered at higher risk of CVD were not at higher risk to be hypertensive. This result is probably due to the very low ratio of tobacco use in Benin. Also, alcohol consumption, a well-known risk factor of

HTN (14) was not significantly reported in our analyses. In other sub-Saharan African studies results are also controversial, showing either association, or no significant association between HTN and moderate to heavy alcohol consumption(27, 28). Furthermore, our study suggests a slightly protective effect of the light alcohol consumption in men, probably as the reflect of a healthy lifestyle. These discrepancies in hardly comparable results might be explained by the use of different settings, and assessments of alcohol consumption from one study to another.

Recent data show that salt consumption around the world is much higher than is physiologically necessary(29), even more in males than females at any age(30, 31). In a systematic review and meta-regression in 2016 no gender difference was reported in African studies(32). In our study, high salt intake, was independently associated with a higher likelihood of prevalent hypertension. Dietary estimates of sodium intake might not be accurate due to recall bias, reporting errors, erroneous food composition tables (especially if not country-specific or outdated), and/or difficulty in quantifying added salt (e.g. salt added during cooking but discarded in cooking water, etc). However, the fact that there might be a difference between men and women, suggests that context-specific research is needed to establish whether patterns of sodium intake in the SSA settings differ from high-income countries or other low- or middle-income countries.

Our finding regarding higher ratios of HTN in subjects of animist religion are intriguing and we do not find any rational as declarative questionnaires on lifestyle used in our study were not exhaustive. Previous studies suggested that religiosity could be a factor involved in the management of health and diseases/patient longevity (33). Nevertheless, further specific confirmatory and exploratory studies are needed.

In our study, less than 42% of hypertensive participants were aware of their pathology and less than one-half of them took antihypertensive treatments. Among treated subjects, only one-third had controlled hypertension. Few studies have analysed awareness and treatment ratios of hypertension, as well as the fraction with controlled hypertension in Africa(8, 26).

The ratios of awareness and controlled hypertension were higher than other studies on the continent. These differences might be related to variances in socioeconomic levels, healthcare access and preventive measures to reduce HTN. The findings of our study also suggest that independently of the socio-cultural status, women could be more receptive to treatment and more regular during the follow up. This could be explained by a better medical survey for women during and after pregnancy.

Our study has several limitations. Although the definition of hypertension respects the current guidelines, no ambulatory pressures were available. Furthermore, the cross-sectional design of the study, do not allow the assessment of the temporal nature of the associations and survival bias cannot be totally excluded. We also cannot exclude recall bias, especially for nutritional factors, but this it is limited since informants were interviewed alongside participants. Also, the 24-hour urinary salt excretion was not available. Assessment of dietary sodium intake used a food frequency questionnaire which might underestimate the sodium intake by 30% to 50%. Data on dyslipidaemia were lacking and thus were not adjusted for in this study.

Despite these limitations, our study remains the largest study in general African population using a rigorous method of HTN assessment according to the latest guidelines and brings important epidemiological data on HNT in developing countries.

More studies are needed in the future to identify reasons behind increased prevalence and poor blood pressure control and examine trends in prevalence, awareness, treatment, and control. TAHES cohort will go on during the years to come in order to analyse the impact of an educational program, based on local culture and economic characteristics, on the management and control of HTN in this population.

The role of new tools for hypertension management will also be tested.

The aim of the TAHES cohort is to provide epidemiological data and propose ways of improvement in HTN management in order to promote preventive programmes aiming to increase public awareness and educate physicians.

Conclusions

In terms of public health, this large population-based study confirms the high burden of HTN in sub-Saharan Africa and highlights the urgent need for HTN screening, treatment, and control. Less than half of the population with hypertension are aware of their hypertension, indicating the burden of undiagnosed and uncontrolled high blood pressure in these populations(34). Future studies in the same setting would enable to assess the evolution of hypertension and its management in this specific setting.

List Of Abbreviations

BMI: body mass index, CVDs: cardiovascular diseases, DBP: diastolic blood pressure, HTN: hypertension, KH: knee height, SBP:systolic blood pressure, SSA: sub-Sahara Africa, TAHES: TAnve HEalth Study

Declarations

Ethics approval and consent to participate

The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki. All participants and/or their families gave informed consent prior to inclusion in the study. Written consent was obtained whenever feasible. For illiterate people, the study's objectives were verbally explained and consent was obtained by thumbprint.

The study and its consentment method had prior approval of the Benin National Health's Research Ethics Committee and the "Comité de Protection des Personnes du Sud-Ouest et d'Outre-Mer 4 in France".

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request

Competing interests

The authors declare that they have no competing interests

Funding

This study was supported by the APREL Fund from CHU Dupuytren, Limoges (2016) and the research grant of SFHTA (Société Française d'Hypertension Artérielle) and FRHTA (Fondation de Recherche sur l'Hypertension Artérielle) 2018. The sponsors had no role in the design, methods, subject recruitment, data collection, analysis and preparation of this manuscript.

Authors' contributions

Authors I.D, S.A.A, Y.C.H, S.D.H, G.N.G, V.A, P.M.P, Ph.L, designed the study, directed its implementation, including quality assurance and control, and prepared its analytic strategy. Authors S.A.A, Y.C.H, S.D.H, G.N.G, V.A, P.M.P helped conduct the literature review and prepare the Methods and the Discussion sections of the text. Authors I.D and S.A.A did the statistical analysis and drafted the manuscript. Author V.A was a major contributor in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank for their assistance the participants in the TAHES study, as well as the students and nurses who were involved. Thanks also to the participants to the regional health directorate of Zou, the Mayor of Agbangnizoun and his staff, Tanve village's chief ministry and the community health workers in Tanve.

Authors' information

Authors I.D, S.A.A, S.D.H, V.A, P.M.P, Ph.L, members of the Research team INSERM 1094 are actively implicated in epidemiological research of non-communicable diseases in tropical countries, with a particular focus on cardiovascular risk factors in Africa.

Authors I.D and V.A actively participated at the 2018 European Society of Cardiology/European Society of Hypertension Guidelines for the management of arterial hypertension, as members of the ESC/ESH Task Force.

References

1. Mackay J MGTAoHDSG, World Health Organization (WHO), 2004 (http://www.who.int/cardiovascular_diseases/resources/atlas/en/).

2. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet* (London, England). 2017;389(10064):37-55.
3. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, et al. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *Jama*. 2013;310(9):959-68.
4. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet* (London, England). 2005;365(9455):217-23.
5. Opie LH, Seedat YK. Hypertension in sub-Saharan African populations. *Circulation*. 2005;112(23):3562-8.
6. Steyn K, Bradshaw D, Norman R, Laubscher R. Determinants and treatment of hypertension in South Africans: the first Demographic and Health Survey. *S Afr Med J*. 2008;98(5):376-80.
7. Hendriks ME, Wit FW, Roos MT, Brewster LM, Akande TM, de Beer IH, et al. Hypertension in sub-Saharan Africa: cross-sectional surveys in four rural and urban communities. *PloS one*. 2012;7(3):e32638.
8. Houinato DS, Gbary AR, Houehanou YC, Djrolo F, Amoussou M, Segnon-Agueh J, et al. Prevalence of hypertension and associated risk factors in Benin. *Revue d'epidemiologie et de sante publique*. 2012;60(2):95-102.
9. Houehanou YC, Mizehoun-Adissoda C, Amidou S, Desormais I, Houenassi M, Preux PM, et al. Feasibility of a cardiovascular cohort in a Sub-Saharan Africa community: preliminary report of the pilot project TAHES (Tanve Health Study) in Benin. *Global health action*. 2017;10(1):1270528.
10. World Health Organization. NCDs| The STEPS Instrument and Support Materials [Internet]. WHO. [Cited Feb 8 Aahwwinss.
11. GNI per capita Am, <https://data.worldbank.org/indicator/NY.GNP.PCAP.CD>.
12. World Health Organisation TWSatndrfsIW, Available at: http://www.who.int/ncds/surveillance/steps/STEPS_Manual.pdf?ua=1 2017, Accessed date: 1 February 2018.
13. of WHODaD, Hyperglycemia: DMaI, at: RoaWICA, http://www.who.int/diabetes/publications/Definition%,20and%20diagnosis%20of%20diabetes_new.pdf.
14. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension. *European heart journal*. 2018;39(33):3021-104.
15. McLean RM, Farmer VL, Nettleton A, Cameron CM, Cook NR, Campbell NRC. Assessment of dietary sodium intake using a food frequency questionnaire and 24-hour urinary sodium excretion: a systematic literature review. *Journal of clinical hypertension* (Greenwich, Conn). 2017;19(12):1214-30.
16. Guideline:Sodium intake for adults and children. https://www.who.int/nutrition/publications/guidelines/sodium_intake_printversion.pdf.
17. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation*. 2001;104(22):2746-53.

18. Dalal S, Beunza JJ, Volmink J, Adebamowo C, Bajunirwe F, Njelekela M, et al. Non-communicable diseases in sub-Saharan Africa: what we know now. *Int J Epidemiol*. 2011;40(4):885-901.
19. Mensah GA, Roth GA, Sampson UK, Moran AE, Feigin VL, Forouzanfar MH, et al. Mortality from cardiovascular diseases in sub-Saharan Africa, 1990-2013: a systematic analysis of data from the Global Burden of Disease Study 2013. *Cardiovascular journal of Africa*. 2015;26(2 Suppl 1):S6-10.
20. Ogah OS, Madukwe OO, Chukwuonye, II, Onyeonoro UU, Ukegbu AU, Akhimien MO, et al. Prevalence and determinants of hypertension in Abia State Nigeria: results from the Abia State Non-Communicable Diseases and Cardiovascular Risk Factors Survey. *Ethnicity & disease*. 2013;23(2):161-7.
21. Pires JE, Sebastiao YV, Langa AJ, Nery SV. Hypertension in Northern Angola: prevalence, associated factors, awareness, treatment and control. *BMC public health*. 2013;13:90.
22. Wamala JF, Karyabakabo Z, Ndungutse D, Guwatudde D. Prevalence factors associated with hypertension in Rukungiri district, Uganda—a community-based study. *African health sciences*. 2009;9(3):153-60.
23. Helelo TP, Gelaw YA, Adane AA. Prevalence and associated factors of hypertension among adults in Durame Town, Southern Ethiopia. *PloS one*. 2014;9(11):e112790.
24. Adeloye D, Basquill C. Estimating the prevalence and awareness rates of hypertension in Africa: a systematic analysis. *PloS one*. 2014;9(8):e104300.
25. Hamid S, Groot W, Pavlova M. Trends in cardiovascular diseases and associated risks in sub-Saharan Africa: a review of the evidence for Ghana, Nigeria, South Africa, Sudan and Tanzania. *The aging male : the official journal of the International Society for the Study of the Aging Male*. 2019:1-8.
26. Pilleron S, Aboyans V, Mbelesso P, Ndamba-Bandzouzi B, Desormais I, Lacroix P, et al. Prevalence, awareness, treatment, and control of hypertension in older people in Central Africa: the EPIDEMCA study. *J Am Soc Hypertens*. 2017;11(7):449-60.
27. Zatu MC, Van Rooyen JM, Kruger A, Schutte AE. Alcohol intake, hypertension development and mortality in black South Africans. *European journal of preventive cardiology*. 2016;23(3):308-15.
28. Chin JH, Twinobuhungiro A, Sandhu A, Hootsmans N, Kayima J, Kalyesubula R. Determinants of Raised Blood Pressure in Urban Uganda: A Community-Based Case-Control Study. *Ethnicity & disease*. 2017;27(1):15-20.
29. Elliott P. Sodium intakes around the world. Background document prepared for the Forum and Technical meeting on Reducing Salt Intake in Populations (Paris 5–7 October 2006). Geneva W.
30. Powles J, Fahimi S, Micha R, Khatibzadeh S, Shi P, Ezzati M, et al. Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. *BMJ open*. 2013;3(12):e003733.
31. Brown IJ, Tzoulaki I, Candeias V, Elliott P. Salt intakes around the world: implications for public health. *Int J Epidemiol*. 2009;38(3):791-813.
32. Oyebo O, Oti S, Chen YF, Lilford RJ. Salt intakes in sub-Saharan Africa: a systematic review and meta-regression. *Population health metrics*. 2016;14:1.

33. Mishra SK, Togneri E, Tripathi B, Trikamji B. Spirituality and Religiosity and Its Role in Health and Diseases. J Relig Health. 2017;56(4):1282-301.

34. Guwatudde D, Nankya-Mutyoba J, Kalyesubula R, Laurence C, Adebamowo C, Ajayi I, et al. The burden of hypertension in sub-Saharan Africa: a four-country cross sectional study. BMC public health. 2015;15:1211.

Tables

Table 1. Characteristics of study participants:

		Total Population n (%)	Men n (%)	Women n (%)	<i>p</i> Men vs. Women
Total sample size		1777	695 (39.1)	1082 (60.9)	<i>0.2732</i>
Sociodemographic variables					
Age (y)		42.5±16.5	43±16.8	42.1±16.4	<i>0.2732</i>
Marital status					<i><0.0001</i>
	Married/in couple	1336 (75.2)	554 (79.7)	782 (72.3)	
	Single	213 (12.0)	116 (16.4)	97 (9.0)	
	Widowed/divorced/separated	228 (12.8)	25 (3.6)	203 (18.7)	
Education					<i><0.0001</i>
	None	1175 (66.1)	348 (50.1)	827 (76.4)	
	Primary	406 (22.9)	216 (31.1)	190 (17.6)	
	Higher	196 (11.0)	131 (18.8)	65 (6.0)	
Occupation					<i><0.0001</i>
	Farmer/breeder/fisherman	337 (18.9)	181 (26.0)	156 (14.4)	
	Craftsman/storekeeper	1170 (65.8)	385 (55.4)	785 (72.6)	
	Employee/government employee	130 (7.3)	93 (13.4)	37 (3.4)	
	Homemade/retired	50 (2.8)	6 (0.9)	44 (4.1)	
	Jobless	90 (5.1)	30 (4.3)	60 (5.5)	
Religion*					<i><0.0001</i>
	Christians	1009 (61.6)	374 (56.4)	635 (65.1)	
	Animists	529 (32.3)	254 (38.3)	275 (28.2)	
	Other	101 (6.2)	35 (5.3)	66 (6.7)	
Average household income / month					
	Low	1011 (56.9)	237 (34.1)	774 (71.5)	<i><0.0001</i>

	Middle	748 (42.1)	447 (64.3)	301 (27.8)	
	High	18 (1.0)	11 (1.6)	7 (0.7)	
Depression		171 (9.7)	54 (7.8)	117 (10.8)	<i>0.0400</i>
Anxiety		249 (14.0)	63 (9.1)	186 (17.2)	<i><0.0001</i>
Cardiovascular risk factors					
Body mass index: kg/m ²		24.2±11.9	23.2±6.5	24.8±5.6	<i><0.0001</i>
	<18.5	210 (11.9)	97 (14.0)	114 (10.5)	
	18.5-25	1032 (58.3)	436 (63.0)	595 (55.1)	
	25-30	374 (21.1)	129 (18.6)	245 (22.7)	
	≥30	155 (8.7)	30 (4.4)	125 (11.7)	
Tobacco use		152 (8.6)	93 (13.4)	59 (5.5)	<i><0.0001</i>
	Smoked tobacco	100 (5.6)	78 (11.2)	22 (2.0)	<i><0.0001</i>
	Chewed tobacco	76 (4.9)	29 (4.2)	47 (4.3)	<i>0.0658</i>
Diabetes**		55 (3.1)	21 (3.1)	34 (3.2)	<i>0.9168</i>
Physical activity <150 min/week		1037(58.4)	394 (56.7)	643 (59.4)	<i>0,2182</i>
Hypertension		584 (32.9)	228 (32.8)	356 (33.0)	<i>0.9342</i>
Nutritional variables					
Fruits and vegetables <5/day		1603 (90.2)	618 (88.9)	985 (91.0)	<i>0.1512</i>
Alcohol consumption					<i><0.0001</i>
	Abstainers	847 (47.7)	196 (28.2)	651 (60.2)	
	Light	513 (28.9)	200 (28.8)	310 (28.7)	
	Moderate to heavy	417 (23.5)	299 (43.0)	118 (10.9)	
Salt intake (often or very often)***		1029 (58.6)	344 (49.7)	685 (63.3)	<i><0.0001</i>

Table 2. Factors associated with hypertension

Factors	Univariable analysis			Multivariable analysis					
				Model 1*		Model 2*		Model 3*	
	OR (95% CI)	<i>p</i>		OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>
Age (y)	1.04 (1.03- 1.04)	<0.0001		1.04 (1.03- 1.05)	<0.0001	1.04 (1.03- 1.05)	<0.0001	1.04 (1.03- 1.05)	<0.0001
Sex									
	Males	Ref. 1		Ref. 1		Ref. 1		Ref. 1	
	Females	1.00 (0.82- 1.23)	0.9558	1.07 (0.83- 1.38)	0.5977	0.94 (0.72- 1.22)	0.6305	1.01 (0.77- 1.33)	0.9460
Education									
	None	Ref. 1		Ref. 1		Ref. 1		Ref. 1	
	Primary	0.59 (0.46- 0.76)	<0.0001	0.95 (0.71- 1.27)	0.7203	0.92 (0.68- 1.24)	0.5762	0.89 (0.66- 1.22)	0.4908
	Higher	0.72 (0.52- 1)	0.0541	0.70 (0.93- 2.28)	0.1024	0.81 (0.83- 2.12)	0.2345	0.80 (0.86- 2.19)	0.1896
Occupation									
Employee/government employee		Ref. 1		Ref. 1		Ref. 1		Ref. 1	
Craftsman/storekeeper		1.38 (0.90- 2.12)	0.1356	1.37(0.80- 2.34)	0.2478	1.29 (0.75- 2.24)	0.3564		0.3987
Farmer/breeder/fisherman		2.2 (1.38- 3.49)	0.0008	1.44 (0.80- 2.59)	0.2216	1.56 (0.85- 2.84)	0.1501	1.49(0.81- 2.72)	0.1971
Homemade/retired		2.84 (1.42- 5.66)	0.0030	2.31 (1.02- 5.22)	0.0451	2.79 (1.19- 6.53)	0.0173	2.81 (1.19- 6.67)	0.0190
Jobless		5.00 (2.78- 8.99)	<0.0001	1.52 (0.72- 3.20)	0.2736	1.73 (0.81- 3.70)	0.1578	1.61 (0.75- 3.46)	0.2222
Religion									

	Christians	Ref. 1		Ref. 1		Ref. 1			
	Animists	1.97 (1.58- 2.46)	<i><0.0001</i>	1.43 (1.12- 1.82)	<i>0.0042</i>	1.57 (1.22- 2.02)	<i>0.0005</i>	1.52 (1.17- 2.62)	<i>0.0048</i>
	Other	1.22 (0.78- 1.89)	<i>0.3815</i>	1.06 (0.67- 1.69)	<i>0.8021</i>	1.12 (0.68- 1.77)	<i>0.6934</i>	1.16 (0.58- 1.89)	0.7834

Marital status

	Single	Ref. 1		Ref. 1	
	Married/in couple	1 (0.73- 1.38)	<i>0.9621</i>	0.89 (0.63- 1.26)	<i>0.5102</i>
Widowed/divorced/separated		2.84 (1.92- 4.21)	<i><0.0001</i>	1.10 (0.69- 1.77)	<i>0.6800</i>

Average household income/month

	Low	Ref. 1	
	Middle	0.97 (0.81- 1.21)	<i>0.8896</i>
	High	0.78 (0.27- 2.20)	<i>0.6368</i>

Depression

	No	Ref. 1	
	Yes	1.1 (0.79- 1.53)	<i>0.5790</i>

Anxiety

	No	Ref. 1		Ref. 1	
	Yes	1.21	<i>0.1895</i>	0.97	<i>0.8536</i>

		(0.91- 1.59)		(0.71- 1.33)				
Body mass index (kg/m²)		1.02 (1.00- 1.03)	<i>0.0002</i>	1.02 (1.01- 1.04)	<i>0.0001</i>	1.05 (1.02- 1.07)	<i>0.0001</i>	1.04 (1.01- 1.07) <i>0.001</i>
	18.5-25	Ref. 1				Ref. 1		Ref. 1
	<18.5	1.48 (1.08- 2.03)	<i>0.0141</i>			1.05 (0.73- 1.51)	<i>0.7940</i>	1 (0.7- 1.46) <i>0.9598</i>
	25-30	1.92 (1.50- 2.46)	<i><0.0001</i>			2.12 (1.60- 2.79)	<i><0.0001</i>	2.10 (1.59- 2.18) <i><0.0001</i>
	≥30	2.78 (1.97- 3.92)	<i><0.0001</i>			3.57 (2.43- 5.25)	<i><0.0001</i>	3.32 (2.25- 4.89) <i><0.0001</i>
Tobacco Use								
(smoked/chewed)								
	No	Ref. 1				Ref. 1		
	Yes (past/current)	1.54 (1.10- 2.17)	<i>0.0119</i>			0.84 (0.57- 1.26)	<i>0.4021</i>	
Diabetes								
	No	Ref. 1						
	Yes	1.09 (0.62- 1.93)	<i>0.7533</i>					
Physical activity								
	≥150 min/week	Ref. 1				Ref. 1		Ref. 1
	<150 min/week	1.37 (1.11- 1.67)	<i>0.0027</i>			1.17 (0.93- 1.48)	<i>0.1755</i>	1.18 (0.93- 1.49) <i>0.1725</i>
Fruits and vegetables								
	≥5/day	Ref. 1						

<5/day 1.03 0.8363
(0.74-
1.45)

Alcohol consumption

Abstainers	Ref. 1		Ref. 1	
Light	0.77 0.0313 (0.60- 0.98)		0.75 0.0500 (0.57- 0.99)	
Moderate to heavy	1.19 0.1487 (0.94- 1.53)		1.11 0.5162 (0.82- 1.50)	

Salt consumption

Rarely	Ref. 1		Ref. 1	
Often and very often	1.23 0.0512 (1- 1.50)		1.11 0.3801 (0.89- 1.41)	

*Model 1: adjusted for demographic and socio-economic factors

**Model 2: adjusted for demographic, socio-economic and cardiovascular risk factors

***Model 3: adjusted for demographic, socio-economic, cardiovascular risk factors and nutritional factors

Table 3. Factors associated with hypertension in male and female population

Factors	FEMALES		MALES	
	Multivariable analysis*			
	OR (95% CI)	<i>p</i>	OR (95% CI)	<i>p</i>
Age (y)	1.05 (1.04-1.06)	<0.0001	1.03 (1.01-1.04)	<0.0001
Marital status				
Single	Ref. 1		Ref. 1	
Married/in couple	0.92 (0.54-1.55)	0.7499	0.86 (0.52-1.45)	0.5752
Widowed/divorced/separated	0.98 (0.52-1.85)	0.9547	1.38 (0.50-3.81)	0.5280
Education				
None	Ref. 1		Ref. 1	
Primary	0.87 (0.56-1.37)	0.5583	0.91 (0.59-1.41)	0.6763
Higher	1.32 (0.58-3.00)	0.5038	1.21 (0.65-2.34)	0.5519
Occupation				
Employee/government employee	Ref. 1		Ref. 1	
Craftsman/storekeeper	0.98 (0.33-2.93)	0.9745	1.21 (0.63-2.34)	0.5706
Farmer/breeder/fisherman	1.30 (0.41-4.16)	0.6550	1.37 (0.64-2.90)	0.4169
Homemade/retired	2.06 (0.55-7.73)	0.2895	7.19 (0.75-69.29)	0.0876
Jobless	1.46 (0.40-5.34)	0.5638	0.85 (0.26-0.74)	0.7804
Religion				
Christians	Ref. 1		Ref. 1	
Animists	1.52 (1.06-2.16)	0.0211	1.75 (1.17-2.62)	0.0068
Other	0.94 (0.51-1.73)	0.8408	1.72 (0.76-3.89)	0.1921
Depression				
No	Ref. 1		Ref. 1	
Yes	1.94 (0.54-1.66)	0.8444	1.55 (2.56-1.19)	0.1277
Anxiety				
No	Ref. 1		Ref. 1	
Yes	0.87 (0.57-1.40)	0.5789	1.77 (0.91-3.42)	0.0922
Body mass index (kg/m ²)				
18.5-25	Ref. 1			
<18.5	0.56 (0.31-0.99)	0.0482	1.29 (0.75-2.22)	0.3593
25-30	2.16 (1.49-3.12)	<0.0001	1.89 (1.19-3.02)	0.0069
≥30	3.02 (1.89-4.81)	<0.0001	5.28 (2.36-11.77)	<0.0001
Tobacco				
No	Ref. 1			
Yes	0.75 (0.39-1.46)	0.4032	0.83 (0.57-1.19)	0.4916
Diabetes				
No	Ref. 1		Ref. 1	

	Yes	0.83 (0.34-2.00)	0.6723	1.43 (0.53-3.86)	0.4761
Physical activity					
	≥150 min/week	Ref. 1		Ref. 1	
	<150 min/week	1.44 (1.05-1.95)	0.0245	0.99 (0.68-1.44)	0.9520
	≥5/day	Ref. 1		Ref. 1	
	<5/day	0.95 (0.55-1.65)	0.8567	1.13 (0.61-2.10)	0.6993
Alcohol consumption					
	Abstainers	Ref. 1		Ref. 1	
	Light	0.73 (0.51-1.04)	0.0838	0.74 (0.45-0.98)	0.0467
	Moderate to heavy	1.05 (0.64-1.73)	0.6723	1.14 (0.73-1.78)	0.5732
Salt consumption					
	Rarely	Ref. 1		Ref. 1	
	Often or very often	1.42 (1.02-1.97)	0.0365	0.83 (0.57-1.20)	0.3168
<i>*Adjusted for demographic and socio-economic, cardiovascular risk factors and nutritional factors</i>					

Table 4. Factors associated with controlled hypertension

Factors	Multivariable analysis	
	OR (95% CI)	<i>p</i>
Age (y)	0.98 (0.96-1.01)	<i>0.2992</i>
Sex		
	Males	Ref. 1
	Females	2.90 (1.11-5.57) <i>0.0296</i>
Education		
	None	Ref. 1
	Primary	0.57 (0.18-1.79) <i>0.3358</i>
	Higher	1.39 (0.37-5.2) <i>0.6273</i>
Occupation		
	Employee/government employee	Ref. 1
	Craftsman/storekeeper	0.67 (0.14-3.18) <i>0.6177</i>
	Farmer/breeder/fisherman	0.54 (0.09-3.36) <i>0.5103</i>
	Homemade/retired	0.54 (0.05-6.69) <i>0.6279</i>
	Jobless	0.40 (0.03-5.90) <i>0.5040</i>
Religion		
	Christians	Ref. 1
	Animists	0.90 (0.38-2.10) <i>0.7997</i>
	Other	1.81 (0.47-7.01) <i>0.3886</i>
Marital status		
	Single	Ref. 1
	Married/in couple	1.14 (0.36-3.60) <i>0.8216</i>
	Widowed/divorced/separated	0.43 (0.08-2.31) <i>0.3197</i>
Average household income/month		
	Low	Ref. 1
	Middle/high	1.24 (0.56-2.73) <i>0.5899</i>

Figures




Figure 1

Venn diagram representing the prevalence and the proportions of awareness, treatment and controlled hypertension in study population. For each category, values of systolic (SBP) and diastolic (DBP) blood pressures in both sexes.




Figure 2

Venn diagrams representing the prevalence and the proportions of awareness, treatment and controlled hypertension in males and females