

# Implementing a Digital Mental Health Intervention for Individuals With Psychosis - A Multi-country Qualitative Study

**Tamara Pemovska**

Queen Mary University of London, Unit for Social and Community Psychiatry

**Aliriza Arënlju** (✉ [aliriza.arenliu@uni-pr.edu](mailto:aliriza.arenliu@uni-pr.edu))

University of Prishtina, Hasan Prishtina <https://orcid.org/0000-0002-7016-4188>

**Jon Konjufca**

Department of Psychology, University of Prishtina, Hasan Prishtina

**Fitim Uka**

Department of Psychology, University of Prishtina, Hasan Prishtina

**Jennifer Hunter**

School of Health Sciences, City University of London

**Stojan Bajraktarov**

Saints Cyril and Methodius University in Skopje Medical Faculty

**Lidija Injac Stevović**

Psychiatry Clinic, University of Montenegro, Podgorica

**Stefan Jerotić**

Faculty of Medicine, University of Belgrade, Serbia

**Alma Džubur Kulenović**

Department of Psychiatry, Clinical Centre of the Sarajevo

**Antoni Novotni**

University Clinic of Psychiatry, Ss. Cyril and Methodius University in Skopje

**Ljubisha Novotni**

University Clinic of Psychiatry, Ss. Cyril and Methodius University in Skopje

**Tamara Radojičić**

Psychiatry Clinic, University of Montenegro, Podgorica

**Selman Repišti**

Psychiatry Clinic, University of Montenegro, Podgorica

**Emina Ribić**

Department of Psychiatry, Clinical Centre of the University of Sarajevo

**Ivan Ristić**

Faculty of Medicine, University of Belgrade, Serbia

**Eldina Smajić Mešević**

Department of Psychiatry, Clinical Centre of the University of Sarajevo

**Mirjana Zebić**

Faculty of Medicine, University of Belgrade, Serbia

**Nikolina Jovanović**

Unit for Social and Community Psychiatry, Queen Mary University of London

---

## Research

**Keywords:** implementation science, psychosis, mental health care, psychosocial intervention, digital intervention, Southeast Europe, low- and middle-income setting, multi-country qualitative research

**Posted Date:** July 29th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-47980/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Psychiatry on September 25th, 2021. See the published version at <https://doi.org/10.1186/s12888-021-03466-x>.

# Abstract

**Background:** Implementation of psychosocial interventions in mental health services has the potential to improve the treatment of psychosis spectrum disorders (PSD) in low- and middle-income countries (LMICs) where care is predominantly focused on pharmacotherapy. The first step is to understand the views of key stakeholders. We conducted a multi-language qualitative study to explore the contextual barriers and facilitators to implementation of a cost-effective, digital psychosocial intervention, called DIALOG+, for treating PSD. DIALOG+ builds on existing clinician-patient relationships without requiring development of new services, making it well-fitting for healthcare systems with scarce resources.

**Methods:** The focus group study included 174 participants (patients, clinicians, policymakers and carers), who were familiarized with DIALOG+ through a presentation. The Southeast European LMICs included in this research were: Bosnia and Herzegovina, Kosovo[1], North Macedonia, Montenegro and Serbia. Framework analysis was used to analyse the participants' accounts.

**Results:** Six major themes were identified. Three themes (Intervention characteristics; Carers' involvement; Patient and organisational benefits) were interpreted as perceived implementation facilitators. The theme Attitudes and perceived preparedness of potential adopters comprised of subthemes that were interpreted as both perceived implementation facilitators and barriers. Two other themes (Frequency of intervention delivery; Suggested changes to the intervention) were more broadly related to the intervention's implementation. Participants were exceedingly supportive of the implementation of a digital psychosocial intervention such as DIALOG+. The attractive intervention's characteristics, efficient use of scarce resources for its implementation and potential to improve mental health services were seen as the main implementation facilitators. The major implementation barrier identified was psychiatrists' time constraints.

**Conclusions:** This study yielded important insights regarding implementation of digital psychosocial interventions for people with PSD in low-resource settings by including perspectives from four stakeholder groups in five LMICs in Southeast Europe – a population and region rarely explored in the literature. The perceived limited availability of psychiatrists could be potentially resolved by increased inclusion of other mental health professionals in service delivery for PSD. These findings will be used to inform the implementation strategy of DIALOG+ across the participating countries. The study also offers insights into multi-country qualitative research.

[1] By United Nations resolution 1244

## Contributions To The Literature (100/100)

- 1. Treatment for psychosis has been centred on medication, while psychosocial aspects have often been neglected. Insights into factors likely to impact the implementation of psychosocial interventions in low-resource settings are essential to achieve a comprehensive mental health care globally.

- 2. This research offers unique insights into implementation factors identified by clinicians, patients, policymakers and carers, emphasizing the importance of including multiple stakeholders' perspectives in implementation science.
- 3. The study was conducted across five Southeast European countries – a region often considered a blind spot on the global mental health map.
- 4. The study offers unique insights into multi-language, cross-country qualitative research in implementation science.

## Background

Effective treatment and management of psychotic spectrum disorders (PSD) is multi-faceted and expensive [1, 2]. PSD frequently cause severe disability, more often than other mental health disorders, resulting in significant health, social and economic burden [1].

In many countries such as the United Kingdom (UK), evidence-based treatment of PSD involves pharmacotherapy, family education and psychosocial interventions [3]. However, care for PSD in low- and middle-income countries (LMICs) in Southeast Europe (SEE) is characterized by meetings between clinicians and patients dominated by pharmacotherapy while psychosocial aspects tend to be overlooked, reportedly because of the lack of qualified staff and sufficient funding of mental health services [4, 5]. This facilitates additional inequalities of an already highly vulnerable population.

Cost-effective and easily deliverable digital psychosocial interventions embedded on a device, such as an application on a tablet [6], offer an opportunity to advance mental health care to people affected by PSD in LMICs and to alleviate the burden of psychosis on patients and their families.

One such intervention is DIALOG+, which is based on patient-centred communication, quality of life research and solution-focused therapy and designed to make routine mental health clinical meetings more therapeutically effective. The intervention is available as an app and uses a tablet computer. Each session begins with the patient rating their satisfaction with eight life areas and three treatment areas on a DIALOG scale of 1 (totally dissatisfied) to 7 (totally satisfied). The ratings are displayed on the tablet screen, allowing for comparisons with ratings from previous sessions. Each rating is reviewed through positive reinforcement and particular areas of concern are addressed in more detail in a four-step, solution-focused approach which aims to identify and utilize the patient's existing resources [7]. Additional file 1 and 2 contain visual representations of DIALOG+ and the Template for Intervention Description and Replication Checklist [8], respectively.

DIALOG+ builds on the existing clinician-patient relationships which do not require formation of new services or referrals to other clinicians, making it a good fit for healthcare systems with scarce resources. After three hours of training, a range of clinicians have been shown to be able to use the intervention [9]. DIALOG+ has proved effective in decreasing clinical symptoms, increasing quality of life and reducing treatment costs of patients with psychosis in the UK [9]. A pilot study in Germany also showed significant

amelioration of schizophrenia and depressive symptoms as well as quality of life [10]. A process evaluation of the intervention identified the following likely 'active ingredients': 1) solution-focused structure; 2) patient empowerment; 3) mobilizing social resources; and 4) self-reflexion and therapeutic expression [11]. DIALOG+ has been implemented as a care planning tool and patient-reported outcome measure in health institutions in England and Wales [12–14]. It is also being researched in implementation and effectiveness studies across four continents [10, 15, 16], emphasizing the flexible use of the intervention.

Translating evidence-based interventions from high-income countries (HICs) to LMICs requires an in-depth understanding of contextual barriers and facilitators to implementation [17], factors which should be considered before implementation of any novel intervention to ensure the greatest chance of a successful outcome [18, 19]. Therefore, this study aims to identify the perceived implementation facilitators and barriers of an evidence-based, digital, psychosocial intervention for patients with psychosis in five LMICs in SEE. The findings could provide evidence about the provision of digital psychosocial treatment for PSD in low-resource mental health settings and facilitate successful uptake of such interventions.

## Methods

### Design

This is a multi-country, multi-language qualitative study of the perceived implementation feasibility of DIALOG+. It involved primary data collection using qualitative methods in the form of semi-structured focus groups to explore perceived implementation facilitators and barriers of DIALOG+ from the perspective of key stakeholders, patients with psychosis, carers, clinicians and policymakers, in five SEE countries (Bosnia and Herzegovina, Kosovo[1], North Macedonia, Montenegro and Serbia). Local Lived Experience Advisory Panels (LEAP) were formed in each country to involve patients and carers as experts by experience in the research. Additional file 3 contains SRQR reporting guidelines [20].

The study is part of the larger research project "Implementation of an effective and cost-effective intervention for patients with psychotic disorders in low and middle-income countries in Southeast Europe" - IMPULSE, which is funded by the European Commission [21].

### Data Collection

#### *Participants*

The research included 174 participants who participated in 31 focus groups and one interview. The participants were a mix of patients, carers, clinicians and policymakers (Table 1). There was a higher proportion of female participants (n = 108; 62%) and most participants were patients (n = 59; 34%) followed by clinicians (n = 47; 27%), caregivers (n = 40; 23%) and policymakers (n = 28; 16%).

<sup>a</sup>By United Nations resolution

Clinicians, patients and carers were purposively recruited from mental health services, service user organisations and mental health non-governmental organisations. Policymakers and service managers were purposively recruited from Ministries of Health, National Psychiatric Associations and mental health medical boards. Patients' inclusion criteria were: 18 to 65 years of age; in psychiatric treatment for at least three months; clinical diagnosis of psychosis or related disorder (i.e., ICD-10 F20-29, F31); and capable of giving informed consent. The inclusion criteria for clinicians were: professional qualification working with mental health services and more than six months working experience in mental health. Patients' carers were included if they were: at least 18 years of age; a carer for someone with mental health difficulties or providing support outside of clinical context; and capable of giving informed consent. The inclusion criteria for policymakers and services providers were managerial and/or policymaking experience in health care, local or national authorities.

31 focus groups were conducted: eight with clinicians, seven with policy makers, nine with patients and seven with carers. One policymaker was interviewed because they were unable to attend the focus groups due to other commitments (Table 1).

### ***Topic guides***

The topic guides explored participants' views on barriers, facilitators and perceived benefits of implementing DIALOG+. They were based on the same set of topics for patients/family members and clinicians/policymakers, but with probes specific for each group (Additional file 4). The same topic guide was used during the interview with a policymaker.

The topic guides were developed in English in an iterative process, and draft versions were circulated among all collaborating centres for feedback. The final versions of the topic guides were translated into the national languages.

### ***Procedure***

The focus groups lasted 59.4 minutes on average and used similar methodology in the following sequence: 1) introduction and explanation of the aim of the focus group; 2) standardized presentation about DIALOG+, lasting 10 minutes on average; 3) focus group discussion related to barriers and facilitators to implementation of DIALOG+.

Focus groups took place between July and November 2018, in private rooms at the collaborating SEE mental health centres by male and female facilitators, who were local medical doctors and/or psychologists. Additional training in facilitating focus groups prior to data collection was provided to ensure a standardized approach.

All focus groups were audiotaped; transcribed verbatim and de-identified. The transcripts were translated to English either by local researchers proficient in the English language or by professional translators.

## Data Analysis

We analysed data using thematic framework analysis [22] suitable for the large breadth of our dataset, collected in five different languages. Our research question was classified within the “diagnostic” and “strategic” typologies defined by Ritchie and Spencer [22] because we were interested in identifying both perceived barriers and facilitators to DIALOG+ implementation (diagnostic) and how they could potentially be addressed (strategic) [22, 23].

The data analysis was conducted by four researchers with background in psychology and public health in Kosovo[2] (AA, JK) and in the UK (TP, JH). First, we familiarized ourselves with all the transcripts and each researcher created a preliminary coding list in light of our research question to identify perceived facilitators and barriers as well as any other codes developed and interpreted from the data. The individual preliminary coding lists were combined in a single one created by consensus through face-to-face discussions. The coding list was distributed to the focus group facilitators to check for accuracy.

Next, we grouped the preliminary codes to develop broad framework categories based on both concerns related to our research question and other emerging ones from the familiarization stage. Upon agreement of the framework, two researchers coded focus groups from clinicians and policy makers, and two other researchers coded focus groups from patients and carers using ATLAS.ti qualitative analysis software. Same transcripts were coded by researchers until there were no discrepancies between them.

After organizing the data according to the framework categories, they were summarized for each category in a chart form. This was followed by mapping and interpretation of the data to find patterns. This process was on-going until no additional patterns were found.

The LEAP provided feedback on the findings from focus groups with patients and carers.

## Results

The results present the major themes related to the research aim to identify perceived facilitators and barriers to DIALOG+ implementation by four key groups of stakeholders. Two developed themes (Frequency of intervention delivery; Suggested changes to the intervention) are more broadly associated with DIALOG+ implementation. Differences between participant groups are explored within the identified themes. Views from the interviewed person were in line with views elicited in focus groups.

Six major themes were developed: Intervention characteristics; Attitudes and perceived preparedness of potential adopters; Carers’ involvement; Perceived potential patient and organisational benefits; Frequency of intervention delivery; and Suggested changes to the intervention (Figure 1). countries in Southeast Europe (SEE).

Each major theme was developed from subthemes, summarized in Table 2.

## Intervention Characteristics

The intervention's characteristics, likely the starting point of adopters' engagement with the intervention, were largely interpreted as facilitators to the implementation across participant groups. Clinicians, patients and policymakers viewed DIALOG+ as clear and easy to use. Clinician45 from Serbia expressed "[DIALOG+] couldn't be simpler". Yet, a few patients expressed ambiguity when asked about their understanding of the DIALOG+ rating scale and its items. Previous familiarity with Likert scales and quality of life measures is likely to influence how easily patients can understand the rating scale that is part of the intervention.

Patient31: Yes, [DIALOG+ rating scale] is a classic Likert scale.

Patient30: [About the rating scale] when there are such gradations, one gets confused. I think there is no need for more than five [options]. It should be simpler.

Clinicians, patients and policymakers viewed DIALOG+ as adding structure to routine clinician-patient meetings while involving the psychosocial needs of the patient. Clinicians spoke about DIALOG+ creating a workflow for patients and clinicians that could lead to more comprehensive therapeutic approaches. Regularly using DIALOG+ would make it less likely for clinicians to omit topics that might be relevant to treatment.

Clinician9: I like that this intervention is well-structured, involving all the most important areas of one's life and visual approach that doesn't let us forget some domain.

Clinicians, patients and carers found the ability of the intervention to track patients and their treatment progress as particularly attractive.

Carer11: [With DIALOG+] there will be a follow up of [the patient's] condition, which is the most important thing for the patient, number one issue (...) Simply, his condition is monitored, how is it today, is it getting worse or better, are there any improvements.

## Attitudes and Perceived Preparedness of Potential Adopters

Some patients, policymakers and carers expressed that the resources to implement DIALOG+ in their facilities are readily usable and available.

Patient19: The medical personnel are sufficient [to implement DIALOG+]. Policymaker3: [DIALOG+] is basically free. There are lots of things that I would like to implement here, but they are expensive (...) But this can be delivered by nurses as well.

However, when asked about resources that might be lacking, other participants expressed serious concerns about the limited resources available, particularly regarding time, staff, funding and space. Conflicting views regarding the existing resources to implement the intervention may reflect resource differences in the explored services.

Patient2: Medical staff of this clinic, they are great, but there just aren't enough of them.

Carer39: I will give a specific example [of a usual clinical meeting] in my experience as both a caregiver and a patient. The doctor asks me – “Are things better? Are life's problems worse than before?” but I try to make my conversations with the doctor as brief as possible. I can see there are so many patients waiting to be examined. I can see that patients come from all over the country with their bags, waiting to be examined. So with the doctor, I speak shortly and concretely. [DIALOG+] is something else; this is likely to last longer.

Policymaker26: In Serbia, it is practically impossible in the conditions of usual ambulatory controls, we do not have time for it, really.

Clinician34: You have to give tablet computers to every clinician and pay for tablets and software and everything else. If there is no money for it, how...

Clinicians, patients and policymakers viewed DIALOG+ as consistent with existing norms, policies and procedures in their services. However, participants from Serbia and North Macedonia questioned the suitability of DIALOG+ with the current funding policies that prioritize the quantity of patients examined. Furthermore, some policymakers expressed that implementing DIALOG+ might lead to “loss of individualized approach” to patients. This is likely motivated by the difference between the current, largely unstructured clinical meetings in mental health care compared to the structured procedure of using DIALOG+ supported by technology. It is interesting that the same characteristics of the digital intervention which are considered to dehumanize the intervention by some are also highly appreciated by others.

Policymaker25: DIALOG+ would fit well, I think, very well in the current practice, because, it is actually a modification of the practice that is already being done, but it is more effective and faster.

Policymaker2: I am just afraid of robotizing sessions after certain period of time.

Opposing views were expressed regarding the motivation across participants and peers to implement DIALOG+. Some participants expect that there will be clinicians' resistance to innovation. Such views likely reflect the context of mental health services in the countries, where digital interventions are not a common practice.

Clinician9: I think that [using DIALOG+] is related to the [clinician's] individual approach; surely some of our colleagues would be thrilled.

Policymaker4: As a member of a management board, I strongly believe that this can be implemented in our routine practice when we do all the legal work needed for something to become a routine intervention... certification, coding, and setting up the price. I am definitely in for implementing this.

Patient5: I could use it every time... I need to talk more, to tell what's on my mind, but the doctor never has enough time

Polycymaker2: I think opinions would be divided. You cannot expect all [clinicians] to accept something; some people have resistance towards technology...

For most clinicians in the study it is vital that their service managers support them in the implementation of DIALOG+. Clinician14 stated "I think that we must first have the support of management".

Additionally, all participant groups reported that skills and experience to implement DIALOG+ by key stakeholders are already existent. Some clinicians and patients asked for more information on how to use DIALOG+. Participants only received a short presentation describing the intervention, which is likely the motivation for such accounts. A few patients reported the need for training due to their lack of familiarity with smart phones/tablets.

Clinician3: We would have to get some instructions, this presentation was great, but we would need detailed training.

Polycymaker8: No, [DIALOG+] won't be different [in the skills required] (...) all of these [clinicians] are trained and they talk to patients – some for therapy, some for supportive psychotherapy – and each of them has his own way.

Patient3: I am good with computers, this is too easy.

Patient10: I don't use a smart phone... so I would need [training].

An important implementation facilitator interpreted from the data is the participants' perceived need for DIALOG+ implementation, particularly as it supports innovation in mental health care. Patients and carers express need for the intervention because current clinical meetings are perceived to last too short to fully understand the patient and limiting in the scope of conversation.

Patient45: [Clinicians] also pay little attention to the patient during the appointments, only 10-15 minutes or 5, irrelevant, but I think [DIALOG+] is already better.

Clinician9: I think that [DIALOG+] is very useful, that is exactly what we need.

Polycymaker17: In the new National Strategy that is under preparation, we have included a mental health category. So this project comes at the right time so that we can think about which services we can develop. We are very thin in this field. It should start inter-sector cooperation at the local level. This project can help us a lot...

Carer26: I think it will succeed, [implementing DIALOG+] is a step forward in medicine, having in mind how patients are treated, all is done in the old fashioned way, and nothing has improved until now.

Participants expressed that poor mental health of patients can be a barrier to DIALOG+ implementation. Polycymaker2 said that "[DIALOG+ areas are significant when patients reach remission, when they can

handle real-life situations.” This result highlights that the intervention may not be suitable for all patients and some selection criteria may be necessary.

### **Carers’ Involvement in Intervention**

Participants expressed varying views regarding involvement of carers in DIALOG+ implementation, although all participant groups saw carers to have a facilitating role. Clinicians and carers proposed carers to be involved in the planning and execution of actions agreed from the sessions. Clinicians expressed that DIALOG+ could offer carers psycho-education. However, patients and clinicians also showed reluctance about the extent to carers’ involvement during sessions. The accounts suggest that the level of carers’ involvement with such an intervention requires adjustment to the patient’s needs.

Clinician10: We should just keep caregivers away from assessing areas instead of patients.

Policymaker3: I think caregivers should take part in this. Whether they take part on the sessions or afterwards... it depends on what the patient wants...

Patient8: I would like [family members to be involved] in that part where doctor gives tasks to family members and me. I wouldn’t like them there all the time.

Patient56: [Carers] could encourage and stimulate clients, and at the same time they could help clinicians be better informed about their clients’ problems – from a different point of view. I think carers could understand and explain certain issues more objectively than clients themselves.

### **Perceived Potential Patient and Organisational Benefits**

Clinicians, patients and carers reported patient empowerment and an improved therapeutic relationship as potential benefits of the intervention to the patient. Other potential benefits at an organisational level were also reported: DIALOG+ would help shift the mental health services away from a typical medical approach to care by reinforcing a psychosocial care approach (clinicians, patients, carers); care could become more patient-centred (patients and carers); the structure and follow-up of care could be improved (all participant groups), and DIALOG+ could offer an opportunity for care to be standardized across services (by policymakers). These subthemes reflect how DIALOG+ could add to the routine service delivery and the elicited views are likely related to what participants are missing from their current care.

Carer36: The clinician gets closer to the patient, and the patient increases his confidence in the clinician [with DIALOG+]. If this intervention goes on for a longer time, this can only deepen and expand the relationship

Patient3: I like that we have freedom, that we can follow this app and that we have information from previous sessions. We see if we have made progress or not, we can see if the doctor’s therapy, advice, or drug were adequate.

Patient56: It would motivate me to work harder on myself and spend less time doing nothing but wandering around.

Policymaker2: [DIALOG+] would be good because we would standardize appointments in time, content, and approach.

Clinician3: It increases functionality in every aspect, what is important for the patient, engages patients in decision-making process, motivates them. That increases self-confidence and empowers the patient.

Clinician45: The mere fact that conversations with my patients would cover eight life areas would bring change, since that was not part of my routine clinical practice so far. I think that structured conversation would allow us to discuss more things in less time and to be sure that we didn't miss any important life area

### **Frequency of Intervention Delivery**

Some participants provided brief answers about the frequency of DIALOG+ delivery. Once per month was the most common view, particularly among clinicians and policymakers. Some patients and carers requested more frequent delivery, for instance weekly or bi-weekly. Less frequent delivery, such as "once in two months" and "at least twice a year", was favoured by some other patients and carers. Carer38 proposed DIALOG+ be delivered every six months because changes to patients' life satisfaction take time. Carer37 expressed that DIALOG+ "should be applied regularly during outpatient appointments". These accounts suggest that the frequency of DIALOG+ sessions should be determined by the individual patient's needs.

### **Suggested Changes to the Intervention**

Some participants' accounts contained broad recommendations about changes to DIALOG+, however these was not explored in detail. Suggestions about the DIALOG scale were made, such as adding a miscellaneous field so patients can discuss topics not included in the scale, and reducing it to a 5-point Likert scale. Clinician46 expressed that "it might be practical and easy to use smartphones for [delivering DIALOG+]". About the area 'job situation', policymaker4 said "it should be clearer that it also refers to education, or how the patient is dealing with being unemployed." Clinician36 spoke about the longer time needed to see changes in patients' satisfaction with areas such as 'job situation' and 'accommodation' than that with their mental health, thus "perhaps these should be assessed only at a 6-month interval."

## **Discussion**

Our study's findings offer important implications for implementation of digital psychosocial intervention for PSD in low-resource settings.

Our results show that resources regarding time, staff, funding and space for the implementation of a novel intervention in the explored services are perceived as limited. Lack of financial and human

resources have been found to be key implementation roadblocks for digital interventions for people with psychosis [24]. The cost of tablets overall was not considered to be a major burden in the implementation of DIALOG+ in the context of SEE. Rather, participants' accounts focused more on the lack of human resources. Special consideration should be provided to psychiatrists' time limitations that arise as a result of limited human resources for the number of patients, because this barrier was mentioned across stakeholders and countries. To maximize its sustainability, DIALOG+ should not require much longer clinical meetings than usual. The findings suggest inclusion of other mental health professionals in the implementation process of DIALOG+ to address the limited availability of psychiatrists in most of the participating countries. Such suggestions were also given by the LEAP and support the stepped care approach in mental health services [25], especially in contexts with limited human resources. Digital support tools have been reported to facilitate a diverse set of mental health workers in delivering the care needed by patients with severe mental illnesses in the community, which is of particular necessity in LMICs [26].

Attractive intervention characteristics, such as clarity and easiness of use, and ability to provide structure to patient-clinician meetings, involve patient's psychosocial needs and track patient's progress were implementation facilitators considered important by the key stakeholders. 'Perceptions of the innovation' has been identified as one of the basic determinants associated with the rate of implementation of the intervention [27], thus it is paramount that key stakeholders have positive perceptions of the intervention. The findings also support previous reports that the comprehensive structure enabled by DIALOG+ is one of its mechanisms of action [11]. Automated data collection offered by digital psychosocial interventions is useful to systematically measure the quality of psychosocial care, a shortcoming identified with current psychosocial therapies [26].

Despite the identified implementation facilitators, the findings show the need to consider the patients' level of existent familiarity with life and treatment satisfaction questionnaires using Likert scales when introducing the intervention, as a few patients found the DIALOG scale unclear. Patients who used DIALOG+ in the UK also reported 'difficulty to understand or cope with questions', although this was not explored in detail [11]. Similarly, a systematic review of implementation determinants for digital interventions for people with psychosis recognized the complexity of the intervention as an impediment for those with psychiatric symptoms, premorbid intelligence quotient and low technological competence [24]. Some clinicians and patients directly expressed a need to receive more training, largely explained by receiving only a short presentation of the intervention prior to the focus group discussions and some patient's lack of familiarity with smart phones. Thus, besides clinicians, patients may also need training to use the intervention as planned. Such training could be delivered by the clinicians who can explain key principles and procedures to the patient during their first sessions.

Additionally, many positive attitudes and sufficient level of preparedness for DIALOG+ implementation across potential intervention adopters was reported, particularly related to the perceived existence of the required skills and high stakeholders' motivation. Such implementation facilitators are of significance

because negative attitudes and scepticism have been shown to reduce motivation to adopt the interventions for both clinicians and patients with psychosis (35).

Some participants raised concerns about potential resistance to innovation in mental health care by clinicians, particularly to the technological aspect of the intervention as use of digital interventions is not part of routine mental health care in the SEE countries that participated in this study. A study exploring the experiences of using DIALOG+ in the UK also identified 'initial apprehension to the technology' as a barrier [11]. The LEAP similarly stressed that clinicians need to be adequately trained, motivated and willing to use DIALOG+, while not having any resistance. Thus, the potential resistance to innovation should be considered when developing training for digital interventions. Moreover, carers and policymakers expressed concerns that poor mental health would limit patients' engagement and thus negatively impact DIALOG+ implementation, further highlighting that the intervention may not be suitable for all patients and clinicians. Accessibility and adaptability of digital interventions for people with psychosis and treating clinicians are reported as main implementation facilitators of digital interventions identified in previous studies [24] and need to be carefully considered.

Despite the views of potential resistance to innovation, key stakeholders perceived the intervention as needed to improve the scope of the conversations during routine clinical meetings. This is promising because stakeholders not perceiving that a change is needed is an important barrier to innovation implementation in mental health care (29).

DIALOG+ was mostly seen as consistent with existing norms and practices, an aspect that is often reported among strong enablers of innovation implementation across health care [28, 29]. However, some important opposing views regarding the incompatibility of DIALOG+ with the mental healthcare funding policy in North Macedonia and Serbia and the lack of intervention flexibility should be considered. Similarly, the LEAP highlighted that asking questions in a mechanical way could make the sessions 'boring and overwhelming'. Previous research in the UK similarly identified repetitiveness of DIALOG+ as a barrier [11]. This result highlights the importance of considering individual patient's needs regarding how frequently the intervention is used to avoid repetitiveness. The UK trial of DIALOG+ suggested that the intervention be used monthly for three months and flexibly afterwards [9]. However, the initial perception among participants in this study was that the intervention should be delivered once a month.

Clinicians expressed that institutional support is a strong determinant for successful DIALOG+ implementation. This insight adds to the limited evidence of contextual and organisational factors regarding implementation of digital interventions for PSD (35).

Different opinions on the way and the extent to which carers should be involved were expressed. The findings together with the LEAP feedback indicate that it is important to involve carers in the implementation process and develop their specific role through discussion between clinicians, patients and carers.

Increased patient empowerment and strengthened clinician-patient relationship were considered to be important potential individual benefits of DIALOG+ by all key stakeholders. This is in line with results from the UK study researching mechanisms of action of DIALOG+ [11]. The opportunity of DIALOG+ to shift the typical medical approach to a more psychosocial approach – an approach that was recognized by all stakeholders as often neglected – was expressed as a potential organisational benefit. In this approach, patients would be at the centre of their treatment and DIALOG+ would offer improved structure and follow up of sessions. These views were also shared by the LEAP. The identified potential benefits from participants' accounts present important facilitators to further the implementation process of digital psychosocial interventions in the explored context in SEE.

Digital interventions are becoming an integral part of the transformation of mental health care because of their potential to advance current services and to enable new pathways for provision of innovative psychosocial therapies [30, 31]. The process of implementing a digital intervention can be time consuming as it requires changes to routine services at an individual and an organisational level that frequently do not occur successively [32]. The collaboration between key stakeholders in the development and implementation of digital interventions is a crucial promoter of implementation of digital interventions for people with psychosis [24]. In order for technology to meaningfully 'disrupt' the current approach to mental health care, it needs to be inclusive of people with severe mental illnesses, consider how mental health providers can use such technologies most adequately and develop digital interventions with having low-resource settings in mind.

## **Strengths and Limitations**

A major strength of this study was the inclusion of perspectives from four key stakeholder groups and feedback from LEAP across five LMICs in SEE, which is rarely seen in the literature. Early communication with key stakeholders is crucial to innovation development and successful implementation as it enables an increased understating of the context in which intervention implementation occurs [33]. The findings from this study contribute to the field of implementation science because they provide insight into currently limited knowledge of perceived facilitators and barriers of implementing a digital psychosocial health intervention in low-resource settings for PSD.

The study has several limitations. Participants' responses about changes to DIALOG+ and its delivery frequency were not explored in detail during the focus groups, limiting the subsequent data analysis. Nonetheless, these data were included since they were related to the study's research aim; however future research should explore the related themes further. Although all focus groups facilitators received training and had topic guides, the quality and depth of information varied across focus groups. This might have occurred because of various styles of individual facilitators and differences in their background (e.g., medical vs. psychological), as well as because of the varied motivation of participants to contribute to a theme. This is often reported as a challenge when conducting cross-national and cross-disciplinary qualitative research as each team member has their own combination of theoretical, national and linguistic underpinnings, as well as varying levels of methodological experience [34]. Additionally, all

transcripts were translated into English from five different languages, which introduced a risk of losing contextual-specific meaning and traits [34]. However, this was a pragmatic decision as early stage translation enabled collaboration in the analysis process within the research team. The transcripts were not back-translated or checked with focus group participants; thereby, the process of translating transcripts could have introduced a bias that might have influenced the analysis of transcripts. However, the analysis team was in regular contact with focus group facilitators to clarify any ambiguous meaning of words. The analysis team consisted of researchers that were both proficient in English and some of the SEE languages and were able to go back to original transcripts where necessary. To future cross-national, multi-disciplinary research teams conducting qualitative studies, we recommend in-depth training in qualitative methods, involving researchers from all countries in the analysis process and maintaining transcripts in their original languages. We hope that our approach and lessons learned can contribute to the literature on conducting such qualitative research.

## Conclusion

This study provides increased understanding from different stakeholders' perspectives regarding the implementation of a digital psychosocial intervention for PSD in five LMICs in SEE. Implementation of DIALOG+ is supported by the key stakeholders, who overwhelmingly reported implementation facilitators compared to barriers, and as such is promising to advance care for PSD patients in SEE. DIALOG+ fits well with the organisational goal of the mental health centres of interest, but would require transformation of the nature of routine clinical meetings in mental health care, which can be challenging for both clinicians and patients. Identification of potential barriers and facilitators promotes successful uptake of the intervention, minimizes waste of scarce resources and provides evidence about the provision of digital psychosocial treatment for PSD in LMICs. Working towards implementation of DIALOG+ is an opportunity to move beyond lip service that mental health services need to pay more attentions to psychosocial aspects. These findings, together with the existing evidence base, provide support for further examination of the effectiveness and implementation of DIALOG+ in LMICs in SEE with the expected impact of improving management of PSD in this region.

## Abbreviations

PSD: psychosis spectrum disorders

SEE: Southeast Europe

LMICs: low- and middle-income countries

LEAP: Lived Experience Advisory Panel

## Declarations

*Ethics approval and consent to participate*

Research activities followed current legislation and regulations in the participating SEE centres where the research was carried out. Each SEE collaborating centre obtained ethical approval from their local ethical committee prior to study beginning: Ethics committee of the Clinical Centre of the University of Sarajevo, Bosnia and Herzegovina (Ref: 03-02-47500, date approved: 13/09/2018); Komisioni Etiko Profesional, Hospital and University Clinical Service of Kosovo, University Clinical Centre of Kosovo, Kosovo[1] (Ref: 904, date approved: 08/06/2018); Ethical Committee for Research with Humans, Medical Faculty at the University of Cyril and Methodius in Skopje, North Macedonia (Ref: 03-2237/12, date approved: 21/05/2018); Ethics Committee for the Clinical Centre of Montenegro, Montenegro (Ref: 03/01-11066/1, date approved: 19/07/2018); Ethical Committee of the University of Belgrade Faculty of Medicine (Ref: 2650/V1-3, date approved: 26/06/2018).

### ***Consent for publication***

Not applicable.

### ***Availability of data and materials***

The dataset (which includes individual transcripts) used and analysed during the current study is not publicly available due to confidentiality policies.

### ***Competing interest***

All authors declare that they have no competing interests.

### ***Funding***



The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded as part of the IMPULSE project under the European Union's Horizon 2020 research and innovation programme [grant number 779334]. The IMPULSE project has received funding through the "Global Alliance for Chronic Diseases (GACD) prevention and management of mental disorders" (SCI-HCO-07-2017) funding call.

### ***Author's contributions***

NJ, chief coordinator of the IMPULSE project, and AA, co-grant holder of the project, wrote the study design. AA, JK, FU, ST, LS, SJ, AK, AN, LN, TR, SR, ER, IR, EM and MZ were involved in facilitating the focus groups, transcription and translation of transcripts. AA, TP, JK, and JH were involved in data analysis. TP and AA were involved in drafting the paper. NJ provided oversight of the project and has reviewed the different versions of the article. All authors read and approved the final submitted manuscript.

### ***Acknowledgements***

We would like to acknowledge the study participants who gave their valuable time to participate in this study as well as the members of the Local Lived Experience Advisory Panels for their helpful feedback.

## References

- [1] Eaton WW, Martins SS, Nestadt G, et al. The burden of mental disorders. *Epidemiol Rev* 2008; 30: 1–14.
- [2] Chong HY, Teoh SL, Wu DB, et al. Global economic burden of schizophrenia: a systematic review. *Neuropsychiatr Dis Treat* 2016; 12: 357–373.
- [3] NICE Treatment Guidelines for Managing Psychosis, <https://www.nice.org.uk/guidance/CG178> (2014, accessed 4 June 2020).
- [4] World Health Organization. Mental Health Atlas 2017, [https://www.who.int/mental\\_health/evidence/atlas/mental\\_health\\_atlas\\_2017/en/](https://www.who.int/mental_health/evidence/atlas/mental_health_atlas_2017/en/) (2017, accessed 4 June 2020).
- [5] World Health Organisation. Mental Health Atlas 2014, [https://www.who.int/mental\\_health/evidence/atlas/mental\\_health\\_atlas\\_2014/en/](https://www.who.int/mental_health/evidence/atlas/mental_health_atlas_2014/en/) (2014, accessed 4 June 2020).
- [6] Murray E, Hekler EB, Andersson G, et al. Evaluating Digital Health Interventions: Key Questions and Approaches. *American Journal of Preventive Medicine* 2016; 51: 843–851.
- [7] Priebe S, Golden E, Kingdon D, et al. Effective patient-clinician interaction to improve treatment outcomes for patients with psychosis: a mixed methods design (DIALOG+). *NIHR Journals Libr Heal Technol Assess Progr Grants Appl Res* 2017; 5: 6.
- [8] Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: Template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014; 348.
- [9] Priebe S, Kelley L, Omer S, et al. The Effectiveness of a Patient-Centred Assessment with a Solution-Focused Approach (DIALOG+) for Patients with Psychosis: A Pragmatic Cluster-Randomised Controlled Trial in Community Care. *Psychother Psychosom* 2015; 84: 304–313.
- [10] Fichtenbauer I, Priebe S, Schrank B. The German Version of DIALOG+ for Patients with Psychosis-A Pilot Study. *Psychiatr Prax* 2019; 46: 376–380.
- [11] Omer S, Golden E, Priebe S. Exploring the mechanisms of a patient-centred assessment with a solution focused approach (DIALOG+) in the community treatment of patients with psychosis: a process evaluation within a cluster-randomised controlled trial. *PLoS One* 2016; 11: 2.

- [12] DIALOG - Healthy London Partnership, <https://www.healthylondon.org/resource/dialog/> (accessed 27 May 2020).
- [13] ELFT - Services - Dialog, <https://www.elft.nhs.uk/service/347/Dialog> (accessed 27 May 2020).
- [14] Dialog+ - Home, <https://dialog.elft.nhs.uk/Home> (accessed 27 May 2020).
- [15] NIHR Global Health Research Group - NIHR Global Health Research Group, <https://www.qmul.ac.uk/nhr-ghrg/> (accessed 27 May 2020).
- [16] Jovanovic N, Francis J, Maric NP, et al. Implementing a psychosocial intervention DIALOG+ for patients with psychotic disorders in low and middle income countries in South Eastern Europe: protocol for a hybrid effectiveness-implementation cluster randomized clinical trial (IMPULSE). *Glob Psychiatry* 2019; 3: 83–96.
- [17] Brooke-Sumner C, Petersen I, Asher L, et al. Systematic review of feasibility and acceptability of psychosocial interventions for schizophrenia in low and middle income countries. *BMC Psychiatry* 2015; 15: 19.
- [18] Devlin AM, McGee-Lennon M, O'Donnell CA, et al. Delivering digital health and well-being at scale: lessons learned during the implementation of the dallas program in the United Kingdom. *J Am Med Informatics Assoc* 2015; 23: 48–59.
- [19] Wisdom JP, Chor KHB, Hoagwood KE, et al. Innovation Adoption: A review of theories and constructs. *Adm Policy Ment Heal* 2014; 41: 480–502.
- [20] O'Brien B, Harris I, Beckman T, et al. Standards for reporting qualitative research: a synthesis of recommendations. *Acad Med* 2014; 89: 1245–1251.
- [21] Implementation of an effective and cost-effective intervention for patients with psychotic disorders in low and middle income countries in South Eastern Europe | IMPULSE Project | H2020 | CORDIS | European Commission, <https://cordis.europa.eu/project/id/779334> (accessed 27 May 2020).
- [22] Ritchie J, Spencer L. *Qualitative data analysis for applied policy research*. Routledge, London and New York, 1994.
- [23] Parkinson S, Eatough V, Holmes J, et al. Framework analysis: a worked example of a study exploring young people's experiences of depression. *Qual Res Psychol* 2016; 13: 109–129.
- [24] Aref-Adib G, McCloud T, Ross J, et al. Factors affecting implementation of digital health interventions for people with psychosis or bipolar disorder, and their family and friends: a systematic review. *The Lancet Psychiatry* 2019; 6: 257–266.

- [25] Singla DR, Raviola G, Patel V. Scaling up psychological treatments for common mental disorders: a call to action. *World Psychiatry* 2018; 17: 226–227.
- [26] Ben-Zeev D, Atkins DC. Bringing digital mental health to where it is needed most. *Nat Hum Behav* 2017; 1: 849–851.
- [27] Berwick D. Disseminating innovations in health care. *J Am Med Assoc* 2003; 289: 1969–1975.
- [28] Greenhalgh T, Robert G, MacFarlane F, et al. Diffusion of Innovations in Service Organizations: Systematic Review and Recommendations. *Milbank Q* 2004; 82: 581–629.
- [29] Rogers EM. *Diffusion of innovations*. Free Press, 2003.
- [30] Ben-Zeev D, Davis KE, Kaiser S, et al. Mobile technologies among people with serious mental illness: Opportunities for future services. *Adm Policy Ment Heal Ment Heal Serv Res* 2013; 40: 340–343.
- [31] Heron KE, Smyth JM. Ecological momentary interventions: Incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010; 15: 1–39.
- [32] O'Connor S, Mair FS, McGee-Lennon M, et al. Engaging in large-scale digital health technologies and services. What factors hinder recruitment? *Stud Health Technol Inform* 2015; 210: 306–10.
- [33] Grol R, Wensing M, Eccles M, et al. *Improving patient care: the implementation of change in health care*. John Wiley & Sons, 2013.
- [34] Haak M, Himmelsbach I, Granbom M, et al. Cross-national and multi-language qualitative research: challenges and recommendations. *Br J Occup Ther* 2013; 76: 333–336.

## Tables

**Table 1.** Number of focus groups (FGs) and study participants per country.

|                               |              | Clinicians | Policymakers         | Patients | Carers   | Total number (%)      |
|-------------------------------|--------------|------------|----------------------|----------|----------|-----------------------|
| <b>Bosnia and Herzegovina</b> | FGs          | 2          | 2                    | 2        | 2        | 8                     |
|                               | Participants | 10         | 5                    | 10       | 10       | 35 (20%)              |
| <b>Kosovo<sup>a</sup></b>     | FGs          | 1          | 1                    | 2        | 1        | 5                     |
|                               | Participants | 8          | 6                    | 16       | 8        | 38 (22%)              |
| <b>North Macedonia</b>        | FGs          | 2          | 1                    | 2        | 1        | 6                     |
|                               | Participants | 13         | 6                    | 15       | 6        | 40 (23%)              |
| <b>Montenegro</b>             | FGs          | 2          | 1                    | 2        | 2        | 7                     |
|                               | Participants | 10         | 7                    | 12       | 11       | 40 (23%)              |
| <b>Serbia</b>                 | FGs          | 1          | 2 FGs<br>1 interview | 1        | 1        | 5 FGs<br>1 interview  |
|                               | Participants | 6          | 4                    | 6        | 5        | 21 (12%)              |
| <b>TOTAL</b>                  | FGs          | 8          | 7 FGs<br>1 interview | 9        | 7        | 31 FGs<br>1 interview |
|                               | Participants | 47 (27%)   | 28 (16%)             | 59 (34%) | 40 (23%) | 174                   |

**Table 2.** Summary of themes and subthemes.

| Theme   | Subthemes   |
|---|---|
| <b>Intervention characteristics</b>                               | <ul style="list-style-type: none"> <li data-bbox="760 159 1550 222">Clarity and easiness of use</li> <li data-bbox="760 222 1550 285">Involving patients' psychosocial needs</li> <li data-bbox="760 285 1550 348">Structuring routine clinician-patient meetings</li> <li data-bbox="760 348 1550 411">Ability to track patients' progress</li> </ul>  |
| <b>Attitudes and perceived preparedness of potential adopters</b> | <ul style="list-style-type: none"> <li data-bbox="760 422 1550 516">Implementation resources (staff, funding, time, space)</li> <li data-bbox="760 516 1550 611">Consistency with existing norms, policies and procedures</li> <li data-bbox="760 611 1550 684">Motivation to implement across potential adopters</li> <li data-bbox="760 684 1550 747">Potential adopters' resistance to innovation</li> <li data-bbox="760 747 1550 842">Level of existent skills and relevant experience by potential adopters</li> <li data-bbox="760 842 1550 905">Perceived need for the intervention</li> <li data-bbox="760 905 1550 968">Institutional support</li> <li data-bbox="760 968 1550 1041">Influence of poor mental health of patients</li> </ul> |
| <b>Carers' involvement</b>  | <ul style="list-style-type: none"> <li data-bbox="760 1041 1550 1104">Role of carers in intervention implementation</li> <li data-bbox="760 1104 1550 1167">Extent to carers involvement</li> </ul>   |
| <b>Perceived potential patient and organisational benefits</b>    | <ul style="list-style-type: none"> <li data-bbox="760 1167 1550 1230">Patient empowerment</li> <li data-bbox="760 1230 1550 1293">Improved therapeutic relationship</li> <li data-bbox="760 1293 1550 1356">Reinforcement of psychosocial care approach</li> <li data-bbox="760 1356 1550 1419">Standardization of care across services</li> <li data-bbox="760 1419 1550 1482">Reinforcing patient-centred care</li> <li data-bbox="760 1482 1550 1566">Improving structure and follow-up of care</li> </ul>   |
| <b>Frequency of intervention delivery</b>                         | <ul style="list-style-type: none"> <li data-bbox="760 1566 1550 1629">Monthly delivery</li> <li data-bbox="760 1629 1550 1692">More frequent delivery than monthly</li> <li data-bbox="760 1692 1550 1755">Less frequent delivery than monthly</li> </ul>   |
| <b>Suggested changes to the intervention</b>                      | <ul style="list-style-type: none"> <li data-bbox="760 1755 1550 1818">Changes to the DIALOG scale</li> <li data-bbox="760 1818 1550 1881">Changes to interpretation of life areas</li> <li data-bbox="760 1881 1550 1957">Changes to mode of intervention delivery</li> </ul>   |

## Figures




Figure 1

Major themes related to the implementation of DIALOG+ across participating.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile4.docx](#)
- [AdditionalFile3.docx](#)
- [AdditionalFile2.docx](#)
- [AdditionalFile1.docx](#)