

Genome Analyses Revealed Genetic Admixture and Selection Signatures in *Bos indicus*

S P Dixit (✉ dixitsp@gmail.com)

ICAR - National Bureau of Animal Genetic Resources

A K Bhatia

ICAR - National Bureau of Animal Genetic Resources

Indrajit Ganguly

ICAR - National Bureau of Animal Genetic Resources

Sanjeev Singh

ICAR - National Bureau of Animal Genetic Resources

Soumya Dash

ICAR - National Bureau of Animal Genetic Resources

Anurodh Sharma

ICAR - National Bureau of Animal Genetic Resources

N. Anand Kumar

ICAR - National Bureau of Animal Genetic Resources

A K Dang

ICAR - National Dairy Research Institute Karnal

S Jayakumar

ICAR - National Bureau of Animal Genetic Resources

Research Article

Keywords: India, Native cattle breeds, SNP, BovineHD BeadChip, Diversity, Selection signatures

Posted Date: May 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-484105/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The genomic diversity and relationship among seven diverse cattle breeds viz. Sahiwal, Tharparkar, Gir, Vechur, Ongole, Kangayam and Hariana were investigated in 132 random samples based on high density SNP array comprising > 777K SNPs. A total of 1993 SNPs (0.25% of the total) having greater power ($F_{st} \geq 0.20$) to differentiate these cattle populations were identified, and utilized to partition genome of each animal into a predefined number of clusters. The structure of these cattle indicated shared ancestry of dairy breeds viz. Gir, Tharparkar and Sahiwal. Most of animals (> 76%) of the different populations under study except Vechur clustered into their own group of animals called breed. Vechur population contained highest rate of admixture, consistent with its crossing with other breeds. Ongole, Kangayam and Hariana shared comparatively less of their genome ($\leq 15\%$) with other breeds. The study indicated that all seven breeds evolved from their independent ancestry but there was intermixing of these breeds in the recent past. The selection signatures identified between draft (Kangayam) and dairy breeds included several genes like FAM19A2, RAB31P, BEST3, DGKA, AHCY, PIGU and PFKP which are involved in immune response, metabolic pathway, transportation of glucose and sugars, signaling pathways, cellular processes, cell division and glycolysis regulation, respectively. Moreover, these genomic regions also harbour QTLs affecting milk performance traits. The signatures were also identified even between the dairy breeds. There were also significant differences in number of QTLs affecting production (body weight, growth rate etc.) and morphological traits (height etc.) of short stature, Vechur, compared to large sized cattle. The HMGA2 gene was present in the selection signature on chromosome 5, which may explain the differences in stature of these cattle.

Introduction

India has 50 registered cattle breeds adapted to different agro-climatic conditions, breeding and management practices (<https://nbagr.icar.gov.in/en/registered-cattle/>) and are likely to differ for a number of traits. These breeds thrive in humid subtropical, semi-arid & arid and tropical wet/dry climatic regions, catering to a variety of specialized functions such as dairy, draft and dual (Dairy & draft) use. Therefore, these cattle would have gene content unique to such roles and adaptations in their genome¹. Hence, they can serve as a great reservoir of genetic pool for identifying genes under selection for different traits that have evolved, as well as for determining genetic diversity.

In India, cattle account for approximately 58% of the country's 303.76 million bovines, but share only around 48% of the total milk produced, due to their low dairy productivity and a higher proportion of draught and dual-purpose breeds than dairy breeds². With the exception of Africa, where output remained stable, global milk production increased 2.0% from 2019 to nearly 906 million tonnes in 2020, owing to rises in all geographical regions. Milk production in India reached to 195 million tonnes in 2020, up 2.0% from the previous year³ and accounting for around 21% of global milk production. In addition to their dairy usefulness, draft/dual cattle breeds in India are also used for agricultural and transport operations. Considering their importance in Indian agriculture, several government schemes, such as National

Programme for Bovine Breeding and Dairy Development, National Dairy Plan and Dairy Entrepreneurship Development scheme have been launched to boost their future productivity.

The use of single nucleotide polymorphism (SNP) array and next-generation sequencing technologies in genomics and population genetics has resulted significant progress in deciphering the genetic structures⁴, genome diversity and selection footprints in cattle^{5-11,1}. Understanding the genomic diversity of native cattle breeds aids in improving their productivity, fitness, fertility and even the behaviour. Additionally, using high-density SNP arrays can increase detection limits of positive selection and lower the false discovery rate¹²⁻¹⁵. Only a few reports using Illumina high-density SNP arrays on Indian native cattle breeds are available^{1,16,17}.

To better understand the genetic mechanisms underlying the local adaptation and functional characteristics of Indian zebu cattle, we investigated three dairy breeds (Sahiwal, Gir, Tharparkar) from sub-tropical and hot arid regions, two dual breeds (Hariana, Ongole) from sub-tropical and hot humid regions, and one draught breed (Kangayam) from the country's hot humid region. We have also included Vechur, a short statured cattle breed native to hot humid climates. This array of cattle breeds, each adapted to a different ecological niche and functional attribute, will help to uncover genomic diversity and natural and artificial selection footprints over centuries. Previously, we identified and characterized genome wide runs of homozygosity (ROH) signatures in these cattle breeds using Illumina BovineHD BeadChip¹⁷. The objectives of the present study were to: 1) assess genomic diversity and effective population size; 2) analyse admixture and structuring; 3) identify diversified selection signatures among breeds using SNP data.

Results And Discussion

Genomic diversity within and among the breeds

Twenty of the 132 animals were excluded due to low genotyping ($MIND > 0.1$), and the average genotyping rate for the remaining 112 animals was 0.99. The final data on 112 cattle samples belonging to Sahiwal (13), Tharparkar (17), Gir (15), Ongole (17), Hariana (18), Kangayam (16) and Vechur (16) breeds were achieved after quality control measures (Table 1). A sample size of ≥ 13 per breed was adequate for the diversity analyses, which was in consonance with other studies^{18,19}. Out of 735293 autosomal SNPs genotyped in these cattle, 60% (438,176) were in Hardy Weinberg equilibrium and revealed higher degree of polymorphism ($MAF > 0.05$). It was also stated that a sample size of 4-6²⁰ and polymorphic SNP filtration^{19,21,17} could mitigate the effect of ascertainment bias if the number of markers is sufficiently large enough, as in the current study. The SNPs in strong LD ($R = 0.5$) were also filtered out to minimize the bias in estimating the genomic diversity. We were left with 165,021 informative SNPs after pruning for further genome diversity analyses. Minor allele frequency varied between breeds, ranging from 0.23 (Kangayam) to 0.26 (Vechur), with an average observed heterozygosity of 0.35 in all samples examined (Table 1). The similar estimates of MAF were also reported in Nellore, Holstein, Iranian, Ethiopian and South African cattle²²⁻²⁶, which varied from 0.21 to

0.25 in *Bos indicus* and *Bos Taurus* cattle breeds. However, higher/lower estimates were also reported in other cattle breeds^{27–29}. The reported estimates of observed heterozygosity in several other zebu cattle^{25,30,31} were in agreement with the present estimates. However, higher heterozygosity was reported in Hanwoo, Rwanda and other Taurine cattle breeds^{25,30,29}. The lower genetic variability estimates in *Bos indicus* relative to *Bos taurus* were consistent with earlier studies^{32,33,29}. The lower genetic diversity in *Bos Indicus* may be due to ascertainment bias of mainly taurine derived SNPs in SNP chip. Out of Bovine HD and 54K SNP chips of Illumina, only 40–50% SNPs were found to be informative for genetic diversity of Zebu cattle breeds of India¹⁶ but, it was 90% in *Bos taurus* after quality control¹⁴.

Table 1
Number of animals, mean of expected and observed heterozygosity (He, Ho), minor allele frequency (MAF) and coefficient of inbreeding (F_{IS}).

Breed	n	He	Ho	MAF	F_{IS} ($p > 0.40$)
Sahiwal	13	0.34	0.35	0.25	-0.009
Tharparkar	17	0.33	0.34	0.24	-0.003
Gir	15	0.32	0.33	0.24	0.010
Ongole	17	0.33	0.34	0.24	0.006
Haryana	18	0.33	0.34	0.25	-0.001
Kangayam	16	0.30	0.33	0.23	-0.062
Vechur	16	0.34	0.35	0.26	-0.009
overall	112	0.36	0.35	0.27	-0.009

The genome diversity of seven cattle breeds viz. Tharparkar, Sahiwal, Gir, Vechur, Ongole, Kangayam and Haryana, was determined first using all filtered SNPs (4 38,176) and then using the 1993 most informative SNPs having F_{ST} value ≥ 0.20 . The genetic differentiation (F_{ST}) of the breeds based on all SNPs was just 0.05 and based on 1993 most differentiating loci with an F_{ST} value of ≥ 0.20 was 0.23. Hence, out of 777K, 1993 loci (0.25% of the total) were identified as having greater power to distinguish these cattle populations, and these loci were further assessed for diversity and structure analysis. The overall estimates of observed heterozygosity (Ho), gene diversity within population (Hs), gene diversity in total population (Ht), gene diversity among breeds (Dst), and Dest, a measure of population differentiation³⁴, across these loci were 0.28 0.27, 0.35, 0.08 and 0.13, respectively.

Manhattan graph (Fig. 1) depicts the distribution of F_{ST} values across the chromosomes. The spread of F_{ST} showed that there are few loci (16 markers) having the higher degree of genetic differentiation ($F_{ST} \geq 0.40$). The inbreeding coefficient (identical by state) was zero, suggesting that the cattle from which the

samples were taken were randomly mated. The genetic differentiation power of those informative SNP loci ($n = 1,993$) in cattle breeds under study ranged from 0.20 to 0.51, with an average value of 0.23, indicating that these loci account for 23% of the genetic variation between the breeds. Table 2 and Fig. 2 showed the pair-wise estimates of F-statistics (F_{ST}) and Nei's genetic distance³⁵. Both estimates revealed higher genetic differentiation between Kangayam and rest of the cattle breeds (F_{ST} : 0.08 to 0.10), followed by Ongole and rest of the breeds (F_{ST} : 0.05 to 0.07). Dairy cattle breeds viz., Tharparkar, Sahiwal and Gir showed the least genetic differentiation (F_{ST} : 0.04 to 0.06). A moderate genetic differentiation³⁶ where F_{ST} ranged from 0.05 to 0.15 was also recorded in *Bos taurus* dairy and beef cattle with an F_{ST} value of 0.08 across all SNPs¹⁴, as well as in African cattle breeds (F_{ST} : 0.04 to 0.08). Many SNPs with higher genetic differentiation power ($F_{ST} > 0.5$) were also reported in *Bos taurus* dairy and beef cattle¹⁴. The degree of genetic differentiation among Indian Zebu cattle was comparable to that of African Zebu cattle³⁰, but lower than that of *Bos taurus* and *Bos indicus*^{30,33}, which was anticipated given their historical divergence.

Table 2
The pair-wise Nei's F_{ST} (above diagonal) and genetic distance (below diagonal) among the cattle breeds

Breed	TP	SW	GR	VC	OG	KG	HR
TP	-	0.044	0.059	0.066	0.064	0.094	0.049
SW	0.028	-	0.051	0.051	0.052	0.081	0.037
GR	0.032	0.032	-	0.072	0.072	0.101	0.056
VC	0.034	0.031	0.037	-	0.069	0.096	0.061
OG	0.033	0.031	0.037	0.035	-	0.092	0.058
KG	0.044	0.042	0.048	0.046	0.043	-	0.088
HR	0.027	0.026	0.031	0.032	0.031	0.043	-
SW-Sahiwal, TP-Tharparkar, GR-Gir, VC-Vechur, OG-Ongole, KG-Kangayam, HR-Haryana							

Effective Population Size

Ancestral and recent effective population sizes (N_e) for seven Indian cattle breeds are presented in Fig. 3. Estimated N_e showed a downward trend in recent generations across the populations. The most rapidly declining recent N_e was found in the KG and GR, whereas HR and TP showed a slowly declining trend. The estimated N_e at 13 generations ago in seven Indian native cattle breeds ranged from 48 to 74 (Supplementary Table S1).

Over the last 999 years, the N_e has displayed a decreasing pattern across Indian breeds, with a steeper slope since about 200 generations ago. Lower N_e had been found in KG and SW with estimated values of 48 and 51, respectively, 13 generations ago (N_{e13}), due to the intensive selection pressure or artificial insemination used for developing these breeds. In Kangayam cattle, we have observed maximum autozygosity attributable to both recent and ancient inbreeding¹⁷.

Recently, a Bovine HD-SNPs array based screening of Chinese native cattle populations revealed a similar trend, with N_{e13} values ranging from 85 to 132³⁷. The genotyping of Italian local beef breeds (Calvana-CAL, Mucca Pisana-MUP, Pontremolese-PON) and Italian Limousin (LIM) using the GeneSeek GGP-LDv4 33k SNP chip containing 30,111 SNPs showed an average estimated historical effective population size (N_{e13}) of 45–310 (CAL-79, MUP-65, PON-45 and LIM-310)³⁸. The application of LD-based N_e estimation in developing countries for local breeds without pedigree information could offer new perspectives for the assessment of the actual gene pool available and the respective decision-making in conservation and management.

Genetic Structuring And Classification Of The Cattle Breeds

The principal component analysis (PCA) based on genomic relationship (IBS-Identity by state) matrix using 165,021 LD pruned autosomal SNPs was undertaken to assess breed composition of the animals. The first and second principal components accounted for 29% and 12 % of the total variation, respectively. Figure 4A, B presented the first, second and third principal components where Vechur, Kangayam and Ongole clustered separately from other breeds under investigation, and rest of the breeds (Gir, Sahiwal, Tharparkar and Hariana) grouped together. The structure and PCA both revealed that dairy breeds clustered apart from the dual (Ongole) except Hariana, and draft breeds. PCA based on SNP data clearly separated the breeds by their utility and size, which was in consonance with analysis of molecular variation (Table 3). Kangayam, a draft breed and Vechur, a small statured breed were quite distinct from the dairy and dual breeds and was in agreement with structuring of these cattle based on runs of homozygosity (ROH)¹⁷. Moreover, the structure and genomic relationship among these cattle was also studied using a small number of highly differentiated loci (180, 50, 10 markers) and it was interesting to note that even 10 markers also revealed the same level of differentiation as with large sized battery of markers. Therefore, the structure and relationship between the cattle breeds were not significantly influenced by ascertainment bias, which was consistent with a previous study in sheep³⁹ and cattle¹⁷.

Table 3
Hierarchical analysis of variance based on pairwise differences*

Source of variance Variance	d.f.	Sum of squares	variation components	Percentage of variation	p value
Among groups	3	314971.297	486.39137	1.64	0.029
Among populations within groups	3	234688.652	1613.69233	5.44	0.00
Among individuals within populations	105	2870271.011	-243.80843	-0.82	0.645
Within individuals	112	3116235.500	27823.53125	93.75	0.001
*Significance tests based on 1023 permutations					

The Hierarchical F-statistics computed across different groups indicated significant differences in milk production categories (High: dairy breeds; Moderate: Dual breeds; Low: Draft and small statured breeds) and hence differentiated dairy (Sahiwal, Gir and Tharparkar), dual (Ongole, Hariana), and draft (Kangayam) and small sized cattle (Vechur) (Table 4). PCA based on genotype displayed similar categorization of these cattle¹⁷. The analysis of molecular variance also revealed significant ($p < 0.05$) differences in the proportion of variation (1.64%) due to their functional characteristics viz. dairy, dual, draft and small size. However, there were significant differences (5.44%) between the breeds having a particular characteristic like dairy and dual (Table 3). ROH regions observed in these cattle breeds were also able to differentiate dairy and draft breeds as well as small stature cattle¹⁷.

Table 4

Hierarchical F-statistics computed over geographical distribution, body size and level of milk production of the 7 cattle breeds

Hierarchical Level	Geographical Region	Body Size	Level of dairy performance	Individual
Geographical Region	0.00	0.19	0.44	0.43
Body Size	0.00	0.00	0.30**	0.29
Level of dairy performance (high, medium and low)	0.00	0.00	0.00	-0.02
**p = 0.001				

Admixture Analyses Of The Cattle Breeds

The population structure of these cattle was studied using admixture model based clustering implemented in Structure⁴⁰ to partition genome of each animal into a predefined number of clusters. The

breeds were clearly grouped into draft and dairy/dual breeds (Fig. 5) for pre-defined $K = 2$, indicating shared ancestry of dairy and dual purpose breeds (Gir, Tharparkar, Sahiwal, Vechur, Ongole, Haryana) very similar to sharing of paternal lineages⁴¹. More than 90% of genome of dairy /dual cattle except Ongole (86%) clustered together and 90% of the genome of draft cattle Kangayam grouped into separate cluster. At $K = 3$, Sahiwal, Tharparkar, Gir, Vechur and Haryana (> 72%) clustered together. Kangayam and Ongole clustered into their own group (89%). All major dairy breeds (Tharparkar, Gir and Sahiwal) share > 80% of their genome. However, at $K = 4$, Vechur, was clustered apart from other breeds and at $K = 5$, Gir also clustered separately. At $K = 7$, most of animals (> 76%) of the different populations under study except Vechur clustered into their own group of animals called breed (Fig. 5). Vechur population displayed highest rate of admixture (Table 5), consistent with its crossing with other breeds and the same was also revealed based on its ROH analysis¹⁷. Ongole, Kangayam and Haryana shared comparatively less of their genome ($\leq 15\%$) with other breeds. The Fig. 6 revealed the increasing mean value of log likelihood across the inferred clusters and indicated the all seven breeds evolved from their independent ancestry. However, the structure at different values of K indicated intermixing of these breeds during the recent past. The admixture analysis ($k = 2$) revealed the structure of these cattle except Vechur similar to that unraveled by principal component analysis.

Table 5
Proportion of membership of each pre-defined population in each of the 7 clusters

Population	Given Inferred Clusters							Population size
	1	2	3	4	5	6	7	
	OG	HR	KG	GR	TH	VC	SW	
TH	0.013	0.107	0.014	0.034	0.766	0.016	0.049	17
SW	0.014	0.095	0.014	0.035	0.047	0.026	0.768	13
GR	0.014	0.073	0.014	0.764	0.064	0.011	0.06	15
VC	0.033	0.07	0.052	0.026	0.028	0.733	0.058	16
OG	0.854	0.082	0.01	0.002	0.015	0.011	0.027	17
KG	0.026	0.024	0.889	0.002	0.027	0.01	0.022	16
HR	0.014	0.89	0.005	0.02	0.026	0.01	0.035	18
SW-Sahiwal, TP-Tharparkar, GR-Gir, VC-Vechur, OG-Ongole, KG-Kangayam, HR-Haryana								

Selection Signatures Among The Breeds

The genomic regions showing most differentiation among breed pairs based on F_{ST} value were identified. The most differentiated breed pairs based on their performance were chosen for the identification of selection signatures. Therefore, Kangayam, a draft breed, was used as control breed for the analysis of

dairy breeds while, Vechur, a small sized breed for the rest of large sized breed to search for signatures that may be associated with stature. Genome wide significance level was set to 0.001 to represent a selection signatures. Out of the significantly differentiated genomic regions, the top most differentiating signatures with $F_{ST} \geq 0.50$ among different breed pairs is presented in Table 6.

Table 6

The selection signatures ($F_{ST} \geq 0.50$) value among the different cattle breeds and their annotation with Bovine QTL database

Breed** pair	Top F_{ST} value	Marker	Chromosome	QTLs*
SW/TP	0.67	BovineHD0400009897	BTA4	MY,MC,R,M/C,H,P,M
SW/GR	0.57	BovineHD0500015340	BTA5	MC,R,M/C,H
SW/VC	0.69	BovineHD2900011510	BTA29	ALL TRAITS
SW/HR	0.53	BovineHD0300002536	BTA3	MC,P
SW/KG	0.80	BovineHD2400003366	BTA24	R,M/C,H,M
SW/OG	0.64	BovineHD0800001684	BTA8	R,M/C,P
TP/GR	0.59	BovineHD0300009588	BTA3	ALL TRAITS EXCEPT M
TP/VC	0.73	BovineHD1900008163	BTA19	ALL TRAITS EXCEPT MY
TP/HR	0.50	BovineHD2800011107	BTA28	M/C
TP/KG	0.81	BovineHD1300018382	BTA13	MC,R,M
TP/OG	0.62	BovineHD0200032623	BTA2	ALL TRAITS
GR/VC	0.81	BovineHD1900018844	BTA19	ALL EXCEPT MY
GR/HR	0.60	BovineHD1000011926	BTA10	ALL EXCEPT MY,M
GR/KG	0.81	BovineHD0500019637	BTA5	ALL TRAITS
GR/OG	0.65	BovineHD0500012675	BTA5	ALL TRAITS
VC/HR	0.80	BovineHD1900008163	BTA19	ALL EXCEPT MY
VC/KG	0.76	BovineHD0200022436	BTA2	R,M/C,H,P
VC/OG	0.61	BovineHD1200026864	BTA12	ALL EXCEPT MY
HR/KG	0.80	BovineHD1300013278	BTA13	ALL TRAITS
HR/OG	0.60	BovineHD1700005865	BTA17	ALL EXCEPT MY,H
KG/OG	0.78	BovineHD2100009715	BTA21	R,M/C,H,P
*MY-Milk Yield, MC-Milk Composition, R-Reproduction, M/C-Meat & Carcass				
H-Health, P-Production, M-Morphology				
**SW-Sahiwal, TP-Tharparkar, GR-Gir, VC-Vechur, OG-Ongole, KG-Kangayam, HR-Hariana				

The annotation of 5 top most genomic regions among these cattle to Bovine QTL database revealed that these regions harbored QTL affecting milk yield, milk composition, reproduction, production, Health status and morphological traits (Supplementary Table S2). The top divergent regions among most divergent breed pairs may be putative selection signature for diversifying traits between the breeds. For example, the marker BovineHD0500014902 lies in the FAM19A2 gene and is the most differentiating region between Kangayam and Sahiwal. This gene is thought to produce brain-specific chemokines or neurokinins, which function as immune and nervous cell regulators (<https://www.genecards.org/cgi-bin/carddisp.pl?gene=TAF2>), and thus may affect Kangayam and Sahiwal's health in different ways. Kangayam is a hardy draft breed compared to Sahiwal. There were significant differences in number of QTLs affecting milk yield and production traits when Kangayam paired with any of the dairy breed. For example, between Hariana and Kangayam, the marker BovineHD0500012581 lies in RAB31 gene which is involved in metabolic pathway (<https://www.genecards.org/cgi-bin/carddisp.pl?gene=RAB31>) and has also been annotated with QTLs affecting milk performance traits (Cattle QTL data base). Hence, may be responsible for variations in these traits.

The other selection signatures identified between Kangayam and dairy breeds included several genes like Bestrophin 3 (BEST3) [<https://www.genecards.org/cgi-bin/carddisp.pl?gene=BEST3&keywords=BEST3>], Diacylglycerol Kinase Alpha (DGKA) [<https://www.genecards.org/cgi-bin/carddisp.pl?gene=DGKA&keywords=DGKA>], Adenosylhomocysteinase (AHCY) [<https://www.genecards.org/cgi-bin/carddisp.pl?gene=AHCY&keywords=AHCY>], Phosphatidylinositol Glycan Anchor Biosynthesis Class U (PIGU) [<https://www.genecards.org/cgi-bin/carddisp.pl?gene=PIGU&keywords=PIGU>] and Phosphofructokinase, Platelet (PFKP) [<https://www.genecards.org/cgi-bin/carddisp.pl?gene=PFKP&keywords=Phosphofructokinase>] which are involved in transportation of glucose and sugars, signaling pathways, cellular processes, cell division and glycolysis regulation, respectively. Moreover, these genomic regions also harbour QTLs affecting milk performance traits (Cattle QTL data base). The signatures were also identified even between the dairy breeds. For example, the genomic region, having BovineHD1300006213 locus, revealing high differentiation between Tharparkar and Gir ($F_{ST}=0.585$) harbours QTLs affecting all the traits under study except milk yield (Cattle QTL data). There were significant differences in number of QTLs affecting production (body weight, growth rate etc.) and morphological traits (height etc.) of Vechur when paired with large sized cattle (Supplementary Table S3). The genomic region having BovineHD0500013882 locus starting from 48008400 to 48069099 on chromosome 5 is having HMGA2 gene which is responsible for explaining the variation in stature both in cattle⁴² and human beings⁴³.

Implications Of Genomic Analyses For Breed Management And Conservation

The genomic analyses detailed herein revealed that all the high yielders dairy breeds viz. Sahiwal, Tharparkar and Gir shared common ancestry but are admixed due to gene flow among them through crossing, migration and /or grading up with other breeds. Hariana, traditionally a dual purpose breed but was more closely associated with dairy breeds. Ongole, Kangayam and Vechure were quite distinct from rest of the breeds under study. The genomic analyses represented the unique gene pool of these cattle

genetic resources, befitting their breed's status. The Haryana cattle was under intensive selection for high milk yield in the past and was being used as improver breed for many other breeds/populations for augmenting milk production in the country, and same was also revealed here through genomic characterization.

In conclusion, BovineHD BeadChip genotyping of Indian cattle breeds is promising, as it can not only structure these cattle but also detect distinct signatures of selection. As a result, it could be used for a wider range of studies, such as genome wide association studies and genomic selection involving larger populations of the breeds. Multi-breed genomic selection may be feasible in dairy breeds due to their shared genome. This study reveals a trend toward shrinking effective population sizes in native Indian cattle breeds, indicating that a long-term breeding strategy is needed to prevent further reductions in N_e , as well as genetic improvement and potential conservation. In future, whole genome sequencing information on these breeds may be useful for pinpointing the genomic regions linked to polygenic productivity, health, fertility, and behavioural traits that evolved under Indian ecological and farming systems.

Methods

Animal Resources, SNP Genotyping and Quality Control. A total of 132 samples of Sahiwal (SW, $n = 19$), Tharparkar (TR, $n = 17$), Gir (GR, $n = 16$), Ongole (OG, $n = 24$), Haryana (HR, $n = 18$), Kangayam (KG, $n = 18$) and Vechur (VC, $n = 20$) breeds of cattle were incorporated. Random blood samples were obtained from various farms across the country in accordance with the regulations and guidelines of the Institutional Animal Ethics Committee (IAEC), National Bureau of Animal Genetics Resources (ICAR-NBAGR), Karnal. Genomic DNA was extracted from the whole blood using HiPurATM SPP Blood DNA isolation kit. The quality of the genomic DNA was tested using an agarose gel electrophoresis, and the quantity of DNA was measured using a Nanodrop Spectrophotometer (Nanodrop ND-1000). The DNA samples were genotyped at Sandor Lifesciences Pvt. Ltd. in Hyderabad, India, using an Illumina BovineHD BeadChip with 777962 SNPs and following the manufacturer's standard procedures. The data files including MAP and PED files were retrieved using Genome Studio. The analysis of the SNP data revealed the call rate ranged from 95 to 99%. The number of SNPs scored in a given sample / the number of SNPs available on chip * 100 was used to calculate the call rate of SNPs. The call rate indicated that the Illumina BovineHD BeadChip is useful in scoring SNP/genotypes in the Indian cattle population and could be used to assess breed signatures and diversity.

The quality control procedure was carried out by using PLINK^{44,45}. The unmapped SNPs and SNPs present on X, Y chromosomes, and on mitochondrial DNA were removed and only the SNPs located on autosomes were considered for analysis. SNPs with call rate (CR) $\leq 95\%$, minor allele frequency (MAF) ≤ 0.05 , and HWE ($P \leq 0.001$) were excluded. Samples that had more than 10% missing genotypes were also excluded. The quality of SNPs genotyped were assessed based on Gene Call Score (< 0.2) and Gene Train Score (< 0.55) using Genome Studio. The gene train score, in general, was > 0.55 for the SNPs genotyped in these samples and hence good quality SNPs were obtained.

Genomic diversity analyses among the breeds. For handling and managing as well as analyzing the large size data on 777 K bovine Bead Chip, several in house computer scripts were written for making the suitable data formats for further downstream analyses using different genetic software including HierFstat in R and Structure. LD was measured for each breed as correlation between adjacent SNPs (r^2) which depends upon the frequencies of the alleles at the loci under consideration. The r^2 values were calculated using PLINK v 1.9^{44,45} keeping the window size limit of 500 kb between pair-wise SNPs. Further, autosomal SNPs were pruned out with an r^2 value of 0.5 using PLINK^{44,45}. The minor allele frequency, heterozygosity and inbreeding in different breeds were also estimated using PLINK. Hierarchical F-statistics were computed to access genomic differences in different groups using Hierfstat in R (<http://www.r-project.org>, <http://github.com/jgx65/hierfstat>): I) geographical distribution (North & South), II) body size (Large & Small), and III) Milk production (High, Medium & Low). The genomic differences among the breeds were also calculated in terms of F_{ST} value⁴⁶ as well as through analysis of molecular variance (AMOVA) using ARLEQUIN⁴⁷.

Past effective population size (N_e).

The historical and recent effective population size (N_e) was estimated using the SNeP v1.1 software as described earlier⁴⁸. It inferred N_e based on linkage disequilibrium (LD) against past t generations, where $t = 1/2c$ and c is the distance between SNPs in Morgans (100 Mb = 1 Morgan was assumed)⁴⁹. The estimation was performed on the SNP data with correction of sample size, phasing and recombination rate.

Structuring and admixture analyses of the breeds. The population structure was studied using admixture model based clustering implemented in Structure⁴⁰ to partition genome of each animal into a predefined number of clusters (K). The optimum number of K was determined based on mean value of log likelihood across the inferred clusters. The principal component analysis (PCA) based on genomic relationship (IBS-Identity by state) matrix using plink tools and R script was also undertaken to assess breed composition of the animals.

Detection of diversified selection signatures and the underlying genes. The diversifying selection signatures were identified among the breeds using F-statistics ($P < 0.0001$). F_{ST} value of ≥ 0.25 was considered as selection sweep between two contrasting groups (dairy/dual verses draft, small verses large stature). The highly significant windows between two groups were annotated for their gene content. NCBI map viewer of the bovine UMD3.1.1 (<https://www.ncbi.nlm.nih.gov/genome/gdv/>) was used to identify the genes underlying the selection sweeps. Genes underlying ± 1 MB of SNPs with F_{ST} value of ≥ 0.25 were identified.

The effect of the top 5 signatures on the underlying QTLs was assessed by exploring cattle QTL database (<https://www.animalgenome.org/QTLdb/cattle>). Test of two proportions was worked out to find the significant differences between the numbers of QTLs affecting the two contrasting groups (dairy versus Draft) for six different traits using XLSTAT.

Declarations

Ethics statement. Random blood samples were obtained from various farms across the country with written informed consent from the owner by qualified Veterinarian in accordance with the guidelines issued by the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA; <http://cpcsea.nic.in/WriteReadData/userfiles/file/Compendium%20of%20CPCSEA.pdf>) and approved by the Institutional Animal Ethics Committee (IAEC) of ICAR-National Bureau of Animal Genetics Resources (ICAR-NBAGR), Karnal.

Data availability

We have uploaded the data on ICAR-Krishi portal and is in public domain with the URL <http://krishi.icar.gov.in/jspui/handle/123456789/31167>.

Acknowledgments

The authors acknowledge the funding support received from the Department of Biotechnology (DBT), Govt. of India. Authors also gratefully acknowledge the assistance of Mr. Ramesh Rana (ICAR-NBAGR) for sample collection and processing. The authors wish to express thanks to Divya Jyoti Jagrati Sansthan, Noormahal, Jalandhar (Punjab), Rajasthan University of Veterinary and Animal Sciences, Bikaner (Rajasthan) and Dr. Jayadevan Narayanan, Kerala for providing necessary samples

Author Contributions

S.P.D conceived and designed the experiments. S.P.D., A.S, A.K and S.D. performed experiments. S.P.D., A.K.B., I.G., and S.S. analyzed data. S.P.D. wrote the main manuscript. I.G., S.S., A.K.D, and J.K assisted with the writing of the manuscript. All the authors read and approved the final manuscript.

Competing Interests: The authors declare no competing interests.

References

1. Bhardwaj, S. *et al.* Genome-wide diversity analysis for signatures of selection of *Bos indicus* adaptability under extreme agro-climatic conditions of temperate and tropical ecosystems. *Anim. Gene* **20**, 200115 , <https://doi.org/10.1016/j.angen.2021.200115> (2021).
2. *Department of Animal Husbandry and Dairy (DAHD), Government of India (GOI) Annual Report 2019/20.* [http://dahd.nic.in/sites/default/files/Annual Report 2019-20.pdf](http://dahd.nic.in/sites/default/files/Annual%20Report%202019-20.pdf).
3. *FAO. 2021. Dairy Market Review: Overview of global dairy market developments in 2020, April 2021.* Rome. <http://www.fao.org/3/cb4230en/cb4230en.pdf>.

4. Gibbs, R. A. *et al.* Genome-Wide Survey of SNP Variation Uncovers the Genetic Structure of Cattle Breeds. *Science (80-)*. **324**, 528–532 , <https://doi.org/10.1126/science.1167936> (2009).
5. Utsunomiya, Y. T. *et al.* Detecting Loci under Recent Positive Selection in Dairy and Beef Cattle by Combining Different Genome-Wide Scan Methods. *PLoS One* **8**, e64280 , <https://doi.org/10.1371/journal.pone.0064280> (2013).
6. Cañas-Álvarez, J. J. *et al.* Genetic diversity and divergence among Spanish beef cattle breeds assessed by a bovine high-density SNP chip1. *J. Anim. Sci.* **93**, 5164–5174 , <https://doi.org/10.2527/jas.2015-9271> (2015).
7. Stothard, P. *et al.* A large and diverse collection of bovine genome sequences from the Canadian Cattle Genome Project. *Gigascience* **4**, 49 , <https://doi.org/10.1186/s13742-015-0090-5> (2015).
8. Gao, Y. *et al.* Species composition and environmental adaptation of indigenous Chinese cattle. *Sci. Rep.* **7**, 16196 , <https://doi.org/10.1038/s41598-017-16438-7> (2017).
9. Kim, S. *et al.* Genetic diversity and divergence among Korean cattle breeds assessed using a BovineHD single-nucleotide polymorphism chip. *Asian-Australasian J. Anim. Sci.* **31**, 1691–1699 , <https://doi.org/10.5713/ajas.17.0419> (2018).
10. Chen, N. *et al.* Whole-genome resequencing reveals world-wide ancestry and adaptive introgression events of domesticated cattle in East Asia. *Nat. Commun.* **9**, 2337 , <https://doi.org/10.1038/s41467-018-04737-0> (2018).
11. Liu, Y. *et al.* Discovery of Genomic Characteristics and Selection Signatures in Southern Chinese Local Cattle. *Front. Genet.* **11**, , <https://doi.org/10.3389/fgene.2020.533052> (2020).
12. Pérez O'Brien, A. M. *et al.* Assessing signatures of selection through variation in linkage disequilibrium between taurine and indicine cattle. *Genet. Sel. Evol.* **46**, 19 , <https://doi.org/10.1186/1297-9686-46-19> (2014).
13. Xu, L. *et al.* Genomic Signatures Reveal New Evidences for Selection of Important Traits in Domestic Cattle. *Mol. Biol. Evol.* **32**, 711–725 , <https://doi.org/10.1093/molbev/msu333> (2015).
14. Zhao, F., McParland, S., Kearney, F., Du, L. & Berry, D. P. Detection of selection signatures in dairy and beef cattle using high-density genomic information. *Genet. Sel. Evol.* **47**, 49 , <https://doi.org/10.1186/s12711-015-0127-3> (2015).
15. González-Rodríguez, A. *et al.* On the performance of tests for the detection of signatures of selection: a case study with the Spanish autochthonous beef cattle populations. *Genet. Sel. Evol.* **48**, 81 , <https://doi.org/10.1186/s12711-016-0258-1> (2016).
16. Dash, S. *et al.* Evaluation of Bovine High-Density SNP Genotyping Array in Indigenous Dairy Cattle Breeds. *Anim. Biotechnol.* **29**, 129–135 , <https://doi.org/10.1080/10495398.2017.1329150> (2018).
17. Dixit, S. P. *et al.* Genome-Wide Runs of Homozygosity Revealed Selection Signatures in *Bos indicus*. *Front. Genet.* **11**, , <https://doi.org/10.3389/fgene.2020.00092> (2020).
18. Upadhyay, M. *et al.* Distribution and Functionality of Copy Number Variation across European Cattle Populations. *Front. Genet.* **8**, , <https://doi.org/10.3389/fgene.2017.00108> (2017).

19. Colli, L. *et al.* New Insights on Water Buffalo Genomic Diversity and Post-Domestication Migration Routes From Medium Density SNP Chip Data. *Front. Genet.* **9**, , <https://doi.org/10.3389/fgene.2018.00053> (2018).
20. Willing, E.-M., Dreyer, C. & van Oosterhout, C. Estimates of Genetic Differentiation Measured by FST Do Not Necessarily Require Large Sample Sizes When Using Many SNP Markers. *PLoS One* **7**, e42649 , <https://doi.org/10.1371/journal.pone.0042649> (2012).
21. Utsunomiya, Y. T. *et al.* Genomic clues of the evolutionary history of *Bos indicus* cattle. *Anim. Genet.* **50**, 557–568 , <https://doi.org/10.1111/age.12836> (2019).
22. Espigolan, R. *et al.* Study of whole genome linkage disequilibrium in Nellore cattle. *BMC Genomics* **14**, 305 , <https://doi.org/10.1186/1471-2164-14-305> (2013).
23. Mustafa H, Heather HJ, EuiSoo K, Ahmad N, Ali A, Khan WA, Pasha TN, Farooq MZ, Javed K, Ajmal A, S. T. Comparative analysis of genome wide difference in Red Sindhi and Holstein cattle breeds using dense SNP marker. *Int J Adv Res* **2**, (2014).
24. Karimi, K., Esmailzadeh Koshkoiyeh, A. & Gondro, C. Comparison of linkage disequilibrium levels in Iranian indigenous cattle using whole genome SNPs data. *J. Anim. Sci. Technol.* **57**, 47 , <https://doi.org/10.1186/s40781-015-0080-2> (2015).
25. Edea, Z., Dadi, H., Kim, S. W., Dessie, T. & Kim, K.-S. Comparison of SNP Variation and Distribution in Indigenous Ethiopian and Korean Cattle (Hanwoo) Populations. *Genomics Inform.* **10**, 200 , <https://doi.org/10.5808/GI.2012.10.3.200> (2012).
26. QWABE, S.O.; VANMARLE-KOSTER, E.; MAIWASHE, A. and MUCHADEYI, F. C. . E. of the B. genotyping array in four S. A. cattle populations. Evaluation of the BovineSNP50 genotyping array in four South African cattle populations. *S. Afr. j. anim. sci.* **41**, 64–67 (2013).
27. Uzzaman, M. R. *et al.* Evaluation of Single Nucleotide Polymorphisms (SNPs) Genotyped by the Illumina Bovine SNP50K in Cattle Focusing on Hanwoo Breed. *Asian-Australasian J. Anim. Sci.* **25**, 1381–1386 , <https://doi.org/10.3390/ani9030083> (2019).
28. Dadi, H., Kim, J.-J., Yoon, D. & Kim, K.-S. Evaluation of Single Nucleotide Polymorphisms (SNPs) Genotyped by the Illumina Bovine SNP50K in Cattle Focusing on Hanwoo Breed. *Asian-Australasian J. Anim. Sci.* **25**, 28–32 , <https://doi.org/10.5713/ajas.2011.11232> (2011).
29. Chagunda, M. G. G. *et al.* Use of High Density Single Nucleotide Polymorphism (SNP) Arrays to Assess Genetic Diversity and Population Structure of Dairy Cattle in Smallholder Dairy Systems: The Case of Girinka Programme in Rwanda. *Front. Genet.* **9**, , <https://doi.org/10.3389/fgene.2018.00438> (2018).
30. Makina, S. O., Muchadeyi, F. C., van Marle-Koester, E., MacNeil, M. D. & Maiwashe, A. Genetic diversity and population structure among six cattle breeds in South Africa using a whole genome SNP panel. *Front. Genet.* **5**, , <https://doi.org/10.3389/fgene.2014.00333> (2014).
31. Alshawi, A., Essa, A., Al-Bayatti, S. & Hanotte, O. Genome Analysis Reveals Genetic Admixture and Signature of Selection for Productivity and Environmental Traits in Iraqi Cattle. *Front. Genet.* **10**, , <https://doi.org/10.3389/fgene.2019.00609> (2019).

32. LIN, B. Z., SASAZAKI, S. & MANNEN, H. Genetic diversity and structure in *Bos taurus* and *Bos indicus* populations analyzed by SNP markers. *Anim. Sci. J.* **81**, 281–289 , <https://doi.org/10.1111/j.1740-0929.2010.00744.x> (2010).
33. Edea, Z. *et al.* Genetic diversity, population structure and relationships in indigenous cattle populations of Ethiopia and Korean Hanwoo breeds using SNP markers. *Front. Genet.* **4**, , <https://doi.org/10.3389/fgene.2013.00035> (2013).
34. JOST, L. G ST and its relatives do not measure differentiation. *Mol. Ecol.* **17**, 4015–4026 , <https://doi.org/10.1111/j.1365-294X.2008.03887.x> (2008).
35. Nei, M. *Molecular Evolutionary Genetics*. (Columbia University Press, 1987). doi:10.7312/nei-92038 , <https://doi.org/10.7312/nei-92038> (1987).
36. Wright S. *Evolution and Genetics of Populations, Volume 4. Variability Within and Among Natural Populations*. (Univ. Chicago Press, Chikago, 1978). (1978).
37. Xu, L. *et al.* Evaluation of Linkage Disequilibrium, Effective Population Size and Haplotype Block Structure in Chinese Cattle. *Animals* **9**, 83 , <https://doi.org/10.3390/ani9030083> (2019).
38. Fabbri, M. C., Dadousis, C. & Bozzi, R. Estimation of Linkage Disequilibrium and Effective Population Size in Three Italian Autochthonous Beef Breeds. *Animals* **10**, 1034 , <https://doi.org/10.3390/ani10061034> (2020).
39. Kijas, J. W. *et al.* Genome-Wide Analysis of the World's Sheep Breeds Reveals High Levels of Historic Mixture and Strong Recent Selection. *PLoS Biol.* **10**, e1001258 , <https://doi.org/10.1371/journal.pbio.1001258> (2012).
40. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of Population Structure Using Multilocus Genotype Data. *Genetics* **155**, 945–959 , <https://doi.org/10.1093/genetics/155.2.945> (2000).
41. Ganguly, I. *et al.* Y-chromosome genetic diversity of *Bos indicus* cattle in close proximity to the centre of domestication. *Sci. Rep.* **10**, 9992 , <https://doi.org/10.1038/s41598-020-66133-3> (2020).
42. Pryce, J. E., Hayes, B. J., Bolormaa, S. & Goddard, M. E. Polymorphic Regions Affecting Human Height Also Control Stature in Cattle. *Genetics* **187**, 981–984 , <https://doi.org/10.1534/genetics.110.123943> (2011).
43. Ligon, A. H. *et al.* Constitutional Rearrangement of the Architectural Factor HMGA2: A Novel Human Phenotype Including Overgrowth and Lipomas. *Am. J. Hum. Genet.* **76**, 340–348 , <https://doi.org/10.1086/427565> (2005).
44. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 , <https://doi.org/10.1086/519795> (2007).
45. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 , <https://doi.org/10.1186/s13742-015-0047-8> (2015).
46. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution (N. Y.)* **38**, 1358 , <https://doi.org/10.2307/2408641> (1984).

47. EXCOFFIER, L. & LISCHER, H. E. L. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567 , <https://doi.org/10.1111/j.1755-0998.2010.02847.x> (2010).
48. Barbato, M., Orozco-terWengel, P., Tapio, M. & Bruford, M. W. SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Front. Genet.* **6** , <https://doi.org/10.3389/fgene.2015.00109> (2015).
49. Barbato, M. *et al.* Genomic signatures of adaptive introgression from European mouflon into domestic sheep. *Sci. Rep.* **7**, 7623 , <https://doi.org/10.1038/s41598-017-07382-7> (2017).

Figures




Figure 1

Distribution of marker FST values across the chromosomes using Manhattan plot




Figure 2

Pair-wise Genetic distance (Nei's) among breeds. The abbreviations are defined below: TP- Tharparkar, SW- Sahiwal, GR- Gir, VC- Vechur, OG- Ongole, KG- Kangayam, HR- Hariana




Figure 3

Effective population size (Ne) of Indian cattle breeds for a number of generations. X and Y axis represents generation and Ne, respectively.





Figure 4

A. Multi-dimensional (2D) scaling plot of genome-wide IBS pairwise distances. B. Multi-dimensional (3D) scaling plot of genome-wide IBS pairwise distances




Figure 5

Population structure of seven Indian cattle breeds inferred by using the STRUCTURE software. Each animal is represented by a single vertical line divided into K colors, where K is the number of assumed ancestral clusters, that ranged from 2 to 7, the color segments shows the individual's estimated membership proportions in a given clusters: Breed abbreviations are defined as below: 1. Tharparkar, 2. Sahiwal, 3. Gir, 4. Vechur, 5. Ongole, 6. Kangayam, 7. Hariana




Figure 6

Graphical presentation of mean value of log likelihood across the inferred clusters

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTableS1.docx](#)
- [supplementaryTableS2.xlsx](#)
- [supplementaryTableS3.docx](#)