

# Coulomb Stress Change in Ahar-Varzaghan (20121108) Earth Quakes

Omid Memarian Sorkhabi (✉ [omidmemaryan@gmail.com](mailto:omidmemaryan@gmail.com))

University of Isfahan

---

## Research Article

**Keywords:** Coulomb Stress Change, Ahar-Varzaghan, Earthquakes, Okada

**Posted Date:** May 3rd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-484748/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# **Coulomb Stress Change in Ahar-Varzaghan (20121108) Earth Quakes**

Omid Memarian Sorkhabi  
(omidmemaryan@gmail.com)

Department of Geomatics Engineering, Faculty of Civil Engineering and Transportation, University of Isfahan,  
Isfahan, Iran.

## **Abstract**

Understanding how the movement of faults and deformation affects such as motion-induced surface stress and strain, which is very important in seismic regions. The best way to learn about the effects of fault movement is modeled. For example, the modeling of surface displacement or deformation and the amount of damage earthquake can be estimated by the model. Coulomb stress changes can be modeled or predicted earthquake aftershocks or future Earthquakes. we employ assumptions on the orientations, rupture lengths and average slip associated with each earthquake to calculate stress changes. Using this model, we displacement, stress and strain at any depth in the Earth's surface acquired. In this study the modeling of earthquakes  $M_w=6.5$ ,  $M_w=6.3$  Ahar-Varzaghan. The earthquakes induced displacements, strains and stresses were calculated at the surface at an average depth and its aftershocks for 10-km Ahar and 4 km Varzaghan.

Keyword: Coulomb Stress Change, Ahar-Varzaghan, Earthquakes, Okada

## **INTRODUCTION**

On the late afternoon of Saturday, August 11, 2012, the northwest of Iran was shaken by two of the strong earthquakes in Iranian history (Figure 1). First was hit by 6.5 Mw at 16:54 local time (12:23 GMT), and about 11 minutes later, a 6.3 Mw struck 10 km to the west.

## **COULOMB FAILURE**

For the calculation, information of Ahar-Varzaghan earthquake used. This information was extracted from the Iran seismological center that is visible in Table 1 and Figure 1.

Tabel 1 Information of 20120811 Ahar-Varzaghan Earthquake extracted from the Iran seismological center

Centroid Lat 38.394 Lon 46.856	Centroid Lat 38.423 Lon 46.825
Centroid Depth : 9.0	Centroid Depth : 9.0
<b>Strike Dip Rake</b>	<b>Strike Dip Rake</b>
173 61 -4	7 57 21
<b>Strike Dip Rake</b>	<b>Strike Dip Rake</b>
265 87 -151	265 72 146
Origin Time 20120811 12:23:15	Origin Time 20120811 12:34:33




Figure 1 The Study area

To calculate the length and width of the rupture zone of the earthquake were used empirical relationships (Table 2).

Table 2 Geometrical characteristics of earthquake faulting obtained from empirical relationships

Earthquake	RW(rupture width(km))	RLD(subsurface rupture length(km))	AD(average displacement(m))	Rake (degree)
Ahar	9.885	28.840	0.407	-4
varzaghan	8.729	21.667	0.286	21

## Results

Ahar vertical and horizontal displacement after  $M_w= 6.5$  mainshock showed in Figures 2 and 3. Varzaghan vertical and horizontal displacement after  $M_w= 6.3$  mainshock showed on Figures 4 and 5.




Figure 2 Ahar vertical displacement by meter after  $M_w= 6.5$  Earthquake




Figure 3 Ahar horizontal displacement by meter after  $M_w= 6.5$  Earthquake




Figure 4 Varzaghan vertical displacement by meter after  $M_w=6.5$  Earthquake




Figure 5 Varzaghan horizontal displacement after  $M_w=6.5$  Earthquake

The Coulomb stress changes after Ahar- Varzaghan Earthquakes are shown in Figure 6 and 7. We computed the Coulomb stress change in an elastic halfspace [1] by assuming a shear modulus of  $3 \times 10^{10}$  Pa, Poisson's ratio 0.25 and an effective coefficient of friction,  $\mu' = 0.4$  at depth of 10 km and 4 km [2]. Although preliminary, these results highlight the need to consider static stress interactions when estimating the seismic hazard in Azerbaijan. Past earthquakes have occurred within sufficient proximity of mainshocks for static stress interactions to affect their timing.




Figure 6 Coulomb stress change After Ahar Earthquake




Figure 7 Coulomb stress change After Varzaghan Earthquake

### Conclusion

Coulomb stress changes can be modeled or predicted earthquake aftershocks or future Earthquakes. we employ assumptions on the orientations, rupture lengths and average slip associated with each earthquake to calculate stress changes. Using this model, we displacement, stress and strain at any depth in the Earth's surface acquired. In this study the modeling of earthquakes  $M_w=6.5$ ,  $M_w=6.3$  Ahar-Varzaghan. The earthquakes induced displacements, strains and stresses were calculated at the surface at an average depth and its aftershocks for 10-km Ahar and 4 km Varzaghan. Geodesy with wavelet tools and deep learning can be helpful in this area [3-11].

### **Competing interests:**

The authors declare no competing interests.

### **References**


- [1] Okada A, Watanabe M, Sato H, JUN MS, Jo WR, Kim SK, Jeon JS, Chi HC, Oike K. Active fault topography and trench survey in the central part of the Yangsan fault, Southeast Korea. *Journal of Geography (Chigaku Zasshi)*. 1994 Apr 25;103(2):111-26.
- [2] Gombert J, Beeler N, Blanpied M. On rate-state and Coulomb failure models. *Journal of Geophysical Research: Solid Earth*. 2000 Apr 10;105(B4):7857-71.
- [3] Memarian Sorkhabi O, Asgari J, Amiri-Simkooei A. Wavelet decomposition and deep learning of altimetry waveform retracking for Lake Urmia water level survey. *Marine Georesources & Geotechnology*. 2021 Mar 10:1-1
- [4] Memarian Sorkhabi O. Geoid determination based on log sigmoid function of artificial neural networks:(a case study: Iran). *Journal of Artificial Intelligence in Electrical Engineering*. 2015 Mar 1;3(12):18-24.
- [5] Sorkhabia OM, DJAMOUR Y. Wavelet Transform Analysis of Ionospheric Electron Content Changes before Large Earthquakes. *Journal of Geomatics Science and Technology*, 2015.
- [6] Sorkhabi OM, Naderi A, Emadi R. Terrain Effect on Geoid Determination Case Study: NW Iran. *Journal of Geomatics Science and Technology*. 2014 Nov 10;4(2):139-48.
- [7] Sorkhabi OM. Deep learning of dynamic sea-level variability to investigate the relationship with the floods in Gothenburg. 2021. DOI: 10.21203/rs.3.rs-422515/v1
- [8] Memarian Sorkhabi O, Asgari J, Amiri-Simkooei A. Monitoring of Caspian Sea-level changes using deep learning-based 3D reconstruction of GRACE signal. *Measurement*. 2021 Apr 1;174:109004.
- [9] Sorkhabia OM, Asgaria J, Amiri-Simkooeia A. Caspian Sea Level Survey with GRACE and GRACE-FO. 2020.
- [10] Memarian Sorkhabi O, Djamour Y. Estimation of Geodetic Virtual Velocity Based On Back Propagation Artificial Neural Networks (Case Study: NW Iran). *Journal of Geoscience*. 2015 Jun 1;24(95):69-76.
- [11] Memarian Sorkhabi O, Asgari J, Amiri Simkooei A. Analysis of Greenland mass changes based on GRACE four-dimensional wavelet decomposition. *Remote Sensing Letters*. 2021 May 4;12(5):499-509.

# Figures



Figure 1


The Study area






**Figure 2**

Ahar vertical displacement by meter after Mw= 6.5 Earthquake



**Figure 3**

Ahar horizontal displacement by meter after Mw= 6.5 Earthquake



**Figure 4**

Varzaghan vertical displacement by meter after Mw= 6.5 Earthquake




Figure 5

Varzaghan horizontal displacement after Mw= 6.5 Earthquake




Figure 6

Coulomb stress change After Ahar Earthquake




Figure 7

Coulomb stress change After Varzaghan Earthquake